

Frontiers in Pharmacology, 2018, vol.9, NJUN

Autoregulation of acetylcholine release and micro-pharmacodynamic mechanisms at neuromuscular junction: Selective acetylcholinesterase inhibitors for therapy of myasthenic syndromes

Petrov K., Nikolsky E., Masson P.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Petrov, Nikolsky and Masson. Neuromuscular junctions (NMJs) are directly involved into such indispensable to life processes as respiration and locomotion. However, motor nerve forms only one synaptic contact at each muscle fiber. This unique configuration requires specific properties and constrains to be effective. The very high density of acetylcholine receptors (AChRs) of muscle type in synaptic cleft and an excess of acetylcholine (ACh) released under physiological conditions make this synapse extremely reliable. Nevertheless, under pathological conditions such as myasthenia gravis and congenital myasthenic syndromes, the safety factor can be markedly reduced. Drugs used for short-term symptomatic therapy of these pathological states, cause partial inhibition of cholinesterases (ChEs). These enzymes catalyze the hydrolysis of ACh, thus terminate its action on AChRs. Extension of the lifetime of ACh molecules compensates muscular AChRs abnormalities and, consequently, rescues muscle contractions. In this mini review, we will first outline the functional organization of the NMJ, and then, consider the concept of the safety factor and how it may be changed. This will be followed by a look at autoregulation of ACh release that influences the safety factor of NMJs. Finally, we will consider the morphological features of NMJs as a putative reserve to increase effectiveness of pathological muscle weakness therapy by ChEs inhibitors due to opportunity to use micro-pharmacodynamic mechanisms.

<http://dx.doi.org/10.3389/fphar.2018.00766>

Keywords

Cholinesterases, Congenital myasthenic syndromes, Micro-pharmacodynamic mechanisms, Myasthenia gravis, Neuromuscular junction

References

- [1] Abbs, E. T., and Joseph, D. N. (1981). The effects of atropine and oxotremorine on acetylcholine release in rat phrenic nerve-diaphragm preparations. *Br. J. Pharmacol.* 73, 481-483. doi: 10.1111/j.1476-5381.1981.tb10446.x
- [2] Anglister, L., Stiles, J. R., and Salpeter, M. M. (1994). Acetylcholinesterase density and turnover number at frog neuromuscular junctions, with modeling of their role in synaptic function. *Neuron* 12, 783-794. doi: 10.1016/0896-6273(94)90331-X

- [3] Arbour, D., VandeVelde, C., and Robitaille, R. (2017). New perspectives on amyotrophic lateral sclerosis: the role of glial cells at the neuromuscular junction. *J. Physiol.* 595, 647-661. doi: 10.1113/JP270213
- [4] Arenson, M. (1991). The effects of muscarine and atropine reveal that inhibitory autoreceptors are present on frog motor nerve terminals but are not activated during transmission. *Naunyn Schmiedebergs Arch. Pharmacol.* 343, 128-133. doi: 10.1007/BF00168599
- [5] Badawi, Y., and Nishimune, H. (2018). Presynaptic active zones of mammalian neuromuscular junctions: nanoarchitecture and selective impairments in aging. *Neurosci. Res.* 127, 78-88. doi: 10.1016/j.neures.2017.11.014
- [6] Bernard, V., Girard, E., Hrabovska, A., Camp, S., Taylor, P., Plaud, B., et al. (2011). Distinct localization of collagen Q and PRiMA forms of acetylcholinesterase at the neuromuscular junction. *Mol. Cell. Neurosci.* 46, 272-281. doi: 10.1016/j.mcn.2010.09.010
- [7] Blotnick-Rubin, E., and Anglister, L. (2018). Fine localization of acetylcholinesterase in the synaptic cleft of the vertebrate neuromuscular junction. *Front. Mol. Neurosci.* 11:123. doi: 10.3389/fnmol.2018.00123
- [8] Cavalcante, P., Bernasconi, P., and Mantegazza, R. (2012). Autoimmune mechanisms in myasthenia gravis. *Curr. Opin. Neurol.* 25, 621-629. doi: 10.1097/WCO.0b013e328357a829
- [9] Ciani, S., and Edwards, C. (1963). The effect of acetylcholine on neuromuscular transmission in the frog. *J. Pharmacol. Exp. Ther.* 142, 21-23
- [10] Copeland, R. (2011). Conformational adaptation in drug-target interactions and residence time. *Future Med. Chem.* 3, 1491-1501. doi: 10.4155/fmc.11.112
- [11] Copeland, R., Pompliano, D., and Meek, T. (2006). Drug-target residence time and its implications for lead optimization. *Nat. Rev. Drug Discov.* 5, 730-739. doi: 10.1038/nrd2082
- [12] Correia-de-Sá, P., Timóteo, M., and Ribeiro, J. (1996). Presynaptic A1 inhibitory/A2A facilitatory adenosine receptor activation balance depends on motor nerve stimulation paradigm at the rat hemidiaphragm. *J. Neurophysiol.* 76, 3910-3919. doi: 10.1152/jn.1996.76.6.3910
- [13] Davis, R., and Koelle, G. (1967). Electron microscopic localization of acetylcholinesterase and nonspecific cholinesterase at the neuromuscular junction by the gold-thiocholine and gold-thiolacetic acid methods. *J. Cell Biol.* 34, 157-171. doi: 10.1083/jcb.34.1.157
- [14] Dudel, J. (2007). The time course of transmitter release in mouse motor nerve terminals is differentially affected by activation of muscarinic M1 or M2 receptors. *Eur. J. Neurosci.* 26, 2160-2168. doi: 10.1111/j.1460-9568.2007.05770.x
- [15] Duncan, C., and Publicover, S. (1979). Inhibitory effects of cholinergic agents on the release of transmitter at the frog neuromuscular junction. *J. Physiol.* 294, 91-103. doi: 10.1113/jphysiol.1979.sp012917
- [16] Dvir, H., Silman, I., Harel, M., Rosenberry, T., and Sussman, J. (2010). Acetylcholinesterase: from 3D structure to function. *Chem. Biol Interact.* 187, 10-22. doi: 10.1016/j.cbi.2010.01.042
- [17] Engel, A. (2007). The therapy of congenital myasthenic syndromes. *Neurotherapeutics* 4, 252-257. doi: 10.1016/j.nurt.2007.01.001
- [18] Engel, A., Shen, X., Selcen, D., and Sine, S. (2015). Congenital myasthenic syndromes: pathogenesis, diagnosis, and treatment. *Lancet Neurol.* 14, 420-434. doi: 10.1016/S1474-4422(14)70201-7
- [19] Garcia, N., Santafé, M., Salon, I., Lanuza, M., and Tomàs, J. (2005). Expression of muscarinic acetylcholine receptors (M1-, M2-, M3-and M4-type) in the neuromuscular junction of the newborn and adult rat. *Histol. Histopathol.* 20, 733-743. doi: 10.14670/HH-20.733
- [20] Heredia, D., Feng, C., Hennig, G., Renden, R., and Gould, T. (2018). Activity-induced Ca²⁺ signaling in perisynaptic Schwann cells of the early postnatal mouse is mediated by P2Y1 receptors and regulates muscle fatigue. *eLife* 7:e30839. doi: 10.7554/eLife.30839
- [21] Howard, J. (2018). Myasthenia gravis: the role of complement at the neuromuscular junction. *Ann. N. Y. Acad. Sci.* 1412, 113-128. doi: 10.1111/nyas.13522
- [22] Jones, R., Harrison, C., Eaton, S., Llaveró Hurtado, M., Graham, L., Alkhamash, L., et al. (2017). Cellular and molecular anatomy of the human neuromuscular junction. *Cell Rep.* 21, 2348-2356. doi: 10.1016/j.celrep.2017.11.008
- [23] Katz, B., and Miledi, R. (1975). The nature of the prolonged endplate depolarization in anti-esterase treated muscle. *Proc. R. Soc. Lond. B Biol. Sci.* 192, 27-38. doi: 10.1098/rspb.1975.0149
- [24] Keogh, M., Sedehizadeh, S., and Maddison, P. (2011). Treatment for Lambert-Eaton myasthenic syndrome. *Cochrane Database Syst. Rev.* 2:CD003279. doi: 10.1002/14651858.CD003279.pub3
- [25] Kesner, V., Oh, S., Dimachkie, M., and Barohn, R. (2018). Lambert-eaton myasthenic syndrome. *Neurol. Clin.* 36, 379-394. doi: 10.1016/j.ncl.2018.01.008
- [26] Kharlamova, A., Lushchekina, S., Petrov, K., Kots, E., Nachon, F., Villard-Wandhammer, M., et al. (2016). Slow-binding inhibition of acetylcholinesterase by an alkylammonium derivative of 6-methyluracil: mechanism and possible advantages for myasthenia gravis treatment. *Biochem. J.* 473, 1225-1236. doi: 10.1042/BCJ20160084

- [27] Ko, C., and Robitaille, R. (2015). Perisynaptic Schwann cells at the neuromuscular synapse: adaptable, multitasking glial cells. *Cold Spring Harb. Perspect. Biol.* 7:a020503. doi: 10.1101/cshperspect.a020503
- [28] Komloova, M., Musilek, K., Dolezal, M., Gunn-Moore, F., and Kuca, K. (2010). Structure-activity relationship of quaternary acetylcholinesterase inhibitors-outlook for early myasthenia gravis treatment. *Curr. Med. Chem.* 17, 1810-1824. doi: 10.2174/092986710791111198
- [29] Kovyazina, I., Tsentsevitsky, A., and Nikolsky, E. (2015). Identification of the muscarinic receptor subtypes involved in autoregulation of acetylcholine quantal release from frog motor nerve endings. *Dokl. Biol. Sci.* 460, 5-7. doi: 10.1134/S0012496615010081
- [30] Kovyazina, I., Tsentsevitsky, A., Nikolsky, E., and Bukharaeva, E. (2010). Kinetics of acetylcholine quanta release at the neuromuscular junction during high-frequency nerve stimulation. *Eur. J. Neurosci.* 32, 1480-1489. doi: 10.1111/j.1460-9568.2010.07430.x
- [31] Kusner, L., Sengupta, M., and Kaminski, H. (2018). Acetylcholine receptor antibody-mediated animal models of myasthenia gravis and the role of complement. *Ann. N. Y. Acad. Sci.* 1413, 136-142. doi: 10.1111/nyas.13555
- [32] Lee, J., and Jander, S. (2017). Myasthenia gravis: recent advances in immunopathology and therapy. *Expert Rev. Neurother.* 17, 287-299. doi: 10.1080/14737175.2017.1241144
- [33] Legay, C. (2018). Congenital myasthenic syndromes with acetylcholinesterase deficiency, the pathophysiological mechanisms. *Ann. N. Y. Acad. Sci.* 1413, 104-110. doi: 10.1111/nyas.13595
- [34] Malomouzh, A., Petrov, K., Nurullin, L., and Nikolsky, E. (2015). Metabotropic GABAB receptors mediate GABA inhibition of acetylcholine release in the rat neuromuscular junction. *J. Neurochem.* 135, 1149-1160. doi: 10.1111/jnc.13373
- [35] Masson, P., and Lockridge, O. (2010). Butyrylcholinesterase for protection from organophosphorus poisons: catalytic complexities and hysteretic behavior. *Arch. Biochem. Biophys.* 494, 107-120. doi: 10.1016/j.abb.2009.12.005
- [36] Masson, P., and Lushchekina, S. (2016). Slow-binding inhibition of cholinesterases, pharmacological and toxicological relevance. *Arch. Biochem. Biophys.* 593, 60-68. doi: 10.1016/j.abb.2016.02.010
- [37] Minic, J., Chatonnet, A., Krejci, E., and Molgó, J. (2003). Butyrylcholinesterase and acetylcholinesterase activity and quantal transmitter release at normal and acetylcholinesterase knockout mouse neuromuscular junctions. *Br. J. Pharmacol.* 138, 177-187. doi: 10.1038/sj.bjp.0705010
- [38] Minic, J., Molgó, J., Karlsson, E., and Krejci, E. (2002). Regulation of acetylcholine release by muscarinic receptors at the mouse neuromuscular junction depends on the activity of acetylcholinesterase. *Eur. J. Neurosci.* 15, 439-448. doi: 10.1046/j.0953-816x.2001.01875.x
- [39] Nascimento, F., Sebastião, A., and Ribeiro, J. (2017). Presymptomatic and symptomatic ALS SOD1(G93A) mice differ in adenosine A1 and A2A receptor-mediated tonic modulation of neuromuscular transmission. *Purinergic Signal.* 11, 471-480. doi: 10.1007/s11302-015-9465-4
- [40] Nicole, S., Azuma, Y., Bauché, S., Eymard, B., Lochmüller, H., and Slater, C. (2017). Congenital myasthenic syndromes or inherited disorders of neuromuscular transmission: recent discoveries and open questions. *J. Neuromuscul. Dis.* 4, 269-284. doi: 10.3233/JND-170257
- [41] Ohno, K., Tsujino, A., Brengman, J., Harper, C., Bajzer, Z., Udd, B., et al. (2001). Choline acetyltransferase mutations cause myasthenic syndrome associated with episodic apnea in humans. *Proc. Natl. Acad. Sci. U.S.A.* 98, 2017-2022. doi: 10.1073/pnas.98.4.2017
- [42] Oliveira, L., Timóteo, M., and Correia-de-Sá, P. (2002). Modulation by adenosine of both muscarinic M1-facilitation and M2-inhibition of [3H]-acetylcholine release from the rat motor nerve terminals. *Eur. J. Neurosci.* 15, 1728-1736. doi: 10.1046/j.1460-9568.2002.02020.x
- [43] Oliveira, L., Timóteo, M., and Correia-de-Sá, P. (2009). Negative crosstalk between M1 and M2 muscarinic autoreceptors involves endogenous adenosine activating A1 receptors at the rat motor endplate. *Neurosci. Lett.* 459, 127-131. doi: 10.1016/j.neulet.2009.05.001
- [44] Petrov, K., Girard, E., Nikitashina, A., Colasante, C., Bernard, V., Nurullin, L., et al. (2014). Schwann cells sense and control acetylcholine spillover at the neuromuscular junction by $\alpha 7$ nicotinic receptors and butyrylcholinesterase. *J. Neurosci.* 34, 11870-11883. doi: 10.1523/JNEUROSCI.0329-14.2014
- [45] Petrov, K., Kharlamova, A., Lenina, O., Nurtudinov, A., Sitdykova, M., Ilyin, V., et al. (2018a). Specific inhibition of acetylcholinesterase as an approach to decrease muscarinic side effects during myasthenia gravis treatment. *Sci. Rep.* 8:304. doi: 10.1038/s41598-017-18307-9
- [46] Petrov, K., Kovyazina, I., Zobov, V., Bukharaeva, E., Nikolsky, E., and Vyskocil, F. (2006). Different sensitivity of miniature endplate currents of the rat extensor digitorum longus, soleus and diaphragm muscles to a novel acetylcholinesterase inhibitor C-547. *Physiol. Res.* 55, 585-589
- [47] Petrov, K., Kovyazina, I., Zobov, V., Bukharaeva, E., Nikolsky, E., and Vyskocil, F. (2009). Different sensitivity of miniature endplate currents in rat external and internal intercostal muscles to the acetylcholinesterase inhibitor C-547 as compared with diaphragm and extensor digitorum longus. *Physiol. Res.* 58, 149-153

- [48] Petrov, K., Malomouzh, A., Kovyazina, I., Krejci, E., Nikitashina, A., Proskurina, S., et al. (2013). Regulation of acetylcholinesterase activity by nitric oxide in rat neuromuscular junction via N-methyl-D-aspartate receptor activation. *Eur. J. Neurosci.* 37, 181-189. doi: 10.1111/ejn.12029
- [49] Petrov, K., Yagodina, L., Valeeva, G., Lannik, N., Nikitashina, A., Rizvanov, A., et al. (2011). Different sensitivities of rat skeletal muscles and brain to novel anti-cholinesterase agents, alkylammonium derivatives of 6-methyluracil (ADEMS). *Br. J. Pharmacol.* 163, 732-744. doi: 10.1111/j.1476-5381.2011.01211.x
- [50] Petrov, K., Zueva, I., Kovyazina, I., Sedov, I., Lushchekina, S., Kharlamova, A., et al. (2018b). C-547, a 6-methyluracil derivative with long-lasting binding and rebinding on acetylcholinesterase: pharmacokinetic and pharmacodynamic studies. *Neuropharmacology* 131, 304-315. doi: 10.1016/j.neuropharm.2017.12.034
- [51] Plomp, J. J. (2018). "Neuromuscular junction physiology and pathophysiology, " in *Myasthenia Gravis and Related Disorders*, eds H. Kaminski, and L. Kusner (Cham: Humana Press), 1-12
- [52] Pope, C., and Brimijoin, S. (2018). Cholinesterases and the fine line between poison and remedy. *Biochem. Pharmacol.* 153, 205-216. doi: 10.1016/j.bcp.2018.01.044
- [53] Pope, C., Karanth, S., and Liu, J. (2005). Pharmacology and toxicology of cholinesterase inhibitors: uses and misuses of a common mechanism of action. *Environ. Toxicol. Pharmacol.* 19, 433-446. doi: 10.1016/j.etap.2004.12.048
- [54] Renew, J. R., Brull, S. J., and Naguib, M. (2018). "Clinical pharmacology of drugs acting at the neuromuscular junction, " in *Basic Sciences in Anesthesia*, eds E. Farag, M. Argalious, J. Tetzlaff, D. Sharma (Cham: Springer), 139-159
- [55] Robitaille, R. (1995). Purinergic receptors and their activation by endogenous purines at perisynaptic glial cells of the frog neuromuscular junction. *J. Neurosci.* 15, 7121-7131. doi: 10.1523/JNEUROSCI.15-11-07121.1995
- [56] Robitaille, R., Jahromi, B., and Charlton, M. (1997). Muscarinic Ca²⁺ responses resistant to muscarinic antagonists at perisynaptic Schwann cells of the frog neuromuscular junction. *J. Physiol.* 504, 337-347. doi: 10.1111/j.1469-7793.1997.337be.x
- [57] Rochon, D., Rousse, I., and Robitaille, R. (2001). Synapse-glia interactions at the mammalian neuromuscular junction. *J. Neurosci.* 21, 3819-3829. doi: 10.1523/JNEUROSCI.21-11-03819.2001
- [58] Ruff, R. L. (2011). Endplate contributions to the safety factor for neuromuscular transmission. *Muscle Nerve* 44, 854-861. doi: 10.1002/mus.22177
- [59] Santafé, M., Lanuza, M., Garcia, N., and Tomàs, J. (2006). Muscarinic autoreceptors modulate transmitter release through protein kinase C and protein kinase A in the rat motor nerve terminal. *Eur. J. Neurosci.* 23, 2048-2056. doi: 10.1111/j.1460-9568.2006.04753.x
- [60] Santafé, M., Lanuza, M., Garcia, N., Tomàs, M., and Tomàs, J. (2007). Coupling of presynaptic muscarinic autoreceptors to serine kinases in low and high release conditions on the rat motor nerve terminal. *Neuroscience* 148, 432-440. doi: 10.1016/j.neuroscience.2007.06.017
- [61] Santafé, M., Salon, I., Garcia, N., Lanuza, M., Uchitel, O., and Tomàs, J. (2003). Modulation of ACh release by presynaptic muscarinic autoreceptors in the neuromuscular junction of the newborn and adult rat. *Eur. J. Neurosci.* 17, 119-127. doi: 10.1046/j.1460-9568.2003.02428.x
- [62] Santafé, M., Salon, I., Garcia, N., Lanuza, M., Uchitel, O., and Tomàs, J. (2004). Muscarinic autoreceptors related with calcium channels in the strong and weak inputs at polyinnervated developing rat neuromuscular junctions. *Neuroscience* 123, 61-73. doi: 10.1016/j.neuroscience.2003.09.012
- [63] Silinsky, E., and Redman, R. (1994). ATP released together with acetylcholine as the mediator of neuromuscular depression at frog motor nerve endings. *J. Physiol.* 477, 117-127. doi: 10.1113/jphysiol.1994.sp020176
- [64] Sine, S. M. (2002). The nicotinic receptor ligand binding domain. *J. Neurobiol.* 53, 431-446. doi: 10.1002/neu.10139
- [65] Slater, C. (2009). "Reliability of neuromuscular transmission and how it is maintained, " in *Neuromuscular Junction Disorders: Handbook of Clinical Neurology*, Vol. 91, ed. A. G. Engel (New York, NY: Elsevier), 27-101
- [66] Slater, C. (2015). The functional organization of motor nerve terminals. *Prog. Neurobiol.* 134, 55-103. doi: 10.1016/j.pneurobio.2015.09.004
- [67] Slater, C. (2017). The structure of human neuromuscular junctions: some unanswered molecular questions. *Int. J. Mol. Sci.* 18:E2183. doi: 10.3390/ijms18102183
- [68] Slutsky, I., Silman, I., Parnas, I., and Parnas, H. (2001). Presynaptic M(2) muscarinic receptors are involved in controlling the kinetics of ACh release at the frog neuromuscular junction. *J. Physiol.* 536(Pt 3), 717-725. doi: 10.1111/j.1469-7793.2001.00717.x
- [69] Taylor, P., Camp, S., and Radic, Z. (2009). "Acetylcholinesterase, " in *Encyclopedia of Neuroscience*, Vol. 1, eds R. Larry and L. R. Squire (New York, NY: Elsevier), 5-7. doi: 10.1016/B978-008045046-9.01132-3
- [70] Todd, K., Auld, D., and Robitaille, R. (2007). Neurotrophins modulate neuron-glia interactions at a vertebrate synapse. *Eur. J. Neurosci.* 25, 1287-1296. doi: 10.1111/j.1460-9568.2007.05385.x

- [71] Todd, K., Darabid, H., and Robitaille, R. (2010). Perisynaptic glia discriminate patterns of motor nerve activity and influence plasticity at the neuromuscular junction. *J. Neurosci* 30, 11870-11882. doi: 10.1523/JNEUROSCI.3165-10.2010
- [72] Tonge, P. (2018). Drug-target kinetics in drug discovery. *ACS Chem. Neurosci.* 9, 29-39. doi: 10.1021/acchemneuro.7b00185
- [73] Vauquelin, G. (2010). Rebinding: or why drugs may act longer in vivo than expected from their in vitro target residence time. *Expert Opin. Drug Discov.* 5, 927-941. doi: 10.1517/17460441.2010.512037
- [74] Vauquelin, G. (2015). On the 'micro'-pharmacodynamic and pharmacokinetic mechanisms that contribute to long-lasting drug action. *Expert Opin. Drug Discov.* 10, 1085-1098. doi: 10.1517/17460441.2015.1067196
- [75] Vauquelin, G. (2016). Effects of target binding kinetics on in vivo drug efficacy: koff, kon and rebinding. *Br. J. Pharmacol.* 173, 2319-2334. doi: 10.1111/bph.13504
- [76] Vauquelin, G., and Charlton, S. (2010). Long-lasting target binding and rebinding as mechanisms to prolong in vivo drug action. *Br. J. Pharmacol.* 161, 488-508. doi: 10.1111/j.1476-5381.2010.00936.x
- [77] Verschuuren, J., Strijbos, E., and Vincent, A. (2016). "Neuromuscular junction disorders, " in *Autoimmune Neurology: Handbook of Clinical Neurology*, Vol. 133, eds J. Sean, S. J. Pittock, and A. Vincent (New York, NY: Elsevier), 447-466. doi: 10.1016/B978-0-444-63432-0.00024-4
- [78] Walkup, G., You, Z., Ross, P., Allen, E., Daryaei, F., Hale, M., et al. (2015). Translating slow-binding inhibition kinetics into cellular and in vivo effects. *Nat. Chem. Biol.* 11, 416-423. doi: 10.1038/nchembio.1796
- [79] Wang, X., and Rich, M. (2018). Homeostatic synaptic plasticity at the neuromuscular junction in myasthenia gravis. *Ann. N. Y. Acad. Sci.* 1412, 170-177. doi: 10.1111/nyas.13472
- [80] Wessler, I., Karl, M., Mai, M., and Diener, A. (1987). Muscarinic receptors on the rat phrenic nerve, evidence for positive and negative muscarinic feedback mechanisms. *Naunyn Schmiedebergs Arch. Pharmacol.* 1987, 605-612
- [81] Wood, S., and Slater, C. (2001). Safety factor at the neuromuscular junction. *Prog. Neurobiol.* 64, 393-429. doi: 10.1016/S0301-0082(00)00055-1