

Boundary stabilization of quasilinear Maxwell equations

Michael Pokojovy, Roland Schnaubelt

CRC Preprint 2018/48, December 2018

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

ISSN 2365-662X

BOUNDARY STABILIZATION OF QUASILINEAR MAXWELL EQUATIONS

MICHAEL POKOJOVY AND ROLAND SCHNAUBELT

ABSTRACT. We investigate an initial-boundary value problem for a quasilinear nonhomogeneous, anisotropic Maxwell system subject to an absorbing boundary condition of Silver & Müller type in a smooth, bounded, strictly star-shaped domain of \mathbb{R}^3 . Imposing usual smallness assumptions in addition to standard regularity and compatibility conditions, a nonlinear stabilizability inequality is obtained by showing nonlinear dissipativity and observability-like estimates enhanced by an intricate regularity analysis. With the stabilizability inequality at hand, the classic nonlinear barrier method is employed to prove that small initial data admit unique classical solutions that exist globally and decay to zero at an exponential rate. Our approach is based on a recently established local well-posedness theory in a class of \mathcal{H}^3 -valued functions.

1. INTRODUCTION

The goal of this work is to establish global-in-time existence and exponential stability for the quasilinear nonhomogeneous, anisotropic Maxwell system in a bounded, smooth star-shaped domain of \mathbb{R}^3 . Provided the initial data are sufficiently small in the Sobolev \mathcal{H}^3 -norm, we will show that the damping effect from an absorbing boundary condition of Silver & Müller-type can render the system globally well-posed and exponentially stable.

The Maxwell equations, laying the mathematical foundation of the theory of electro-magnetism, establish a relation between the electric fields E , D and the magnetic fields B , H via Ampère’s circuital law and Faraday’s law of induction. These basic continuity equations read as

$$\partial_t D = \operatorname{curl} H \quad \text{and} \quad \partial_t B = -\operatorname{curl} H, \quad t > 0, \quad x \in \Omega,$$

where $\Omega \subset \mathbb{R}^3$ is a bounded domain with a smooth boundary Γ and outer unit normal ν . In our work, we take (E, H) as the state variables and postulate the instantaneous nonlinear material laws

$$D = \varepsilon(\cdot, E)E \quad \text{and} \quad B = \mu(\cdot, H)H \quad \text{in } \Omega$$

with nonlinear, nonhomogeneous, anisotropic tensor-valued permittivity ε and permeability μ . Imposing a generalized linear Silver & Müller boundary condition with a tensor-valued λ , we arrive at the quasilinear Maxwell system

$$\begin{aligned} \partial_t(\varepsilon(x, E(t, x))E(t, x)) &= \operatorname{curl} H(t, x), & t \geq 0, \quad x \in \Omega, \\ \partial_t(\mu(x, H(t, x))H(t, x)) &= -\operatorname{curl} E(t, x), & t \geq 0, \quad x \in \Omega, \\ \operatorname{div}(\varepsilon(x, E)E) &= 0, \quad \operatorname{div}(\mu(x, H)H) = 0, & t \geq 0, \quad x \in \Omega, \\ H(t, x) \times \nu(x) + (\lambda(x)(E(t, x) \times \nu(x))) \times \nu(x) &= 0, & t \geq 0, \quad x \in \Gamma, \\ E(0, x) &= E^{(0)}(x), \quad H(0, x) = H^{(0)}(x), & x \in \Omega. \end{aligned} \tag{1.1}$$

The absorbing boundary condition in Equation (1.1) above emerges as a first-order approximation of the more comprehensive “transparent” boundary condition. But, while being dissipative in its

Date: December 14, 2018.

2000 Mathematics Subject Classification. 35Q61, 35L60, 35L50, 35A02, 35B40, 35B65.

Key words and phrases. Quasilinear Maxwell equations; Silver-Müller boundary conditions; nonhomogeneous anisotropic materials; global existence; exponential stability.

We gratefully acknowledge financial support from the Deutsche Forschungsgemeinschaft (DFG) through CRC 1173.

nature, in contrast to the latter condition, Silver & Müller-type boundary conditions (sometimes [48] also – somewhat inadequately [5] – referred to as “impedance boundary conditions”) account for the waves being reflected back into the domain (cf. [17, p. 136]). From the physical point of view, such boundary conditions model the process of how electromagnetic waves are scattered by an obstacle Ω under the assumption that the underlying medium does not allow for deep penetration of the wave (cf. [9, p. 20]).

The solenoidality condition or the divergence-freeness of E and H express the absence of electric and magnetic charges in the medium, which is true in any physical environment for the magnetic charges. This condition excludes certain non-zero equilibrium states of the system. Because of the absence of interior conductivity or currents, the solenoidality conditions in Equation (1.1) are uniquely related to the so-called “initial solenoidality”

$$\operatorname{div}(\varepsilon(\cdot, E^{(0)})E^{(0)}) = 0, \quad \operatorname{div}(\mu(\cdot, H^{(0)})H^{(0)}) = 0 \quad \text{in } \Omega. \quad (1.2)$$

Provided the system (1.1) possesses a regular solution, the solenoidality condition in the fourth equation of (1.1) trivially yields the initial solenoidality (1.2). Vice versa, applying the div-operator to the first two equations in (1.1) and recalling the distributional identity $\operatorname{div} \operatorname{curl} = 0$, the solenoidality conditions of (1.1) follow (cf. [58, Lemma 7.25]). Therefore, in the following, we will be freely switching between the two equivalent conditions.

Since we are interested in classical \mathcal{H}^3 -valued solutions, it is natural to expect that both regularity and compatibility conditions are to be imposed. Indeed, given a smooth solution on a closed time-space cylinder, an obvious requirement is that all response tensors (i.e., ε , μ and λ), the initial data $(E^{(0)}, H^{(0)})$ and the boundary Γ are smooth. Further, the higher-order time derivatives of the solution at $t = 0$ are expected to satisfy the boundary conditions translating into compatibility conditions on the initial data. These have been described in the recent well-posedness study [52]. Unless the nonlinear response tensors ε and μ are globally positive definite, the smallness of the solution and thus the initial data, is also necessary to preserve the “hyperbolicity” of the system (1.1). Last but not least, appropriate symmetry conditions on the response tensors along with a “tangentiality” condition (cf. Section 2) on the tensor field λ are required.

Turning to global solutions, a uniform stabilization mechanism in the system (1.1) needs to be discovered to pave the way for a reasonable long-time analysis. In contrast to semilinear problems, uniform stability is often the only way to global existence, especially for hyperbolic problems. In bounded domains, it is the exponential stability that furnishes the global existence of smooth solutions originating in a vicinity of a stable equilibrium [21, 30, 31, 32, 38, 42]. In the full space \mathbb{R}^d , dispersive estimates are the path to the long-time existence [36, 37, 40, 41, 49, 50, 56]. In case a (too) weak solution concept is employed [13, 57] or there is little to no dissipation (even in a smooth solution class) [1, 2, 3, 4, 22], blow-ups are known to occur in finite time – both for large and small data, often independently of whether the initial data are smooth or compactly supported, etc.

In view of the dissipation present in the Maxwell system (1.1) at the basic energy level and rooted in its absorbing boundary condition, the question arises whether this dissipative mechanism is sufficiently strong to drive the system to the zero equilibrium. In this paper, we give a positive answer to this question in strongly star-shaped bounded C^5 -domains Ω of \mathbb{R}^3 – provided the initial data are small in $\mathcal{H}^3(\Omega)$ and satisfy the compatibility conditions (2.9) introduced in Section 2 to follow. Our stability proof will heavily rely on the local well-posedness theory, in particular, the *a priori* estimates, recently established in [52] and a mixture of energy methods and control-theoretic techniques such as Rellich multipliers for proving observability/stabilizability, etc. The core step of our reasoning is a refined regularity analysis based on div-curl “elliptic” theory.

We continue with a brief literature review accompanied by a general discussion. Maxwell equations (1.1) with symmetric and positive definite $\partial_E(\varepsilon(\cdot, E)E)$ and $\partial_H(\mu(\cdot, H)H)$ can be viewed as a first-order quasilinear symmetric hyperbolic system. Due to their generality and ubiquity in mathematical physics, these systems have drawn major attention in the literature. In the full space

$\Omega = \mathbb{R}^d$, a well-posedness theory due to Kato [23] is known. Applied to Equation (1.1), it furnishes the existence of a local-in-time \mathcal{H}^3 -valued solution. As for the global existence, for the quasilinear Maxwell system, we are only aware of very few results [37, 49, 56] on the full space $\Omega = \mathbb{R}^3$ that establish global existence and uniform stability of smooth solutions to small initial data. These studies have at their heart dispersive estimates which are not available on bounded domains. On the other hand, a blow-up behavior in $\mathcal{H}(\text{curl})$ under various boundary conditions has recently been shown in [13], whereas blow-up in $W^{1,\infty}(\Omega)$ has been known since a long time, cf. [1].

Turning to domains with boundary, numerous solution theories are available [7, 20, 24, 46, 51, 54, 55]. However, due to the characteristicity of the boundary symbol associated with (1.1) (cf. [58, 60] for the case of a perfect conductor), a regularity loss in the normal direction possibly occurs making only few of these theories (e.g., [20, 55]) applicable. The additional price to pay is that a cumbersome framework of weighted Sobolev spaces of very high order needs to be employed greatly reducing the practical applicability of these results. While the existence (without uniqueness or continuous dependence on the data) for Maxwell system (1.1) has been established in the early work [48], the full Hadamard well-posedness in \mathcal{H}^3 for perfectly conducting boundary conditions [58, 59, 60], conservative interface conditions [53] or absorbing boundary conditions [52] as in Equation (1.1) have only been proved very recently. We will strongly rely on [52] in this paper.

In our main Theorem 2.2, we demonstrate that local \mathcal{H}^3 -valued solutions to (1.1) can be globally extended and decay exponentially to zero in the same topology under a smallness assumption on the initial data (in addition to natural compatibility conditions). In addition, we assume that the domain Ω is strictly star-shaped and the space derivatives of $\varepsilon(x, 0)$ and $\mu(x, 0)$ satisfy the lower bound (2.24). The long-time behavior of Maxwell system has been extensively investigated in recent literature (see, e.g., [6, 15, 17, 18, 19, 25, 27, 43, 44, 45, 47] and references therein). In all of these works, the authors restrict their attention to linear permittivity and permeability responses, often homogeneous or isotropic ones, rendering the problem linear or semilinear. In contrast, our problem (1.1) is genuinely quasilinear with the only dissipativity source entering the system through the Silver & Müller-type boundary condition. To the best of our knowledge, apart from our companion paper [30] for the case of interior conductivity and perfectly conducting boundary conditions, no global results on *quasilinear* Maxwell systems in domains with boundary are available.

In the linear situation, the aforementioned asymptotic results are furnished by an dissipativity inequality coupled with an observability-type estimate providing a lower bound on the dissipation. We refer the reader to [25] for the case of scalar, constant $\varepsilon, \mu, \lambda$; [17, 18] for x -dependent scalar ε, μ and nonlinear x -independent scalar λ ; [43] for the case of (t, x) -dependent scalar ε, μ and nonlinear x -dependent scalar λ ; [6] for general nonhomogeneous, anisotropic ε, μ and nonlinear x -independent scalar λ and a nonlinear boundary feedback, etc. The use of Rellich multipliers in the present paper is motivated by the linear observability result [16].

In our nonlinear situation, given the initial data are small, the perturbation terms also remain small till a certain time, but the smallness is only guaranteed in much stronger norms than those controlled by the dissipation. Hence, a rather elaborated regularity theory needs to be developed to close the regularity gap – and, even beyond that, with constants independent of the time interval length. Namely, for $k \in \{0, 1, 2\}$, the $\mathcal{H}^{3-k}(\Omega)$ -norms of $\partial_t^k(E, H)$ need to be estimated by the L^2 -norm of $(\partial_t^l(E, H))_{l \in \{0, 1, 2, 3\}}$, plus superlinear error terms. This fact is established in our core Proposition 4.1 which is proved in Section 6 based on theory developed in Section 5. There we show an “elliptic” regularity result (essentially due to [12]), which allows us to reconstruct the full $\mathcal{H}^1(\Omega)$ -norm of (E, H) from the $L^2(\Omega)$ -norms of $\text{curl}(E, H)$ and $\text{div}(\varepsilon E, \mu H)$ as well as the $\mathcal{H}^{1/2}(\Gamma)$ -norm of the boundary data. In contrast to our earlier work [30], due to the absence of the electric resistance term, solenoidality properties are available both for E and H , which facilitates the application of this fact. However, higher-order normal derivatives cannot be controlled in this way since they destroy the boundary condition. Instead, the required normal regularity has to be deduced from the evolution and divergence equations in (1.1), using ideas as in [30]. Appropriate localization

techniques in a boundary collar also need to be employed. Along this way, both anisotropy and inhomogeneity impose an additional challenge. Paired with the dissipativity/observability estimates of Propositions 3.1 & Corollary 3.5, the regularity boost of Proposition 4.1 furnishes the nonlinear stabilizability estimate of Proposition 4.2. Then our main result from Theorem 2.2 is a direct consequence of the standard barrier method (cf. [30, 31, 32, 38, 42], etc.).

In the next section we discuss basic notation & results and present our main Theorem 2.2. The energy and observability-type estimates are proved in Section 3. In Section 4 we state the crucial regularity result of Proposition 4.1 before we establish Theorem 2.2. The proof of this proposition is based on various auxiliary results discussed in Section 5. The technically most demanding part is Section 6, where Proposition 4.1 is shown.

2. PROBLEM SETTINGS AND MAIN RESULTS

Throughout this paper, let $\Omega \subset \mathbb{R}^3$ be a bounded domain with a C^5 -boundary $\Gamma := \partial\Omega$ and the unit outer normal vector $\nu: \Gamma \rightarrow \mathbb{R}^3$.

For an arbitrary, but fixed $T > 0$, let $J = J_T = [0, T]$ denote the time interval, while $\Omega_T = (0, T) \times \Omega$ and $\Gamma_T = (0, T) \times \Gamma$ stand for the interior and the lateral boundary of the time-space cylinder, respectively. For the sake of brevity and convenience, we will often make no distinction between respective spaces of scalar and vector-valued functions, e.g., $L^2(\Omega)$ and $(L^2(\Omega))^3 \simeq L^2(\Omega, \mathbb{R}^3)$, etc. Further, we will frequently omit the domain of integration and merely write \mathcal{H}^k in lieu of $\mathcal{H}^k(\Omega)$, etc., in case the domain is unambiguously clear from the context. We will always employ the standard Lebesgue measure on open domains of \mathbb{R}^3 and the associated surface measure on smooth oriented surfaces Γ in \mathbb{R}^3 . In both cases, these measures will be denoted as dx in respective volumetric or surface integrals.

Our central assumptions employed throughout the paper are the regularity and uniform positivity conditions of the material tensors

$$\begin{aligned} \varepsilon, \mu \in C^3(\overline{\Omega} \times \mathbb{R}^3, \mathbb{R}_{\text{sym}}^{3 \times 3}), \quad \lambda \in C^3_\tau(\Gamma, \mathbb{R}_{\text{sym}}^{3 \times 3}), \\ \varepsilon(x, 0) \geq 2\eta I, \quad \mu(x, 0) \geq 2\eta I \quad \text{for all } x \in \overline{\Omega} \quad \text{and} \quad \lambda(x) \geq \eta I \quad \text{for all } x \in \Gamma \end{aligned} \quad (2.1)$$

for some constant $\eta > 0$, where I stands for the (3×3) -identity matrix. The space $C^3(\overline{\Omega})$ denotes the space of (here: tensor-valued) functions, which – along with their derivatives up to order three – possess continuous extensions onto $\overline{\Omega}$. Likewise, $C^3_\tau(\Gamma) := C^3(\Gamma) \cap L^2_\tau(\Gamma)$ (cf. Equation (2.5)) is the space of “tangential” tensor-valued functions α such that $\text{im}_{\alpha(x)}(\text{span}\{\nu\}^\perp) \subset \text{span}\{\nu\}^\perp$ for all $x \in \Gamma$.

Next, following [30], we introduce matrix-valued “linearizations” ε^d and μ^d of ε and μ via

$$\varepsilon^d_{jk}(x, \xi) = \varepsilon_{jk}(x, \xi) + \sum_{l=1}^3 \partial_{\xi_k} \varepsilon_{jl}(x, \xi) \xi_l, \quad \mu^d_{jk}(x, \xi) = \mu_{jk}(x, \xi) + \sum_{l=1}^3 \partial_{\xi_k} \mu_{jl}(x, \xi) \xi_l$$

for $x \in \overline{\Omega}$, $\xi \in \mathbb{R}^3$ and $j, k \in \{1, 2, 3\}$, which will be used to express the time-derivatives of the left-hand sides of the first two equations in (1.1). We further impose the following additional regularity and symmetry conditions:

$$\partial_\xi \varepsilon, \partial_\xi \mu \in C^3(\overline{\Omega} \times \mathbb{R}^3, \mathbb{R}^{3 \times 3}), \quad \varepsilon^d = (\varepsilon^d)^\top, \quad \text{and} \quad \mu^d = (\mu^d)^\top, \quad (2.2)$$

where $(\cdot)^\top$ denotes the standard matrix transposition operator.

The continuity of respective functions along with assumptions (2.1) furnish the existence of a (possibly) small radius $\tilde{\delta} \in (0, 1]$ such that the following “local positivity” condition

$$\varepsilon(x, \xi) \geq \eta I, \quad \mu(x, \xi) \geq \eta I, \quad \varepsilon^d(x, \xi) \geq \eta I, \quad \mu^d(x, \xi) \geq \eta I \quad (2.3)$$

holds true for all $\xi \in \mathbb{R}^3$ with $|\xi| \leq \tilde{\delta}$ uniformly in $x \in \overline{\Omega}$.

Remark 2.1. The class of functions satisfying our assumptions (2.1) and (2.2) is non-trivial. Indeed, apart from obvious linear candidates, physically relevant instances of genuinely nonlinear tensor responses – both isotropic and anisotropic – can be found in [30, Example 2.1]. For further examples, we refer the reader to the monograph [39].

Continuing, for a smooth vector field u , let $\text{tr}_n u$ be the trace of the normal component $u \cdot \nu$ on Γ of u , while $\text{tr}_t u$ denotes the tangential trace $u \times \nu$ of u on Γ . The tangential component

$$\nu \times (\text{tr}_t u) = \text{tr} u - (\text{tr}_n u)\nu \quad (2.4)$$

of the full trace $\text{tr} u$ will similarly be denoted as tr_τ . By well-known results (cf. Theorems IX.1.1 and IX.1.2 in [14]), the (linear) mappings

$$\text{tr}_n: \mathcal{H}(\text{div}) \rightarrow \mathcal{H}^{-1/2}(\Gamma) \quad \text{and} \quad \text{tr}_t: \mathcal{H}(\text{curl}) \rightarrow (\mathcal{H}^{-1/2}(\Gamma))^3$$

are continuous. Here, the Hilbert spaces

$$\mathcal{H}(\text{curl}) = \left\{ u \in (L^2(\Omega))^3 \mid \text{curl} u \in (L^2(\Omega))^3 \right\} \quad \text{and} \quad \mathcal{H}(\text{div}) = \left\{ u \in (L^2(\Omega))^3 \mid \text{div} u \in L^2(\Omega) \right\}$$

are equipped with the natural inner products. Similarly, following [45, Equation (2.12)], we define the “tangential” L^2 -space

$$L_\tau^2(\Gamma) = \left\{ v \in (L^2(\Gamma))^3 \mid v \cdot \nu = 0 \quad \text{on } \Gamma \right\} \quad (2.5)$$

endowed with the standard L^2 -inner product.

In this paper, we are interested in classical solutions

$$(E, H) \in \bigcap_{k=0}^3 C^k([0, T_{\max}], (\mathcal{H}^{3-k}(\Omega))^6) =: G^3([0, T_{\max}]) = G^3. \quad (2.6)$$

The space $\mathcal{H}^3(\Omega)$ in the equation above is known to be the optimal integer-order Sobolev space for $E(t), H(t)$ from the quasilinear Maxwell system (1.1), cf. [58]. Provided the C^2 -regularity of solutions at $t = 0$, the boundary conditions in the third equation in (1.1) need to be satisfied by respective time-derivates of (E, H) at time $t = 0$. These facts translate into usual “compatibility conditions” naturally arising when treating quasilinear problems. To express these conditions in terms of the initial data $(E^{(0)}, H^{(0)})$, we let $E^{(0)}, H^{(0)} \in (\mathcal{H}^3(\Omega))^3$ and introduce the vector fields

$$\begin{aligned} E^{(1)} &= (\varepsilon^{\text{d}}(E^{(0)}))^{-1} \text{curl} H^{(0)}, & H^{(1)} &= -(\mu^{\text{d}}(H^{(0)}))^{-1} \text{curl} E^{(0)}, \\ E^{(2)} &= (\varepsilon^{\text{d}}(E^{(0)}))^{-1} [\text{curl} H^{(1)} - (\nabla_E \varepsilon^{\text{d}}(E^{(0)}) E^{(1)}) \cdot E^{(1)}], \\ H^{(2)} &= -(\mu^{\text{d}}(H^{(0)}))^{-1} [\text{curl} E^{(1)} + (\nabla_H \mu^{\text{d}}(H^{(0)}) H^{(1)}) \cdot H^{(1)}] \end{aligned} \quad (2.7)$$

as formal representations of $\partial_t(E(t), H(t))|_{t=0}$ for $k \in \{1, 2\}$, where we used the convention

$$((\nabla A)\xi \cdot \eta)_j = \left(\sum_{i,k} \partial_i A_{jk} \xi_k \eta_i \right)_j. \quad (2.8)$$

With the notation from Equations (2.7), for $k \in \{0, 1, 2\}$ we obtain the following compatibility and initial solenoidality conditions

$$\text{tr}_t H^{(k)} + \text{tr}_t (\lambda \text{tr}_t E^{(k)}) = 0 \quad \text{on } \Gamma, \quad \text{div} (\varepsilon(E^{(0)}) E^{(0)}) = \text{div} (\mu(H^{(0)}) H^{(0)}) = 0 \quad \text{on } \Omega. \quad (2.9)$$

We will proceed by invoking the local well-posedness theory recently developed in [52]. Introducing the constant

$$\delta_0 = \min\{1, \tilde{\delta}/C_S\},$$

where C_S denotes the norm of the Sobolev embedding $\mathcal{H}^2(\Omega) \hookrightarrow L^\infty(\Omega)$, we select numbers $T > 0$ and $\delta \in (0, \delta_0]$. The (small) parameter $\delta > 0$ will be chosen appropriately in subsequent proofs. Under the regularity and positivity conditions (2.1)–(2.3), the local well-posedness result in [52,

Theorem 6.4] furnishes the existence of a (small) maximal radius $r(T, \delta) \in (0, r(T, \delta_0)]$ such that for all radii $r \in (0, r(T, \delta)]$ and initial data $E^{(0)}, H^{(0)} \in (\mathcal{H}^3(\Omega))^3$ satisfying the compatibility conditions (2.9) and the smallness assumption

$$\|E^{(0)}\|_{\mathcal{H}^3(\Omega)}^2 + \|H_0\|_{\mathcal{H}^3(\Omega)}^2 \leq r^2, \quad (2.10)$$

there exists a unique classical solution $(E, H) \in G^3([0, T_{\max}))$ (see (2.6)), to the quasilinear Maxwell system (1.1) up to a maximal existence time $T_{\max} \in (T, \infty]$. Further, the fields (E, H) have tangential traces in $\mathcal{H}^3(\Gamma_{T'})$ for $T' < T_{\max}$ and satisfy the estimate

$$\max_{k \in \{0, 1, 2, 3\}} \max_{t \in [0, T]} \left(\|\partial_t^k E(t)\|_{\mathcal{H}^{3-k}(\Omega)}^2 + \|\partial_t^k H(t)\|_{\mathcal{H}^{3-k}(\Omega)}^2 \right) \leq \delta^2 \leq 1 \quad (2.11)$$

along with the solenoidality condition

$$\operatorname{div}(\varepsilon(\cdot, E(t))E(t)) = 0, \quad \operatorname{div}(\mu(\cdot, H(t))H(t)) = 0 \quad \text{in } \Omega \quad (2.12)$$

for all $t \in [0, T_{\max}) =: J_{\max}$. Here and in the sequel, we often view vector fields $E(t, \cdot)$, etc., as a Hilbert space element and write $E(t)$, etc. As noted in the introduction, condition (2.12), i.e., the third equation in (1.1), easily follows from the initial solenoidality assumption in Equation (1.2) and the first two lines of (1.1), cf. [58, Lemma 7.25].

The *a priori* bound (2.11) will repeatedly be invoked throughout the article either explicitly or implicitly. This inequality will prove essential for treating the nonlinearity as it both provides a uniform bound on the solution and furnishes its smallness. Moreover, in view of assumption (2.11), it guarantees the ‘‘hyperbolicity’’ of the system by preserving the uniform positive definiteness of respective nonlinear tensors $\varepsilon(\cdot, E(t))$ and $\mu(\cdot, H(t))$ everywhere in Ω for $t \in [0, T]$, which is indispensable both for the local existence and the global stability of regular solutions.

Fixing now $T = 1$, we obtain $r(\delta) := r(1, \delta)$. Given arbitrary initial data fields $(E^{(0)}, H^{(0)})$ satisfying the compatibility conditions (2.9) and the smallness assumption (2.10), we introduce the final time

$$T_* = \sup \{T \in [1, T_{\max}) \mid \text{global bound (2.11) is valid for } t \in [0, T]\}. \quad (2.13)$$

Hence, the estimate (2.11) holds true on $[0, T_*) =: J_*$. Unless $T_* < \infty$, the blow-up condition in [52, Theorem 6.4] implies $T_{\max} > T_*$ and, therefore, by continuity provided by G^3 ,

$$z(T_*) := \max_{k \in \{0, 1, 2, 3\}} \left(\|\partial_t^k E(T_*)\|_{\mathcal{H}^{3-k}(\Omega)}^2 + \|\partial_t^k H(T_*)\|_{\mathcal{H}^{3-k}(\Omega)}^2 \right) = \delta^2. \quad (2.14)$$

To study the regularity of (E, H) , we need to proceed to time-differentiated versions of Equation (1.1). To this end, letting

$$\widehat{\varepsilon}_k = \begin{cases} \varepsilon(\cdot, E), & k = 0, \\ \varepsilon^{\text{d}}(\cdot, E), & k \in \{1, 2, 3\}, \end{cases} \quad \widehat{\mu}_k = \begin{cases} \mu(\cdot, H), & k = 0, \\ \mu^{\text{d}}(\cdot, H), & k \in \{1, 2, 3\}, \end{cases} \quad (2.15)$$

we arrive at the non-homogeneous system

$$\begin{aligned} \partial_t(\widehat{\varepsilon}_k \partial_t^k E) &= \operatorname{curl} \partial_t^k H - \partial_t f_k, & t \in J_{\max}, x \in \Omega, \\ \partial_t(\widehat{\mu}_k \partial_t^k H) &= -\operatorname{curl} \partial_t^k E - \partial_t g_k, & t \in J_{\max}, x \in \Omega, \\ \operatorname{tr}_t \partial_t^k H - \operatorname{tr}_t(\lambda \operatorname{tr}_t \partial_t^k E) &= 0, & t \in J_{\max}, x \in \Gamma, \end{aligned} \quad (2.16)$$

for $k \in \{0, 1, 2, 3\}$, with the commutator terms

$$\begin{aligned} f_0 = f_1 = 0, \quad f_2 &= (\partial_t \varepsilon^{\text{d}}(\cdot, E)) \partial_t E, \quad f_3 = (\partial_t^2 \varepsilon^{\text{d}}(\cdot, E)) \partial_t E + 2(\partial_t \varepsilon^{\text{d}}(\cdot, E)) \partial_t^2 E, \\ g_0 = g_1 = 0, \quad g_2 &= (\partial_t \mu^{\text{d}}(\cdot, H)) \partial_t H, \quad g_3 = (\partial_t^2 \mu^{\text{d}}(\cdot, H)) \partial_t H + 2(\partial_t \mu^{\text{d}}(\cdot, H)) \partial_t^2 H. \end{aligned} \quad (2.17)$$

In a similar fashion, differentiating Equation (2.12), the higher-order solenoidality properties

$$\operatorname{div}(\varepsilon^{\text{d}}(\cdot, E) \partial_t^k E) = -\operatorname{div} f_k \quad \text{and} \quad \operatorname{div}(\mu^{\text{d}}(\cdot, H) \partial_t^k H) = -\operatorname{div} g_k \quad \text{for } k \in \{1, 2, 3\} \quad (2.18)$$

emerge. In the latter two equations (2.17) and (2.18), the functions $\partial_t f_k$, $\partial_t g_k$, $\operatorname{div} f_k$ and $\operatorname{div} g_k$ belong to $L^\infty(J_*, L^2(\Omega))$ on the strength of the estimate (2.22) stated below. For $k = 3$, the first equation in (1.1) is interpreted in $\mathcal{H}^{-1}(\Omega_T)$, while the divergence operator in Equation (2.16) is viewed as a map from $(L^2(\Omega))^3$ into $\mathcal{H}^{-1}(\Omega)$. Since the forcing terms belong to $L^2(\Omega)$, the traces in Equation (2.16) exist as $\mathcal{H}^{-1/2}(\Gamma)$ -distributions (cf. [58, Section 2.1]).

To facilitate energy estimates, we also consider the following equivalent version of Equation (2.16):

$$\begin{aligned} \varepsilon^{\mathrm{d}}(\cdot, E) \partial_t (\partial_t^k E) &= \operatorname{curl} \partial_t^k H - \tilde{f}_k, & t \in J_{\max}, x \in \Omega, \\ \mu^{\mathrm{d}}(\cdot, H) \partial_t (\partial_t^k H) &= -\operatorname{curl} \partial_t^k E - \tilde{g}_k, & t \in J_{\max}, x \in \Omega, \\ \operatorname{tr}_t \partial_t^k H - \operatorname{tr}_t (\lambda \operatorname{tr}_t \partial_t^k E) &= 0, & t \in J_{\max}, x \in \Gamma, \end{aligned} \quad (2.19)$$

for $k \in \{0, 1, 2, 3\}$ with the new commutator terms

$$\begin{aligned} \tilde{f}_0 &= 0, & \tilde{f}_k &= \sum_{j=1}^k \binom{k}{j} (\partial_t^j \varepsilon^{\mathrm{d}}(\cdot, E)) \partial_t^{k+1-j} E & \text{for } k \in \{1, 2, 3\}, \\ \tilde{g}_0 &= 0, & \tilde{g}_k &= \sum_{j=1}^k \binom{k}{j} (\partial_t^j \mu^{\mathrm{d}}(\cdot, H)) \partial_t^{k+1-j} H & \text{for } k \in \{1, 2, 3\}. \end{aligned} \quad (2.20)$$

Continuing, we define higher-order ‘‘energies’’

$$\begin{aligned} e_k(t) &= \frac{1}{2} \max_{0 \leq j \leq k} \left(\|\widehat{\varepsilon}_k^{1/2} \partial_t^j E(t)\|_{L^2(\Omega)}^2 + \|\widehat{\mu}_k^{1/2} \partial_t^j H(t)\|_{L^2(\Omega)}^2 \right), & e &:= e_3, \\ d_k(t) &= \max_{0 \leq j \leq k} \|\lambda^{1/2} \operatorname{tr}_t \partial_t^j E(t)\|_{L^2(\Gamma)}^2, & d &:= d_3, \\ z_k(t) &= \max_{0 \leq j \leq k} \left(\|\partial_t^j E(t)\|_{\mathcal{H}^{k-j}(\Omega)}^2 + \|\partial_t^j H(t)\|_{\mathcal{H}^{k-j}(\Omega)}^2 \right), & z &:= z_3 \end{aligned} \quad (2.21)$$

of order $k \in \{0, 1, 2, 3\}$ for $t \in J_{\max}$. The ‘‘weight’’ tensors in the definition of e_k and d_k are introduced in light of the energy and observability-type estimates proved in the next section. The ‘‘natural’’ higher-order energies e_k arise from the basic energy associated with Equation (1.1) and (2.19), respectively, the expression d_k measures the squared norms of boundary damping higher-order temporal derivatives, while $z_k(t)$ combine both time- and space-higher-order squared norms, which induce the topology on G^3 after taking the maximum over t . Having shown a stabilizability estimate for e and d with an error term involving z , the goal will be to bound z through e by carefully exploiting the structure of the underlying dynamics.

We proceed with useful *a priori* estimates on various linear and nonlinear quantities involved in our future energy estimates. To this end, here and in the sequel, let c_k and c denote positive constants independent of $t \in [0, T_*]$, T_* , $\delta \in (0, \delta_0]$, $r \in (0, r(\delta)]$, and the ‘‘admissible’’ initial data $(E^{(0)}, H^{(0)})$ satisfying the conditions (2.9) and (2.10). Adopting the arguments from [52], similar to [30], we can estimate

$$\begin{aligned} \|\widehat{\varepsilon}_k(t)\|_\infty, \|\widehat{\mu}_k(t)\|_\infty, \|\widehat{\varepsilon}_k^{-1}(t)\|_\infty, \|\widehat{\mu}_k^{-1}(t)\|_\infty &\leq c, \\ \|\partial_{(t,x)}^\alpha \widehat{\varepsilon}_j(t)\|_{L^2(\Omega)}, \|\partial_{(t,x)}^\alpha \widehat{\mu}_j(t)\|_{L^2(\Omega)} &\leq c(z_k^{1/2}(t) + \delta_{\alpha_0=0}), \\ \max_{2 \leq k \leq 3, 0 \leq j \leq 1} \left(\|\partial_t^j f_k(t)\|_{\mathcal{H}^{4-j-k}(\Omega)} + \|\partial_t^j g_k(t)\|_{\mathcal{H}^{4-j-k}(\Omega)} \right) &\leq cz(t), \\ \|f_2(t)\|_{L^2(\Omega)}, \|g_2(t)\|_{L^2(\Omega)}, \|f_3(t)\|_{L^2(\Omega)}, \|g_3(t)\|_{L^2(\Omega)} &\leq ce_2^{1/2}(t), \\ \|\tilde{f}_k(t)\|_{\mathcal{H}^{3-k}(\Omega)}, \|\tilde{g}_k(t)\|_{\mathcal{H}^{3-k}(\Omega)} &\leq cz(t) \end{aligned} \quad (2.22)$$

for all $0 \leq j, k \leq 3$, $\alpha \in \mathbb{N}_0 \times \mathbb{N}_0^3$ with $|\alpha| = k > 0$ and $t \in J_*$, where, as stated before, the constants c do not depend on t . The expression $\delta_{\alpha_0=0}$ is the Kronecker delta defined as $\delta_{\alpha_0=0} = 1$ if $\alpha_0 = 0$

and $\delta_{\alpha_0=0} = 0$ if $\alpha_0 > 0$. The “correction” term “+c” on the right-hand side of the second line of Equation (2.22) emerges from applying the ∂^α -operator to the x -variable of $\varepsilon(\cdot, E)$ or $\mu(\cdot, H)$.

As previously announced, the thrust of this paper is to prove that the Maxwell system (1.1) subject to a Silver & Müller-type boundary damping admits a unique classical solution in the solution space G^3 defined in Equation (2.6) provided the initial data are sufficiently small and satisfy appropriate regularity and compatibility conditions. When studying a quasilinear dynamics in bounded domains, it is commonly recognized that the global well-posedness goes hand in hand with the exponential stability of classical solutions (cf. [30, 31, 32, 38], etc.). Hence, a big part of the present paper is devoted to showing the desired stability property.

For our main result we need two more assumptions. First, Ω has to be strictly star-shaped with respect to some point $x_0 \in \mathbb{R}^3$, i.e., there exists a number $\bar{\eta} > 0$ such that

$$\nu \cdot m \geq \bar{\eta} > 0 \quad \text{on } \Gamma \quad \text{with } m(x) = x - x_0. \quad (2.23)$$

Second, we suppose that the permittivity and permeability tensors satisfy the lower bounds

$$\varepsilon(x, 0) + (m(x) \cdot \nabla_x) \varepsilon(x, 0) \geq \kappa \varepsilon(x, 0) \quad \text{and} \quad \mu(x, 0) + (m(x) \cdot \nabla_x) \mu(x, 0) \geq \kappa \mu(x, 0) \quad (2.24)$$

for a constant $\kappa > 0$ and all $x \in \bar{\Omega}$. Variants of this condition have often been used in the nonhomogeneous linear and semilinear cases (see, e.g., [6, 16, 43, 44]). Of course, it is satisfied in the homogeneous case where the tensors do not depend on x .

We state our main result which is proved at the end of Section 4.

Theorem 2.2. *Let $\Omega \subset \mathbb{R}^3$ be a bounded, simply connected domain with a smooth boundary $\Gamma := \partial\Omega \in C^5$ satisfying the strict star-shapedness condition (2.23). Further, let the coefficient tensors ε , μ and λ satisfy the assumptions (2.1), (2.2) and (2.24), and let the initial data $E^{(0)}, H^{(0)} \in (\mathcal{H}^3(\Omega))^3$ fulfill the compatibility conditions (2.9) and the smallness assumption (2.10). Then there exists a radius $r > 0$ entering in assumption (2.10) and stability constants $M, \omega > 0$ such that the classical solution $(E, H) \in G^3$ of the Maxwell system (1.1) uniquely exists for all times $t \geq 0$ and exhibits an exponential decay rate*

$$\max_{0 \leq j \leq 3} \left(\|\partial_t^j E(t)\|_{\mathcal{H}^{3-j}(\Omega)}^2 + \|\partial_t^j H(t)\|_{\mathcal{H}^{3-j}(\Omega)}^2 \right) \leq M e^{-\omega t} \|(E^{(0)}, H^{(0)})\|_{\mathcal{H}^3(\Omega)}^2 \quad \text{for all } t \geq 0.$$

3. ENERGY AND OBSERVABILITY-TYPE ESTIMATES

We first establish the basic dissipation of the system from the energy loss at the boundary through the Silver & Müller-type boundary condition. This property will be expressed via the quantities e_k and d_k at all energy levels $0 \leq k \leq 3$. In the proof we apply standard energy techniques.

Proposition 3.1 (Dissipativity inequality). *Let the assumptions of Theorem 2.2 – with the exception of the simple connectedness of Ω and (2.24) – be satisfied. Then, for $0 \leq k \leq 3$ and $0 \leq s \leq t \leq T_*$, the energy estimate*

$$e_k(t) + \int_s^t d_k(\tau) \, d\tau \leq e_k(s) + c_1 \int_s^t z^{3/2}(\tau) \, d\tau \quad (3.1)$$

holds true with positive constants c_1 and η independent of s, t .

To streamline the exposition, we consider the following linear non-autonomous Maxwell system (motivated by (2.19)). Let $\alpha, \beta \in W^{1,\infty}(J, L^\infty(\Omega, \mathbb{R}_{\text{sym}}^{3 \times 3}))$ with $\alpha, \beta \geq \eta$, $\varphi, \psi \in L^2(\Omega_T)^3$, $u^{(0)}, v^{(0)} \in (L^2(\Omega))^3$. By [52, Proposition 3.1], the non-homogeneous problem

$$\begin{aligned} \alpha \partial_t u &= \text{curl } v + \varphi && \text{in } J \times \Omega, \\ \beta \partial_t v &= -\text{curl } u + \psi && \text{in } J \times \Omega, \\ v \times \nu + (\lambda(u \times \nu)) \times \nu &= 0 && \text{in } J \times \Gamma, \end{aligned} \quad (3.2)$$

$$u(0) = u^{(0)}, \quad v(0) = v^{(0)} \quad \text{in } \Omega$$

possesses a unique weak solution $(u, v) \in C^0([0, T], (L^2(\Omega))^6)$ with $(\text{tr}_\tau u, \text{tr}_\tau v) \in L^2(\Omega_T, \mathbb{R}^6)$. See also the earlier paper [8] for a slight variant. The latter extra regularity of the trace in time and space – in addition to what one would expect from the mapping $\text{tr}_\tau: L^2(\Omega) \rightarrow \mathcal{H}^{-1/2}(\Gamma)$ – is referred as “hidden regularity” and often occurs in hyperbolic problems such as the wave equation [28, 34, 35], the dynamic equations of elasticity [26], fluid/structure interaction systems [11], plate and beam equations [29, 33], etc. Proposition 3.1 in [52] also yields the following energy equality.

Lemma 3.2. *Under the above assumptions, for $0 \leq s \leq t \leq T$ we have*

$$\begin{aligned} & \|\alpha^{1/2}(T)u(T)\|_{L^2(\Omega)}^2 + \|\beta^{1/2}(T)v(T)\|_{L^2(\Omega)}^2 + 2 \int_s^t \|\lambda \text{tr}_t u \cdot \text{tr}_t v\|_{L^2(\Gamma)}^2 d\tau \\ &= \|\alpha^{1/2}(0)u(0)\|_{L^2(\Omega)}^2 + \|\beta^{1/2}(0)v(0)\|_{L^2(\Omega)}^2 + 2 \int_s^t \int_\Omega (u \cdot \varphi + v \cdot \psi) dx d\tau \\ &+ \int_s^t \int_\Omega (u \cdot (\partial_t \alpha)u + v \cdot (\partial_t \beta)v) dx d\tau. \end{aligned}$$

In view of (2.19), Proposition 3.1 for $k \in \{1, 2, 3\}$ is a consequence of the bounds in (2.22) and Lemma 3.2 with $\alpha = \widehat{\varepsilon}_k$, $\beta = \widehat{\mu}_k$, $\varphi = -\tilde{f}_k$, and $\psi = -\tilde{g}_k$. To show Proposition 3.1 for $k = 0$, we can use the ample regularity of our solution $(E, H) \in G^3$ to Maxwell system (1.1). Using the symmetry of the coefficients, inserting Equation (1.1) and integrating by parts, we compute

$$\begin{aligned} & \partial_t \int_\Omega (\widehat{\varepsilon}_0 E \cdot E + \widehat{\mu}_0 H \cdot H) dx \\ &= 2 \int_\Omega (\partial_t(\widehat{\varepsilon}_0 E) \cdot E + \partial_t(\widehat{\mu}_0 H) \cdot H) dx + \int_\Omega (\widehat{\varepsilon}_0 E \cdot (\partial_t \widehat{\varepsilon}_0^{-1}) \widehat{\varepsilon}_0 E + \widehat{\mu}_0 H \cdot (\partial_t \widehat{\mu}_0^{-1}) \widehat{\mu}_0 H) dx \\ &= 2 \int_\Omega (\text{curl } H \cdot E - \text{curl } E \cdot H) dx - \int_\Omega (E \cdot (\partial_t \widehat{\varepsilon}_0) E + H \cdot (\partial_t \widehat{\mu}_0) H) dx \\ &= -2 \int_\Gamma (H \times \nu) \cdot E dx - \int_\Omega (E \cdot (\partial_t \widehat{\varepsilon}_0) E + H \cdot (\partial_t \widehat{\mu}_0) H) dx \\ &= 2 \int_\Gamma ((\lambda(E \times \nu)) \times \nu) \cdot E dx - \int_\Omega (E \cdot (\partial_t \widehat{\varepsilon}_0) E + H \cdot (\partial_t \widehat{\mu}_0) H) dx \\ &= -2 \int_\Gamma |\lambda^{1/2}(E \times \nu)|^2 dx - \int_\Omega (E \cdot (\partial_t \widehat{\varepsilon}_0) E + H \cdot (\partial_t \widehat{\mu}_0) H) dx. \end{aligned}$$

We now integrate in time over (s, t) with respect to τ and apply the estimates in (2.22). Proposition 3.1 also follows for $k = 0$, too.

Next, we aim to prove a nonlinear observability-type estimate involving the higher-order natural energies e_k and the dissipation functionals d_k . To this end, we now have to assume that Ω is strictly star-shaped (viz. Equation (2.23)) and that ε and μ satisfy the lower bound (2.24).

Proposition 3.3 (Observability-like inequality). *Under the assumptions of Theorem 2.2, there is a radius $\delta_1 \in (0, \delta_0]$ such that for all $\delta \in (0, \delta_1]$ in (2.11), the estimate*

$$\int_s^t e_k(\tau) d\tau \leq c_2 \int_s^t d_k(\tau) d\tau + c_3(e_k(t) + e_k(s)) + c_4 \int_s^t z^{3/2}(\tau) d\tau$$

holds true for $k \in \{0, 1, 2, 3\}$, $0 \leq s \leq t < T_*$, and some constants c_j independent of s and t .

To prove Proposition 3.3 for $k \geq 1$, we again look at a non-autonomous Maxwell system. Since now the divergence conditions are involved, it is more convenient to use the system

$$\partial_t(\alpha u) = \text{curl } v + \partial_t \varphi \quad \text{in } J \times \Omega,$$

$$\begin{aligned}
\partial_t(\beta v) &= -\operatorname{curl} u + \partial_t \psi && \text{in } J \times \Omega, \\
v \times \nu + (\lambda(u \times \nu)) \times \nu &= 0 && \text{in } J \times \Gamma, \\
u(0) &= u^{(0)}, \quad v(0) = v^{(0)} && \text{in } \Omega,
\end{aligned} \tag{3.3}$$

which better matches (2.16). We take coefficients $\alpha, \beta \in C^1(J \times \overline{\Omega}, \mathbb{R}_{\text{sym}}^{3 \times 3})$ with $\alpha, \beta \geq \eta$ and data $\varphi, \psi \in C^1([0, T], L^2(\Omega)^3) \cap C([0, T], \mathcal{H}^1(\Omega)^3) = G^1$ and $u^{(0)}, v^{(0)} \in (L^2(\Omega))^3$ such that $\operatorname{div} \varphi, \operatorname{div} \psi \in C(J, L^2(\Omega))$, $\operatorname{div}(\alpha(0)u^{(0)}) = \operatorname{div} \varphi(0)$, and $\operatorname{div}(\beta(0)v^{(0)}) = \operatorname{div} \psi(0)$. Proposition 1.1 of [8] or Proposition 3.1 of [52] provide a unique weak solution $(u, v) \in C^0([0, T], (L^2(\Omega))^6)$ of (3.3) with $(\operatorname{tr}_\tau u, \operatorname{tr}_\tau v) \in L^2(\Omega_T, \mathbb{R}^6)$. We will see below that our assumptions and Theorem 1.2 of [8] imply that the full traces $(\operatorname{tr} u, \operatorname{tr} v)$ belong to $L^2(\Omega_T, \mathbb{R}^6)$. We further assume that

$$\begin{aligned}
\|\alpha\|_{L^\infty}, \|\beta\|_{L^\infty}, \|\lambda\|_{L^\infty} &\leq M, \quad \text{for some } M > 0, \\
\tilde{\alpha} := \alpha + (m \cdot \nabla)\alpha &\geq \frac{\kappa\alpha}{2} \geq \frac{\kappa\eta}{2}, \quad \text{and} \quad \tilde{\beta} := \beta + (m \cdot \nabla)\beta \geq \frac{\kappa\beta}{2} \geq \frac{\kappa\eta}{2}
\end{aligned} \tag{3.4}$$

for some constants $\eta, \kappa > 0$, which are given by (2.2) and (2.24) later on.

Lemma 3.4. *Besides the assumptions stated above, we also require that the domain Ω is strictly star-shaped (viz. Equation (2.23)). Let $(u, v) \in C^0([0, T], L^2(\Omega, \mathbb{R}^6))$ be the weak solution of Equation (3.3). Then the traces $(\operatorname{tr} u, \operatorname{tr} v)$ belong to $L^2(\Omega_T, \mathbb{R}^6)$ and the fields (u, v) fulfill the estimate*

$$\begin{aligned}
&\frac{\kappa}{4} \int_s^t \int_\Omega (\alpha u \cdot u + \beta v \cdot v) \, dx \, d\tau + \frac{\bar{\eta}}{4} \int_s^t \int_\Gamma \operatorname{tr} (\alpha u \cdot u + \beta v \cdot v) \, dx \, d\tau \\
&\leq \frac{\|m\|_{L^\infty}^2}{\bar{\eta}} M(1+M)^2 \int_s^t \int_\Gamma |u \times \nu|^2 \, dx \, d\tau + \frac{M\|m\|_{L^\infty}}{2} (\|(u(t), v(t))\|_{L^2}^2 + \|(u(s), v(s))\|_{L^2}^2) \\
&\quad + \|m\|_{L^\infty} \int_s^t \int_\Omega (M(|\partial_t \varphi| \cdot |v| + |\partial_t \psi| \cdot |v|) + (|\operatorname{div} \varphi| |u| + |\operatorname{div} \psi| |v|)) \, dx \, d\tau
\end{aligned} \tag{3.5}$$

for $0 \leq s \leq t \leq T$.

Proof. We let $s = 0$ without loss of generality.

1) We start by regularizing the data. To this end, we first look at the homogeneous autonomous version of (3.3) with $\varphi = \psi = 0$, time-independent coefficients $\alpha(0)$ and $\beta(0)$, and initial ‘‘charges’’ $\operatorname{div}(\alpha(0)u^{(0)}), \operatorname{div}(\beta(0)v^{(0)}) \in L^2(\Omega)$. In this case the coefficients commute with the time derivative. Theorem 1.2 of [8] implies that the weak solutions (\tilde{u}, \tilde{v}) of (3.3) with these coefficients and data form a C_0 -semigroup on the space

$$X = \{(u_0, v_0) \in L^2(\Omega)^6 \mid \operatorname{div}(\alpha(0)u_0), \operatorname{div}(\beta(0)v_0) \in L^2(\Omega)\}$$

endowed with the norm given by $\|u_0\|_2^2 + \|v_0\|_2^2 + \|\operatorname{div}(\alpha(0)u_0)\|_2^2 + \|\operatorname{div}(\beta(0)v_0)\|_2^2$. (We note that $\operatorname{div}(\alpha(0)\tilde{u}(t)) = \operatorname{div}(\alpha(0)u_0)$ and $\operatorname{div}(\beta(0)\tilde{v}(t)) = \operatorname{div}(\beta(0)v_0)$ for all $t \geq 0$. Moreover, one has to use the remarks at the end of the first section of [8], to extend the results of this paper from scalar λ to positive definite and bounded ones.) This semigroup has a generator A whose domain $D(A)$ is dense in X . For data $(u^{(0)}, v^{(0)}) \in D(A)$ the solution (\tilde{u}, \tilde{v}) is an element of $C^1([0, \infty), X)$, and hence $(\tilde{u}(t), \tilde{v}(t))$ is an element of $\mathcal{H}(\operatorname{curl})^2$ by the evolution equations. Since these fields also satisfy the boundary condition in (3.3), Lemma 5.1 shows that (\tilde{u}, \tilde{v}) is contained in $C([0, \infty), \mathcal{H}^1(\Omega))$. (Of course, the proof of this lemma is independent of the present one.) In particular, the initial values $(\tilde{u}(0), \tilde{v}(0)) \in (A)$ are contained in $\mathcal{H}^1(\Omega)$ and fulfill the boundary condition.

Let now $(u^{(0)}, v^{(0)})$, φ , and ψ be given as in the statement, and let $(u, v) \in C([0, T], L^2(\Omega))$ be the solution of (3.3). In view of the previous paragraph, there are functions $(u_n^{(0)}, v_n^{(0)})$ in $\mathcal{H}^1(\Omega)$ that satisfy the third line of (3.3) and tend to $(u^{(0)}, v^{(0)})$ in X . We can also construct maps $\varphi_n, \psi_n \in C^2(\overline{\Omega_T})$ which tend to φ respectively ψ in G^1 . Theorem 1.3 in [8] then provides solutions $(u_n, v_n) \in G^1$ of (3.3). Theorem 1.2 in [8] now shows that (u_n, v_n) and $(\operatorname{tr} u_n, \operatorname{tr} v_n)$ tend to (u, v)

and $(\operatorname{tr} u, \operatorname{tr} v)$ in $C([0, T], L^2(\Omega))$ respectively $L^2(\Gamma_T)$. Below we establish the asserted estimate for (u_n, v_n) , so that the claims will then follow by approximation.

2) We may thus assume that (u, v) belongs to G^1 and use the given data. We set $\Omega_t := (0, t) \times \Omega$ and $\Gamma_t := (0, t) \times \Gamma$. Now, in view of Equation (3.3), we can write

$$\begin{aligned} J &:= \int_{\Omega} \partial_t((m \times \alpha u) \cdot \beta v) \, dx & (3.6) \\ &= \int_{\Omega} \left((m \times (\operatorname{curl} v + \partial_t \varphi)) \cdot \beta v + (m \times \alpha u) \cdot (-\operatorname{curl} u + \partial_t \psi) \right) \, dx \\ &= - \int_{\Omega} \operatorname{curl} v \cdot (m \times \beta v) \, dx - \int_{\Omega} \operatorname{curl} u \cdot (m \times \alpha u) \, dx + \int_{\Omega} (m \times \partial_t \varphi) \cdot \beta v + (m \times \alpha u) \cdot \partial_t \psi \, dx. \end{aligned}$$

Next, we proceed as in [16, Equations (3.6)–(3.8)]. First, we obtain

$$\int_{\Omega} \operatorname{curl} v \cdot (m \times \beta v) \, dx = \int_{\Omega} v \cdot \operatorname{curl}(m \times \beta v) \, dx + \int_{\Gamma} (\nu \times v) \cdot (m \times \beta v) \, dx$$

and compute

$$\begin{aligned} \operatorname{curl}(m \times \beta v) &= \operatorname{div}(\beta v)m + (\beta v \cdot \nabla)m - \operatorname{div}(m)\beta v - (m \cdot \nabla)(\beta v) \\ &= \operatorname{div}(\beta v)m - 2\beta v - (m \cdot \nabla)(\beta v). \end{aligned}$$

Applying div the second equation in (3.2) and integrating in time, it follows

$$\operatorname{div}(\beta(t)v(t)) = \operatorname{div}(\beta(0)v^{(0)}) + \operatorname{div} \psi(t) - \operatorname{div} \psi(0) = \operatorname{div} \psi(t).$$

Integrating over $(0, t)$ and performing partial integration, we arrive at

$$\begin{aligned} \int_{\Omega_t} \operatorname{curl} v \cdot (m \times \beta v) \, d(\tau, x) &= -\frac{1}{2} \int_{\Omega_t} |\tilde{\beta}^{1/2} v|^2 \, d(\tau, x) - \frac{1}{2} \int_{\Gamma_t} (\nu \cdot m)(\beta v \cdot v) \, d(\tau, x) \\ &\quad + \int_{\Omega_t} m \cdot v \operatorname{div} \psi \, d(\tau, x) + \int_{\Gamma_t} (\nu \times v) \cdot (m \times \beta v) \, d(\tau, x). \end{aligned}$$

(Recall the definition of $\tilde{\alpha}$ and $\tilde{\beta}$ in Equation (3.4).) We proceed analogously with u . Invoking the identity (3.6) and using the boundary condition in (3.3), we get

$$\begin{aligned} &\int_{\Omega} (m \times \alpha(t)u(t))\beta(t)v(t) \, dx - \int_{\Omega} (m \times \alpha(0)u(0))\beta(0)v(0) \, dx = \int_0^t J \, d\tau \\ &= \frac{1}{2} \int_{\Omega_t} (\tilde{\alpha}u \cdot u + \tilde{\beta}v \cdot v) \, d(\tau, x) + \frac{1}{2} \int_{\Gamma_t} (\nu \cdot m) \operatorname{tr}(\alpha u \cdot u + \beta v \cdot v) \, d(\tau, x) \\ &\quad - \int_{\Gamma_t} ((\nu \times u) \cdot (m \times \alpha u) + (\lambda(u \times \nu) \times \nu) \cdot (m \times \beta v)) \, d(\tau, x) \\ &\quad + \int_{\Omega_t} \left((m \times \partial_t \varphi) \cdot \beta v + (m \times \alpha u) \cdot \partial_t \psi - ((m \cdot u) \operatorname{div} \varphi + m \cdot v \operatorname{div} \psi) \right) \, d(\tau, x). \end{aligned}$$

By virtue of the strict star-shapedness condition (2.23) and the assumption (3.4), we now obtain

$$\begin{aligned} &\frac{\kappa}{4} \int_{\Omega_t} (\alpha u \cdot u + \beta v \cdot v) \, d(\tau, x) + \frac{\bar{\eta}}{2} \int_{\Gamma_t} \operatorname{tr}(\alpha u \cdot u + \beta v \cdot v) \, d(\tau, x) \\ &\leq \frac{\|m\|_{L^\infty}}{2} M^2 (\|u(t)\|_{L^2}^2 + \|v(t)\|_{L^2}^2 + \|u(0)\|_{L^2}^2 + \|v(0)\|_{L^2}^2) \\ &\quad + \frac{M\|m\|_{L^\infty}^2}{4} (1+M)^2 \frac{1}{\theta} \int_{\Gamma_t} |u \times \nu|^2 \, d(\tau, x) + \theta \int_{\Gamma_t} \operatorname{tr}(\alpha u \cdot u + \beta v \cdot v) \, d(\tau, x) \\ &\quad + \|m\|_{L^\infty} \int_{\Omega_t} \left[M(|\partial_t \varphi| |v| + |\partial_t \psi| |u|) + (|\operatorname{div} \varphi| |u| + |\operatorname{div} \psi| |v|) \right] \, d(\tau, x), \end{aligned}$$

where we applied Young's inequality. Selecting $\theta = \bar{\eta}/4$, the assertion follows. \square

We are now able to prove Proposition 3.3. First, let $k \in \{1, 2, 3\}$. By continuity and (2.24), we can choose a number $\delta_1 \in (0, \delta_0]$ such that the tensors $\alpha := \widehat{\varepsilon}_k = \varepsilon^d(\cdot, E)$ and $\beta := \widehat{\mu}_k = \mu^d(\cdot, H)$ satisfy the conditions (3.4) for all $\delta \in (0, \delta_1]$ in (2.11). Further, let $\varphi := -f_k$ and $\psi := -g_k$ for the functions f_k and g_k defined in Equation (2.17). As in (2.22), one can see that they belong to G^1 since $(E, H) \in G^3$. Equation (2.18) at $t = 0$ yields $\operatorname{div}(\alpha(0)u^{(0)}) = \operatorname{div} \varphi(0)$ and $\operatorname{div}(\beta(0)v^{(0)}) = \operatorname{div} \psi(0)$. So we can apply Lemma 3.4. Combined with Equations (2.18) and (2.22), it implies the assertion of Proposition 3.3 for $k \geq 1$. The case $k = 0$ can be treated in the same way with $\varphi = \psi = 0$, but here one does not need a regularization step since $(E, H) \in G^3$.

Arguing as in the proof of [30, Corollary 3.5], Proposition 3.1 and 3.3 furnish the following “incomplete” nonlinear stabilizability inequality for time-like higher-order energies e_k .

Corollary 3.5. *Suppose the conditions of Theorem 2.2 hold true and that $\delta \in (0, \delta_1]$ in (2.11) with the number δ_1 from Proposition 3.3. Then we have the estimate*

$$e_k(t) + \int_s^t e_k(\tau) \, d\tau \leq C_1 e_k(s) + C_2 \int_s^t z^{3/2}(\tau) \, d\tau \quad (3.7)$$

for $0 \leq s \leq t < T_*$ and $k \in \{0, 1, 2, 3\}$, where the constants C_k depend neither on t nor on s .

In the linear or the Lipschitzian semilinear situation (cf. [17]), one can restrict oneself to the case $k = 0$ and show that the latter term on the right-hand side of Equation (3.7) vanishes. In this situation, the squared norms $e_0(t)$ and $z_0(t)$ are equivalent. This renders the “incomplete” stabilizability inequality complete and the exponential decay of $e_0(t) \sim z_0(t)$ follows similarly to the proof of Datko & Pazy's theorem (cf. [30, Remark 4.2]). In contrast, in the quasilinear situation we are concerned with, such an argument is impossible at the basic energy level. Hence, a genuinely quasilinear strategy needs to be developed.

4. UNIFORM STABILIZABILITY INEQUALITY AND PROOF OF THEOREM 2.2

The natural higher-order energies e_k in the “incomplete” stabilizability estimate (3.7) in Corollary 3.5 only contain higher-order temporal derivatives of the solution pair (E, H) and, therefore, do not match the topology of the solution space G^3 . For the estimate (3.7) to be complete, the “missing” higher-order time-space derivatives need to be recovered so that both sides of the inequality can be formulated in terms of $z(t)$. Trivially, the squared norms $e(t)$ and $z(t)$ are not equivalent in general. At the same time, along classical solutions to the quasilinear Maxwell system (1.1), we will be able to bound $z(t)$ by a multiple of $e(t)$ plus a quadratic term in $z(t)$. This amounts to a careful higher-order regularity analysis of solutions. Namely, in Section 6, we prove the following regularity boost result.

Proposition 4.1. *Under conditions of Proposition 3.1, there exist constants c_5 and c_6 independent of t such that*

$$z(t) \leq c_5 e(t) + c_6 z^2(t) \quad \text{for } t \in [0, T_*]. \quad (4.1)$$

A result reminiscent of Proposition 4.1 for local-in-time solutions to Equation (1.1) has recently been obtained in [52]. Here, our goal is to improve upon these results by showing the estimate holds uniform in time, i.e., the constants do not blow-up, provided the initial data are sufficiently small. This will result in a uniform stabilizability inequality (4.2) stated in Proposition 4.2 below. Combining Corollary 3.5 and Proposition 4.1, we derive (4.2) in a fashion vaguely reminiscent of the proof of Proposition 4.1 of [30].

Proposition 4.2 (Nonlinear stabilizability inequality). *Suppose the conditions of Theorem 2.2 hold true and that $\delta \in (0, \delta_1]$ in (2.11) with the number δ_1 from Proposition 3.3. For all initial data*

satisfying the smallness condition (2.10) with $r \in (0, r(\delta)]$ the associated classical solution (E, H) fulfills the uniform stabilizability inequality

$$z(t) + \int_s^t z(\tau) d\tau \leq C \left(z(s) + z^2(t) + \int_s^t z^2(\tau) d\tau \right) \quad \text{for} \quad 0 \leq s < t < T_*, \quad (4.2)$$

where the complete (squared) norm z is defined in (2.21) and the constant C is independent of t and the radius $r \in (0, r(\delta)]$.

Proof. Utilizing Equation (3.7) from Corollary 3.5 (summed up over $k \in \{0, 1, 2, 3\}$) and Equation (4.1) from Proposition 4.1, we estimate

$$\begin{aligned} c_5^{-1}z(t) - c_5^{-1}c_6z^2(t) + \int_s^t \left(c_5^{-1}z(\tau) - c_5^{-1}c_6z^2(\tau) \right) d\tau &\leq e(t) + \int_s^t e(\tau) d\tau \\ &\leq C_1e(s) + 4C_2 \int_s^t z^{3/2}(\tau) d\tau \\ &\leq C_1z(s) + 4C_2 \int_s^t z^{3/2}(\tau) d\tau \end{aligned}$$

and, therefore,

$$c_5^{-1}z(t) + c_5^{-1} \int_s^t z(\tau) d\tau \leq C_1z(s) + c_5^{-1}c_6z^2(t) + \int_s^t \left(4C_2z^{3/2}(\tau) + c_5^{-1}c_6z^2(\tau) \right) d\tau. \quad (4.3)$$

Using Young's inequality to estimate $z^{3/2}(\tau) \leq \frac{1}{2}(z(\tau) + z^2(\tau))$, the estimate in (4.3) yields the desired inequality with an appropriate positive constant C . \square

Theorem 2.2 can now easily be proved, cf. [30] or [31].

Proof of Theorem 2.2. Fixing $\delta = \min\{\delta_1, 1/2C\}$, estimate (4.2) furnishes the enhanced linear stabilizability inequality

$$z(t) + \int_s^t z(\tau) d\tau \leq 2Cz(s) \quad \text{for} \quad 0 \leq s \leq t < T_*. \quad (4.4)$$

If we have here $T_* = \infty$, the result follows as in the linear case, see [30, Remark 4.2] or the proofs of [32, Corollary 5.7] or [6, Theorem A.1].

Suppose that $T_* < \infty$. Equation (2.14) then yields $z(T_*) = \delta^2$. On the other hand, (4.4) shows that $z(t) \leq 2Cz(0)$ for $0 \leq t < T_*$ and initial data with $\|(E_0, H_0)\|_{\mathcal{H}^3}^2 \leq r^2$ for all $r \in (0, r(\delta)]$. (The number $r(\delta) > 0$ was introduced before (2.13).) Formulas (2.19) and (2.22) then imply

$$z(t) \leq sCz(0) \leq 2c_0C \|(E_0, H_0)\|_{\mathcal{H}^3}^2 \leq 2c_0Cr^2$$

for a constant $c_0 > 0$. We now fix the radius

$$r := \min \left\{ r(\delta), \frac{\delta}{\sqrt{4c_0C}} \right\}. \quad (4.5)$$

As a result, the quantity $z(t)$ is bounded by $\delta^2/2$ for $t < T_*$ and by continuity also for $t = T_*$. This fact contradicts $z(T_*) = \delta^2$, and hence $T_* = \infty$. \square

5. AUXILIARY RESULTS

5.1. curl-div-estimates. The following lemma is a variant of [12, Lemma 4.5.5] for tensors λ . In a certain sense, it ensures optimal regularity for the ‘‘elliptic part’’ of Equation (3.3).

Lemma 5.1. *Let Ω be simply connected. Assume that $u, v \in \mathcal{H}(\text{curl})$ and $\alpha, \beta \in W^{1,\infty}(\Omega, \mathbb{R}^{3 \times 3}_{\text{sym}})$ satisfy $\alpha, \beta \geq \eta > 0$, $\text{div}(\alpha u), \text{div}(\beta v) \in L^2(\Omega)$ and $v \times \nu + \lambda(u \times \nu) \times \nu =: h \in \mathcal{H}^{1/2}(\Gamma, \mathbb{R}^3)$. Then u and v belong to $\mathcal{H}^1(\Omega, \mathbb{R}^3)$ and*

$$\|u\|_{\mathcal{H}^1} + \|v\|_{\mathcal{H}^1} \leq c(\|\text{curl } u\|_{L^2} + \|\text{curl } v\|_{L^2} + \|\text{div}(\alpha u)\|_{L^2} + \|\text{div}(\beta v)\|_{L^2} + \|h\|_{\mathcal{H}^{1/2}(\Gamma)}).$$

Proof. Proposition IX.1.3 and Remark IX.1.4 of [14] yield a function $w \in \mathcal{H}^1(\Omega, \mathbb{R}^3)$ with $\text{curl } u = \text{curl } w$, $\text{div } w = 0$, and $\nu \cdot w = 0$ on Γ . Moreover, $\|w\|_{\mathcal{H}^1} \leq c\|u\|_{L^2}$. Further, $u - w = \nabla \varphi$ for some $\varphi \in \mathcal{H}^1(\Omega)$ by [14, Proposition IX.1.2]. Hence,

$$\text{div}(\alpha \nabla \varphi) = \text{div}(\alpha u) - \text{div}(\alpha w) \in L^2(\Omega).$$

As in Definition 2.2 of [10] we let $\nabla_{\Gamma} \varphi = \nu \times (\nabla \varphi \times \nu)$ be the tangential trace of $\nabla \varphi$, and we define div_{Γ} by duality. We can write $u \times \nu = Bu$ with

$$B := \begin{pmatrix} 0 & \nu_3 & -\nu_2 \\ -\nu_3 & 0 & \nu_1 \\ \nu_2 & -\nu_1 & 0 \end{pmatrix}.$$

The matrix $\tilde{\lambda} := B^{\top} \lambda B$ is again symmetric, positive definite, and Lipschitzian in x . Observe that

$$\tilde{\lambda}(\nu \times (\nabla \varphi \times \nu)) = B^{\top} \lambda (\nabla \varphi \times \nu) = -(\lambda (\nabla \varphi \times \nu)) \times \nu.$$

Invoking [10, Theorem 2.4] and the boundary condition, we thus deduce

$$\tilde{\lambda} \nabla_{\Gamma} \varphi = (\lambda(w \times \nu)) \times \nu - (\lambda(u \times \nu)) \times \nu = (\lambda(w \times \nu)) \times \nu + v \times \nu - h =: \psi \in \mathcal{H}^{-1/2}(\text{div}_{\Gamma}),$$

where we also use the fact $v \in \mathcal{H}(\text{curl})$ and Theorem 2.4 in [10]. Hence, $\text{div}_{\Gamma}(\tilde{\lambda} \nabla_{\Gamma} \varphi) \in H^{-1/2}(\Gamma)$ and the standard elliptic theory implies that $\text{tr } \varphi$ belongs to $H^{3/2}(\Gamma)$. Again, by elliptic results, it follows that φ is contained in $\mathcal{H}^2(\Omega)$ and satisfies

$$\begin{aligned} \|\varphi\|_{\mathcal{H}^2(\Omega)} &\leq c(\|\text{div}(\alpha u)\|_{L^2(\Omega)} + \|\text{div}(\alpha w)\|_{L^2(\Omega)} + \|\text{div}_{\Gamma} \psi\|_{\mathcal{H}^{-1/2}(\Gamma)}) \\ &\leq c(\|\text{div}(\alpha u)\|_{L^2(\Omega)} + \|w\|_{\mathcal{H}^1(\Omega)} + \|h\|_{\mathcal{H}^{1/2}(\Omega)} + \|\text{curl } v\|_{L^2(\Omega)}) \\ &\leq c(\|\text{div}(\alpha u)\|_{L^2(\Omega)} + \|\text{curl } u\|_{L^2(\Omega)} + \|h\|_{\mathcal{H}^{1/2}(\Omega)} + \|\text{curl } v\|_{L^2(\Omega)}). \end{aligned}$$

We have thus shown the desired estimate for u . The inequality for v is proved similarly, but here one can directly use that $\nu \times (v \times \nu)$ is bounded in $\mathcal{H}^{1/2}(\Gamma)$ by $c\|\text{curl } v\|_{L^2(\Omega)}$. \square

5.2. Tangential and normal derivatives. For a sufficiently small $a > 0$, consider the ‘‘boundary collar’’ $\Gamma_a := \{x \in \bar{\Omega} \mid \text{dist}(x, \Gamma) < a\}$ of Γ with $\text{dist}(x, \Gamma) := \max_{y \in \Gamma} |x - y|$. Fix tangential vectors $\{\tau^1(x), \tau^2(x)\}$ spanning the tangential plane at each $x \in \Gamma$. Since $\Gamma \in C^5$, the vector bundle $\{\tau^1(x), \tau^2(x), \nu(x)\}$ can be extended in C^4 from Γ to Γ_a such that $\{\tau^1(x), \tau^2(x), \nu(x)\}$ form an orthonormal basis of \mathbb{R}^3 for each $x \in \Gamma_a$.

Similar to [30, Section 5.2], we introduce the following differential operators. For $\xi, \zeta \in \{\tau^1, \tau^2, \nu\}$, $u \in \mathbb{R}^3$ and $a \in \mathbb{R}^{3 \times 3}$, let

$$\partial_{\xi} = \sum_{j=1}^3 \xi_j \partial_j, \quad u_{\xi} = u \cdot \xi, \quad u^{\xi} = u_{\xi} \xi, \quad u^{\tau} = \sum_{i=1}^2 u_{\tau^i} \tau^i, \quad a_{\xi\zeta} = \xi^{\top} a \zeta \quad \text{in } \Gamma_a.$$

With this notation, for a smooth function φ , we can write

$$\nabla \varphi = \sum_{\xi \in \{\tau^1, \tau^2, \nu\}} \xi (\xi \cdot \nabla \varphi) = \sum_{\xi \in \{\tau^1, \tau^2, \nu\}} \xi \partial_{\xi} \varphi \quad \text{and} \quad \partial_j \varphi = \sum_{\xi \in \{\tau^1, \tau^2, \nu\}} \xi_j \partial_{\xi} \varphi \quad \text{in } \Gamma_a.$$

Therefore, observing the identity

$$\text{curl } u = \partial_1(0, -u_3, u_2)^{\top} + \partial_2(u_3, 0, -u_1)^{\top} + \partial_3(-u_2, u_1, 0)^{\top},$$

the curl-operator can be expressed as

$$\operatorname{curl} u = \sum_{j=1}^3 J_j \partial_j = \sum_{j=1}^3 \sum_{\xi \in \{\tau^1, \tau^2, \nu\}} J_j \xi_j \partial_\xi = \sum_{\xi \in \{\tau^1, \tau^2, \nu\}} J(\xi) \partial_\xi,$$

where

$$J_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad J_2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad J_3 = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad J(\xi) = \sum_{j=1}^3 \xi_j J_j.$$

(The matrix $-J(\nu)$ was called B in the proof of Lemma 5.1.) Note that

$$\ker(J(\nu)) = \operatorname{span}\{\nu\} \quad \text{and} \quad (\ker(J(\nu)))^\top = \operatorname{span}\{\tau^1, \tau^2\} \quad \text{in } \Gamma_a.$$

In particular, factoring out the null space, we can write $J(\nu)u = J(\nu)u^\tau$. Moreover, the restriction of $J(\nu)$ onto the span of $\{\tau^1, \tau^2\}$ is invertible with an inverse $R(\nu)$.

5.3. Reconstructing the normal derivatives. Typically, when showing the “full” Sobolev spatial regularity for elliptic boundary-value problems, the regularity of the boundary symbol proves very beneficial. Unfortunately, this property is violated by the Maxwell system. Indeed, the associated boundary-value problem is characteristic since the boundary symbol $J(\nu)$ has a nontrivial kernel $\operatorname{span}\{\nu\}$ and, thus, is singular. Hence, additional effort is needed to reconstruct the regularity in the normal component. Following [30, Section 5.2], we employ the so-called “curl-div-strategy” which roots in the observation that the curl-operator stores the information about the normal derivative of the tangential components, whereas the solenoidality condition furnishes an estimate for the normal derivatives of the normal component. We briefly protocol this procedure below. By Subsection 5.2, we can write

$$\operatorname{curl} u = J(\nu)(\partial_\nu u)^\tau + J(\tau^1)\partial_{\tau^1} u + J(\tau^2)\partial_{\tau^2} u.$$

Using the invertibility of $J(\nu)|_{\{\tau^1, \tau^2\}}$, we get

$$\begin{aligned} \partial_\nu u^\tau &= \sum_{i=1}^2 (\partial_\nu \tau^i u_{\tau^i} + \tau^i \partial_\nu \tau^i \cdot u) + R(\nu) \left(\operatorname{curl} u - \sum_{i=1}^2 J(\tau^i) \partial_{\tau^i} u \right) \\ &= R(\nu) \left(\operatorname{curl} u - \sum_{i=1}^2 J(\tau^i) \partial_{\tau^i} u \right) + \text{l.o.t.}(u), \end{aligned} \tag{5.1}$$

where the lower-order terms are a zeroth-order operator determined by Ω .

Proceeding to the divergence, for a smooth matrix-valued α , we obtain

$$\begin{aligned} \operatorname{div}(\alpha u) &= \sum_{\xi, \zeta} \partial_\xi (\alpha_{\xi\zeta} u_\zeta) + \operatorname{div}(\xi) \xi^\top \alpha u \\ &= \sum_{\xi, \zeta} (a_{\xi\zeta} \partial_\xi u_\zeta + \partial_\xi \alpha_{\xi\zeta} u_\zeta) + \sum_{\xi} \operatorname{div}(\xi) \xi^\top \alpha u \end{aligned}$$

with $\alpha_{\xi\zeta} = \xi^\top \alpha \zeta$. Solving this equation for $\partial_\nu u_\nu$ yields

$$\alpha_{\nu\nu} \partial_\nu u_\nu = \operatorname{div}(\alpha u) - \sum_{(\xi, \zeta) \neq (\nu, \nu)} \alpha_{\xi\zeta} \partial_\xi u_\zeta - \sum_{\xi, \zeta} \partial_\xi a_{\xi\zeta} u_\zeta - \sum_{\xi} \operatorname{div}(\xi) \xi^\top \alpha u. \tag{5.2}$$

If $\alpha = \alpha^\top \geq \eta I$, we have $\alpha_{\nu\nu} \geq \eta$ and thus the normal derivative $\partial_\nu u_\nu$ of the normal component of u is represented through $\operatorname{div}(\alpha u)$, the normal derivatives of the tangential components of u , the tangential derivatives of u , and lower-order terms depending on Ω and α along with its derivatives.

In contrast to [30], where the solenoidality condition was violated for the electric field due to the presence of an internally distributed electrical resistance term, both fields in (1.1) satisfy appropriate solenoidality conditions and can be handled via Equation (5.2).

6. REGULARITY BOOST: PROOF OF PROPOSITION 4.1

Proof. We first outline the argument. In Step 1), the desired estimates of $\partial_t^k E$ and $\partial_t^k H$ easily follow from our “elliptic” curl-div-estimate in Lemma 5.1 and the system (2.16). Similarly, one can also bound the higher-order terms away from the boundary in the second step. Near the boundary, one has to distinguish between the tangential and the normal regularity. The former can again be treated by means of Lemma 5.1 since tangential derivatives preserve the boundary condition up to lower-order terms. The normal regularity has to be recovered from the Maxwell equations and the divergence conditions as explained in the previous section. Iteratively, this is repeated in Step 4) for (E, H) in \mathcal{H}^2 , in Step 5) for $\partial_t(E, H)$ in \mathcal{H}^2 and in step 6) for (E, H) in \mathcal{H}^3 .

1) \mathcal{H}^1 -estimates for $\partial_t^k E$ and $\partial_t^k H$. Let $k \in \{0, 1, 2\}$. We first control the L^2 -norms of the curl and divergence of these fields via the time-differentiated Maxwell system (2.16) and the solenoidality conditions in (2.12) and (2.18). Also invoking the *a priori* estimates (2.22), we arrive at

$$\begin{aligned} \|\operatorname{curl} \partial_t^k E(t)\|_{L^2(\Omega)} &\leq ce_{k+1}^{1/2}(t) + cz(t)\delta_{k=2}, & \|\operatorname{curl} \partial_t^k H(t)\|_{L^2(\Omega)} &\leq ce_{k+1}^{1/2}(t) + cz(t)\delta_{k=2}, \\ \|\operatorname{div}(\widehat{\varepsilon}_k(t)\partial_t^k E(t))\|_{L^2(\Omega)} &\leq cz(t)\delta_{k=2}, & \|\operatorname{div}(\widehat{\mu}_k(t)\partial_t^k H(t))\|_{L^2(\Omega)} &\leq cz(t)\delta_{k=2} \end{aligned}$$

with the Kronecker delta $\delta_{k=2} = 1$ for $k = 2$ and $\delta_{k=2} = 0$ for $k \in \{0, 1\}$. Lemma 5.1 thus yields

$$\|\partial_t^k E(t)\|_{\mathcal{H}^1(\Omega)}^2 \leq ce_{k+1}(t) + cz^2(t)\delta_{k=2}, \quad \|\partial_t^k H(t)\|_{\mathcal{H}^1(\Omega)}^2 \leq ce_{k+1}(t) + cz^2(t)\delta_{k=2}. \quad (6.1)$$

(Later on we often omit the argument t .) Note that the right-hand sides $\partial_t f_k$ and $\partial_t g_k$ in Equation (2.16) are (locally) quadratic with respect to (E, H) and can thus be dominated by z according to Equation (2.22). This results in a superlinear remainder term of order 2 in z and will prove to be crucial for our reasoning.

2) *Interior spatio-temporal estimates for E and H .* Let $k \in \{0, 1\}$. For a fixed sufficiently small $a > 0$, we have the boundary “collar” $\Gamma_a = \{x \in \overline{\Omega} \mid \operatorname{dist}(x, \Gamma) < a\}$ from Subsection 5.2. Consider a cut-off function $\chi \in C_c^\infty(\overline{\Omega}, [0, 1])$ with $\operatorname{supp}(\chi) \subset \Gamma_a$ and $\chi = 1$ on $\Gamma_{a/2}$. Then $1 - \chi$ is smooth with $\operatorname{supp}(1 - \chi) \subset \Omega \setminus \Gamma_{a/2}$ and, hence, vanishes near the boundary Γ . Moreover, the maps $(1 - \chi)E$ and $(1 - \chi)H$ are as regular as E and H and satisfy the Maxwell system (2.16) with the new right-hand sides

$$\widehat{f}_k = \nabla \chi \times \partial_t^k H, \quad \widehat{g}_k = -\nabla \chi \times \partial_t^k E \quad (6.2)$$

in place of $-\partial_t f_k = 0$ respectively $-\partial_t g_k = 0$.

Take a multi-index $\alpha \in \mathbb{N}_0^3$ with $1 \leq |\alpha| \leq 2 - k$. Using Equation (2.16), we calculate

$$\operatorname{curl}(\partial_t^k \partial_x^\alpha((1 - \chi)H)) = \partial_t \partial_x^\alpha(\widehat{\varepsilon}_k(1 - \chi)\partial_t^k E) - \partial_x^\alpha(\nabla \chi \times \partial_t^k H).$$

Employing also the estimates of $\widehat{\varepsilon}_k$ in (2.22), we derive

$$\|\operatorname{curl}(\partial_t^k \partial_x^\alpha((1 - \chi)H))\|_{L^2(\Omega)} \leq c \max_{0 \leq j \leq 3 - |\alpha|} \|\partial_t^j E\|_{\mathcal{H}^{|\alpha|}(\Omega)} + c \|\partial_t^k H\|_{\mathcal{H}^{|\alpha|}(\Omega)}.$$

(One also invokes the Sobolev embeddings $\mathcal{H}^1(\Omega) \hookrightarrow L^6(\Omega)$ and $\mathcal{H}^2(\Omega) \hookrightarrow L^\infty(\Omega)$ to bound $\partial_t^j E$.) Note that $\partial_t^k \partial_x^\alpha((1 - \chi)E)$ and $\partial_t^k \partial_x^\alpha((1 - \chi)H)$ trivially satisfy the boundary condition in (2.16). We recall the divergence properties

$$\operatorname{div}(\widehat{\mu}_0 H) = 0 \quad \text{and} \quad \operatorname{div}(\widehat{\mu}_1 \partial_t H) = 0,$$

from Equations (2.12) and (2.18). Applying the product rule, it follows

$$\begin{aligned} \operatorname{div}(\widehat{\mu}_k \partial_t^k \partial_x^\alpha((1-\chi)H)) &= \sum_{0 \leq \beta \leq \alpha} \binom{\alpha}{\beta} \partial_x^{\alpha-\beta} ((\nabla \chi)^\top \widehat{\mu}_k) \partial_t^k \partial_x^\beta H \\ &\quad - \sum_{0 \leq \beta < \alpha} \binom{\alpha}{\beta} \operatorname{div}(\partial_x^{\alpha-\beta} \widehat{\mu}_k \partial_t^k \partial_x^\beta((1-\chi)H)). \end{aligned} \quad (6.3)$$

The inequalities in (2.22) yield

$$\begin{aligned} \|\operatorname{div}(\widehat{\mu}_k \partial_t^k \partial_x^\alpha((1-\chi)H))\|_{L^2(\Omega)} &\leq c \|\partial_t^k H\|_{\mathcal{H}^{|\alpha|}(\Omega)}, \\ \|\operatorname{div}(\widehat{\varepsilon}_k \partial_t^k \partial_x^\alpha((1-\chi)E))\|_{L^2(\Omega)} &\leq c \|\partial_t^k E\|_{\mathcal{H}^{|\alpha|}(\Omega)}, \end{aligned} \quad (6.4)$$

where the latter estimate is shown analogously.

First, let $|\alpha| = 1$. Lemma 5.1 and inequality (6.1) then imply the estimates

$$\|\partial_t^k((1-\chi)E(t))\|_{\mathcal{H}^2(\Omega)}^2, \|\partial_t^k((1-\chi)H(t))\|_{\mathcal{H}^2(\Omega)}^2 \leq ce(t) + cz^2(t). \quad (6.5)$$

For $|\alpha| = 2$, it similarly follows

$$\|(1-\chi)(E(t), H(t))\|_{\mathcal{H}^3(\Omega)}^2 \leq c \max_{j \in \{0,1\}} \{\|\partial_t^j E(t)\|_{\mathcal{H}^2(\Omega)}^2, \|\partial_t^j H(t)\|_{\mathcal{H}^2(\Omega)}^2\}. \quad (6.6)$$

3) *Preparations for boundary collar estimates for E and H .* Let $k \in \{0, 1\}$ and $\tau \in \{\tau^1, \tau^2\}$. We next treat χE and χH . Similar as $((1-\chi)E, (1-\chi)H)$, the fields χE and χH solve the Maxwell equations (2.16) with

$$\check{f}_k = -\nabla \chi \times \partial_t^k H, \quad \check{g}_k = \nabla \chi \times \partial_t^k E$$

in place of $-\partial_t f_k = 0$ respectively $-\partial_t g_k = 0$. An analogue of divergence equation (6.3) remains true. While the boundary condition is preserved (since $\chi = 1$ near Γ), the latter is only invariant under the tangential derivatives $\partial_\tau = \tau \cdot \nabla$ and not for the normal one. Taking the tangential derivative of the boundary condition in Equation (2.16), we can write

$$\begin{aligned} \partial_\tau \partial_t^k H \times \nu + (\lambda(\partial_\tau \partial_t^k E \times \nu)) \times \nu &= -\partial_t^k H \times \partial_\tau \nu - (\partial_\tau \lambda(\partial_t^k E \times \nu)) \times \nu \\ &\quad - (\lambda(\partial_t^k E \times \partial_\tau \nu)) \times \nu - (\lambda(\partial_t^k E \times \nu)) \times \partial_\tau \nu \\ &=: h_k. \end{aligned} \quad (6.7)$$

4) \mathcal{H}^2 -estimate for E and H . An application of the tangential derivative to the first two equations in (1.1) produces

$$\begin{aligned} \partial_t(\widehat{\varepsilon}_0 \partial_\tau(\chi E)) &= \operatorname{curl} \partial_\tau(\chi H) + [\partial_\tau, \operatorname{curl}](\chi H) - \partial_\tau(\nabla \chi \times H) - \partial_t(\partial_\tau \widehat{\varepsilon}_0(\chi E)), \\ \partial_t(\widehat{\mu}_0 \partial_\tau(\chi H)) &= -\operatorname{curl} \partial_\tau(\chi H) - [\partial_\tau, \operatorname{curl}](\chi E) + \partial_\tau(\nabla \chi \times E) - \partial_t(\partial_\tau \widehat{\mu}_0(\chi H)) \end{aligned} \quad (6.8)$$

(cf. Equation (6.2)), where $[A, B] = AB - BA$ stands for the commutator. Thus, in view of Equation (6.1), we can estimate

$$\|\operatorname{curl} \partial_\tau(\chi H(t))\|_{L^2(\Omega)}^2 \leq c(\|H(t)\|_{\mathcal{H}^1(\Omega)}^2 + \|E(t)\|_{\mathcal{H}^1(\Omega)}^2 + \|\partial_t E(t)\|_{\mathcal{H}^1(\Omega)}^2) \leq ce(t).$$

Similarly, observing

$$\operatorname{div}(\widehat{\mu}_0 \partial_\tau(\chi H)) = -[\partial_\tau, \operatorname{div}](\widehat{\mu}_0 \chi H) + \partial_\tau(\nabla \chi \cdot \widehat{\mu}_0 H) - \operatorname{div}(\partial_\tau \widehat{\mu}_0(\chi H)), \quad (6.9)$$

we can bound the divergence by

$$\|\operatorname{div}(\widehat{\mu}_0(t) \partial_\tau(\chi H(t)))\|_{L^2(\Omega)}^2 \leq c \|H(t)\|_{\mathcal{H}^1(\Omega)}^2 \leq ce(t). \quad (6.10)$$

Equations (6.7) with $k = 0$ and (6.1) yield

$$\|\partial_\tau(\chi H(t)) \times \nu + \lambda(\partial_\tau(\chi E(t)) \times \nu) \times \nu\|_{\mathcal{H}^{1/2}(\Gamma)}^2 \leq c(\|H(t)\|_{\mathcal{H}^1(\Omega)}^2 + \|E(t)\|_{\mathcal{H}^1(\Omega)}^2) \leq ce(t).$$

Repeating the procedure for E and invoking Lemma 5.1, we arrive at

$$\|\partial_\tau(\chi E(t))\|_{\mathcal{H}^1(\Omega)}^2, \|\partial_\tau(\chi H(t))\|_{\mathcal{H}^1(\Omega)}^2 \leq ce(t). \quad (6.11)$$

Turning to the normal derivatives, we apply ∂_j to the first equation in (1.1) and obtain

$$\partial_t(\widehat{\varepsilon}_0 \partial_j(\chi E)) = \operatorname{curl}(\partial_j(\chi H)) - \partial_j(\nabla \chi \times H) - \partial_t(\partial_j \widehat{\varepsilon}_0(\chi E))$$

for $j \in \{1, 2, 3\}$. Using (5.1) with $u = \partial_j(\chi H)$ as well as inequalities (6.11) and (6.1), we compute

$$\|\partial_\nu(\partial_j(\chi H(t)))_\tau\|_{L^2}^2 \leq c(\|\partial_\tau \partial_j(\chi H(t))\|_{L^2}^2 + \|H(t)\|_{\mathcal{H}^1}^2 + \|E(t)\|_{\mathcal{H}^1}^2 + \|\partial_t E(t)\|_{\mathcal{H}^1}^2) \leq ce(t). \quad (6.12)$$

An analogous estimate follows for E . As in (6.9), we can write

$$\operatorname{div}(\widehat{\mu}_0 \partial_j(\chi H)) = \partial_j(\nabla \chi \cdot \widehat{\mu}_0 H) - \operatorname{div}(\partial_j \chi \widehat{\mu}_0 H).$$

Therefore, employing (5.2), (6.1) and (6.12), we conclude

$$\|\partial_\nu(\partial_j(\chi H(t)))_\nu\|_{L^2(\Omega)}^2 \leq c(\|\partial_\nu(\partial_j(\chi H(t)))_\tau\|_{L^2(\Omega)}^2 + \|H(t)\|_{\mathcal{H}^1(\Omega)}^2) \leq ce(t). \quad (6.13)$$

Repeating this argumentation for H , we establish

$$\|\chi E(t)\|_{\mathcal{H}^2(\Omega)}^2, \|\chi H(t)\|_{\mathcal{H}^2(\Omega)}^2 \leq ce(t).$$

This inequality and (6.5) with $k = 0$ imply the \mathcal{H}^2 -estimate

$$\|E(t)\|_{\mathcal{H}^2(\Omega)}^2, \|H(t)\|_{\mathcal{H}^2(\Omega)}^2 \leq ce(t) + cz^2(t). \quad (6.14)$$

5) \mathcal{H}^2 -estimate for $\partial_t E$ and $\partial_t H$. We apply the ∂_τ -operator to (2.16) and derive

$$\begin{aligned} \partial_t(\widehat{\varepsilon}_1 \partial_\tau \partial_t(\chi E)) &= \operatorname{curl}(\partial_\tau \partial_t(\chi H)) + [\partial_\tau, \operatorname{curl}](\partial_t(\chi H)) \\ &\quad - \partial_\tau(\nabla \chi \times \partial_t H) - \partial_t(\partial_\tau \widehat{\varepsilon}_1 \partial_t(\chi E)), \\ \partial_t(\widehat{\mu}_1 \partial_\tau \partial_t(\chi H)) &= -\operatorname{curl}(\partial_\tau \partial_t(\chi E)) - [\partial_\tau, \operatorname{curl}](\partial_t(\chi E)) \\ &\quad + \partial_\tau(\nabla \chi \times \partial_t E) - \partial_t(\partial_\tau \widehat{\mu}_1 \partial_t(\chi H)). \end{aligned} \quad (6.15)$$

Recycling the \mathcal{H}^1 -estimate (6.1), it follows

$$\|\operatorname{curl} \partial_\tau \partial_t(\chi E)\|_{L^2(\Omega)} \leq c(\|\partial_t^2 H\|_{\mathcal{H}^1(\Omega)} + \|\partial_t H\|_{\mathcal{H}^1(\Omega)} + \|\partial_t E\|_{\mathcal{H}^1(\Omega)}) \leq ce^{1/2}(t) + cz(t).$$

We take the tangential derivative of the identity

$$\operatorname{div}(\widehat{\varepsilon}_1 \partial_t(\chi E)) = -\nabla \chi \widehat{\varepsilon}_1 \partial_t E$$

and conclude

$$\operatorname{div}(\widehat{\varepsilon}_1 \partial_\tau \partial_t(\chi E)) = -[\partial_\tau, \operatorname{div}](\widehat{\varepsilon}_1 \partial_t(\chi E)) - \operatorname{div}(\partial_\tau \widehat{\varepsilon}_1 \partial_t(\chi E)) - \partial_\tau(\nabla \chi \widehat{\varepsilon}_1 \partial_t E). \quad (6.16)$$

In view of (6.1), this identity furnishes the inequality

$$\|\operatorname{div}(\widehat{\varepsilon}_1(t) \partial_\tau \partial_t(\chi E(t)))\|_{L^2(\Omega)} \leq c\|\partial_t E(t)\|_{\mathcal{H}^1(\Omega)} \leq ce^{1/2}(t).$$

Observe that $\partial_\tau \partial_t(\chi E)$ satisfies the boundary condition (6.7) with

$$\|h_1(t)\|_{H^{1/2}(\Omega)} \leq c(\|\partial_t E(t)\|_{H^1(\Omega)} + \|\partial_t H(t)\|_{H^1(\Omega)}) \leq ce^{1/2}(t),$$

where we again employed the \mathcal{H}^1 -estimate (6.1). Lemma 5.1 thus implies the tangential bound

$$\|\partial_\tau \partial_t(\chi E(t))\|_{\mathcal{H}^1(\Omega)}^2, \|\partial_\tau \partial_t(\chi H(t))\|_{\mathcal{H}^1(\Omega)}^2 \leq ce(t). \quad (6.17)$$

We next look at the normal derivative. An application of the ∂_j -operator to (2.16) yields

$$\partial_t(\widehat{\varepsilon}_1 \partial_j \partial_t(\chi E)) = \operatorname{curl} \partial_j \partial_t(\chi H) - \partial_j(\nabla \chi \times \partial_t H) - \partial_t(\partial_j \widehat{\varepsilon}_1 \partial_t(\chi E)),$$

cf. (6.15). By means of (5.1) with $u = \partial_j \partial_t(\chi H)$, (6.1) and (6.17), we deduce

$$\|\partial_\nu(\partial_j \partial_t(\chi H))_\tau\|_{L^2(\Omega)}^2 \leq c(\|\partial_t H\|_{\mathcal{H}^1(\Omega)}^2 + \|\partial_\tau \partial_t(\chi H)\|_{\mathcal{H}^1(\Omega)}^2 + \|\partial_t(\chi E)\|_{\mathcal{H}^1(\Omega)}^2 + \|\partial_t^2(\chi E)\|_{\mathcal{H}^1(\Omega)}^2)$$

$$\leq ce(t) + cz^2(t). \quad (6.18)$$

On the other hand, (2.18) yields

$$\operatorname{div}(\widehat{\mu}_1 \partial_j \partial_t(\chi H)) = \partial_j(\nabla \chi \cdot \widehat{\mu}_1 \partial_t H) - \operatorname{div}(\partial_j \widehat{\mu}_1 \partial_t(\chi H)).$$

Exploiting (5.2), (6.1), (6.17) and (6.18), we get the inequality

$$\begin{aligned} \|\partial_\nu(\partial_j \partial_t(\chi H(t)))_\nu\|_{L^2(\Omega)}^2 &\leq c(\|\partial_\tau \partial_t(\chi H(t))\|_{\mathcal{H}^1(\Omega)}^2 + \|\partial_\nu(\partial_j \partial_t(\chi H(t)))_\tau\|_{L^2(\Omega)}^2 + \|\partial_t H(t)\|_{\mathcal{H}^1(\Omega)}^2) \\ &\leq ce(t) + cz^2(t). \end{aligned} \quad (6.19)$$

A similar estimate follows in an analogous fashion for E . The bounds (6.17), (6.18) and (6.19) yield the collar estimate

$$\|\chi \partial_t E(t)\|_{\mathcal{H}^2(\Omega)}^2, \|\chi \partial_t H(t)\|_{\mathcal{H}^2(\Omega)}^2 \leq ce(t) + cz^2(t).$$

Combining this with (6.5), we arrive at the \mathcal{H}^2 -inequality for $\partial_t(E, H)$

$$\|\partial_t E(t)\|_{\mathcal{H}^2(\Omega)}^2, \|\partial_t H(t)\|_{\mathcal{H}^2(\Omega)}^2 \leq ce(t) + cz^2(t). \quad (6.20)$$

Hence, (6.6) leads to the interior bound

$$\|(1 - \chi)(E(t), H(t))\|_{\mathcal{H}^3(\Omega)}^2 \leq ce(t) + cz^2(t). \quad (6.21)$$

6) \mathcal{H}^3 -estimate for E and H . As in (6.8), letting $\partial_\tau^2 := \partial_{\tau_i} \partial_{\tau_j}$ for $i, j \in \{1, 2, 3\}$, we can write

$$\partial_t(\widehat{\varepsilon}_0 \partial_\tau^2(\chi E)) = \operatorname{curl} \partial_\tau^2(\chi H) + [\partial_\tau^2, \operatorname{curl}](\chi H) - \partial_\tau^2(\nabla \chi \times H) - 2\partial_t(\partial_\tau \widehat{\varepsilon}_0 \partial_\tau(\chi E)) - \partial_t(\partial_\tau^2 \widehat{\varepsilon}_0 \chi E).$$

Using the inequalities (6.14) and (6.20), we derive

$$\|\operatorname{curl} \partial_\tau^2(\chi H(t))\|_{L^2(\Omega)}^2 \leq ce(t) + cz^2(t).$$

As before, a similar estimate can be shown for E . In the same fashion as in (6.9), we compute

$$\operatorname{div}(\widehat{\mu}_0 \partial_\tau^2(\chi H)) = -[\partial_\tau^2, \operatorname{div}](\widehat{\mu}_0 \chi H) + \partial_\tau^2(\nabla \chi \cdot \widehat{\mu}_0 H) - 2\operatorname{div}(\partial_\tau \widehat{\mu}_0 \partial_\tau(\chi H)) - \operatorname{div}(\partial_\tau^2 \widehat{\mu}_0(\chi H)).$$

Thus, on the strength of (6.14), we get

$$\|\operatorname{div}(\widehat{\mu}_0(t) \partial_\tau^2(\chi H(t)))\|_{L^2(\Omega)}^2 \leq ce(t) + cz^2(t).$$

A modification of (6.7) further yields

$$\|\partial_\tau^2(\chi H) \times \nu + (\lambda(\partial_\tau^2(\chi E) \times \nu)) \times \nu\|_{\mathcal{H}^{1/2}(\Gamma)}^2 \leq c(\|E(t)\|_{\mathcal{H}^2(\Omega)}^2 + \|H(t)\|_{\mathcal{H}^2(\Omega)}^2) \leq ce(t) + cz^2(t).$$

In summary, Lemma 5.1 implies the tangential bound

$$\|\partial_\tau^2(\chi H(t))\|_{\mathcal{H}^1(\Omega)}^2, \|\partial_\tau^2(\chi E(t))\|_{\mathcal{H}^1(\Omega)}^2 \leq ce(t) + cz^2(t). \quad (6.22)$$

We next bound the normal derivative. In spirit of (6.8), we obtain

$$\begin{aligned} \partial_t(\widehat{\varepsilon}_0 \partial_\tau \partial_j(\chi E)) &= \operatorname{curl}(\partial_\tau \partial_j(\chi H)) + [\partial_\tau, \operatorname{curl}]\partial_j(\chi H) - \partial_\tau \partial_j(\nabla \chi \times H) \\ &\quad - \partial_t(\partial_\tau \widehat{\varepsilon}_0 \partial_j(\chi E)) - \partial_t(\partial_j \widehat{\varepsilon}_0 \partial_\tau(\chi E)) - \partial_t(\partial_\tau \partial_j \widehat{\varepsilon}_0(\chi E)) \end{aligned}$$

for $j \in \{1, 2, 3\}$. A joint application of (5.1), (6.14), (6.20) and (6.22) furnishes the estimate

$$\|\partial_\nu(\partial_\tau \partial_j H(t))_\tau\|_{L^2(\Omega)}^2 \leq ce(t) + cz^2(t). \quad (6.23)$$

We can treat E analogously. As in (6.9), we compute

$$\begin{aligned} \operatorname{div}(\widehat{\mu}_0 \partial_\tau \partial_j(\chi H)) &= -[\partial_\tau, \operatorname{div}](\widehat{\mu}_0 \partial_j(\chi H) + \partial_j \widehat{\mu}_0(\chi H)) - \operatorname{div}(\partial_\tau \widehat{\mu}_0 \partial_j(\chi H)) \\ &\quad - \operatorname{div}(\partial_j \widehat{\mu}_0 \partial_\tau(\chi H)) - \operatorname{div}(\partial_\tau \partial_j \widehat{\mu}_0(\chi H)) + \partial_\tau \partial_j(\nabla \chi \cdot \widehat{\mu}_0 H). \end{aligned}$$

Using formula (5.2) and estimates (2.22), (6.14), (6.22) and (6.23), we estimate

$$\|\partial_\nu(\partial_\tau \partial_j(\chi H(t)))_\nu\|_{L^2(\Omega)}^2 \leq ce(t) + cz^2(t). \quad (6.24)$$

The electric field E is controlled similarly. Combining (6.23) and (6.24), we arrive at

$$\|\partial_\nu \partial_\tau \partial_j(\chi E(t))\|_{L^2(\Omega)}^2 + \|\partial_\nu \partial_\tau \partial_j(\chi H(t))\|_{L^2(\Omega)}^2 \leq ce(t) + cz^2(t). \quad (6.25)$$

Replacing ∂_τ by ∂_ν in the calculations above, we treat $\partial_\nu^2 \partial_j(\chi E)$ and $\partial_\nu^2 \partial_j(\chi H)$ in the same way. Together with (6.22) and (6.25), this leads to the interior bound

$$\|\chi E(t)\|_{\mathcal{H}^3(\Omega)}^2 + \|\chi H(t)\|_{\mathcal{H}^3(\Omega)}^2 \leq ce(t) + z^2(t).$$

This inequality and (6.21) imply the final \mathcal{H}^3 -estimate

$$\|E(t)\|_{\mathcal{H}^3(\Omega)}^2 + \|H(t)\|_{\mathcal{H}^3(\Omega)}^2 \leq ce(t) + z^2(t). \quad (6.26)$$

The claim now follows from inequalities (6.1), (6.20), and (6.26). \square

REFERENCES

- [1] S. Alinhac. *Blowup for Nonlinear Hyperbolic Equations*. Birkhäuser, Basel, 1995.
- [2] S. Alinhac. Blowup of small data solutions for a class of quasilinear wave equations in two space dimensions II. *Acta Mat.*, 182:1–23, 1999.
- [3] S. Alinhac. Blowup of small data solutions for a quasilinear wave equation in two space dimensions. *Ann. Math.*, 149:97–127, 1999.
- [4] S. Alinhac. Remarks on the blowup rate of classical solutions to quasilinear multidimensional hyperbolic system. *J. Math. Pure Appl.*, 79:839–854, 2000.
- [5] H. Ammari and J.-C. Nédélec. Generalized impedance boundary conditions for the Maxwell equations as singular perturbations problems. *Comm. Partial Differential Equations*, 24:821–849, 1999.
- [6] A. Anikushyn and M. Pokojovy. Global well-posedness and exponential stability for heterogeneous anisotropic Maxwell’s equations under a nonlinear boundary feedback with delay. <https://arxiv.org/abs/1807.00098>, 2018.
- [7] C. Bardos, A. Y. Le Roux, and J. C. Nédélec. First order quasilinear equations with boundary conditions. *Comm. Partial Differential Equations*, 4:1017–1034, 1979.
- [8] J. Cagnol and M. Eller. Boundary regularity for Maxwell’s equations with applications to shape optimization. *J. Differential Equations*, 250:1114–1136, 2011.
- [9] F. Cakoni, D. Colton, and P. Monk. *The Linear Sampling Method in Inverse Electromagnetic Scattering*. SIAM, Philadelphia, PA, 2011.
- [10] M. Cessenat. *Mathematical Methods in Electromagnetism. Linear Theory and Applications*. World Scientific Publishing, River Edge (NJ), 1996.
- [11] I. Chueshov and I. Lasiecka. Generation of a semigroup and hidden regularity in nonlinear subsonic flow-structure interactions with absorbing boundary conditions. *J. Abstr. Differ. Equ. Appl.*, 3:1–27, 2012.
- [12] M. Costabel, M. Dauge, and S. Nicaise. Corner Singularities and Analytic Regularity for Linear Elliptic Systems. Part I. <http://hal.archives-ouvertes.fr/hal-00453934/en/>, 2010.
- [13] P. D’Ancona, S. Nicaise, and R. Schnaubelt. Blow-up for nonlinear Maxwell equations. *Electron. J. Differential Equations*, Paper No. 73, 2018.
- [14] R. Dautray and J.-L. Lions. *Spectral Theory and Applications*. Springer-Verlag, Berlin, 2000.
- [15] M. Eller. Stability of the anisotropic Maxwell equations with a conductivity term. *Evol. Equ. Control Theory*. To appear.
- [16] M. Eller. Continuous observability for the anisotropic Maxwell system. *Appl. Math. Optim.*, 55:185–201, 2007.
- [17] M. Eller, J. Lagnese, and S. Nicaise. Decay rates for solutions of a Maxwell system with nonlinear boundary damping. *Comput. Appl. Math.*, 21(1):135–165, 2002.
- [18] M. Eller, J.E. Lagnese, and S. Nicaise. Stabilization of heterogeneous Maxwell’s equations by linear and nonlinear boundary feedbacks. *Electronic J. Differential Equations*, 21:1–26, 2002.
- [19] M. Eller and D. Toundykov. Carleman estimates for elliptic boundary value problems with applications to the stabilization of hyperbolic systems. *Evol. Equ. Control Theory*, 1(2):271–296, 2012.
- [20] O. Guès. Problème mixte hyperbolique quasi-linéaire caractéristique. *Comm. Partial Differential Equations*, 15(5):595–645, 1990.
- [21] S. Jiang and R. Racke. *Evolution Equations in Thermoelasticity*. Chapman & Hall/CRC, Boca Raton, 2000.
- [22] F. John. Blowup for quasilinear wave equations in three space dimensions. *Comm. Pure. Appl. Math.*, 34:29–51, 1981.
- [23] T. Kato. The Cauchy problem for quasi-linear symmetric hyperbolic systems. *Arch. Ration. Mech. Anal.*, 58(3):181–205, 1975.

- [24] S. Kawashima, T. Yanagisawa, and Y. Shizuta. Mixed problems for quasi-linear symmetric hyperbolic systems. *Proc. Japan Acad. Ser. A Math. Sci.*, 63(7):243–246, 1987.
- [25] V. Komornik. Boundary stabilization, observation and control of Maxwell’s equations. *Pan-American Math. J.*, 4:47–61, 1994.
- [26] I. Kukavica, A. L. Mazzucato, and A. Tuffaha. Sharp trace regularity for an anisotropic elasticity system. *Proc. Amer. Math. Soc.*, 141(8):2673–2682, 2013.
- [27] J. Lagnese. Exact controllability of Maxwell’s equations in a general region. *SIAM J. Control Optim.*, 27(2):374–388, 1989.
- [28] I. Lasiecka, J.-L. Lions, and R. Triggiani. Non-homogeneous boundary value problems for second order hyperbolic operators. *J. Math. Pures Appl.*, 65:149–192, 1986.
- [29] I. Lasiecka, R. Marchand, and T. McDevitt. Boundary control and hidden trace regularity of a semigroup associated with a beam equation and non-dissipative boundary conditions. *Dynam. Systems Appl.*, 21:467–490, 2012.
- [30] I. Lasiecka, M. Pokojovy, and R. Schnaubelt. Exponential decay of quasilinear Maxwell equations with interior conductivity. <https://arxiv.org/abs/1810.12390>, 2018.
- [31] I. Lasiecka, M. Pokojovy, and X. Wan. Global existence and exponential stability for a nonlinear thermoelastic Kirchhoff-Love plate. *Nonlinear Anal. Real World Appl.*, 38:184–221, 2017.
- [32] I. Lasiecka, M. Pokojovy, and X. Wan. Long-time behavior of quasilinear thermoelastic Kirchhoff-Love plates with second sound. <https://arxiv.org/abs/1811.01138>, 2018.
- [33] I. Lasiecka and D. Toundykov. Semigroup generation and “hidden” trace regularity of a dynamic plate with non-monotone boundary feedbacks. *Commun. Math. Anal.*, 8(1):109–144, 2010.
- [34] I. Lasiecka and R. Triggiani. Sharp regularity theory for second order hyperbolic equations of Neumann type. Part I. L_2 nonhomogeneous data. *Ann. Mat. Pura Appl.*, IV:285–367, 1990.
- [35] I. Lasiecka and R. Triggiani. Regularity theory of hyperbolic equations with non-homogeneous Neumann boundary conditions. II. General boundary data. *J. Differential Equations*, 94(1):112–164, 1991.
- [36] F. Linares and G. Ponce. *Introduction to Nonlinear Dispersive Equations*. Springer-Verlag, New York, 2nd edition, 2015.
- [37] S. Lucente and G. Ziliotti. Global existence for a quasilinear Maxwell system. *Rend. Istit. Mat. Univ. Trieste*, 31(suppl. 2):169–187, 2000.
- [38] S. A. Messaoudi, M. Pokojovy, and B. Said-Houari. Nonlinear damped Timoshenko systems with second sound – global existence and exponential stability. *Math. Methods Appl. Sci.*, 32(5):505–534, 2009.
- [39] J. Moloney and A. Newell. *Nonlinear Optics*. Westview Press. Advanced Book Program, Boulder, CO, 2004.
- [40] J. E. Muñoz Rivera and R. Racke. Smoothing properties, decay, and global existence of solutions to nonlinear coupled systems of thermoelastic type. *SIAM J. Math. Anal.*, 26:1547–1563, 1995.
- [41] J. E. Muñoz Rivera and R. Racke. Large solutions and smoothing properties for nonlinear thermoelastic systems. *J. Differential Equations*, 127:454–483, 1996.
- [42] J. E. Muñoz Rivera and R. Racke. Mildly dissipative nonlinear Timoshenko systems – Global existence and exponential stability. *J. Math. Anal. Appl.*, 276:248–278, 2002.
- [43] S. Nicaise and C. Pignotti. Boundary stabilization of Maxwell’s equations with space-time variable coefficients. *ESAIM Control Optim. Calc. Var.*, 9:563–578, August 2003.
- [44] S. Nicaise and C. Pignotti. Internal stabilization of Maxwell’s equations in heterogeneous media. *Abstr. Appl. Anal.*, 2005(7):791–811, 2005.
- [45] S. Nicaise and C. Pignotti. Partially delayed stabilizing feedbacks for Maxwell’s system. *Adv. Differential Equations*, 12(1):27–54, 2007.
- [46] T. Ohkubo. Well posedness for quasi-linear hyperbolic mixed problems with characteristic boundary. *Hokkaido Math. J.*, 18(1):79–123, 1981.
- [47] K. D. Phung. Contrôle de stabilisation d’ondes électromagnétiques. *ESAIM Control Optim. Calc. Var.*, 5:87–137, 2000.
- [48] R. H. Picard and W. M. Zajaczkowski. Local existence of solutions of impedance initialboundary value problem for nonlinear maxwell equations. *Math. Methods Appl. Sci.*, 18(3):169–199, 1995.
- [49] R. Racke. *Lectures on Nonlinear Evolution Equations: Initial Value Problems*. Birkhäuser, Basel, 2015.
- [50] R. Racke and Y. Ueda. Nonlinear thermoelastic plate equations – Global existence and decay rates for the Cauchy problem. *J. Differential Equations*, 263:8138–8177, 2017.
- [51] J. Rauch. Symmetric positive systems with boundary characteristic of constant multiplicity. *Trans. Amer. Math. Soc.*, 291(1):167–187, 1985.
- [52] R. Schnaubelt and M. Spitz. Local wellposedness of quasilinear Maxwell equations with absorbing boundary conditions. <http://arxiv.org/abs/1812.03803>, 2018.
- [53] R. Schnaubelt and M. Spitz. Local wellposedness of quasilinear maxwell equations with conservative interface conditions. <http://arxiv.org/abs/1811.08714>, 2018.

- [54] S. Schochet. The compressible Euler equations in a bounded domain. Existence of solutions and the incompressible limit. *Comm. Math. Phys.*, 104:49–75, 1986.
- [55] P. Secchi. Well-posedness of characteristic symmetric hyperbolic systems. *Arch. Ration. Mech. Anal.*, 134(2):155–197, 1996.
- [56] J. Speck. The nonlinear stability of the trivial solution to the Maxwell-Born-Infeld system. *J. Math. Phys.*, 53(8):083703, 83, 2012.
- [57] J. Speck. *Shock Formation in Small-Data Solutions to 3D Quasilinear Wave Equations*. Amer. Math. Soc., Providence, RI, 2016.
- [58] M. Spitz. *Local wellposedness of nonlinear Maxwell equations*. PhD thesis, Karlsruhe Institute of Technology, Karlsruhe, 2017.
- [59] M. Spitz. Local wellposedness of nonlinear Maxwell equations with perfectly conducting boundary conditions. *J. Differential Equations*, 2018. Early version online.
- [60] M. Spitz. Regularity theory for nonautonomous Maxwell equations with perfectly conducting boundary conditions. <https://arxiv.org/abs/1805.00671>, 2018.

M. POKOJOVY, DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF TEXAS AT EL PASO, 500 W UNIVERSITY AVE, EL PASO, TX 79968, USA

E-mail address: `mpokojovy@utep.edu`

R. SCHNAUBELT, DEPARTMENT OF MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY, 76128 KARLSRUHE, GERMANY

E-mail address: `schnaubelt@kit.edu`