

PAPER • OPEN ACCESS

QED loop effects in the spacetime background of a Schwarzschild black hole

To cite this article: Viacheslav A. Emelyanov 2017 *J. Phys.: Conf. Ser.* **942** 012009

View the [article online](#) for updates and enhancements.

Related content

- [Fast Track Communication](#)
S A Hayward, R Di Criscienzo, M Nadalini et al.
- [Hawking radiation from decoherence](#)
Claus Kiefer
- [Quantum radiation of general nonstationary black holes](#)
Jia-Chen Hua and Yong-Chang Huang

QED loop effects in the spacetime background of a Schwarzschild black hole

Viacheslav A. Emelyanov

Institute of Theoretical Physics, Karlsruhe Institute of Technology, 76131 Karlsruhe, Germany

E-mail: viacheslav.emelyanov@kit.edu

Abstract. The black-hole evaporation implies that the quantum-field propagators in a local Minkowski frame acquire a correction, which gives rise to this process. The modification of the propagators causes, in turn, non-trivial local effects due to the radiative/loop diagrams in non-linear QFTs. In particular, there should be imprints of the evaporation in QED, if one goes beyond the tree-level approximation. Of special interest in this respect is the region near the black-hole horizon, which, already at tree level, appears to show highly non-classical features, e.g., negative energy density and energy flux into the black hole.

1. Introduction

After many years of theoretical and experimental studies, we have been successful in establishing a couple of models, which incredibly well describe physics at microscopic and macroscopic scales. The basic ideas behind of the Standard Model of Particle Physics and Cosmology, are different, but not inconsistent. It is the Einstein equivalence principle that allows us to go over from the general relativity regime to the particle physics one.

According to the large scale observations, the global geometrical structure of the Universe turns out to be roughly described by de Sitter spacetime. At smaller length scales, the Universe is no longer homogeneous and isotropic as a consequence of the presence of Dark Matter as well as baryonic matter in the form of clusters of galaxies, galaxies and so on. However, Minkowski spacetime appears to be a good approximation to the Universe locally, i.e. in the neighbourhood of any given space-time point.

The geometrical structure relevant for Particle Physics is given by the Minkowski metric of a local Minkowski frame. The Poincaré group $R^4 \times SO(1, 3)$ is, thus, a local isometry group of the Universe. This group is employed to define the notion of particles, which are associated with its unitary and irreducible representations, with a no-particle state being the local Minkowski vacuum. The Lehmann-Symanzik-Zimmermann reduction formula relates, in turn, physical particle states with poles of the quantum-field propagators. We know *a posteriori* that this formalism adequately describes various reactions observed in the particle colliders on Earth.

The notion of particles identified with the unitary, irreducible representations of the Poincaré group and, hence, labeled by values of its two Casimir generators, i.e. mass and spin, is the only one, which has been successfully tested so far. The Wigner particle is, thus, a localized object, that can be defined in a local Minkowski frame only. It seems that if the local curvature in a certain space-time region of the Universe becomes comparable with, e.g., the Compton length λ_c of the electron, then one should not expect that the notion of the electron is physically



The figure shows two Feynman diagrams. The top diagram represents the vacuum polarization tensor $i\Pi^{\mu\nu}(k)$. It consists of a loop of fermions (represented by a circle with arrows) with an incoming photon line (wavy line) on the left and an outgoing photon line (wavy line) on the right. The loop momentum is p , the incoming photon momentum is k , and the outgoing photon momentum is $k+p$. The bottom diagram represents the electron self-energy $-i\Sigma(p)$. It shows a fermion line (solid line with arrows) with an incoming momentum p and an outgoing momentum p . A loop of photons (wavy line) is attached to the fermion line, with loop momentum k and the fermion momentum in the loop being $p-k$.

Figure 1. One-loop diagrams providing the lowest order correction to the photon as well as electron self-energy. The object $\Pi^{\mu\nu}(k)$ is known as the vacuum polarization tensor. The vacuum polarization gives rise to the running of the electric charge and contributes to the Lamb shift.

meaningful there. Nevertheless, the Wigner electrons (i.e. electrons we observe in the particle colliders) might, still, make physical sense if relativistic, i.e. correspond to the de Broglie length $\lambda_e \ll \lambda_c$. Therefore, the Wigner electrons have to be highly relativistic in that region. Although the proof of this conjecture in a lab is impossible, we take it for granted below as a natural extrapolation of physics we have experimentally probed up to now.

The fact that we find ourselves in non-flat spacetime implies that one should bear in mind the curvature length scale l_c . Still, the Minkowski-space approximation employed in Particle Physics is adequate, whenever the length scale of a certain particle reaction is negligible with respect to l_c . Consequently, one does not need to integrate over all space-time points of the Universe in the coordinate representation of vertices in non-linear QFTs. Moreover, the same also appears to hold in the path integral formalism and, furthermore, one can use the powerful technique of the Fourier transform as if the Universe were globally Minkowski spacetime.

Thus, one can imagine that the local curvature at a given point of the Universe is described by the length scale l_c . According to the Einstein equivalence principle, one can introduce a local Minkowski frame in its vicinity. In this frame, we can follow standard methods used in Particle Physics to quantize, e.g., the Maxwell or Dirac field. There is no reason to expect that this procedure is physically meaningless, because otherwise it would be unnatural to assume that the formalism works on Earth only. Of course, one should pay attention to the fact that the Standard Model is not scale invariant. For instance, if $l_c \lesssim \lambda_c \propto 1/m_e$, one should then allow only electrons/positrons as external legs in the Feynman diagrams with momenta $|\vec{p}| \gg 1/l_c$. In this paper, we intend to employ these ideas near a black hole, so that l_c corresponds to the size of the black-hole event horizon, i.e. $l_c = r_H = 2M$, where M is a black-hole mass.

Throughout this paper we assume that $c = G = k_B = \hbar = 1$, unless otherwise stated.

2. Loop effects in Quantum Electrodynamics

A plenty of observable effects in QFT cannot be understood at tree level, i.e. without counting loop diagrams. In quantum electrodynamics, the most remarkable manifestations of the loops are the anomalous magnetic moment of the electron, the running of the electric charge and the Lamb shift. We shall deal in this paper with the one-loop correction to the photon and electron self-energy (see Fig. 1).

2.1. Isotropic electron-positron plasma

In the presence of particles and/or external fields, novel non-trivial manifestations of the loops appear. As an instructive example, we consider first an isotropic and neutral plasma composed of electrons, positrons and photons.

Maxwell- and Dirac-field propagators: To determine how the plasma particles influence, e.g., the photon kinematics, one needs to know the vector- and fermion-field propagators inside the plasma. One can show that

$$\underbrace{\frac{i(\not{p} + m)}{p^2 - m^2 + i\varepsilon}}_{\text{Vacuum}} \implies \underbrace{\frac{i(\not{p} + m)}{p^2 - m^2 + i\varepsilon} - 2\pi \frac{(\not{p} + m)}{e^{\beta|p_0|} + 1} \delta(p^2 - m^2)}_{\text{Isotropic plasma}}, \quad (1)$$

where $\beta \equiv 1/T$ is the inverse plasma temperature, and

$$\underbrace{\frac{-i\eta_{\mu\nu}}{k^2 + i\varepsilon}}_{\text{Vacuum}} \implies \underbrace{\frac{-i\eta_{\mu\nu}}{k^2 + i\varepsilon} - 2\pi \frac{\eta_{\mu\nu}}{e^{\beta|k_0|} - 1} \delta(k^2)}_{\text{Isotropic plasma}}. \quad (2)$$

Since the electron-positron field is massive, the extra term in its propagator is not suppressed by the Boltzmann factor $\exp(-\beta m_e)$ if we consider the case $T \gg m_e$. It is legitimate, then, to neglect the mass parameter m_e , so that the fermion field becomes effectively massless. This limit is known as the hard-thermal-loop approximation [13].

Modified dispersion relation: The one-loop effects originating from non-vanishing temperature are the Debye screening of a point-like charge, emergent propagating degrees of freedom (so-called plasmon and plasmino) as a result of the collective plasma-particles excitations and others (see, e.g., [13]). We focus herein on the photon and electron dispersion relations.

With the propagators given in Eqs. (1) and (2), we find in the on-mass-shell limit in the hot plasma ($T \gg m_e$) that

$$k_0^2 - \vec{k}^2 \approx \frac{2\pi}{3} \alpha T^2, \quad (3)$$

$$p_0^2 - \vec{p}^2 \approx m_e^2 + \pi \alpha T^2 \quad (4)$$

at one-loop level, where $\alpha \equiv e^2/4\pi$ is the fine-structure constant and m_e is a physical electron mass (whereas m is a bare electron mass). Thus, QED photons acquire a thermal gauge-invariant mass, which changes their kinematics [13].

2.2. Anisotropic electron-positron plasma

An anisotropic plasma is characterized by a unit vector \vec{n} , that gives the direction along which the plasma moves. This medium appears to be closely related to the quantum physics near to and far from the black-hole horizon [4, 5, 7, 8].

Maxwell- and Dirac-field propagators: If the anisotropic and neutral electron-positron plasma is held at temperature T , then one needs to implement in Eqs. (1) and (2) the following changes:

$$\delta(k^2) \implies 2\pi |\vec{k}| \delta(\vec{k} - k_0 \vec{n}), \quad (5)$$

$$\delta(p^2 - m^2) \implies 2\pi |\vec{p}| \delta(\vec{p} - (p_0^2 - m^2)^{\frac{1}{2}} \vec{n}), \quad (6)$$

in order to gain the photon- and electron-positron-field propagators in this medium.

Starting with quantum kinetic theory (see, e.g., [6]), it is straightforward to show that the anisotropic plasma is described by the following one-particle distribution:

$$f(x, p) = \frac{g_s}{2\pi^2} \frac{|\vec{p}|^2}{e^{\beta p_0} - (-1)^{2s}} \theta(\vec{p} \cdot \vec{n}) \delta(\vec{p} \times \vec{n}), \quad (7)$$

where p_0 is either $|\vec{p}|$ or $(|\vec{p}|^2 + m_s^2)^{\frac{1}{2}}$, depending on whether the quantum field of spin s is massless or massive and g_s is its number of degrees of freedom. For instance, this implies that the photon number current \vec{j}_γ is given by $n_\gamma \cdot \vec{n}$, where $n_\gamma = 2\zeta(3)T^3/\pi^2$ is the photon number density in the plasma. Therefore, the anisotropy of the plasma manifests itself in the macroscopic local observables.

Modified dispersion relation: It turns out that the modified dispersion relations given in Eqs. (3) and (4) are independent on whether the electron-positron plasma is isotropic or anisotropic. This holds only at one-loop approximation. The photon dispersion relation at two-loop order gets a dependence on the unit vector \vec{n} , implying that the two-loop diagrams do not contribute to the thermal photon mass [9].

2.3. Near-horizon region of small black holes

The near-horizon region of black holes is exotic from a classical point of view. Specifically, the renormalized stress tensor of the quantum fields describes an inward flux of the negative energy density, originating well outside of the black-hole horizon [16, 2, 12]. This explains, however, why the black-hole horizon shrinks, i.e. $2\dot{M} < 0$, where dot refers to the Schwarzschild time coordinate.

This non-classical flux, like the Hawking flux in the far-from-horizon region, is anisotropic. This phenomenon is described by the field propagators of the form given in Eqs. (1) and (2) with the modifications (5) and (6) implemented with an extra multiple factor roughly equaling to $-27/4$ and $\beta = 2/T_H$ [6, 7],¹ where T_H is the Hawking temperature. Thus, the following result holds:

$$k_0^2 - \vec{k}^2 \approx -\frac{9\pi}{8} \alpha T_H^2, \quad (8)$$

$$p_0^2 - \vec{p}^2 \approx m_e^2 - \frac{27\pi}{16} \alpha T_H^2, \quad (9)$$

for black holes with $T_H \gg m_e$ [8].

3. Concluding remarks

We have found above that $k^2 < 0$ and $p^2 < 0$ close to the event horizon. What is the physics of this result?

Einstein acausality vs. instability: First, it is useful to consider a classical field model with a “wrong” sign in the mass-squared term. In general, there are two logically non-excludable ways of quantizing this theory. One can either discard modes with imaginary frequencies or take those into account. In the former case, the Einstein causality is violated, whereas, in the latter one, it is not, but the price is the quantum-vacuum instability [3, 15, 1]. The very reason of the instability are frequency modes with $|\vec{q}| < \mu$, where $\mu > 0$ is the field “mass”. In other words, the modes with $q_0^2 = \vec{q}^2 - \mu^2 < 0$ are exponentially increasing in time. This model, however, is not equivalent to our case, because $m_\gamma^2 < 0$ is a one-loop result obtained within standard QED, but in the non-standard quantum-vacuum state.

It is further instructive to consider a classical field model with the standard mass term, but, e.g., in flat de Sitter spacetime with the Hubble constant H . In this case, the superhorizon modes increase without bound at future time infinity. These correspond to imaginary time-dependent frequencies. This fact does not serve any problem in Particle Physics, because those modes have

¹ The factor 2 in β comes from the Fermi normal coordinates, which are introduced in the near-horizon region.

a physical wavelength larger than the space-time curvature, i.e. no notion of particles can be associated with them [14].

It, now, seems that the tachyonic scalar field model with $\mu \lesssim H$ could be physically (almost) indistinguishable from a massless scalar field in local tests in the particle colliders. Still, the dispersion relation is $q_0^2 = \vec{q}^2 - \mu^2$ with the IR cutoff $l_{IR} \ll 1/H \lesssim 1/\mu$ and, hence, $|\vec{q}| \gg H \gtrsim \mu$. This, in turn, means that $dq_0/d|\vec{q}| \approx 1 + \mu^2/2\vec{q}^2 > 1$ holds for scalar particles. Moreover, any local perturbation (e.g., a particle) is oblivious to the large-scale structure and, thus, described by momenta with $|\vec{q}|$ much larger than $1/l_{IR}$,² implying that this perturbation is described by the group velocity larger than c .

This last example reflects our case, if one takes into account that $m_\gamma^2 < 0$ emerges due to the black-hole evaporation at one-loop approximation. Hence, it does not hold at length scales larger than r_H , because the field propagators, we have employed in the one-loop computations, are accurate in the local Minkowski frame only. Thus, a high-energy QED photon/electron might be able to propagate out of the black hole, bringing information about its internal structure. This appears to provide a semi-classical mechanism for the proposal of [10, 11] towards the resolution of the information-loss problem.

Black-hole explosions: The non-violent Einstein acausality near the event horizon implies that the black-hole explosions happen much earlier, i.e. rather before black holes reach the Planckian size. It might provide the explanation of why these explosions have not been observed so far, because these appear to be less energetic events in the sky than it has been thought.

Acknowledgments

It is a pleasure to thank the participants of the 3rd Karl Schwarzschild Meeting on Gravitational Physics and the Gauge/Gravity Correspondence for interesting discussions and the organizers for making it all happen. I am also thankful to Frans Klinkhamer and José Queiruga for their comments on an early version of this paper.

References

- [1] Aharonov, Y., Komar, A., Susskind L.: Superluminal behavior, causality, and instability. *Phys. Rev.* **182**, 1400 (1969)
- [2] Candelas, P.: Vacuum polarization in Schwarzschild space-time. *Phys. Rev. D* **21**, 2185 (1980)
- [3] Dhar, J., Sudarshan, E.C.G.: Quantum field theory of interacting tachyons. *Phys. Rev.* **174**, 1808 (1968)
- [4] Emelyanov, V.A.: Effective photon mass from black-hole formation. *Nucl. Phys. B* **919**, 110 (2017)
- [5] Emelyanov, V.A.: Quantum electromagnetic phenomena far from small evaporating black holes. In: Gomboc A. (ed.) *New Frontiers in Black Hole Astrophysics*, pp. 279-282. Cambridge University Press (2016)
- [6] Emelyanov, V.A.: Quantum kinetic theory of a massless scalar model in the presence of a Schwarzschild black hole. *Annalen Phys.* **529**, 1700078 (2017)
- [7] Emelyanov, V.A.: Magnetostatic-field screening induced by small black holes. *Nucl. Phys. B* **921**, 796 (2017)
- [8] Emelyanov, V.A.: Can a QED photon propagate out of the black-hole horizon? arXiv:hep-th/1703.05078
- [9] Emelyanov, V.A.: Two-loop QED photon self-energy near the black-hole horizon. In progress
- [10] Giddings, S.B.: Models for unitary black hole disintegration. *Phys. Rev. D* **85**, 044038 (2012)
- [11] Giddings, S.B.: Nonviolent nonlocality. *Phys. Rev. D* **88**, 064023 (2013)
- [12] Giddings, S.B.: Hawking radiation, the Stefan-Boltzmann law, and unitarization. *Phys. Lett. B* **754**, 39 (2016)
- [13] Le Bellac, M.: *Thermal Field Theory*. Cambridge University Press (1996)
- [14] Mukhanov, V.F., Winitzki, S.: *Introduction to Quantum Effects in Gravity*. Cambridge University Press (2007)

² It should be noted that the (standard) Fourier transform is inapplicable at length scales, at which the space-time curvature is non-negligible. This is, however, not a problem in Particle Physics, because particles are localized in coordinate space. The infinite-extent approximation of a local Minkowski frame is, thus, applicable, if particles are characterized by a high-enough energy scale.

- [15] Schroer, B.: Quantization of $m^2 < 0$ field equations. Phys. Rev. D **3**, 1764 (1971)
- [16] Unruh, W.G.: Origin of the particles in black hole evaporation. Phys. Rev. D **15**, 365 (1977)