



KFK-347:2

# KERNFORSCHUNGSZENTRUM KARLSRUHE

Mai 1965

Gesellschaft für Kernforschung m. b. H. KFK 347  
Zentralbibliothek

Institut für Angewandte Kernphysik

The Spin of the 2 sec Isomeric State of In<sup>116</sup>

W. Ponitz



- 7. Okt 1965



GESELLSCHAFT FÜR KERNFORSCHUNG M. B. H.  
KARLSRUHE



1.E.1:  
2.A.1

*Nuclear Physics* 66 (1965) 297–300; © North-Holland Publishing Co., Amsterdam

Not to be reproduced by photoprint or microfilm without written permission from the publisher

Gesellschaft für Kernforschung m.b.H.

THE SPIN OF THE 2 sec ISOMERIC STATE OF In<sup>116m2</sup>

W. PÖNITZ

Institut für Angewandte Kernphysik Kernforschungszentrum Karlsruhe

Received 17 November 1964

**Abstract:** The cross section ratios for the formation of the 2 sec and the 54 min isomeric states of In<sup>116</sup> due to neutron capture in the three lowest resonances of the In<sup>116</sup> compound nucleus were measured. A comparison of the results with the isomeric cross section ratios calculated by the  $\gamma$ -ray cascade statistics yields for the spin of the 2 sec state, a value  $J_{m2} = 5$ . The activation cross section for thermal neutrons was calculated from the isomeric cross section ratio in the 1.46 eV resonance to be  $\sigma_{m2}(0.0253 \text{ eV}) = 81 \pm 8 \text{ b}$ .

NUCLEAR REACTIONS In<sup>115</sup>(n,  $\gamma$ ),  $E = 1.4, 3.9, 9.1 \text{ eV}$ ; measured  $I_\gamma$ .  
In<sup>115</sup> deduced  $\sigma(\text{In}^{116} \text{ isomers})$ . In<sup>116</sup> deduced  $J$ . Natural target.

## 1. Introduction

Two isomeric states of indium 116 are known. The first isomeric state In<sup>116m1</sup> ( $T_{\frac{1}{2}} = 54.12 \text{ min}$ ) is a  $5^+$  state<sup>1,13</sup>) with an excitation energy<sup>2)</sup> of about 108 keV. This state decays by  $\beta$ -radiation to excited states of Sn<sup>116</sup>. The second isomeric state<sup>3,4</sup>) In<sup>116m2</sup> ( $T_{\frac{1}{2}} = 2.16 \text{ sec}$ ) decays by 163 keV  $\gamma$ -transitions to In<sup>116m1</sup>. From the E3 character of this isomeric transition it has been concluded that this is a  $8^-$  state. The spin value 8 was confirmed by the agreement between the excitation cross section of the 2 sec state for thermal neutrons measured by Fettweis<sup>5)</sup> and the calculated one obtained by the cascade statistics theory from Huizenga and Vandebosch<sup>6)</sup>. However, the approximately 20 times larger cross section measured by Alexander *et al.*<sup>4)</sup> is only compatible with smaller spin values. In order to determine which of the two measured cross sections  $\sigma_{m2}$  is more accurate and to get further information on the spin value of the second isomeric state of indium 116, the cross section ratios  $\sigma_{m2}/\sigma_{m1}$  in the three lowest resonances of the indium 116 compound nucleus were measured in the present work. In the first resonance<sup>7)</sup> ( $E = 1.46 \text{ eV}$ ),  $\sigma_{m2}/\sigma_{m1}$  was determined by two different methods. For the second and third resonances<sup>7)</sup> ( $E = 3.9$  and  $9.1 \text{ eV}$ ), the isomeric cross section ratios were measured relative to that in the first resonance.

## 2. Experimental Procedure and Results

Metallic indium foils were irradiated with neutrons of  $\approx 1.4, 3.9$  and  $9.1 \text{ eV}$  using crystal spectrometer at the Karlsruhe reactor FR2. The  $\gamma$ -activity of the foil was

counted in a 10.2 cm × 15.2 cm NaI(Tl) crystal either with a single- or a 256-channel analyser.

### 2.1. DETERMINATION OF $(\sigma_{m2}/\sigma_{m1})_{1st\ res}$

For the determination of  $(\sigma_{m2}/\sigma_{m1})_{1st\ res}$ , the  $\gamma$ -ray spectra after irradiation times of 10 sec or 20 min were measured for several runs. The cross section ratio can be obtained from the counting rates  $N$  in the photopeaks of the 163 keV  $\gamma$ -rays and the  $\gamma$ -rays of one transition in the  $Sn^{116}$  spectrum in the following way:

$$\sigma_{m2}/\sigma_{m1} = \left( \frac{N}{\eta_{\gamma} P_{\gamma} \epsilon f T} \right)_{2\ sec} / \left( \frac{N}{\eta_{\gamma} P_{\gamma} \epsilon f T} \right)_{54\ min} \quad (1)$$

Here  $f$  is the  $\gamma$ -ray intensity per decay for the transition considered,  $\eta_{\gamma}$  the total detection efficiency of the NaI(Tl) crystal and  $P_{\gamma}$  its peak-to-total ratio<sup>8)</sup>. The factor  $\epsilon$  represents the  $\gamma$ -absorption in the foil<sup>9)</sup> and  $T$  is a factor dependent on irradiation and counting time. From the internal conversion coefficients<sup>10)</sup> for the isomeric transition a value of  $f$  equal to  $0.37 \pm 0.03$  was calculated. Eq. (1) was evaluated for the 408, 1085 and 1270 keV  $\gamma$ -peaks of the 54 min spectra. This yields the average value

$$(\sigma_{m2}/\sigma_{m1})_{1st\ res} = 0.52 \pm 0.05.$$

The error is mainly due to the errors of  $f$ .

Another determination of the cross section ratio was made by observing the increase of the  $Sn^{116}$   $\gamma$ -activity in the first few seconds after a short-time irradiation due to the decay of the second to the first isomeric state. The counting rate of the 1085 and 1270 keV peaks after a 2 sec activation was measured. Thus the counting rate is

$$N(t) \approx A e^{-\lambda m1 t} - e^{-\lambda m2 t}, \quad (2)$$

where  $A$  is related in a straightforward manner with  $\sigma_{m2}/\sigma_{m1}$ . The evaluation of eq. (2) yields

$$(\sigma_{m2}/\sigma_{m1})_{1st\ res} = 0.49 \pm 0.05.$$

In this case the error depends mainly on statistical fluctuations of the counting rate.

### 2.2. RELATIVE DETERMINATION OF $(\sigma_{m2}/\sigma_{m1})_{2nd\ res}$ AND $(\sigma_{m2}/\sigma_{m1})_{3rd\ res}$

In order to determine  $\sigma_{m2}/\sigma_{m1}$  in the second and third resonance relative to the first resonance, the  $\gamma$ -rays with energies above 100 keV were measured after an irradiation time of 20 sec. These data allow the deduction of the ratio of  $\gamma$ -ray intensities of the decays of the 2 sec and 54 min isomeric states, respectively. The counting probabilities are eliminated by relating the ratio to that in the first resonance. The results are

$$(\sigma_{m2}/\sigma_{m1})_{2nd\ res}/(\sigma_{m2}/\sigma_{m1})_{1st\ res} = 0.98 \pm 0.06,$$

$$(\sigma_{m2}/\sigma_{m1})_{3rd\ res}/(\sigma_{m2}/\sigma_{m1})_{1st\ res} = 0.93 \pm 0.09.$$

### 3. Discussion

It is possible to calculate the isomeric cross section ratios from the  $\gamma$ -ray cascade statistics<sup>6</sup>). For this cascade model it is assumed that only E1 transitions occur and that the transition probability is proportional to the spin-dependent part of the nuclear level density  $\rho(J)$  only, where

$$\rho(J) = \rho(0)(2J+1)\exp\left(-\left(J+\frac{1}{2}\right)^2/2\sigma^2\right), \quad (3)$$

where  $\rho(0)$  is the level density at the spin  $J = 0$ ,  $\sigma$  is the spin cut-off parameter. One assumes furthermore that after  $\bar{N}_\gamma - 1$  transitions of E1 character the last one follows with the lowest spin differences to the final states ( $\bar{N}_\gamma$  is the average multi-




Fig. 1. Comparison of calculated and measured isomeric cross section ratios.

plicity of the  $\gamma$ -ray cascade). In order to compare the experimental results with those of the cascade theory, the values  $\bar{N}_{\gamma, 1st\ res} = 4.4$  and  $\bar{N}_{\gamma, 2nd\ res} = 5.6$  (ref. <sup>11</sup>), compound state spin values  $J_{1st\ res} = 5$  and  $J_{2nd\ res} = 4$  (ref. <sup>12</sup>) and the final state spin  $J_{m1} = 5$  (refs. <sup>13, 14</sup>) were used. The isomeric cross section ratios were calculated for the two limiting values  $\sigma = 2$  and  $\sigma = 5$  of the spin cut-off parameter<sup>15</sup>).

The ratio  $(\sigma_{m2}/\sigma_{m1})_{2nd\ res}/(\sigma_{m2}/\sigma_{m1})_{1st\ res}$  thus calculated for various spin values of the 2-sec isomer is shown in fig. 1 and it is seen that the experimental value is compatible with a spin value  $J_{m2} = 5$ . The value of  $(\sigma_{m2}/\sigma_{m1})_{1st\ res}$  itself allows us to determine the spin of the 2 sec state only by additional use of the value of  $(\sigma_g/\sigma_{m1})_{1st\ res}$  measured by Albold<sup>16</sup>) where  $\sigma_g$  is the formation cross section<sup>17, 1)</sup> for the ground state In<sup>116g</sup> ( $T_{\frac{1}{2}} = 14.10$  sec). By this alternative method, which was

previously used by Alexander *et al.*<sup>5</sup>),  $J_{m2} = 5$  was found, in agreement with the result of the method discussed above.

The spin value 5 was assigned with the aid of a statistical theory which does not take into account individual properties of states and transitions. It would be possible to consider the influence of the low-lying states of the compound nucleus as was done in calculations of  $\gamma$ -ray spectra<sup>18</sup>), but low-lying states of  $\text{In}^{116}$  and their  $\gamma$ -ray transitions have not been previously investigated. However, agreement between measured and calculated isomeric cross section ratios was observed in many cases<sup>6, 19-21</sup>). Furthermore, in this special case of equal final state spin values the assumption of E1 transitions in the  $\gamma$ -ray cascades is no restriction, as is easy to see.

Due to the overall constancy of the ratio  $\sigma_{m2}/\sigma_{m1}$  within experimental error,  $\sigma_{m2}$  for thermal neutrons can be calculated using  $\sigma_{m1}$  from earlier measurements with thermal neutrons<sup>1</sup>). The result is

$$\sigma_{m2}(0.0253 \text{ eV}) = 81 \pm 8 \text{ b.}$$

This is in agreement with  $\sigma_{m2}(0.0253 \text{ eV}) = 92 \pm 14 \text{ b}$  measured by Alexander *et al.*<sup>4</sup>) and disagrees with  $\sigma_{m2}(0.0253 \text{ eV}) = 4 \pm 1.5 \text{ b}$  measured by Fettweis<sup>5</sup>)

### References

- 1) K. H. Beckurts *et al.*, Nucl. Sci. Eng. **17** (1963) 329
- 2) J. Colard *et al.*, J. Phys. Rad. **21** (1960) 863
- 3) P. H. Heckmann *et al.*, Z. Phys. **163** (1961) 451
- 4) K. F. Alexander *et al.*, Phys. Lett. **4** (1963) 302
- 5) P. Fettweis, Phys. Lett. **3** (1962) 40
- 6) J. R. Huizenga and R. Vandenbosch, Phys. Rev. **120** (1960) 1305
- 7) D. J. Hughes and R. B. Schwartz, BNL 325 (1958) p. 47
- 8) C. Weitkamp, Nucl. Instr. **23** (1963) 13
- 9) G. W. Grodstein, National Bureau of Standards Circular 583 (1957)
- 10) L. A. Sliv and I. M. Band, Coefficients of internal conversion of gamma radiation (USSR Academy of Sciences, Moscow, Leningrad, 1956)
- 11) J. E. Draper and T. E. Springer, Nuclear Physics **16** (1960) 27
- 12) A. Stolovy, Phys. Rev. **118** (1960) 211
- 13) L. S. Goodman and S. Weber, Phys. Rev. **108** (1957) 1524
- 14) P. B. Nutter, Phil. Mag. **1** (1956) 587
- 15) M. Bormann and H. Neuert, Forts. Phys. **11** (1963) 277
- 16) E. Albold, to be published
- 17) A. Ducat and R. H. Thomas, Nuclear Physics **15** (1960) 525
- 18) E. S. Troubetzkoy, Phys. Rev. **122** (1961) 212
- 19) K. F. Alexander and H. F. Brinckmann, Ann. der Phys. **12** (1963) 225
- 20) B. Keisch, Phys. Rev. **129** (1963) 769
- 21) S. K. Mangal and P. S. Gill, Nuclear Physics **41** (1963) 372