



GLOWA

GLOWA-Jordan River Project

Regional Climate Simulations for the Middle East with MM5

Harald Kunstmann, Andreas Heckl, Peter Suppan, Gerhard Smiatek
Forschungszentrum Karlsruhe, Garmisch-Partenkirchen, Germany



**Institute for Meteorology and Climate Research
(IMK-IFU)
Garmisch-Partenkirchen**



GLOWA

Regional Climate Simulations

Objectives

- 1) Estimation of future temporal and spatial distribution of temperature and precipitation
- 2) Provision of climate change information to impact WPs in GLOWA-Jordan
- 3) Estimation of uncertainty bounds of climate change projections





Scientific Challenge

- 1) Changes in the regional climate can differ significantly from the overall trend of global climate change
- 2) Region has sharp climatic gradients: subhumid mediterranean ↔ arid climate
- 3) Resolution of global climate models are too coarse for hydrological & biological impact studies
⇒ Higher resolution information required that account for regional and local geographic features (particularly orography, land use and water bodies)


Approach:

Dynamic downscaling of global climate scenarios





Scientific Challenge




IPCC 4AR, 2007

**Eastern Mediterranean/Near East:
is in between increasing and decreasing dominant
large scale patterns of DJF precipitation change**



Regional Climate Simulations



phase III

⇒ Uncertainty bounds

Emission scenarios: based on different assumptions on future GHG emissions



GLOWA

Regional Climate Simulations

Population Growth Economic Development
Technological Progress



Emission Scenarios
Greenhouse Gas Concentrations



Global Climate Models



Global Climate Scenarios




Downscaling Methods



Regional Climate Scenarios




The Mesoscale Meteorological Model MM5




Land Use Discretization

Soil Discretization



- Dynamic Downscaling of ECHAM4 with MM5
- 3 nests: 54x54 km², 18x18 km², 6x6 km²
- 26 Vertical Layers, Model Top: 100 mbar (ca.17 km)
- Coupled OSU-Land-Surface Model
- Time slices: 1961-1990 & 2070-2099, B2, ECHAM4
- Transient: 1960-2050, A2 & B2, ECHAM4

Regional Climate Modeling



... accounts for soil-vegetation-atmosphere feedbacks



Basic differences between SVAT-based hydrological models and “traditional” hydrological models


- **SVAT-Hydro Models (designed for atmospheric feedback purposes):**
 - full energy balance (soil heat & sensible heat fluxes)
 - 2-way interaction with PBL
- **“Traditional”-Hydro models (designed for pure hydrol. applications):**
 - lateral water fluxes, surface runoff routing
 - deeper soils considered
 - finer vertical & horizontal resolutions
 - often groundwater interaction
 - often extensions for reactive flow & transport, erosion, etc.
 - but: depending on specific model choice**



GLOWA

Regional Climate Simulations

Quality of regional climate simulations: control run vs. long term observation




Simulated annual mean precipitation
(ECHAM4, 18 km², 1961-1990)
VS.
observed long term annual mean
(for selected stations 1961-1990)



How accurate does the downscaled Control Run reproduce observed precipitation?

Mean Annual Precipitation







GLOWA

Regional Climate Simulations

What do we expect from the High Resolution Simulations with **6 km**?

Intermediate results of 6 km runs: mean 1961-1975

Domain 1



Yearly Mean Precipitation 1961-1975

54km

18 km

6 km

... more detailed spatial information: land-sea & orography dependent features



GLOWA


Regional Climate Simulations

First example

ECHAM4, B2, 18km, 2070-99 vs. 1961-90




What are the expected changes in temperature?



Change in annual mean temperature

ECHAM4 & MM5, 18 km, B2, 2070-2099 vs 1961-1990, Jordan Area North of Dead Sea




Change in temporal distribution, averaged over domain 2




GLOWA

Regional Climate Simulations

What are the expected changes in precipitation?



Absolute change in [mm]





Relative Change in [%]

ECHAM4 & MM5, 18 km, B2, 2070-2099 vs 1961-1990



How does seasonal precipitation change depend on the region?

2070-2099 vs. 1961-1990, ECHAM4, B2, 18km




■ Area A ■ Area B ■ Area C ■ Area D

For all subregions: Decreased winter, increased spring precipitation






Are drought risks changing? Analysis of effective drought index EDI



Subregion A: shift towards drier conditions & increased drought risks



Are drought risks changing? Analysis of effective drought index EDI



Subregion A: Increasing drought intensities, but “unchanging” drought durations



GLOWA

Regional Climate Simulations

Second example

ECHAM4, A2 & B2, 54km, 1961-2050 transient




GLOWA

Regional Climate Simulations

2021-50


Change [K]

Control Run 1961-90





Szenario B2


Szenario A2



A2 (2021-2050)






Change (°K)








Regional changes in temperature



■ A2 2021-50 ■ B2 2021-50 ■ B2 2070-99




Regional changes in temperature





Regional Climate Simulations

Control Run 1961-90





Szenario B2

Szenario A2


2021-50

Change [mm]



2070-99

Change [mm]






Increase,
little differences between
A2 + B2 till 2050

**But significant decrease
in 2070-99!**






Changes in seasonality of precipitation






Changes in seasonality of precipitation





Changes in heavy precipitation frequency





Summary & Conclusions

- Increase of temperatures in all scenarios (up to +4°C)
- **Long term** projections of precipitation differ from **medium term** projections:
 - 1) precipitation & intensity increase till 2050 for scenarios A2 & B2 (transient)
 - 2) precipitation & intensity decrease till 2100 for scenario B2 (time slice)
- Little differences between A2 and B2 till 2050 in mean annual precipitation change **but significant differences in monthly changes**

“To Do” in phase II & identified research needs for phase III

⇒ **what happens between 2050 and 2070?**

⇒ continuation of transient simulations in 18km resolution

⇒ extension towards HadCM3 and ECHAM5 (A1B)

A blue-tinted image featuring a crown with a central jewel, set against a background of concentric ripples in water. The crown is positioned in the center, and the ripples create a sense of depth and movement. The overall color scheme is a gradient of blues, from light to dark.

Thank you for your attention



Summary: Available Regional Climate Simulations

Global Model	Scenario	Regional Model	Resolution	Time Slice	Availability
ECHAM4	CT	MM5	54km	1961-90	IMK-IFU
ECHAM4	CT	MM5	18km	1961-90	IMK-IFU
ECHAM4	CT	MM5	6km	1961-75	IMK-IFU
ECHAM4	B2	MM5	54km	2070-99	IMK-IFU
ECHAM4	B2	MM5	18km	2070-99	IMK-IFU
ECHAM4	B2	MM5	6km	2070-85	IMK-IFU end of 2007
ECHAM4	CT+A2	MM5	54km	1961-2050	IMK-IFU
ECHAM4	CT+A2	MM5	18km	1961-2050	IMK-IFU end of 2007
HadAM3P	CT	RegCM3	50km	1961-90	TAU
HadAM3P	A2	RegCM3	50km	2071-2100	TAU
HadAM3P	B2	RegCM3	50km	2071-2100	TAU
NASA FV GCM	CT	RegCM3	50km	1961-90	TAU
NASA FV GCM	A2	RegCM3	50km	2071-2100	TAU
ECHAM5	CT+A1B	RegCM3	50km	1960-2050	TAU
ECHAM4	CT	STAR		1958-1996	PALAST
ECHAM4	A1B	STAR		2007-2040	PALAST




GLOWA


Regional Climate Simulations

ECHAM5 transient vs. ECHAM4 transient

Mean Annual Precipitation



ECHAM4-MM5, $\Delta x=18\text{km}$



ECHAM5-RegCM, $\Delta x=0.5^\circ$

Control runs 1961-90 vs. long term observation at 41 stations in the Near East