

# Improved regional climate model data for climate change impact studies in the VGTB basin of Vietnam

P. Laux<sup>1</sup>, S. Vogl<sup>2</sup>, G. Mao<sup>1</sup>, T. Dang<sup>3,1</sup>, H. Kunstmann<sup>1</sup>

1 Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research IMK-IFU, 82467 Garmisch-Partenkirchen, Germany

2 Technical University Deggendorf, 94469 Deggendorf

3 Institute of Meteorology, Hydrology and Environment (IMHEN), Hanoi, Vietnam

## Introduction

Hydro-meteorological information is crucial for regional and local impact studies. Regional Climate Models (RCMs) are suitable to downscale large-scale climate information to regional (SE Asia, Fig. 1) and local (VGTB basin, Fig. 2) scales. After downscaling, there remains still a mismatch between observation data and high resolution RCM results. This holds especially true for precipitation, which is highly variable in space and time, and thus difficult to correct.




Fig. 1: Mean daily precipitation bias (WRF-ERA40 minus APHRODITE) for 1971-2000, (left). WRF-ERA40 (@Domain2: 15x15 km) is re-gridded onto the APHRODITE grid (0.25°) using bilinear interpolation. Digital elevation model (DEM) as used in the WRF-ERA40 simulations (right).

- Wet bias of WRF-ERA40 @Domain2 is observed, which is higher for complex terrain (Fig. 1)
- Problems to capture seasonality (Fig. 2, top)




Fig. 2: Mean monthly precipitation amount obtained by WRF-ERA40 for Domain 1, 5x5 km (top) and Domain 2, 15x15 km (bottom) at grid cells, 15x15 km corresponding to 23 rain gauges in the Vu Gia-Thu Bon basin (1971-2000).

For WRF-ERA40 @Domain3:

- Still wet bias, but significantly reduced during the rainy period (SOND)
- Seasonality of precipitation well captured (Fig. 2, bottom)

## Bias correction methods:

### 1. Mean correction:

- Linear approach based on long-term monthly correction factors
- Easy applicable for future climate projections

### 2. Quantile mapping:

- Corrects for the statistical distribution (here: performed for extreme and normal part of distribution separately)
- Easy applicable to future climate projections, but historical (daily) values cannot be exceeded

### 3. Copula-based correction (still experimental stage):


- Dynamic approach (performed for each time step separately) and based on the concept of Copulas.
- Models dependence structure (Copula) between variables (here: RCM and observed precipitation, but can be extended to multivariate)
- Generates a probability density function (PDF) for each time step → allows uncertainty estimation
- Spatial extension: Estimate time series for location between the observation stations (based on concept of similarity)
- New and promising approach (Laux et al., 2011; Vogl et al., 2012), but not yet applicable to correct future climate projections

## Objectives

- Analyze **skill** of different bias correction methods
- Provide a **set of corrected data** to local stakeholders in the VGTB basin and to LUCCI consortium

## Procedure & Results

- Split series into Calibration/Validation period



Observations: Precipitation data obtained by IMHEN  
WRF-ERA40: WRF simulation driven by ERA40 reanalysis (ECMWF)

- Calculate performance measures for the validation period




Fig. 3: RMSE and NSE of bias corrected precipitation compared to observed precipitation for 23 observation locations (top). RMSE (NSE) > (<) 60 (-1) are set to NaN for illustration. RMSE for TamKy (bottom, left) and DaNang (bottom, right). The black line shows the WRF raw data (without any bias correction).

- Lowest RMSE for the q-q approach (global value), cop bc (mode) better for most quantiles
- Cop bc (median) performed best by cop bc (median) (higher NSE values)
- Combination of bc mode (normal part) and bc mean (extremes) could improve the results

## Towards hydrological impact studies ...




Fig. 4: First simulations using hydrological model Mike-SHE for Domain 2 (15 km, left) after bias correction and for Domain 3 (5 km) after bias correction.