

HYLA	VOL. 2011.	No. 2	Str. 37-38	PHOTO NOTE
------	------------	-------	------------	------------

## A field observation of common buzzard predating on common toad

### Terensko opažanje predacije običnog škanjca na sneđu krastaču

OLGA JOVANOVIĆ\*<sup>1</sup>, GORAN ŠAFAREK<sup>1</sup>, MIROSLAV SAMARDŽIĆ<sup>2</sup>

<sup>1</sup>Society for Research and Popularization of Science Baobab, Trg k. Krešimira 10, 48000 Koprivnica, Croatia, corresponding author: [jovanovic.olga@googlemail.com](mailto:jovanovic.olga@googlemail.com)

<sup>2</sup> Ecological Society Koprivnica, Hrvatske državnosti 10, 48 000 Koprivnica, Croatia




Figure 1. Skinned toad in the breeding pond near Đelekovec  
Slika 1. Koža smeđe krastače na mrijestilištu u blizini Đelekovca

Common toad *Bufo bufo* (Linnaeus, 1758), similarly to many species of bufonid family, has parotoid glands that release a secretion that is toxic to many predators and serve as defence mechanism (WILBER, CARTOLL, 1940, TRONCHET, 1952, TOLEDO, JARED, 1995). In that context, predation on bufonid frogs is not very common, and it mostly happens under specific conditions (e.g. during the breeding season; OLSON, 1989, SLATER, 2002). We observed common buzzard *Buteo buteo* (Linnaeus, 1758)

on 16<sup>th</sup> March 2008 in the dusk, skinning the common toads that were injured in the traffic during their spring migration to the breeding site, on the county road from Koprivnica to Đelekovec. On the next morning, in the nearby pond which is the main breeding site of *Bufo bufo* in this area, we observed completely skinned individual of common toad, which was skinned and eaten by common buzzard. The same predation pattern was also observed on the road near the gravel excavation pit Gabajeva Greda. It seems that this pattern is here even more common due to lower frequency of traffic. Skinning of common toads was previously reported by Eurasian otter *Lutra lutra* (Linnaeus, 1758) (SLATER, 2002), while the same pattern has also been reported for other bufonid frogs (e.g. *B. boreas* predated by *Corvus corax*; OLSON, 1989).

#### REFERENCES

- OLSON, D. H. (1989): Predation on breeding western toads (*Bufo boreas*). *Copeia*: 1989 (2): 391-397.
- SLATER, F. (2002): Progressive skinning of toads (*Bufo bufo*) by the Eurasian otter (*Lutra lutra*). *IUCN Otter Specialist Group Bulletin* 19 (1): 25-29.
- TOLEDO, R. C. & JARED, C. (1995): Cutaneous granular glands and amphibian venoms. *Comparative Biochemistry and Physiology Part A: Physiology*, 111 (1): 1-29.
- TRONCHET, J. (1952): Contribution a l'étude histochimique des glandes paratoides du crapaud commun *Bufo bufo* (Linne). *Bulletin de la Societe Zoologique de France* 71: 204-215.
- WILBER, C. G. & CARTOLL, P. L. (1940): Studies on the histology of the glands in the skin of Anura. I. The parotoid gland of *Bufo americanus*. *Holbrook. Transactions of the American Microscopical Society* 59: 123-128.