INSECTA MUNDI A Journal of World Insect Systematics

0334

On the southeastern United States distributions of *Stictoleptura* canadensis (Olivier, 1795), Leptura subhamata Randall, 1838 (Coleoptera: Cerambycidae: Lepturinae) and Heterosternuta cocheconis (Fall, 1917) (Coleoptera: Dytiscidae: Hydroporinae)

> John M. Leavengood, Jr. University of Kentucky Department of Entomology S-225 Agricultural Science Center Building – North Lexington, KY 40546-0091

Eric G. Chapman University of Kentucky Department of Entomology Agricultural Science Center Building – North Lexington, KY 40546-0091

Date of Issue: January 24, 2014

John M. Leavengood, Jr. and Eric G. Chapman

On the southeastern United States distributions of *Stictoleptura canadensis* (Olivier, 1795), *Leptura subhamata* Randall, 1838 (Coleoptera: Cerambycidae: Lepturinae) and *Heterosternuta cocheconis* (Fall, 1917) (Coleoptera: Dytiscidae: Hydroporinae) Insecta Mundi 0334: 1-3

ZooBank Registered: urn:lsid:zoobank.org:pub:557BF8B7-CD0A-4D63-AD31-CE2031703822

Published in 2014 by Center for Systematic Entomology, Inc. P. O. Box 141874 Gainesville, FL 32614-1874 USA http://www.centerforsystematicentomology.org/

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology. Manuscript preparation guidelines are available at the CSE website.

Managing editor: Eugenio H. Nearns, e-mail: gino@nearns.com
Production editor: Michael C. Thomas, Paul E. Skelley, Brian Armitage, Ian Stocks
Editorial board: J. H. Frank, M. J. Paulsen
Subject editors: G.B. Edwards, J. Eger, A. Rasmussen, G. Steck, I. Stocks, A. Van Pelt, J. Zaspel, N.P. Lord, A. Brunke
Spanish editors: Julieta Brambila, Angélico Asenjo

Printed copies (ISSN 0749-6737) annually deposited in libraries: CSIRO, Canberra, ACT, Australia Museu de Zoologia, São Paulo, Brazil Agriculture and Agrifood Canada, Ottawa, ON, Canada The Natural History Museum, London, Great Britain Muzeum i Instytut Zoologii PAN, Warsaw, Poland National Taiwan University, Taipei, Taiwan California Academy of Sciences, San Francisco, CA, USA Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA Field Museum of Natural History, Chicago, IL, USA National Museum of Natural History, Smithsonian Institution, Washington, DC, USA Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico. Florida Virtual Campus: http://purl.fcla.edu/fcla/insectamundi University of Nebraska-Lincoln, Digital Commons: http://digitalcommons.unl.edu/insectamundi/ Goethe-Universität, Frankfurt am Main: http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/

Author instructions available on the Insecta Mundi page at: http://www.centerforsystematicentomology.org/insectamundi/

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. http://creativecommons.org/ licenses/by-nc/3.0/

On the southeastern United States distributions of *Stictoleptura* canadensis (Olivier, 1795), *Leptura subhamata* Randall, 1838 (Coleoptera: Cerambycidae: Lepturinae) and *Heterosternuta cocheconis* (Fall, 1917) (Coleoptera: Dytiscidae: Hydroporinae)

John M. Leavengood, Jr. University of Kentucky Department of Entomology S-225 Agricultural Science Center Building – North Lexington, KY 40546-0091 tokay@ufl.edu

Eric G. Chapman University of Kentucky Department of Entomology Agricultural Science Center Building – North Lexington, KY 40546-0091 ericgchapman@gmail.com

Abstract. We present new southeastern United States distributional records for *Stictoleptura canadensis* (Olivier, 1795), *Leptura subhamata* Randall, 1838 (Coleoptera: Cerambycidae: Lepturinae), and *Heterosternuta cocheconis* (Fall, 1917) (Coleoptera: Dytiscidae: Hydroporinae) which extend the southernmost limits of the known distributions for these species. We present new state records for these species.

Introduction

The recent collection of a series of specimens of *Stictoleptura canadensis* (Olivier, 1795), *Leptura subhamata* Randall, 1838 (Coleoptera: Cerambycidae: Lepturinae) and *Heterosternuta cocheconis* (Fall, 1917) (Coleoptera: Dytiscidae: Hydroporinae) from various southeastern localities in West Virginia, Kentucky and Georgia (by the authors) queued an interest in the known distribution of these three species. Current literature records describe the distribution of *S. canadensis* as southeastern Canada and the northeastern United States for subspecies *S. c. canadensis* (Olivier), western Canada to California for the subspecies *S. canadensis cribripennis* (LeConte, 1859), and Arizona and New Mexico for the subspecies *S. canadensis arizonensis* Linsley and Chemsak, 1976 (Monné et al. 2007). The distribution of *L. subhamata* ranges from southeastern Canada through the northeastern United States and south to Tennessee (Linsley and Chemsak 1976; Monné et al. 2007; Lingafelter 2007). These distributions are in agreement with Linsley and Chemsak (1972, 1976) regarding eastern North America. The distribution of *H. cocheconis* ranges from southern Quebec and central New Brunswick to central New York and Connecticut (Matta and Wolfe 1981; Larson et al. 2000).

List of Depository Abbreviations

Distribution records were collected from the following private and institutional depositories:

- CUIC Cornell University Insect Collection (Ithaca, NY)
- EDNC North Carolina Department of Agriculture (Raleigh, NC)
- EGCC Eric G. Chapman, private collection (Lexington, KY)
- ENPC Eugenio H. Nearns, private collection (Gaithersburg, MD)
- FSCA Florida State Collection of Arthropods (Gainesville, FL)
- GCIM Georgia College Insect Museum (Milledgeville, GA)
- JMLC John M. Leavengood, Jr., private collection (Lexington, KY)
- JPHC Jeffrey P. Huether, private collection (Geneva, NY)

2 •	INSECTA	MUNDI	0334,	January	2014
-----	---------	-------	-------	---------	-------------

MCZ —	_	Museum of Comparative Zoology, Harvard University (Cambridge, MA)
OSUC —	_	The Ohio State University (Columbus, OH)
UMIC —	_	University of Mississippi (Oxford, MI)
NMNH —	_	National Museum of Natural History, Smithsonian Institution (Washington, DC)
WVDA —	_	West Virginia Department of Agriculture (Charlestown, WV)

Stictoleptura c. canadensis (Olivier)

Distribution Records for *Stictoleptura c. canadensis* (Olivier). North Carolina: Yancey Co.: Mount Mitchell, 11-VIII (NMNH). **Tennessee**: Sevier County, 7-IX (CUIC). **Virginia**: (Chesterfield/ Henrico Co.:) Richmond, 5-VIII (Jeff Huether, personal communication); Chelsea, 6-VIII (NMNH). **West Virginia**: Hancock Co.: Tomlinson Run State Park, 15-VIII-2007, M. Harriman (WVDA); Pocahontas Co.: Droop, State Hwy 39/55, @ junction Hill Creek Falls Road, 8 August 2009, 3390' elev., 38°10.86'N, 80°20.39'W, EG Chapman & JM Leavengood (EGCC); Pocahontas Co.: Droop, State Hwy 39/55, @ junction Hill Creek Falls Road, 8-VIII-2009 elevation 3390', 38°10.51'18 N/80°20.21'55 W, Coll: J. M. Leavengood, Jr. & E. Chapman (JMLC); Preston Co.:r: Aurora, 20-VIII (NMNH); Cranesville, 14-VII (NMNH); Tucker Co.:r Dolly Sods, 6-VIII-1986, M. C. Thomas (WVDA).

Discussion on *Stictoleptura c. canadensis.* While not indicated on the label data, the majority of JMLC and EGCC specimens from West Virginia were collected on Queen Anne's Lace (Apiaceae: *Daucus carota* Linnaeus) in the presence of milkweeds (Apocynaceae: *Asclepias* sp.). Records from North Carolina, **new state record**, and Tennessee, **new state record**, represent the southernmost known distribution of the species.

Leptura subhamata Randall

Distribution Records for Leptura subhamata Randall. Georgia: Lumpkin Co.: Chattahoochee National Forest, near Boggs Creek, 9-VI-2001, E.H. Nearns (ENPC); Murray County (FSCA); Towns Co.: on flowers along Left Mill Creek Rd, ~2 mi east of SR 17, 34°51.84'N, 83°41.38'W, 20-VI-2010, EG Chapman & JM Leavengood (EGCC); Kentucky: Laurel Co.: along Craigs Creek Road, on milkweed, 5-VI-2010, 36°59.17'N, 84°15.09'W, Eric G Chapman (EGCC); Laurel Co.: Daniel Boone National Forest, Craig's Creek road, 5-VI-2009, ca. 36°59.173'/ 84°'15.086', J. M. Leavengood, Jr., EG chapman & MV Spelman (JMLC); Pulaski Co.: Bee Rock Campground, east end at trail head, 37.0297°N, 84.3195°W, Eric G Chapman (EGCC). North Carolina: Haywood County (FSCA, UMIC); Macon County (FSCA); Macon Co.: Clear Creek, Highlands, 1-VII (CUIC); Highlands, 18-VI (MCZ); Swain County: Andrew's Bald, Great Smoky Mountains National Park, 28-VI (MCZ); Watauga Co.: Zionville, V (EDNC); South Carolina: Oconee Co.: Mount Rest (Jeff Huether, personal communication); Pickens Co.: Rocky Bottom (Jeff Huether, personal communication). Tennessee: Blount County (FSCA); Sevier Co.: Great Smoky Mountains National Park, on dead fraser fir (JPHC); Gatlinburg, 23-VII (OSUC); West Virginia: Jackson Co.: Ripley, 25-VI (CUIC); Morgan Co.: Paw Paw, 23-VI (MCZ); Nicholas Co.: Richwood, 4/VII, on Hydrangea arborescens (FSCA); Pendleton Co.: 5 mi. NW of Ruddle; 16 June 1993, J. D. Hacker, Malaise trap (WVDA); Pocahontas Co.: Droop, State Hwy 39/55, @ junction Hill Creek Falls Road, 8-VIII-2009 elevation 3390', 38°10.51'18 N/80°20.21'55 W, Coll: J. M. Leavengood, Jr. & E. Chapman (JMLC); Pocahontas Co.: Monongahela National Forest, Tea Creek Campground, on flowers, 25-26-VI-2010 elevation 3350', 38°10.51'18 N/80°20.21'55 W, Coll: J. M. Leavengood, Jr. w/ EG chapman & MV Spelman (JMLC); Pocahontas Co.: Monongahela Natl. For., Cranberry Glades, 8 July 1986, M. C. Thomas (WVDA); Spruce, 8 August 1971, P. D. Hardwood (WVDA); along Williams River Road, from Tea Creek Campground to SR 150, 26-VI-2010, 38°20.99'N, 80°12.76'W, EG Chapman, JM Leavengood and MV Spelman (EGCC); Webster Co.: Monongahela Natl. For., 10 mi. NW Richwood, 9 July 1986, H. V. Weems Jr. (WVDA).

Discussion of *Leptura subhamata*. Records from Georgia, **new state record**, represent the southernmost known limits of this species. This is not to suggest the "discovery" or expansion of the species in these areas, but rather that the lack of a recent revision of this group has resulted in incomplete published distributions.

Heterosternuta cocheconis (Fall)

Distribution Records for *Heterosternuta cocheconis* (Fall). West Virginia: Pocahontas Co.: tributary to Tea Creek at Tea Creek campground, east of Williams River Road, 38°20.99'N, 80°12.76'W, 2988 m elev., 8 August 2009 (EGCC); tributary to Tea Creek at Tea Creek campground, east of Williams River Road, 38°20.99'N, 80°12.76'W, 2988 m elev., 27 June 2010 (GCIM).

Discussion of *Heterosternuta cocheconis*. These collections from West Virginia, **new state record**, extend this species' southern distribution at least 350 miles south-southwest. This is not to suggest the "discovery" or expansion of the species in these areas, but rather that the lack of a recent revision of this group has resulted in incomplete published distributions.

Acknowledgments

We are grateful to David Serrano (Broward College) and David T. Almquist (Florida Natural Areas Inventory) for preliminary reviews of the manuscript. We offer many thanks to Michael C. Thomas (FSCA), Laura T. Miller (WVDA), Jeff P. Huether (JPHC) and Eugenio Nearns (ENPC) for providing locality information from collections in their care. Most notably, Jeff Huether provided all records from the following museums: CUIC, EDNC, MCZ, OSUC, UMIC and NMNH. We thank G. W. Wolfe (Georgia State College and University) for confirming the identification of *Heterosternuta cocheconis*.

Literature Cited

- Larson, D. J., Y. Alarie, and R. E. Roughley. 2000. Predaceous diving beetles (Coleoptera: Dytiscidae) of the Nearctic Region, with emphasis on the fauna of Canada and Alaska. NRC Research Press; Ottawa., XIV + 982 p.
- Lingafelter, S. W. 2007. Illustrated key to the long-horned woodboring beetles of the eastern United States. Coleopterists Society, Special Publication No. 3: 1–206.
- Linsley, E. G., and J. A. Chemsak. 1972. Cerambycidae of North America Part VI, No. 1. Taxonomy and classification of the Subfamily Lepturinae. University of California Publications in Entomology 69: 1–138 p.
- Linsley, E. G., and J. A. Chemsak. 1976. Cerambycidae of North America. Part VI, No.2. Taxonomy and classification of the Subfamily Lepturinae. University of California Publications in Entomology 80e: 1–186 p.
- Matta, J. F., and G. W. Wolfe. 1981. A revision of the subgenus *Heterosternuta* Strand of *Hydroporus* Clairville (Coleoptera: Dytiscidae). Pan-Pacific Entomologist 57: 176–219.
- Monné, M. A., L. G. Bezark, and F. T. Hovore [Compilers]. 2007. Checklist of the Cerambycidae, or long-horned beetles (Coleoptera) of the Western Hemisphere. BioQuip Products; Rancho Dominguez, California. 417 p.

Received July 23, 2013; Accepted October 21, 2013.