

INSECTA MUNDI

A Journal of World Insect Systematics

0338

The centipede *Scolopendra morsitans* L., 1758,
new to the Hawaiian fauna, and potential representatives of the
“*S. subspinipes* Leach, 1815, complex”
(Scolopendromorpha: Scolopendridae: Scolopendrinae)

Rowland M. Shelley
Research Laboratory
North Carolina State Museum of Natural Sciences
MSC #1626
Raleigh, NC 27699-1626 USA

William D. Perreira
P.O. Box 61547
Honolulu, HI 96839-1547 USA

Dana Anne Yee
1717 Mott Smith Drive #904
Honolulu, HI 96822 USA

Date of Issue: January 31, 2014

Rowland M. Shelley, William D. Perreira, and Dana Anne Yee
The centipede *Scolopendra morsitans* L., 1758, new to the Hawaiian fauna, and potential representatives of the “*S. subspinipes* Leach, 1815, complex” (Scolopendromorpha: Scolopendridae: Scolopendrinae)
Insecta Mundi 0338: 1-4

ZooBank Registered: urn:lsid:zoobank.org:pub:E3C358D2-298F-4029-96B7-E4595E36A9B4

Published in 2014 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology. Manuscript preparation guidelines are available at the CSE website.

Managing editor: Eugenio H. Nearn, e-mail: gino@nearns.com

Production editors: Michael C. Thomas, Paul E. Skelley, Brian Armitage, Ian Stocks, Eugenio H. Nearn

Editorial board: J. H. Frank, M. J. Paulsen

Subject editors: G.B. Edwards, Joe Eger, A. Rasmussen, Gary Steck, Ian Stocks, A. Van Pelt, Jennifer M. Zaspel, Nathan P. Lord, Adam Brunke

Spanish editors: Julieta Brambila, Angélico Asenjo

Website coordinator: Eugenio H. Nearn

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and Agrifood Canada, Ottawa, ON, Canada
The Natural History Museum, London, Great Britain
Muzeum i Instytut Zoologii PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/>

Author instructions available on the *Insecta Mundi* page at:

<http://centerforsystematicentomology.org/insectamundi/>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

The centipede *Scolopendra morsitans* L., 1758, new to the Hawaiian fauna, and potential representatives of the “*S. subspinipes* Leach, 1815, complex” (Scolopendromorpha: Scolopendridae: Scolopendrinae)

Rowland M. Shelley
Research Laboratory
North Carolina State Museum of Natural Sciences
MSC #1626
Raleigh, NC 27699-1626 USA
rowland.shelley@naturalsciences.org

William D. Perreira
P.O. Box 61547
Honolulu, HI 96839-1547 USA
24snow@hawaii.rr.com

Dana Anne Yee
1717 Mott Smith Drive #904
Honolulu, HI 96822 USA
dana1@hawaii.rr.com

Abstract. *Scolopendra morsitans* L., 1758, is documented from Honolulu, Oahu, Hawaiian Islands, the first record of this anthropochoric chilopod from both the archipelago and state. Hawaii thus becomes the second American state to harbor the species, the other being Florida, where an individual has been taken in Jacksonville, Duval County. Meristic and morphological data are presented for three Hawaiian specimens. At least two other species of *Scolopendra*, both introduced, occur on these islands: *S. polymorpha* Wood, 1861, known only from one specimen from Oahu, and one or more representatives of the “*S. subspinipes* Leach, 1815, complex,” which is widespread and even inhabits Midway Atoll.

Key Words. *Scolopendra subspinipes*, *S. polymorpha*, Scolopendrinae, introduction, Oahu.

Introduction

In the review of Hawaiian scolopendromorph centipedes, all of which possess 21 leg-bearing segments, Shelley (2000) reported two species, both anthropochores, in the nominate subfamily of Scolopendridae - *Scolopendra subspinipes* Leach, 1815, occurring throughout the archipelago including Midway Atoll, and *S. polymorpha* Wood, 1861, represented by one individual from an Oahu pineapple plantation. The latter derives from North America, where it occupies an irregular area from the Central Plains westward to the Pacific Ocean, extending northward to Washington and Montana, USA, and southward to Baja California Sur and Guerrero, Mexico (Shelley 2002). Shelley (2000, 2002) combined the subspecies of *S. subspinipes* under the nominate (Kraepelin 1903; Attems 1914, 1930; Chamberlin 1944; Chamberlin and Wang 1952; Wang 1962; Schileyko 2007), but Kronmüller (2012) showed them to be full species that can be collectively referenced as the “*S. subspinipes* complex.” The dominant scolopendrines in southeast Asia, three have been widely introduced in Oceania (Shelley 2004), and as *S. subspinipes*, the assemblage was first documented from Hawaii by Gervais (1847), suggesting transport there and throughout the Pacific by migrating Polynesians (Buxton and Hopkins 1927, Shelley 2000). Not recalling which representatives have been encountered in the Hawaiian Islands, we incorporate the three most widely introduced species into the ensuing key.

While sampling in Sand Island State Recreation Area at the entrance to Honolulu Harbor, Oahu, in August 2013, WDP and DAY discovered three moderate-size scolopendrines beneath trunks of coconut palms (*Cocos nucifera* L.) that were shipped to RMS for deposit in the North Carolina State Museum of

Natural Sciences (NCSM). They are *S. morsitans* L., 1758, considered native to Africa, Australia, and Asia from India to the Philippines/Papua New Guinea (Koch 1983, Khanna 2001, Shelley et al. 2005, Shelley 2006), which constitutes a third anthropochoric Hawaiian scolopendrine. A fourth specimen, not shipped but considered conspecific, was discovered at the same locality three weeks later in beach sand beneath driftwood at the high water mark. Although one of the first three individuals is a subadult and markedly smaller, all match published diagnoses of *S. morsitans*, with the cephalic plate overlapping the first tergite that lacks the anterior transverse sulcus (terminology per Lewis et al. [2005] and Bonato et al. [2010]). An ultimate leg on one larger specimen is slightly shorter and exhibits different spination; apparently the original appendage was lost and a new one regenerated. Meristic and morphological data are provided in the following table; illustrations are available in Attems (1930, fig. 38), Lewis (2001, figs 2-4), and Shelley (2002, figs. 57-60). Shelley (2004) cited *S. morsitans* from the following Oceanian countries and territories: Cook Islands, Federated States of Micronesia, French Polynesia, Guam, Kiribati, New Caledonia, Northern Marianna Islands, Solomon Islands (Papua New Guinea), Republic of the Marshall Islands, Tonga, and Tuvalu. Hawaii becomes the second US state where the centipede has been documented, the other being Florida, where an individual from Jacksonville, Duval County, constituted the first authentic North American record (Shelley et al. 2005). Sample data are as follows:

USA, Hawaii, Oahu, Sand Island State Recreation Area at entrance to Honolulu Harbor (N 21° 18' 05.23", W 157° 52' 55.99"), 3 specimens, 26 August 2013, WD Pereira, DA Yee (NCSM) and 1 individual, 4 September 2013, WD Pereira (Bishop Museum, Honolulu).

***Scolopendra morsitans* from Hawaii.** L, left; R, right; PFP, Prefemoral Process. Measurements are in mm.

Length	Width	L Anten- nomeres	R Anten- nomeres	L Setal Transi- tion	R Setal Transi- tion	L Ventral Prefemoral Spines/Rows	R Ventral Prefemoral Spines/Rows	L PFP Spines	R PFP Spines
62.9	6.3	20	19	6	6 ^{1/2}	3/3/3	3/3/3	3	4
67.4	7.1	20	20	6	6	3/4/3	2/1/1/2	4	8
48.4	5.3	21	21	5 ^{3/4}	6 ^{1/4}	3/3/3	3/3/3	4	2

Key to known and potential Hawaiian Scolopendromorpha. In the spination figures in couplet 6, normal conditions are not within parentheses and rare ones are; thus (1)2(3) means normally with two spines, occasionally with one or three.

1. With four ocelli on each side of the cephalic plate. **2**
— Ocelli absent **Cryptopidae**
2. Spiracles rounded and without valves, those on segment 3 typically larger.
..... **Scolopendridae: Otostigminae**
— Spiracles triangular, valvular, and subequal in size (Scolopendridae: Scolopendrinae). **3**
3. Tergite 1 with conspicuous anterior transverse sulcus. ***Scolopendra polymorpha* Wood, 1861**
— Without this character. **4**
4. Prefemora of ultimate legs with 0-3 ventral spines arranged in one longitudinal row; large-
bodied, aggressive species, adults >15 cm (6") long. **5**
— Ultimate prefemora typically with 9-10 ventral spines arranged in three longitudinal rows;
smaller species, adults <7.5 cm (3") long. ***S. morsitans* L., 1758**
5. Ultimate prefemora without ventral spines. ***S. dehaani* Brandt, 1840**
— Ultimate prefemora with ventral spines. **6**

6. Ultimate prefemora with (1)2(3) ventral spines, coxopleural process with (1)2(3) spines.
 *S. subspinipes* Leach, 1815
 — Ultimate prefemora with (2)3 ventral spines, coxopleural process with 2-3 spines.
 *S. japonica* L. Koch, 1878

Acknowledgments

We thank F.G. Howarth and J.G.E. Lewis for beneficial pre-submission reviews.

Literature Cited

- Attems, C. 1914.** Die indo-australischen myriopoden. Archiv für Naturgeschichte. 80: 1–398.
- Attems, C. 1930.** Myriapoda 2. Scolopendromorpha. Das Tierreich 54: 1–308.
- Bonato, L., G. D. Edgecombe, J. G. E. Lewis, A. Minelli, L. A. Pereira, R. M. Shelley, and M. Zapparoli. 2010.** A common terminology for the external anatomy of centipedes (Chilopoda). ZooKeys 69: 17–51.
- Buxton P. A., and G. H. Hopkins. 1927.** Researches in Melanesia and Polynesia. An account of investigations in Samoa, Tonga, Ellice group and the New Hebrides in 1924, 1925 (1–4). London School of Tropical Medicine, London, UK, 260 p.
- Chamberlin, R. V. 1944.** Some chilopods from the Indo-Australian Archipelago. Notulae Naturae of the Academy of Natural Sciences of Philadelphia. 147: 1–14.
- Chamberlin, R. V., and Y. M. Wang. 1952.** Some records and descriptions of chilopods from Japan and other oriental areas. Proceedings of the Biological Society of Washington 65: 177–188.
- Gervais, P. 1847.** Myriapodes. *In*: Walckenaer, C. A. and P. Gervais (eds.), Histoire Naturelle des Insectes. Aptères 4: 1–333, 577–595.
- Khanna, V. 2001.** A check-list of the Indian species of the centipedes (Chilopoda: Scolopendromorpha). Annals of Forestry 9(2): 199–219.
- Koch, L. E. 1983.** Morphological characters of Australian scolopendrid centipedes, and the taxonomy and distribution of *Scolopendra morsitans* L. (Chilopoda: Scolopendridae: Scolopendrinae). Australian Journal of Zoology 31: 79–91.
- Kraepelin, K. 1903.** Revision der Scolopendriden. Mitteilungen aus dem Naturhistorischen Museum in Hamburg 20: 1–276.
- Kronmüller, C. 2012.** Review of the subspecies of *Scolopendra subspinipes* Leach, 1815 with the new description of the South Chinese member of the genus *Scolopendra* Linnaeus, 1758 named *Scolopendra hainanum* spec. nov. Spixiana 35(1): 19–27.
- Lewis, J. G. E. 2001.** The scolopendrid centipedes in the collection of the National Museum of Natural History in Sofia (Chilopoda: Scolopendromorpha: Scolopendridae). Historia Naturalis Bulgarica 13: 5–51.
- Lewis, J. G. E., G. D. Edgecombe, and R. M. Shelley. 2005.** A proposed standardised terminology for the external taxonomic characters of the Scolopendromorpha (Chilopoda). Fragmenta Faunistica 48: 1–8.
- Schileyko, A. A. 2007.** The scolopendromorph centipedes (Chilopoda) of Vietnam, with contributions to the faunas of Cambodia and Laos. Part 3. Arthropoda Selecta 16(2): 71–95.
- Shelley, R. M. 2000.** The centipede order Scolopendromorpha in the Hawaiian Islands (Chilopoda). Bishop Museum Occasional Papers 64: 39–48.
- Shelley, R. M. 2002.** A synopsis of the North American centipedes of the order Scolopendromorpha (Chilopoda). Virginia Museum of Natural History Memoir 5: 1–108.
- Shelley, R. M. 2004.** Occurrences of the centipedes *Scolopendra subspinipes* Leach and *S. morsitans* L. on Pacific islands (Chilopoda: Scolopendromorpha: Scolopendridae). Entomological News 115(2): 95–100.
- Shelley, R. M. 2006.** A chronological catalog of the New World species of *Scolopendra* L., 1758 (Chi-

lopoda: Scolopendromorpha: Scolopendridae). *Zootaxa* 1253: 1–50.

Shelley, R. M., G. B. Edwards, and A. Chagas, Jr. 2005. Introduction of the centipede *Scolopendra morsitans* L., 1758, into northeastern Florida, the first authentic North American record, and a review of its global occurrences (Scolopendromorpha: Scolopendridae: Scolopendrinae). *Entomological News* 116(1): 39–58.

Wang, Y. M. 1962. The Chilopoda of the Philippine Islands. *Quarterly Journal of the Taiwan Museum* 15: 79–106.

Received October 28, 2013; Accepted January 22, 2014.