



You have downloaded a document from
RE-BUŚ
repository of the University of Silesia in Katowice

Title: Identification of iron-bearing minerals in basalts and pillow lavas of the Kaczawa Mountains using ^{57}Fe Mössbauer spectroscopy

Author: Dariusz Malczewski, Martyna Jeleń, Jerzy Żaba, Artur Błachowski, Krzysztof Ruebenbauer, Maria Dziurawicz

Citation style: Malczewski Dariusz, Jeleń Martyna, Żaba Jerzy, Błachowski Artur, Ruebenbauer Krzysztof, Dziurawicz Maria. (2017). Identification of iron-bearing minerals in basalts and pillow lavas of the Kaczawa Mountains using ^{57}Fe Mössbauer spectroscopy. " Nukleonika " (Vol. 62, Iss. 2, s. 145-148). doi 10.1515/nuka-2017-0021



Uznanie autorstwa - Użycie niekomercyjne - Bez utworów zależnych Polska - Licencja ta zezwala na rozpowszechnianie, przedstawianie i wykonywanie utworu jedynie w celach niekomercyjnych oraz pod warunkiem zachowania go w oryginalnej postaci (nie tworzenia utworów zależnych).





Identification of iron-bearing minerals in basalts and pillow lavas of the Kaczawa Mountains using ^{57}Fe Mössbauer spectroscopy

Dariusz Malczewski,
Martyna Jeleń,
Jerzy Żaba,
Artur Błachowski,
Krzysztof Ruebenbauer,
Maria Dziurawicz

Abstract. The Kaczawa Mountains along with the Kaczawa foothill comprise a complicated geological unit that is called the Kaczawa metamorphic (Sudetes, SW Poland). The aim of our work was to identify the iron-bearing minerals in samples of basalts and pillow lavas from the Kaczawa metamorphic using ^{57}Fe Mössbauer spectroscopy. Based on the preliminary results, the $\text{Fe}^{3+}/\text{Fe}^{2+}$ ratio in the samples was determined.

Keywords: Kaczawa Mountains • Mössbauer spectroscopy • basalts • pillow lavas

Introduction

The Kaczawa metamorphic is located in the Sudetes and covers an area of 700 km². The unit is built of igneous, metamorphic and sedimentary rocks from the Cambrian to the Neogene period. The geological structure is extremely complicated [1]. The unit is divided into two zones – the oldest and deepest one that includes rocks formed in the early Paleozoic era and the younger zone that lies over the first zone and is built from rocks of the Carboniferous to the Cretaceous period. Cenozoic (the Miocene Epoch) basalts occur between these two zones.

The Miocene basalt samples were collected from four locations – Wilcza Góra, Małe Organy Myśluborskie, Klecza and Czartowska Skała. Basalt is a mafic, extrusive igneous rock that was formed from the rapid cooling of basaltic lava exposed to the surface. Basalts generally contain plagioclase, pyroxene, olivine, hornblende and magnetite as important accessory mineral [2].

Pillow lavas were collected from two locations – Wąwóz Myśluborski and Wleński Gródek near Klecza. The pillow lavas contain characteristic pillow-shaped structures that are attributed to the extrusion of lava under the water of Carboniferous ocean floor. Pillow lavas are of a basaltic chemical composition [2].

Materials and methods

The rock samples were crushed and powdered in an agate ball mill and then prepared in the shape of a thin disk absorber. The Mössbauer transmission spectra were recorded at room temperature using RENOM MsAa-3 spectrometer and linear arrangement of a $^{57}\text{Co}/\text{Rh}$ source (50 mCi) absorber and de-

D. Malczewski[✉], M. Jeleń, J. Żaba, M. Dziurawicz
Faculty of Earth Sciences,
University of Silesia,
60 Będzińska Str., 41-200 Sosnowiec, Poland,
E-mail: dariusz.malczewski@us.edu.pl

A. Błachowski, K. Ruebenbauer
Mössbauer Spectroscopy Laboratory,
Pedagogical University,
2 Podchorążych Str., 30-084 Kraków, Poland

Received: 14 August 2016

Accepted: 11 December 2016




Fig. 1. ^{57}Fe Mössbauer spectra at room temperature of basalt samples. Solid dots – experimental data; thick solid line – fitted curve; thin solid lines – fitted doublets and sextets.

tector (proportional counter). All Mössbauer spectra were numerically analysed using the Recoil software.

Results and discussion

Basalts

The Mössbauer spectra of the four basalt samples are shown in Fig. 1. The hyperfine parameters derived from the fitting procedure are given in Table 1. The Mössbauer spectra of basalts show

doublets representing both Fe^{2+} (labelled with no. 1 and no. 2 in Table 1) and Fe^{3+} (labelled with no. 3) iron ions in octahedral coordination. The Fe^{2+} doublets (no. 1, Fig. 1) with isomer shift (IS) values of 1.11–1.25 mm/s and quadrupole splitting (QS) of 2.5–3.0 mm/s are characteristic for olivine [3]. Olivine is a group of nesosilicate minerals that have the general formula of $(\text{Mg,Fe})_2\text{SiO}_4$. The chemical composition of an olivine specimen falls between forsterite (Mg_2SiO_4) and fayalite (Fe_2SiO_4) [4]. The dominated Fe^{2+} doublets with IS values of ~ 1.1 – 1.24 and $\text{QS} \sim 2$ mm/s (Table 1) can be assigned to augite

Table 1. Parameters for the ^{57}Fe Mössbauer spectra (shown in Fig. 1) for investigated basalt samples. Isomer shift values are given relative to the α -Fe standard

Sample	Line no.	χ^2	IS [mm/s]	QS [mm/s]	B [T]	Γ^* [mm/s]	Valency	A^{**} [%]	Mineral
Wilcza Góra (WG)	1	1.1	1.11	2.90	38	0.17	Fe^{2+}	14	Olivine
	2		1.10	2.02		0.35	Fe^{2+}	22	Augite
	3		0.36	0.85		0.32	Fe^{3+}	26	Hornblende
	4		0.44	-0.07		1.05	Fe^{3+}	38	Goethite
Małe Organy Myśluborskie (MOM)	1	1.8	1.14	2.85	38	0.18	Fe^{2+}	14	Olivine
	2		1.14	2.00		0.41	Fe^{2+}	21	Augite
	3		0.43	0.88		0.32	Fe^{3+}	17	Hornblende
	4		0.50	-0.04		1.00	Fe^{3+}	48	Goethite
Wleński Gródek near Klecza (WGK)	1	1.6	1.25	3.00		0.16	Fe^{2+}	9	Olivine
	2		1.24	2.08		0.40	Fe^{2+}	51	Augite
	3		0.47	0.88		0.38	Fe^{3+}	40	Hornblende
Czartowska Skała (CZS)	1	1.4	1.24	2.50	37	0.20	Fe^{2+}	9	Olivine
	2		1.09	1.98		0.33	Fe^{2+}	22	Augite
	3		0.34	0.99		0.30	Fe^{3+}	34	Hornblende
	4		0.42	-0.05		1.00	Fe^{3+}	35	Goethite

* Half width at half maximum. ** Relative contribution.




Fig. 2. Fe³⁺/Fe²⁺ ratio for investigated basalts.

(Al,Ca,Fe,Mg,Ti)₂(Al,Si)₂O₆. Augite is an essential mineral in basalts and belongs to the pyroxenes [2, 4]. Both olivine and augite were identified in all of the basalt samples. It is worth noting that similar components and contributions have been recorded in the spectrum of basalt from Wilcza Góra in the only paper dealing with Mössbauer spectroscopy in basalts from the Kaczawa Mountains [5].

According to the hyperfine parameters, Fe³⁺ component (doublet no. 3), which is also identified in each basalt sample, most probably may be assigned to hornblende [3, 6]. Hornblende (Ca,K,Na)₂₋₃(Mg,Fe²⁺,Fe³⁺,Al)₅(Si,Al)₈O₂₂(OH)₂ represents the amphiboles that is an important group of inosilicate rock-forming minerals, which occur in igneous and metamorphic rocks [2, 4].

The visible difference in the spectra of basalts that were measured is connected with the magnetic fraction (no. 4, Fig. 1). A Zeeman sextet with magnetic field (*B*) of 38 T, was observed for WG, MOM and CZS samples. Based on the hyperfine parameters (Table 1), the magnetic phase can be unambiguously assigned to goethite – FeO(OH) [7]. Goethite is a common iron hydroxide in materials formed in the weathering environment. Consequently, the presence of goethite may be an indication of an early stage of the weathering process. On the other hand, goethite may be a result of hydrothermal processes during the formation of these basalts.

According to the hyperfine parameters listed in Table 1, a relatively high ratio of Fe³⁺/Fe²⁺ in the range of 1.8 to 2.2 is observed for WG, MOM and CZS basalts. It is due to the occurrence of goethite in these samples. If goethite is a result of the oxidation process, the high ratio of Fe³⁺/Fe²⁺ may reflect a stage of weathering. In WGK basalt, we have not observed goethite. Consequently, the Fe³⁺/Fe²⁺ ratio of 0.7 is lower than those calculated for other basalts (Table 1, Fig. 2).

Pillow lavas

The Mössbauer spectra of the two pillow lava samples are shown in Fig. 3. The hyperfine parameters from the fitting procedure are summarized in Table 2. As can be seen in Fig. 3, despite the fact that both samples represent the same type of rock they show




Fig. 3. ⁵⁷Fe Mössbauer spectra at room temperature of pillow lava samples. Solid dots – experimental data; thick solid line – fitted curve; thin solid lines – fitted doublets and sextet.

Table 2. Parameters for the ⁵⁷Fe Mössbauer spectra (shown in Fig. 3) for investigated pillow lava samples. Isomer shift values are given relative to the α-Fe standard

Sample	Line no.	χ ²	IS [mm/s]	QS [mm/s]	B [T]	Γ* [mm/s]	Valency	A** [%]	Mineral	
Wąwóz Mysliborski (WM)	1	1.2	1.11	2.60	0.19	0.19	Fe ²⁺	45	Chlorite	
	2		1.34	2.87			Fe ²⁺	15	Hornblende	
	3		0.23	1.45			Fe ³⁺	40	Chlorite	
Wleński Gródek near Klecza (WGK)	1	1.6	1.17	2.61	52	0.20	Fe ²⁺	38	Chlorite	
	2		0.40	1.97			0.24	Fe ³⁺	4	Ferrihydrite
	3		0.46	0.67			0.31	Fe ³⁺	11	Chlorite
	4		0.42	-0.11			0.19	Fe ³⁺	47	Hematite

* Half width at half maximum. ** Relative contribution.




Fig. 4. Fe³⁺/Fe²⁺ ratio for investigated pillow lavas.

significant differences. The Mössbauer spectrum of the pillow lava from Wleński Gródek near Klecza (Fig. 3) is dominated by the sextet which can be unambiguously identified as hematite – Fe₂O₃ (labelled with no. 4 in Table 2). Hematite may be regarded as a secondary mineral in this type of rock and can be formed by the weathering of primary iron-bearing minerals or regional metamorphism [1, 5]. The presence of ferrihydrite Fe₅(O₄H₃)₃ (doublet no. 2) and hematite in the sample may be a result of advanced weathering.

The spectrum of the pillow lava sample from Wąwóz Myśluborski is characterized by the presence of two Fe²⁺ doublets (Fig. 3) representing chlorite (no. 1) and hornblende (no. 2), and one Fe³⁺ doublet (no. 3) also associated with chlorite [6, 8]. Chlorite (Mg,Al,Fe²⁺)₁₂[(Si,Al)₈O₂₀](OH)₈, a micaceous greenish mineral, is common in low-grade schist or as the alteration product of pyroxenes or amphiboles [5]. Chlorite phase was also identified in the WGK sample (doublets no. 1 and 3).

Surprisingly, unlike the WGK basalt, the sample of WGK lava from nearly the same location (Wleński Gródek near Klecza) is characterized by Fe³⁺/Fe²⁺ ratio of 1.6 (Fig. 4). It is a result of the occurrence of hematite in this sample; its contribution covers about 47% of the spectrum. In the WM sample, we have not observed hematite phase and the Fe³⁺/Fe²⁺ ratio is 0.7 (Fig. 4).

Conclusions

Augite, olivine and hornblende phases were observed in all the Miocene basalts investigated. Using ⁵⁷Fe Mössbauer spectroscopy, magnetic phases associated with goethite and hematite were identified in basalts and pillow lava. The high ratio of Fe³⁺/Fe²⁺ determined by the Mössbauer measurements for three basalts is a result of the occurrence of goethite in these samples. The present work shows that ⁵⁷Fe Mössbauer spectroscopy is a useful method to identify iron-bearing minerals in igneous rocks. Such measurements of rock samples from other locations in the Sudetes are planned.

References

1. Kryza, R. (2008). *The Variscan Kaczawa Complex: fitness od Palaeozoic basin Variscan orogeny. Geo-educational potential of the Sudety Mts.* University of Wrocław, Poland. Available from: <http://geopark.org.pl/Rozne/geoeducationalpotential.pdf>.
2. Lapidus, D. F., Winstanley, I., MacDonald, J., & Burton, C. (2006). *Geology*. London: HarperCollins.
3. Stevens, J. G., Khasanov, A. M., Miller, J. W., Pollak, H., & Li, Z. (2005). *Mössbauer mineral handbook*. Asheville, NC, USA: Mössbauer Effect Data Center, University of North Carolina. Available from <https://www.mtholyoke.edu/courses/mdyar/data/Mineral-Handbook.pdf>.
4. Bolewski, A. (1982). *Mineralogia szczegółowa*. Warszawa: Wydawnictwo Geologiczne.
5. Komraus, J. L., Adamczyk, Z., Popiel, E. S., & Malczewski, D. (1996). Identification of ferruginous minerals in basalt from Kaczawa Mountains Region by Mössbauer spectroscopy. In: All-Polish Seminar on Mössbauer Spectroscopy, 27–29 May 1996, Lublin, Poland (pp. 51–56).
6. Hawthorne, F. C. (1988). Spectroscopic methods in mineralogy and geology. *Rev. Mineral.*, 18, 255–340. DOI: 10.1017/S0016756800022962.
7. Murad, E., & Cashion, J. (2004). *Mössbauer spectroscopy of environmental materials and their industrial utilization*. Boston, USA: Kluwer Academic Publishers.
8. Malczewski, D., & Popiel, S. (2008). ⁵⁷Fe Mössbauer study of the oxidation state of iron in stilpnomelane from granite pegmatites in Poland. *Am. Mineral.*, 93, 1404–1411. DOI: 10.2138/am.2008.2719.