



# Antigen receptor signalling: a distinctive role for the p110 $\delta$ isoform of PI3K

Klaus Okkenhaug<sup>1\*</sup>, Khaled Ali<sup>2,3</sup> and Bart Vanhaesebroeck<sup>2,3\*</sup>

<sup>1</sup>Laboratory of Lymphocyte Signalling and Development, Babraham Institute, Cambridge, UK, CB2 4AT

<sup>2</sup>Ludwig Institute for Cancer Research, London, UK, W1W 7BS

<sup>3</sup>Department of Biochemistry and Molecular Biology, University College London, London, UK, WC1E 6BT

**The activation of antigen receptors triggers two important signalling pathways originating from phosphatidylinositol(4,5)-bisphosphate [PtdIns(4,5)P<sub>2</sub>]. The first is phospholipase C $\gamma$  (PLC $\gamma$ )-mediated hydrolysis of PtdIns(4,5)P<sub>2</sub>, resulting in the activation of Ras, protein kinase C and Ca<sup>2+</sup> flux. This culminates in profound alterations in gene expression and effector-cell responses, including secretory granule exocytosis and cytokine production. By contrast, phosphoinositide 3-kinases (PI3Ks) phosphorylate PtdIns(4,5)P<sub>2</sub> to yield phosphatidylinositol (3,4,5)-trisphosphate, activating signalling pathways that overlap with PLC $\gamma$  or are PI3K-specific. Pathways that are PI3K-specific include Akt-mediated inactivation of Foxo transcription factors and transcription-independent regulation of glucose uptake and metabolism. The p110 $\delta$  isoform of PI3K is the main source of PI3K activity following antigen recognition by B cells, T cells and mast cells. Here, we review the roles of p110 $\delta$  in regulating antigen-dependent responses in these cell types.**

## Introduction

Leukocytes express a repertoire of receptors to recognize and bind to extracellular antigen. This binding can be direct, as in the case of the B-cell antigen receptor (BCR) and the T-cell antigen receptor (TCR), or indirect, namely through receptors that interact with the Fc portion of immunoglobulins (Ig). The latter include the high affinity receptor for IgE (termed Fc $\epsilon$ RI) on mast cells and basophils, and the Fc $\gamma$ R for IgG on phagocytes, NK cells and B cells. PI3K activity has been implicated in the signalling of all types of antigen receptors.

## PI3K isoforms

Mammals have eight isoforms of phosphoinositide 3-kinase (PI3K), divided into three classes [1]. Class IA PI3Ks signal downstream of tyrosine kinases and Ras [1]. Class IA p110 catalytic subunits (p110 $\alpha$ , p110 $\beta$  and p110 $\delta$ ) are constitutively bound to an SH2 domain-containing adaptor protein, of which there are five species in mammals (p85 $\alpha$ , p55 $\alpha$ , p50 $\alpha$ , p85 $\beta$  and p55 $\gamma$ ; often referred to as 'p85s'). None of the

distinct p85s has been shown to be enriched in leukocytes and, *in vitro*, each p85 can interact with each p110 species. The Src homology 2 (SH2) domains of p85s are thought to bind preferentially to tyrosine (Tyr)-based motifs known as Y<sub>p</sub>xxM (Y<sub>p</sub>, phosphoTyr; M, methionine; x, any amino acid).

The p110 $\gamma$  PI3K, which belongs to the class IB subset of PI3Ks, also phosphorylates phosphatidylinositol(4,5)-bisphosphate [PtdIns(4,5)P<sub>2</sub>] but is activated by G protein-coupled receptors (GPCRs) instead of through tyrosine kinases. p110 $\gamma$  is also regulated by Ras [2]. In immune cells, p110 $\gamma$  is activated mainly by chemokines and by adenosine [3–5]. Therefore, p110 $\gamma$  might generate a distinct pool of phosphatidylinositol (3,4,5)-trisphosphate [PtdIns(3,4,5)P<sub>3</sub>] in activated cells.

The class II and class III PI3Ks have not been implicated in immune signalling, and are not considered further here.

## PI3K isoforms and antigen receptor signalling

Of the three class IA PI3K catalytic isoforms (p110 $\alpha$ , p110 $\beta$  and p110 $\delta$ ), p110 $\delta$  seems to have evolved to regulate PI3K-dependent processes in immune cells, most probably in part related to its high expression in these cells compared with most other cell types. Lymphocyte and mast-cell antigen receptor-dependent PI3K signalling is compromised in mice in which p110 $\delta$  has been inactivated by gene deletion [6,7], point mutation (p110 $\delta$ <sup>D910A</sup>) [8] or small-molecule p110 $\delta$  inhibitors [9–11]. p110 $\delta$  seems to be less crucial in IgG-based antigen receptor (Fc $\gamma$ R)-mediated phagocytosis in macrophages, where the p110 $\beta$  isoform seems to be more important [12]. It should be noted that p110 $\alpha$  and p110 $\beta$  are also expressed in leukocytes, and together can contribute up to 50% of the total p85-associated PI3K activity in some leukocytes [8–10,14]. Therefore, although we emphasize the predominant role of p110 $\delta$  in antigen receptor signalling in this review, it is possible that roles for p110 $\alpha$  and p110 $\beta$  in this signalling context will be uncovered with the ongoing development of conditional knockout mice and selective inhibitors for these PI3K isoforms.


## Conversion of PtdIns(4,5)P<sub>2</sub> downstream of antigen receptors

Antigen receptor stimulation initiates the activation of Src and Syk family Tyr kinases, resulting in Tyr

Corresponding authors: Okkenhaug, K. ([klaus.okkenhaug@bbsrc.ac.uk](mailto:klaus.okkenhaug@bbsrc.ac.uk)); Vanhaesebroeck, B. ([bartvanh@ludwig.ucl.ac.uk](mailto:bartvanh@ludwig.ucl.ac.uk)).

\* K.O. and B.V. are consultants for PIRamed, UK.

Available online 5 January 2007



**Figure 1.** Metabolism of PtdIns(4,5) $P_2$  by PLC $\gamma$  and PI3K. PLC $\gamma$  hydrolyses PtdIns(4,5) $P_2$  to yield Ins(1,4,5) $P_3$  and DAG, both of which function as signalling molecules. Ins(1,4,5) $P_3$  stimulates the release of Ca $^{2+}$  from the ER into the cytosol, which triggers the nuclear translocation of NFAT. DAG binds to and activates RasGRP, which stimulates Ras and the Erk pathway, leading to AP-1-dependent transcription. Ras also binds to p110 and contributes to optimal PI3K activation. DAG binds to and activates PKC, which activates NF- $\kappa$ B through CARMA1, BCL10 and MALT1. By contrast, PI3K phosphorylates PtdIns(4,5) $P_2$  at position 3 to produce the membrane phosphoinositol lipid PtdIns(3,4,5) $P_3$ . PtdIns(3,4,5) $P_3$  functions as an anchor and cofactor for proteins with PtdIns(3,4,5) $P_3$ -binding PH domains such as Akt, Tec family kinases, and various GEFs and GAPs. Pdk1 is required to co-activate Akt. Akt phosphorylates and inactivates Foxo and GSK3. GSK3 can phosphorylate and inactivate NFAT. Akt stimulates mTOR through Tsc1 and Tsc2. Tec kinases can phosphorylate PLC $\gamma$  and contribute to its optimal activity. PI3K signalling is antagonised by the Pten phosphoinositide phosphatase, which removes the 3-phosphate, and the SHIP phosphatase, which removes the 5-phosphate. The role of PI(3,4) $P_2$ -binding proteins is still unknown. Although PLC $\gamma$  and PI3K generate mutually exclusive second messenger signalling molecules, several of the pathways activated by these second messengers interact, and the signals are further integrated by the cell to promote gene transcription, cell growth and differentiation. p110 $\delta$  seems to be the principal PI3K isoform in the context of antigen receptor signaling; however, p110 $\alpha$  and p110 $\beta$  are also expressed in lymphocytes but their roles in antigen receptor signalling are unknown. Abbreviations: BCL10, B cell lymphoma 10; CARMA1, caspase recruitment domain (CARD)-containing membrane-associated guanylate kinase (MAGUK) protein 1; IP $_3$ , inositol(1,4,5)-trisphosphate; MALT1, mucosa-associated lymphoid tissue lymphoma translocation protein 1; PI, phosphatidylinositol.

phosphorylation of adaptor proteins and the activation of two PtdIns(4,5) $P_2$ -based signalling pathways (Figure 1).

The first of these is mediated by phospholipase C (PLC) $\gamma$ , which hydrolyses PtdIns(4,5) $P_2$  to generate diacylglycerol (DAG) and inositol(1,4,5)-trisphosphate [Ins(1,4,5) $P_3$ ] (Figure 1). The water-soluble Ins(1,4,5) $P_3$  triggers a biphasic Ca $^{2+}$  response, initially by inducing Ca $^{2+}$  release from the endoplasmic reticulum, followed by Ca $^{2+}$ -dependent opening of plasma membrane channels to enable sustained Ca $^{2+}$  influx and the nuclear translocation of NFAT (nuclear factor of activated T cells) transcription factors [15]. DAG activates Ras guanyl nucleotide-releasing proteins (RasGRPs, which bind to Ca $^{2+}$  and DAG) and isoforms of protein kinase C (PKC), which initiate the Ras–Erk–AP-1 and nuclear factor (NF)- $\kappa$ B signalling pathways, respectively [16–20]. Hence, the hydrolysis of PtdIns(4,5) $P_2$  by PLC $\gamma$  is sufficient to initiate the activation of three transcription factor families (Figure 1) that coordinately regulate the expression of a vast number of genes involved in cytokine production, cell division and differentiation [21,22]. Antigen receptor cross-linking also activates PLC $\gamma$ -dependent pathways involved in promoting the release of preformed mast-cell secretory granules, which promote the early symptoms of allergic hypersensitivity reactions [19].

Class I PI3Ks mediate an alternative conversion of PtdIns(4,5) $P_2$ , namely by phosphorylating this lipid to PtdIns(3,4,5) $P_3$  (Figure 1). In contrast to its precursor, PtdIns(3,4,5) $P_3$  is resistant to PLC-mediated hydrolysis and, instead, signals at the plasma membrane by functioning as docking sites for pleckstrin homology (PH) domains that are present in several proteins [1]. These PtdIns(3,4,5) $P_3$  targets include protein kinases (such as Pdk1, Akt and Tec kinases), adaptor proteins (such as Gab2), and GTPase activating proteins (GAPs) and guanine nucleotide exchange factors (GEFs) for small GTPases (such as P-Rex, ARAP, SWAT-70, IBP and Vav) [1,23,24]. PtdIns(3,4,5) $P_3$  binding induces the rapid recruitment of these proteins to the membrane in response to PI3K activation and/or alteration of their conformation or activity [25,26]. This abundance of downstream targets link PI3K to its well-established roles in cell cycle progression, growth, prevention of apoptosis, cell migration, differentiation and secretory granule exocytosis.

PtdIns(3,4,5) $P_3$  is a substrate for lipid phosphatases, amongst which the 3-phosphatase Pten and the 5-phosphatase SHIP are most widely studied and have important roles in antagonising PI3K signalling [27,28]. SHIP converts PtdIns(3,4,5) $P_3$  to PtdIns(3,4) $P_2$ , which can bind to a limited set of PH domains, including those of tandem PH

domain-containing protein (TAPP)1 and TAPP2 [29] (Figure 1).

### PI3K coupling to the antigen receptor

The details of how PI3Ks are linked to antigen receptor-associated signalling complexes are still vague [30–32]. What is clear is that PtdIns(3,4,5) $P_3$  accumulation occurs extremely rapidly following antigen recognition, which suggests that the association between the antigen receptor signalling complexes and PI3K is tightly coupled to the initial tyrosine kinase signals [29,33–38]. Several of the adaptor molecules in the antigen receptor complex contain canonical YxxM motifs, although the role of some of these, including B-cell adaptor protein (BCAP) and T-cell receptor interacting molecule (TRIM), is still unclear [39,40] (Figure 2). An intriguing recent study suggested that p85 can bind to Syk and zeta-chain associated protein kinase of 70 kDa (ZAP-70) directly through non-canonical Tyr-based motifs [41]. There is also evidence that p85 can bind to SH2 domain-containing leukocyte phosphoprotein of 76 kD (SLP76), again through noncanonical motifs [42]. The SH3 domains and proline-rich regions present in p85s also offer scope for phosphorylation-independent binding to signalling proteins. If these interactions can be confirmed under physiologically relevant conditions, then we might need to cast the net wider in considering potential players in the recruitment of p85–p110 heterodimers to antigen receptor signalling complexes.

Given that p85 species seem to have no binding preference for specific p110 PI3K isoforms, it is anticipated that this type of recruitment would not favour p110 $\delta$  over p110 $\alpha$  and p110 $\beta$  for association with the antigen receptor. However, in addition to binding to p85, the p110 subunits can also bind to Ras-GTP [1]. Ras signalling is important in antigen receptor signalling [16]. Evidence is accumulating that each p110 isoform has a distinct binding capacity to Ras, or shows a binding preference for specific Ras isoforms (reviewed in Ref. [1]). In one study, p110 $\alpha$  was found to become activated by most Ras isoforms, whereas p110 $\delta$  became activated selectively by R-Ras and Tc21; p110 $\beta$  did not become activated by Ras at all [43]. Hence, the differential usage of Ras isoforms could contribute to selective recruitment and/or activation of class IA PI3K isoforms. Therefore, the potential role of Ras in regulating the activity of different PI3K isoforms in the context of antigen receptor signalling warrants further investigation.

### p110 $\delta$ in B-cell development and function


p110 $\delta$  deficiency<sup>†</sup> does not have a major impact on the early development of B cells in the bone marrow [6–8]. By contrast, the development of mature B cells in the spleen and in pleural cavities is strongly affected. Thus, the number of follicular B2 cells is reduced to <50% of normal numbers. In addition, the development of peritoneal B1 cells and marginal-zone B cells is almost completely blocked [30,31,44].

The *in vitro* proliferation of B cells triggered by antibody-mediated crosslinking (through anti-IgM) of the BCR crucially depends on p110 $\delta$  activity [6–9]. Deletion or inactivation of p110 $\delta$  or p85 $\alpha$  largely ablates BCR-induced phosphorylation of Akt, Foxo and protein kinase D, and results in reduced Ca<sup>2+</sup> flux, impaired cell cycle progression and reduced glucose metabolism [9,45–50]. Interleukin (IL)-4-dependent survival is also compromised in the absence of p110 $\delta$  activity [9]. CD40 and lipopolysaccharide signalling is less dependent on PI3K signalling, so activated T cells and selected pathogens might still stimulate B-cell responses even in the context of strongly attenuated BCR signalling [8,51]. PI3K had initially been suggested to be part of a BCR-associated signalosome based on the similar phenotypes of Btk, p85 $\alpha$ , p110 $\delta$ , and PLC $\gamma$  knockout mice. In this model, the principal role for PI3K would be to promote PLC $\gamma$  activity [52]. Although the role of PI3K in regulating PLC $\gamma$  is well established, recent evidence shows that PLC $\gamma$  and PI3K pathways also function in parallel (Figure 1). Thus, p85 $\alpha$ –Btk and p85 $\alpha$ –PLC- $\gamma$ 2 double-knockout mice show much more dramatic phenotypes than any of the single knockout mice and provide evidence for PLC $\gamma$ -independent functions for PI3K [53,54].

PI3K activation in B cells can be enhanced by the coordinated engagement of the BCR and its co-receptor CD19, which contains YxxM recruitment motifs for PI3K [55]. This enables added sensitivity of the B cells to antigens that are coated with complement. In this context, Vav is required for optimal PI3K responses, perhaps reflecting a role for Vav in coordinating crosstalk between these receptors by assembling larger signalling complexes [56,57]. The B-cell phenotypes observed in p85 $\alpha$ -knockout and p110 $\delta$ -deficient mice largely overlap with those observed in CD19-deficient mice. The key role for PI3K in the context of CD19 function is further evidenced by the failure to rescue a normal B-cell phenotype in CD19 knockout mice by the transgenic expression of a CD19 mutant that cannot bind to PI3K [58]. Moreover, Pten-deficiency can partially revert the effect of CD19 deficiency, presumably by lowering the signalling threshold required to initiate PI3K signalling [59]. Co-ligation of the BCR and Fc $\gamma$ RIIB, a low affinity receptor for IgG, also forms the basis for modulating PtdIns(3,4,5) $P_3$  signals downstream of the BCR through the recruitment of SHIP, which converts PtdIns(3,4,5) $P_3$  into PtdIns(3,4) $P_2$ , thereby inhibiting BCR responses [60].

Antigen challenge triggers a primary humoral immune response that is characterized by clonal expansion and B-cell differentiation into IgM-secreting plasma cells. Alternatively, B cells can be recruited into the T-cell-rich areas of the splenic follicles, where the Ig loci undergo class-switch recombination (CSR) and somatic hypermutation leading to the production of higher affinity antibodies of the IgG or IgE subclasses. Immunized p110 $\delta$ -deficient mice show reduced germinal centre (GC) formation and impaired T-cell-dependent and T-cell-independent immune responses, suggesting a positive role for p110 $\delta$  in the GC reactions [6,8]. However, there is also evidence for a negative role of p110 $\delta$  in the GC reaction: Pten-deficient B cells fail to induce activation-induced cytidine deaminase, an essential regulator of CSR

<sup>†</sup> We have previously commented on the relative merits of different gene-targeting strategies [13] and in this review, we will refer collectively to p110 $\delta$ <sup>-/-</sup> and p110 $\delta$ <sup>D910A</sup> mice as p110 $\delta$ -deficient.



**Figure 2.** Antigen receptor complexes and p110 $\delta$  antigen receptor signalling in different cell types has key commonalities, the most crucial of which is the phosphorylation of ITAM motifs found in proteins that are noncovalently associated with the polypeptides that bind to the antigen or antibody–antigen complexes. Src-family kinases phosphorylate these ITAM motifs, thus providing docking sites for Syk kinases. Syk kinases phosphorylate PLC $\gamma$ , resulting in its activation, in addition to the recruitment of various cellular and membrane-bound adaptor proteins, such as LAT, Gab2, SLP-76 (in T cells and mast cells) and SLP-65 (in B cells) that nucleate larger signalling complexes [18,19]. The upstream activators of PI3K in the context of antigen receptor signalling have not been definitively defined, and possible links with the SH2 domains of p85 are indicated by dashed arrows. Phosphorylation-independent interactions of PI3K with upstream signalling molecules are not shown. Note that the antigens are shown as monomers for illustrative purposes. In reality, dimers or oligomers of the ligands and receptors are required to trigger the signalling cascades shown. **(a)** BCR signalling. Lyn-dependent phosphorylation of Ig $\alpha$  and Ig $\beta$  ITAM motifs results in the recruitment of Syk and the phosphorylation of SLP-65. Several proteins in the BCR receptor complex have been implicated in binding to p85. BCAP is a transmembrane adaptor protein with YxxM motifs which becomes phosphorylated by Syk upon BCR activation. BCAP has been shown to regulate PI3K signalling in DT40 chicken cells but was found to be not required for PI3K signalling in primary mouse B cells [40,87]. Similarly, Vav had been proposed to lie upstream of PI3K signalling [56]; however, BCR crosslinking of primary Vav-deficient B cells results in normal Akt phosphorylation (although Akt phosphorylation in response to BCR and CD19 coligation was Vav-dependent) [51]. Other candidates that link PI3K to the activated BCR include Gab1, non T-cell activation linker (NTAL) and LAT. However, many of these interactions have only been identified in cell lines, and in several cases, their roles in PI3K signalling downstream of the BCR have not been confirmed in mice [88,89]. **(b)** TCR signalling. Lck and Fyn phosphorylate ITAM motifs resulting in the recruitment of ZAP-70, which

[61]. Moreover, Akt phosphorylation and the inactivation of Foxo proteins by p110 $\delta$  was shown to suppress CSR, favouring the formation of IgM-secreting antibody effectors [62]. How these results correlate with the diminished IgG-mediated responses in p110 $\delta$ -deficient mice is unclear. One possibility is that p110 $\delta$ -deficient T cells cannot support full GC reactions. In addition, an earlier developmental lesion in p110 $\delta$ -deficient B cells might reduce dramatically the number of B cells poised for participation in the GC reaction. Therefore, although p110 $\delta$  has a positive role in mitogenic signalling through the BCR, p110 $\delta$  might also negatively regulate differentiation programmes that lead to CSR and secondary immune responses.

### p110 $\delta$ in T-cell development and function

T-cell development in the thymus progresses through three checkpoints:  $\beta$ -selection, where CD4<sup>-</sup>CD8<sup>-</sup> double-negative T cells are examined for pre-TCR $\beta$  expression; positive selection, where CD4<sup>+</sup>CD8<sup>+</sup> double-positive TCR $\alpha\beta$ <sup>+</sup> T cells are selected to become CD4<sup>+</sup> or CD8<sup>+</sup> single-positive T cells; and negative selection, where autoreactive T cells are eliminated.

PtdIns(3,4,5) $P_3$  signalling is both sufficient and necessary for  $\beta$ -selection. Lck-Cre-mediated conditional deletion of Pten in the T-cell lineage enables the development of double-positive thymocytes in the absence of pre-TCR expression [63,64]. Similarly, in the presence of artificially high expression of its ligand CD86, CD28 can promote the development of double-positive T cells in the absence of pre-TCR, but only if the PI3K-binding motif is intact [65].

$\beta$ -selection seems normal in p110 $\delta$ -deficient mice [8,66], suggesting the involvement of other class I PI3Ks. One of these might be p110 $\gamma$ , as T cells lacking this PI3K have a partial defect at this selection stage, resulting in a reduced number of double-positive T cells [4]. p110 $\gamma$ <sup>-/-</sup> p110 $\delta$ <sup>-/-</sup> double-knockout mice have a profound block at the pre-TCR selection step, with a dramatic reduction of double-positive cells [66,67]. This is a surprising observation, given that GPCRs have not previously been shown to be required at this selection step. It is presently unclear whether there is a chemokine receptor or another GPCR that functions in concert with the pre-TCR at this stage, or whether p110 $\gamma$  can be activated by non-GPCR receptors. Positive selection is unaffected in p110 $\delta$ -deficient thymocytes; however, negative selection of autoreactive T cells is partially impaired [68].

In peripheral T cells, PI3Ks can be activated by the TCR, by costimulatory receptors such as CD28 and ICOS, and by receptors for cytokines and chemokines [30,69]. p110 $\delta$  seems to be the main PI3K isoform that generates PtdIns(3,4,5) $P_3$  downstream of the TCR and CD28, although p110 $\alpha$  and/or p110 $\beta$  are also likely to contribute [8,14,70]. In

the immune synapse, PtdIns(3,4,5) $P_3$  accumulation is observed as one of the earliest traceable signals and is sustained for hours, as long as the conjugate formation between antigen-presenting cells and T cells is maintained [33–35]. The relative contributions made by the TCR and CD28 to PI3K activity in the synapse are still unknown; however, synapse accumulation can occur under conditions where CD28 costimulation is absent [33,37]. A key effect of PI3K signalling in T cells is the activation of Akt, which phosphorylates Foxo transcription factors that then become excluded from the nucleus [14,38,71]. In addition, Akt can contribute to glycolysis and protein synthesis; however, PI3K-independent pathways can also contribute to at least some of these effects [38,72–74].

In T cells with inactive p110 $\delta$ , defects in proliferation and cytokine secretion were most clearly revealed when TCR-transgenic T cells were stimulated with cognate antigen [8,14]. In particular, T-helper (Th)1 and Th2 cytokine production was reduced dramatically [14]. As a consequence of reduced Th2 responses, p110-deficient mice were protected from experimentally induced airway inflammation [75]. *In vitro*, the reduction in Th1 and Th2 cytokine production could not be rescued by providing an excess of exogenous cytokines and was still defective among T cells that had undergone several rounds of division [14]. Therefore, there seems to be a block in the genetic programme that enables T cells to open the IL-4 or interferon (IFN) $\gamma$  gene loci. We hypothesize that this block reflects the capacity of PI3Ks to relieve T cells from the blocks imposed by transcription factors of the Foxo family. Unless the suppression by Foxo transcription factors is lifted by PI3K activation, the cell is unable to differentiate further. In this scenario, PI3Ks and, more specifically, p110 $\delta$ , function in parallel to the canonical TCR signalling pathway initiated by PLC $\gamma$  (Figure 1). That is, initial activation events that lead to clonal expansion and IL-2 secretion do occur but further differentiation is blocked. Consistent with this notion, mice that lack Foxo3a, one of the Foxo transcription factors, suffer from exaggerated Th1 and Th2 responses and autoimmune syndromes [76].

A third lineage of CD4<sup>+</sup> T cells, referred to as regulatory T (Treg) cells, restrict the expansion and function of Th cells [77]. Mice that lack Treg cells die young as a consequence of T cell-mediated multi-organ destruction. Mice that have partial defects in Treg-cell development tend to develop colitis as Treg cells have a key role in suppressing immune responses against the gut flora [78]. p110 $\delta$ -deficient mice also show subclinical signs of colitis as detected by histological examination [8]. Moreover, p110 $\delta$ -deficient Treg cells show attenuated capacity to suppress Th cells *in vitro* and fail to protect against experimentally induced colitis *in vivo* [68]. Interestingly, p85 $\beta$  knockout mice with T-cell-specific deletion of p85 $\alpha$  have

(Figure 2 Legend Continued) phosphorylates LAT. TRIM is a transmembrane adaptor protein that associates with the TCR and becomes phosphorylated on YxxM motifs. However, TRIM knockout mice show enhanced instead of impaired Akt phosphorylation [39]. Vav has an important role in regulating Akt and Foxo phosphorylation. Although the exact biochemical link between Vav and PI3K is not clearly defined, it may reflect a more general role for Vav in assembling LAT complexes [71,90,91]. CD28 can bind to the SH2 domains of p85 directly; however, CD28 can provide potent costimulatory signals independently of its association with PI3K [30]. (c) Fc $\epsilon$ R1 signalling. Lyn phosphorylates ITAM motifs in the  $\beta$  and  $\gamma$  chains of the Fc $\epsilon$ R1 resulting in the recruitment of Syk, which phosphorylates GAB2 and LAT, inducing two parallel pathways through GAB2-PI3K and LAT-PLC $\gamma$ . The link shown between NTAL and GAB2 is hypothetical. c-Kit can bind to p85 directly and can potentiate Fc $\epsilon$ R1-stimulated degradation. (d) Chemokine receptor signalling. The p110 $\gamma$  heterodimer binds to the G $\beta\gamma$  subunit released from G $\alpha$  following GPCR stimulation with agonists such as chemokines. Despite the potent activation of p110 $\gamma$  by chemokines, p110 $\gamma$  seems to have a minor role in promoting lymphocyte chemotaxis [3]. Instead, p110 $\gamma$  might promote the survival of developing thymocytes and memory T cells [92].

reduced proportions of Treg cells, show signs of colitis and develop an autoimmune disease that is reminiscent of Sjögren's syndrome [79]. The latter experiments also demonstrated that although p85 $\alpha$ -deficiency is sufficient to suppress PI3K signalling in B cells, both p85 $\alpha$  and p85 $\beta$  need to be deleted to uncover a PI3K-deficient phenotype in T cells [93].

### p110 $\delta$ in mast cell development and function

Mast cells are amongst a select group of cells (including basophils and eosinophils) that express the Fc $\epsilon$ RI (Figure 2). Antigen-specific IgE from the plasma binds to Fc $\epsilon$ RI with high affinity, enabling the mast cells to participate in the adaptive immune response. Antigen-induced aggregation of Fc $\epsilon$ RI-bound IgE activates a series of intracellular signalling events, culminating in secretory-granule exocytosis and the release of pro-inflammatory mediators that promote the allergic cascade [19].

The activation of Fc $\epsilon$ RI triggers a tyrosine kinase cascade involving Lyn, Fyn and Syk, resulting in the activation of the linker for activation of T cells (LAT)–PLC $\gamma$  and GRB2-associated binding protein 2 (GAB2)–PI3K pathways (Figure 2) [19]. These pathways recruit PI3K and interact with effectors that drive Ca<sup>2+</sup> mobilization and PKC activation, which are both a prerequisite for mast-cell exocytosis [80].

Pan-PI3K inhibitors can attenuate IgE–antigen-dependent secretory-granule exocytosis severely [81]. Conversely, genetic interference with SHIP and Pten enhances Fc $\epsilon$ RI responses [82,83]. Akt might control IL-2 and TNF- $\alpha$  production by regulating NF- $\kappa$ B [84]. Two other potential PI3K effectors downstream of Fc $\epsilon$ RI in mast cells are Btk and phospholipase D1 (PLD<sub>1</sub>), which both influence Ca<sup>2+</sup> mobilization and PKC activation strongly. Btk enhances the activities of PLC $\gamma$  and PKC $\beta$ 1 (the latter directly, or indirectly through the regulation of PLC $\gamma$ ) [19]. Deletion or mutation of the Btk PH domain (as observed in Xid mice) results in reduced Fc $\epsilon$ RI-dependent Ca<sup>2+</sup> signalling, degranulation and cytokine release [19]. PLD<sub>1</sub> contains a PtdIns(3,4,5)P<sub>3</sub>-interacting PX domain [85], and its activation leads to the production of phosphatidic acid, which is metabolized further into an important source of DAG that can activate PKC and might provide an additional link between the PI3K and the PLC $\gamma$  pathway in augmenting the PKC activation [86].

Most PI3K-dependent activity downstream of the Fc $\epsilon$ RI is mediated by p110 $\delta$ , and antigen receptor-induced Akt activation is almost completely eliminated following p110 $\delta$  inactivation [10]. Mast cells derived from p110 $\delta$ -deficient mice, or wild-type mast cells treated with a p110 $\delta$ -selective inhibitor, have a substantial reduction in IgE–antigen-dependent exocytosis and release of the pro-inflammatory cytokines TNF- $\alpha$  and IL-6 [10].

Another important pathway for regulating mast-cell proliferation, migration and adhesion is through the c-kit receptor Tyr kinase, which is activated by the SCF ligand. c-kit has a YxxM motif that can recruit p85–p110 complexes. Interestingly, similar to the Fc $\epsilon$ RI, PI3K-dependent responses downstream of the c-kit receptor are almost completely p110 $\delta$ -dependent [10]. Allergic responses occur in an SCF-rich environment, which

potentiates Fc $\epsilon$ RI responses. The observation that both the Fc $\epsilon$ RI and its key modulating receptor c-kit depend heavily on p110 $\delta$  activity is an important rationale for developing p110 $\delta$  inhibitors for allergy indications [10]. Indeed, inactivating p110 $\delta$  in mice results in a substantial reduction in the allergic response [10,11]. However, a residual PI3K-dependent allergic response is observed in p110 $\delta$ -deficient mast cells and mice [10]. The identity of this PI3K(s) is still elusive and could either be another class 1A PI3K isoform or it could be p110 $\gamma$ . Indeed, GPCR-coupled p110 $\gamma$  is thought to operate downstream of activated Fc $\epsilon$ RI through a GPCR-dependent autocrine or paracrine amplification mechanism [5].

### Concluding remarks

p110 $\delta$  seems to have evolved as the main source of PtdIns(3,4,5)P<sub>3</sub> production following antigen recognition by B cells, T cells and mast cells. Although the p110 $\alpha$  and p110 $\beta$  isoforms might also contribute to antigen receptor signalling, their contributions seem less important than those of p110 $\delta$ , for reasons that are not understood. In this respect, the role of p110 $\delta$  is unique: its attenuation affects antigen receptor signalling in B cells, T cells and mast cells, yet p110 $\delta$  is dispensable for the function of most tissues and organs. No other kinase associated with the antigen receptor signalling machinery shares this profile. Instead, the expression of Src, Syk and Tec family kinase isoforms differs between B cells, T cells and mast cells. Hence, p110 $\delta$  presents a unique opportunity to modulate antigen receptor signalling using small-molecule inhibitors.

It will also be important to investigate the contribution of each of the four class I PI3K isoforms to signalling by other immune receptors. This should be facilitated through the availability of a range of PI3K gene-targeted mice and PI3K isoform-selective inhibitors that are now becoming available. These studies should pave the way towards the clinical development of small-molecule inhibitors against PI3K isoforms, in particular p110 $\delta$ , which could alleviate harmful immune responses against self antigens, transplantation antigens and innocuous foreign antigens in allergy.

### Acknowledgements

We thank Elena Vigorito and Robert Salmond for critical reading of the manuscript. K.O. is supported by a BBSRC David Phillips Fellowship. K.A. is supported by the Biotechnology and Biological Science Research Council, UK (grant BB/C505659/1). Work in B.V.'s laboratory is supported by the Ludwig Institute for Cancer Research and the European Union MAIN consortium (FP6–502935).

### References

- 1 Vanhaesebroeck, B. *et al.* (2001) Synthesis and function of 3-phosphorylated inositol lipids. *Annu. Rev. Biochem.* 70, 535–602
- 2 Suire, S. *et al.* (2006) G $\beta$  $\gamma$ s and the Ras binding domain of p110 $\gamma$  are both important regulators of PI3K $\gamma$  signalling in neutrophils. *Nat. Cell Biol.* 8, 1303–1309
- 3 Ward, S.G. (2006) T lymphocytes on the move: chemokines, PI 3-kinase and beyond. *Trends Immunol.* 27, 80–87
- 4 Sasaki, T. *et al.* (2000) Function of PI3K $\gamma$  in thymocyte development, T cell activation, and neutrophil migration. *Science* 287, 1040–1046
- 5 Laffargue, M. *et al.* (2002) Phosphoinositide 3-kinase  $\gamma$  is an essential amplifier of mast cell function. *Immunity* 16, 441–451

- 6 Clayton, E. *et al.* (2002) A crucial role for the p110 $\delta$  subunit of phosphatidylinositol 3-kinase in B cell development and activation. *J. Exp. Med.* 196, 753–763
- 7 Jou, S.T. *et al.* (2002) Essential, nonredundant role for the phosphoinositide 3-kinase p110 $\delta$  in signaling by the B-cell receptor complex. *Mol. Cell. Biol.* 22, 8580–8591
- 8 Okkenhaug, K. *et al.* (2002) Impaired B and T cell antigen receptor signaling in p110 $\delta$  PI 3-kinase mutant mice. *Science* 297, 1031–1034
- 9 Bilancio, A. *et al.* (2006) Key role of the p110 $\delta$  isoform of PI3K in B-cell antigen and IL-4 receptor signaling: comparative analysis of genetic and pharmacologic interference with p110 $\delta$  function in B cells. *Blood* 107, 642–650
- 10 Ali, K. *et al.* (2004) Essential role for the p110 $\delta$  phosphoinositide 3-kinase in the allergic response. *Nature* 431, 1007–1011
- 11 Lee, K.S. *et al.* (2006) Inhibition of phosphoinositide 3-kinase  $\delta$  attenuates allergic airway inflammation and hyperresponsiveness in murine asthma model. *FASEB J.* 20, 455–465
- 12 Leverrier, Y. *et al.* (2003) Class I phosphoinositide 3-kinase p110 $\beta$  is required for apoptotic cell and Fc $\gamma$  receptor-mediated phagocytosis by macrophages. *J. Biol. Chem.* 278, 38437–38442
- 13 Vanhaesebroeck, B. *et al.* (2005) Signalling by PI3K isoforms: insights from gene-targeted mice. *Trends Biochem. Sci.* 30, 194–204
- 14 Okkenhaug, K. *et al.* (2006) The p110 $\delta$  isoform of phosphoinositide 3-kinase controls clonal expansion and differentiation of Th cells. *J. Immunol.* 177, 5122–5128
- 15 Gallo, E.M. *et al.* (2006) Lymphocyte calcium signaling from membrane to nucleus. *Nat. Immunol.* 7, 25–32
- 16 Samelson, L.E. (2002) Signal transduction mediated by the T cell antigen receptor: the role of adapter proteins. *Annu. Rev. Immunol.* 20, 371–394
- 17 Niiro, H. and Clark, E.A. (2002) Decision making in the immune system: regulation of B-cell fate by antigen-receptor signals. *Nat. Rev. Immunol.* 2, 945–956
- 18 Koretzky, G.A. *et al.* (2006) SLP76 and SLP65: complex regulation of signalling in lymphocytes and beyond. *Nat. Rev. Immunol.* 6, 67–78
- 19 Gilfillan, A.M. and Tkaczyk, C. (2006) Integrated signalling pathways for mast-cell activation. *Nat. Rev. Immunol.* 6, 218–230
- 20 Thome, M. (2004) CARMA1, BCL-10 and MALT1 in lymphocyte development and activation. *Nat. Rev. Immunol.* 4, 348–359
- 21 Diehn, M. *et al.* (2002) Genomic expression programs and the integration of the CD28 costimulatory signal in T cell activation. *Proc. Natl. Acad. Sci. U. S. A.* 99, 11796–11801
- 22 Riley, J.L. *et al.* (2002) Modulation of TCR-induced transcriptional profiles by ligation of CD28, ICOS, and CTLA-4 receptors. *Proc. Natl. Acad. Sci. U. S. A.* 99, 11790–11795
- 23 Welch, H.C. *et al.* (2003) Phosphoinositide 3-kinase-dependent activation of Rac. *FEBS Lett.* 546, 93–97
- 24 Vanhaesebroeck, B. and Alessi, D.R. (2000) The PI3K–PDK1 connection: more than just a road to PKB. *Biochem. J.* 346, 561–576
- 25 Milburn, C.C. *et al.* (2003) Binding of phosphatidylinositol 3,4,5-trisphosphate to the pleckstrin homology domain of protein kinase B induces a conformational change. *Biochem. J.* 375, 531–538
- 26 Lemmon, M.A. and Ferguson, K.M. (2000) Signal-dependent membrane targeting by pleckstrin homology (PH) domains. *Biochem. J.* 350, 1–18
- 27 Krystal, G. (2000) Lipid phosphatases in the immune system. *Semin. Immunol.* 12, 397–403
- 28 Sasaki, T. *et al.* (2002) Phosphoinositide 3-kinases in immunity: lessons from knockout mice. *J. Biochem. (Tokyo)* 131, 495–501
- 29 Krahn, A.K. *et al.* (2004) Two distinct waves of membrane-proximal B cell antigen receptor signaling differentially regulated by Src homology 2-containing inositol polyphosphate 5-phosphatase. *J. Immunol.* 172, 331–339
- 30 Okkenhaug, K. and Vanhaesebroeck, B. (2003) PI3K in lymphocyte development, differentiation and activation. *Nat. Rev. Immunol.* 3, 317–330
- 31 Deane, J.A. and Fruman, D.A. (2004) Phosphoinositide 3-kinase: diverse roles in immune cell activation. *Annu. Rev. Immunol.* 22, 563–598
- 32 Koyasu, S. (2003) The role of PI3K in immune cells. *Nat. Immunol.* 4, 313–319
- 33 Harriague, J. and Bismuth, G. (2002) Imaging antigen-induced PI3K activation in T cells. *Nat. Immunol.* 3, 1090–1096
- 34 Costello, P.S. *et al.* (2002) Sustained and dynamic inositol lipid metabolism inside and outside the immunological synapse. *Nat. Immunol.* 3, 1082–1089
- 35 Huppa, J.B. *et al.* (2003) Continuous T cell receptor signaling required for synapse maintenance and full effector potential. *Nat. Immunol.* 4, 749–755
- 36 Astoul, E. *et al.* (1999) The dynamics of protein kinase B regulation during B cell antigen receptor engagement. *J. Cell Biol.* 145, 1511–1520
- 37 Kane, L.P. *et al.* (2004) A proline-rich motif in the C terminus of Akt contributes to its localization in the immunological synapse. *J. Immunol.* 172, 5441–5449
- 38 Fabre, S. *et al.* (2005) Stable activation of phosphatidylinositol 3-kinase in the T cell immunological synapse stimulates Akt signaling to FoxO1 nuclear exclusion and cell growth control. *J. Immunol.* 174, 4161–4171
- 39 Kolsch, U. *et al.* (2006) Normal T-cell development and immune functions in TRIM-deficient mice. *Mol. Cell. Biol.* 26, 3639–3648
- 40 Yamazaki, T. *et al.* (2002) Essential immunoregulatory role for BCAP in B cell development and function. *J. Exp. Med.* 195, 535–545
- 41 Moon, K.D. *et al.* (2005) Molecular basis for a direct interaction between the Syk protein-tyrosine kinase and phosphoinositide 3-kinase. *J. Biol. Chem.* 280, 1543–1551
- 42 Shim, E.K. *et al.* (2004) Association of the Src homology 2 domain-containing leukocyte phosphoprotein of 76 kD (SLP-76) with the p85 subunit of phosphoinositide 3-kinase. *FEBS Lett.* 575, 35–40
- 43 Rodriguez-Viciania, P. *et al.* (2004) Signaling specificity by Ras family GTPases is determined by the full spectrum of effectors they regulate. *Mol. Cell. Biol.* 24, 4943–4954
- 44 Donahue, A.C. and Fruman, D.A. (2004) PI3K signaling controls cell fate at many points in B lymphocyte development and activation. *Semin. Cell Dev. Biol.* 15, 183–197
- 45 Yusuf, I. *et al.* (2004) Optimal B-cell proliferation requires phosphoinositide 3-kinase-dependent inactivation of FOXO transcription factors. *Blood* 104, 784–787
- 46 Glassford, J. *et al.* (2005) Phosphatidylinositol 3-kinase is required for the transcriptional activation of cyclin D2 in BCR activated primary mouse B lymphocytes. *Eur. J. Immunol.* 35, 2748–2761
- 47 Donahue, A.C. and Fruman, D.A. (2003) Proliferation and survival of activated B cells requires sustained antigen receptor engagement and phosphoinositide 3-kinase activation. *J. Immunol.* 170, 5851–5860
- 48 Vigorito, E. *et al.* (2006) Synergistic activation of PKD by the B cell antigen receptor and CD19 requires PI3K, Vav1 and PLC $\gamma$ . *Cell. Signal.* 18, 1455–1460
- 49 Doughty, C.A. *et al.* (2006) Antigen receptor-mediated changes in glucose metabolism in B lymphocytes: role of phosphatidylinositol 3-kinase signaling in the glycolytic control of growth. *Blood* 107, 4458–4465
- 50 Piatelli, M.J. *et al.* (2004) Phosphatidylinositol 3-kinase-dependent mitogen-activated protein/extracellular signal-regulated kinase 1/2 and NF- $\kappa$ B signaling pathways are required for B cell antigen receptor-mediated cyclin D2 induction in mature B cells. *J. Immunol.* 172, 2753–2762
- 51 Hebeis, B.J. *et al.* (2004) The p110 $\delta$  subunit of phosphoinositide 3-kinase is required for the lipopolysaccharide response of mouse B cells. *Biochem. Soc. Trans.* 32, 789–791
- 52 Fruman, D.A. *et al.* (2000) Xid-like phenotypes: a B cell signalosome takes shape. *Immunity* 13, 1–3
- 53 Suzuki, H. *et al.* (2003) PI3K and Btk differentially regulate B cell antigen receptor-mediated signal transduction. *Nat. Immunol.* 4, 280–286
- 54 Dai, X. *et al.* (2006) Distinct roles of phosphoinositide-3 kinase and phospholipase C $\gamma$ 2 in B-cell receptor-mediated signal transduction. *Mol. Cell. Biol.* 26, 88–99
- 55 Del Nagro, C.J. *et al.* (2005) CD19 function in central and peripheral B-cell development. *Immunol. Res.* 31, 119–131
- 56 Inabe, K. *et al.* (2002) Vav3 modulates B cell receptor responses by regulating phosphoinositide 3-kinase activation. *J. Exp. Med.* 195, 189–200
- 57 Vigorito, E. *et al.* (2004) Vav-dependent and vav-independent phosphatidylinositol 3-kinase activation in murine B cells determined by the nature of the stimulus. *J. Immunol.* 173, 3209–3214
- 58 Wang, Y. *et al.* (2002) The physiologic role of CD19 cytoplasmic tyrosines. *Immunity* 17, 501–514

- 59 Anzelon, A.N. *et al.* (2003) Pten inactivation alters peripheral B lymphocyte fate and reconstitutes CD19 function. *Nat. Immunol.* 4, 287–294
- 60 Brauweiler, A.M. and Cambier, J.C. (2003) Fc $\gamma$ RIIB activation leads to inhibition of signalling by independently ligated receptors. *Biochem. Soc. Trans.* 31, 281–285
- 61 Suzuki, A. *et al.* (2003) Critical roles of Pten in B cell homeostasis and immunoglobulin class switch recombination. *J. Exp. Med.* 197, 657–667
- 62 Omori, S.A. *et al.* (2006) Regulation of class-switch recombination and plasma cell differentiation by phosphatidylinositol 3-kinase signaling. *Immunity* 25, 545–557
- 63 Suzuki, A. *et al.* (2001) T cell-specific loss of Pten leads to defects in central and peripheral tolerance. *Immunity* 14, 523–534
- 64 Hagenbeek, T.J. *et al.* (2004) The loss of PTEN allows TCR  $\alpha\beta$  lineage thymocytes to bypass IL-7 and pre-TCR-mediated signaling. *J. Exp. Med.* 200, 883–894
- 65 Williams, J.A. *et al.* (2005) Regulated costimulation in the thymus is critical for T cell development: dysregulated CD28 costimulation can bypass the pre-TCR checkpoint. *J. Immunol.* 175, 4199–4207
- 66 Webb, L.M. *et al.* (2005) Cutting edge: T cell development requires the combined activities of the p110 $\gamma$  and p110 $\delta$  catalytic isoforms of phosphatidylinositol 3-kinase. *J. Immunol.* 175, 2783–2787
- 67 Swat, W. *et al.* (2006) Essential role of PI3K $\delta$  and PI3K $\gamma$  in thymocyte survival. *Blood* 107, 2415–2422
- 68 Patton, D.T. *et al.* (2006) Cutting edge: The phosphoinositide 3-kinase p110 $\delta$  is critical for the function of CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> regulatory T cells. *J. Immunol.* 177, 6598–6602
- 69 Rudd, C.E. and Schneider, H. (2003) Unifying concepts in CD28, ICOS and CTLA4 co-receptor signalling. *Nat. Rev. Immunol.* 3, 544–556
- 70 Vanhaesebroeck, B. *et al.* (1999) Autophosphorylation of p110 $\delta$  phosphoinositide 3-kinase: a new paradigm for the regulation of lipid kinases *in vitro* and *in vivo*. *EMBO J.* 18, 1292–1302
- 71 Charvet, C. *et al.* (2006) Vav1 promotes T cell cycle progression by linking TCR/CD28 costimulation to FOXO1 and p27kip1 expression. *J. Immunol.* 177, 5024–5031
- 72 Frauwirth, K.A. *et al.* (2002) The CD28 signaling pathway regulates glucose metabolism. *Immunity* 16, 769–777
- 73 Wu, L.X. *et al.* (2005) CD28 regulates the translation of Bcl-xL via the phosphatidylinositol 3-kinase/mammalian target of rapamycin pathway. *J. Immunol.* 174, 180–194
- 74 Fox, C.J. *et al.* (2005) The Pim kinases control rapamycin-resistant T cell survival and activation. *J. Exp. Med.* 201, 259–266
- 75 Nashed, B.F. *et al.* Role of the phosphoinositide 3-kinase p110 $\delta$  in the generation of type 2 cytokine responses and allergic airway inflammation. *Eur. J. Immunol.* (in press)
- 76 Lin, L. *et al.* (2004) Regulation of NF- $\kappa$ B, Th activation, and autoinflammation by the forkhead transcription factor Foxo3a. *Immunity* 21, 203–213
- 77 Sakaguchi, S. (2004) Naturally arising CD4<sup>+</sup> regulatory T cells for immunologic self-tolerance and negative control of immune responses. *Annu. Rev. Immunol.* 22, 531–562
- 78 Coombes, J.L. *et al.* (2005) Regulatory T cells and intestinal homeostasis. *Immunol. Rev.* 204, 184–194
- 79 Oak, J.S. *et al.* (2006) Sjogren's syndrome-like disease in mice with T cells lacking class IA phosphoinositide-3-kinase. *Proc. Natl. Acad. Sci. U. S. A.* 103, 16882–16887
- 80 Wilson, B.S. *et al.* (2002) Fc $\epsilon$ RI signaling observed from the inside of the mast cell membrane. *Mol. Immunol.* 38, 1259–1268
- 81 Tkaczyk, C. *et al.* (2003) The phospholipase C $\gamma$ 1-dependent pathway of Fc $\epsilon$ RI-mediated mast cell activation is regulated independently of phosphatidylinositol 3-kinase. *J. Biol. Chem.* 278, 48474–48484
- 82 Furumoto, Y. *et al.* (2006) Cutting edge: lentiviral short hairpin RNA silencing of PTEN in human mast cells reveals constitutive signals that promote cytokine secretion and cell survival. *J. Immunol.* 176, 5167–5171
- 83 Rivera, J. and Gilfillan, A.M. (2006) Molecular regulation of mast cell activation. *J. Allergy Clin. Immunol.* 117, 1214–1225
- 84 Kitaura, J. *et al.* (2000) Akt-dependent cytokine production in mast cells. *J. Exp. Med.* 192, 729–740
- 85 Lee, J.S. *et al.* (2005) Phosphatidylinositol (3,4,5)-trisphosphate specifically interacts with the phox homology domain of phospholipase D1 and stimulates its activity. *J. Cell Sci.* 118, 4405–4413
- 86 Peng, Z. and Beaven, M.A. (2005) An essential role for phospholipase D in the activation of protein kinase C and degranulation in mast cells. *J. Immunol.* 174, 5201–5208
- 87 Okada, T. *et al.* (2000) BCAP: the tyrosine kinase substrate that connects B cell receptor to phosphoinositide 3-kinase activation. *Immunity* 13, 817–827
- 88 Itoh, S. *et al.* (2002) Adapter molecule Grb2-associated binder 1 is specifically expressed in marginal zone B cells and negatively regulates thymus-independent antigen-2 responses. *J. Immunol.* 168, 5110–5116
- 89 Wang, Y. *et al.* (2005) Single and combined deletions of the NTAL/LAB and LAT adaptors minimally affect B-cell development and function. *Mol. Cell. Biol.* 25, 4455–4465
- 90 Wood, J.E. *et al.* (2006) TcR and TcR-CD28 engagement of protein kinase B (PKB/AKT) and glycogen synthase kinase-3 (GSK-3) operates independently of guanine nucleotide exchange factor VAV-1. *J. Biol. Chem.* 281, 32385–32394
- 91 Reynolds, L.F. *et al.* (2002) Vav1 transduces T cell receptor signals to the activation of phospholipase C- $\gamma$ 1 via phosphoinositide 3-kinase-dependent and -independent pathways. *J. Exp. Med.* 195, 1103–1114
- 92 Barber, D.F. *et al.* (2006) Class IB-phosphatidylinositol 3-kinase (PI3K) deficiency ameliorates IA-PI3K-induced systemic lupus but not T cell invasion. *J. Immunol.* 176, 589–593
- 93 Deane J.A. *et al.* T cell function is partially maintained in the absence of class IA phosphoinositide 3-kinase signaling *Blood* (in press)

## The ScienceDirect collection

ScienceDirect's extensive and unique full-text collection covers more than 1900 journals, including titles such as *The Lancet*, *Cell*, *Tetrahedron* and the full suite of *Trends*, *Current Opinion* and *Drug Discovery Today* journals. With ScienceDirect, the research process is enhanced with unsurpassed searching and linking functionality, all on a single, intuitive interface.

The rapid growth of the ScienceDirect collection is a result of the integration of several prestigious publications and the ongoing addition to the Backfiles – heritage collections in a number of disciplines. The latest step in this ambitious project to digitize all of Elsevier's journals back to volume one, issue one, is the addition of the highly cited *Cell Press* journal collection on ScienceDirect. Also available online for the first time are six *Cell* titles' long-awaited Backfiles, containing more than 12,000 articles that highlight important historic developments in the field of life sciences.

For more information, visit [www.sciencedirect.com](http://www.sciencedirect.com)