



Chaotic distributions for relativistic particles

Downloaded from: <https://research.chalmers.se>, 2023-01-21 01:11 UTC

Citation for the original published paper (version of record):

Mustafa, D., Wennberg, B. (2016). Chaotic distributions for relativistic particles. *Kinetic and Related Models*, 9(4): 749-766. <http://dx.doi.org/10.3934/krm.2016014>

N.B. When citing this work, cite the original published paper.

CHAOTIC DISTRIBUTIONS FOR RELATIVISTIC PARTICLES

DAWAN MUSTAFA AND BERNT WENNBORG

Department of Mathematical sciences
Chalmers University of Technology and the University of Gothenburg
412 96 GÖTEBORG, Sweden

(Communicated by Reinhard Illner)

ABSTRACT. We study a modified Kac model where the classical kinetic energy is replaced by an arbitrary energy function $\phi(v)$, $v \in \mathbb{R}$. The aim of this paper is to show that the uniform density with respect to the microcanonical measure is $Ce^{-z_0\phi(v)}$ -chaotic, $C, z_0 \in \mathbb{R}_+$. The kinetic energy for relativistic particles is a special case. A generalization to the case $v \in \mathbb{R}^d$ which involves conservation momentum is also formally discussed.

1. Introduction. In 1956, Mark Kac published a paper [9], in which he answered some fundamental questions concerning the derivation of the spatially homogeneous Boltzmann equation. Kac considered a stochastic model consisting of N identical particles, and obtained an equation like the spatially homogeneous Boltzmann equation as a mean-field limit when the number of particles tends to infinity. The key ingredient in his paper was the notion of *chaos*, which goes back to Boltzmann's *stosszahlansatz*. Loosely speaking, chaos is related to asymptotic independence. We give a precise mathematical definition later. He showed that if the probability distribution of the initial state of the particles is chaotic then this property is propagated in time, i.e., the probability distribution at time $t > 0$ is also chaotic. This is referred to as propagation of chaos.

Kac's model is a jump process on $\mathbb{S}^{N-1} = \{(v_1, \dots, v_N) \in \mathbb{R}^N \mid v_1^2 + \dots + v_N^2 = N\}$. This corresponds to an N -particle system where a particle is represented by its velocity $v \in \mathbb{R}$, and the kinetic energy of a particle is given by v^2 . The total energy is conserved in this process, which is ergodic and hence has a unique invariant measure. In the original model, the stationary distribution is the uniform measure on \mathbb{S}^{N-1} . It is easy to prove that this uniform measure is $(2\pi)^{-1/2}e^{-v^2/2}$ -chaotic according to the definition given below, and Kac also provides a method of constructing other families of chaotic measures on \mathbb{S}^{N-1} which may serve as suitable initial distributions for the process.

In this paper we consider a generalization of Kac's model, where the energy of a particle is given by a large class of functions $\phi(v)$, which includes $\phi(v) = v^2$. The motivation for studying this problem comes from the theory of special relativity,

2010 *Mathematics Subject Classification.* Primary: 60K35, 82B40, 82C40; Secondary: 60J75, 83A05.

Key words and phrases. Chaotic distributions, master equation, relativistic Kac model, propagation of chaos, microcanonical measure.

where v represents the momentum of a particle rather than its velocity, and where the kinetic energy is given by

$$\phi(v) = \sqrt{1 + v^2} - 1.$$

The kinetic theory for relativistic particles is far less developed than for classical particles, but there is a growing interest in this kind of problems (see below for references), and this paper is a contribution to this work.

The first result of this paper is a proof that the uniform distribution on the manifold represented by the total kinetic energy of the particles (depends on ϕ) with respect to the microcanonical measure is chaotic by using the approach of Kac. We also consider a jump process on this manifold, which is a direct generalization of Kac's model, and prove that propagation of chaos holds for this process. We also consider a generalization to a model where the particles have momenta $v \in \mathbb{R}^d$.

One of the most influential equations in the mathematical kinetic theory of gases is the Boltzmann equation, which describes the time evolution of the density of a single particle in a gas consisting of a large number of particles.

The Boltzmann equation is

$$\begin{cases} \frac{\partial}{\partial t} f(x, v, t) + \tilde{v} \cdot \nabla_x f(x, v, t) = Q(f, f)(x, v, t), & v \in \mathbb{R}^3, t > 0, \\ f(v, 0) = f_0, \end{cases} \quad (1)$$

where in the classical case $\tilde{v} = v$, and in the relativistic case $\tilde{v} = v/\sqrt{1 + |v|^2}$. The collision operator on the right hand side is a quadratic operator acting only in the velocity variables. In the classical case it is given by

$$Q(f, f) = \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} (f(v', t)f(v'_*, t) - f(v, t)f(v_*, t)) B(v - v_*, \sigma) dv_* d\sigma.$$

The function B is called the collision kernel and is derived from the physics of the model. The corresponding collision term is considerably more complicated in the relativistic case, which is thoroughly investigated in [15], for example. In both cases, (v, v_*) and (v', v'_*) represent the velocities of a pair of particles before and after an elastic collision, and satisfy the conservation of momentum,

$$v + v_* = v' + v'_*, \quad (2)$$

and energy

$$\phi(|v|) + \phi(|v_*|) = \phi(|v'|) + \phi(|v'_*|). \quad (3)$$

The derivation of the Boltzmann equation from a classical (Newtonian) many particle system is still an important research topic. The classical result in this direction is by Lanford [10], however, this result is valid only over a fraction of the mean free time between collisions. Up to this date the result by Lanford has been improved upon only in details, see e.g., [12] and the recent book [7]. For the study of the relativistic Boltzmann equation, see e.g., [4].

To obtain an equation like (1) as a mean-field limit, Kac considered the *master equation*, which describes the evolution of the probability density on state space. Assuming that velocities of the particles is distributed according to a probability density F_0 initially, its time evolution is given by the following linear master equation

$$\begin{cases} \frac{\partial}{\partial t} F_N(v_1, \dots, v_N, t) = \mathcal{K}F_N(v_1, \dots, v_N, t), \\ F_N(v_1, \dots, v_N, 0) = F_0(v_1, \dots, v_N). \end{cases} \quad (4)$$

where \mathcal{K} is the generator of a Markov jump process,

$$\mathcal{K}F_N(v_1, \dots, v_N) = N[Q - I]F_N(v_1, \dots, v_N),$$

with I being the identity operator and

$$QF_N(v_1, \dots, v_N) = \frac{2}{N(N-1)} \sum_{i < j} \int_{-\pi}^{\pi} F_N(v_1, \dots, v'_i(\theta), \dots, v'_j(\theta), \dots, v_N) \frac{d\theta}{2\pi}.$$

Hence \mathcal{K} can be written as a sum of generators acting on only two variables, v_i and v_j , say, which corresponds to binary collisions in a real gas. The pair of velocities $v'_i(\theta), v'_j(\theta)$ is the outcome of the velocities v_i, v_j undergoing a collision, and in the classical case, as proposed by Kac they are given by

$$v'_i(\theta) = v_i \cos \theta + v_j \sin \theta, \quad v'_j(\theta) = -v_i \sin \theta + v_j \cos \theta. \tag{5}$$

The energy of a pair of particles is always conserved in a collision, i.e.,

$$v'_i(\theta)^2 + v'_j(\theta)^2 = v_i^2 + v_j^2.$$

In a realistic model the momentum should also be conserved, but since the particles have one-dimensional velocities, imposing conservation of momentum would lead to either the particles keeping the velocities during a collision or exchanging velocities. Momentum conservation is therefore sacrificed in this case.

The fact that particles are assumed to be indistinguishable corresponds to the initial distribution F_0 being symmetric with respect to permutations, and this property is preserved by \mathcal{K} , so that F_N is symmetric for all $t > 0$.

In order to obtain a mean-field limit of the master equation (4) as $N \rightarrow \infty$, Kac introduced the notion of chaos (in [9], he referred to it as the Boltzmann property).

Definition 1.1. Let f be probability density on \mathbb{R} . For each $N \in \mathbb{N}$, let F_N be a probability density on \mathbb{R}^N with respect to a measure $m^{(N)}$. The sequence $\{F_N\}_{N \in \mathbb{N}}$ of probability densities on \mathbb{R}^N is said to be f -chaotic if the following two conditions are satisfied:

1. Each F_N is a symmetric function of v_1, \dots, v_N .
2. For each fixed $k \in \mathbb{N}$, the k -th marginal $f_k^N(v_1, \dots, v_k)$ of F_N converges to $\prod_{i=1}^k f(v_i)$, as $N \rightarrow \infty$, where $f(v) = \lim_{N \rightarrow \infty} f_1^N(v)$. The convergence is to be taken in the weak sense, that is, if $\varphi(v_1, \dots, v_k)$ is a bounded continuous function of k variables, $v_1, \dots, v_k \in \mathbb{R}$, then

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_{\mathbb{R}^N} \varphi(v_1, \dots, v_k) F_N(v_1, \dots, v_N) dm^{(N)} \\ = \int_{\mathbb{R}^k} \varphi(v_1, \dots, v_k) \prod_{i=1}^k f(v_i) dv_1 \dots dv_k. \end{aligned} \tag{6}$$

More generally, this can be formulated for probability densities on products E^N where E may be an arbitrary Polish space. The setting of this paper is also slightly different from the Definition 1.1 because the dynamics take place on a submanifold of \mathbb{R}^N .

A chaotic family of probability densities on $\mathbb{S}^{N-1}(\sqrt{N})$ is the following:

Example 1.2. It is a well known fact that the surface area of the sphere $\mathbb{S}^{N-1}(\sqrt{E})$ in \mathbb{R}^N is given by

$$|\mathbb{S}^{N-1}(\sqrt{E})| = \frac{2\pi^{\frac{N}{2}} E^{\frac{N-1}{2}}}{\Gamma(\frac{N}{2})}.$$

Let

$$F_N(v_1, \dots, v_N) = \frac{1}{|\mathbb{S}^{N-1}(\sqrt{N})|}$$

be the symmetric uniform density on $\mathbb{S}^{N-1}(\sqrt{N})$ with respect to surface measure $\sigma^{(N)}$. Let φ be continuous function on \mathbb{R} . It is easy to see that

$$\int_{\mathbb{S}^{N-1}(\sqrt{E})} \varphi(v_1) d\sigma^{(E)} = \int_{-\sqrt{E}}^{\sqrt{E}} \varphi(v_1) \left| \mathbb{S}^{N-2} \left(\sqrt{E - v_1^2} \right) \right| dv_1.$$

Replacing E with N we have

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{\int_{\mathbb{S}^{N-1}(\sqrt{N})} \varphi(v_1) d\sigma^{(N)}}{|\mathbb{S}^{N-1}(\sqrt{N})|} \\ &= \lim_{N \rightarrow \infty} \frac{\Gamma\left(\frac{N}{2}\right)}{\pi^{\frac{1}{2}} N^{\frac{1}{2}} \Gamma\left(\frac{N-1}{2}\right)} \int_{-\sqrt{N}}^{\sqrt{N}} \varphi(v_1) \left(1 - \frac{v_1^2}{N}\right)^{\frac{N-3}{2}} dv_1 \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \varphi(v_1) e^{-\frac{v_1^2}{2}} dv_1. \end{aligned}$$

Taking φ to be a function on \mathbb{R}^k , the same calculations show that, the family of uniform densities on $\mathbb{S}^{N-1}(\sqrt{N})$, with respect to the surface measure, is $\frac{1}{\sqrt{2\pi}} e^{-\frac{v^2}{2}}$ -chaotic.

Kac showed by using a combinatorial argument that the master equation (4) propagates chaos, that is, if $\{F_N(v_1, \dots, v_N, 0)\}_{N \in \mathbb{N}}$ is f_0 -chaotic, then the solutions to (4), $\{F_N(v_1, \dots, v_N, t)\}_{N \in \mathbb{N}}$ is $f(v, t)$ -chaotic. The density $f(v, t)$ satisfies the Boltzmann-Kac equation

$$\begin{aligned} \frac{\partial}{\partial t} f(v, t) &= 2 \int_{\mathbb{R}} \int_{-\pi}^{\pi} [f(v(\theta), t) f(w(\theta), t) - f(v, t) f(w, t)] \frac{d\theta}{2\pi} dw, \\ f(v, 0) &= f_0, \end{aligned} \tag{7}$$

with $v(\theta), w(\theta)$ given by (5).

Kac described also a method to construct other chaotic probability densities on $\mathbb{S}^{N-1}(\sqrt{N})$: Let

$$F_N(v_1, \dots, v_N) = \frac{\prod_{i=1}^N f(v_i)}{Z(\sqrt{N})}, \tag{8}$$

where

$$Z(\sqrt{E}) = \int_{\mathbb{S}^{N-1}(\sqrt{E})} \prod_{i=1}^N f(v_i) d\sigma^{(E)}, \tag{9}$$

with $\sigma^{(E)}$ being the surface measure on $\mathbb{S}^{N-1}(\sqrt{E})$. Kac showed that, under some conditions on the function f , the family of probability densities $\{F_N\}_{N \in \mathbb{N}}$ is $h(v)$ -chaotic, where

$$h(v) = \frac{e^{-z_0 v^2} f(v)}{\int_{\mathbb{R}} e^{-z_0 v^2} f(v) dv}, \tag{10}$$

and z_0 is a positive constant. Note that the case $f(v) = 1$ corresponds to the previous example.

The purpose of the present article is to study chaotic probability densities on other surfaces than $\mathbb{S}^{N-1}(\sqrt{N})$. Define

$$\Omega^{N-1}(\sqrt{E}) = \left\{ (v_1, \dots, v_N) \mid \sum_{i=1}^N \phi(v_i) = E \right\}, \tag{11}$$

where $\phi(v)$ represents the energy of a particle with velocity v , and

$$\begin{aligned} \phi &\in C^1(\mathbb{R}, \mathbb{R}^+), & \phi &\text{ is even and convex,} \\ \phi(0) &= 0. \end{aligned} \tag{12}$$

We show that the uniform density on $\Omega^{N-1}(\sqrt{N})$ with respect to the microcanonical measure is $Ce^{-z_0\phi(v)}$ -chaotic, where C is a normalisation constant and $z_0 > 0$ is the unique solution to a specific equation.

Chaotic probability densities on spheres have been considered by many authors. In a paper by Carlen, Carvalho, Le Roux, Loss and Villani, [1], the authors show the following theorem, which differs from Kac’ result both in the exact definition of chaos (the notion of entropic chaos is introduced), and in the method of proof:

Theorem 1.3. *Let f be a probability density on \mathbb{R} such that $f \in L^p(\mathbb{R})$ for some $p > 1$, $\int_{\mathbb{R}} v^2 f(v)dv = 1$ and $\int_{\mathbb{R}} v^4 f(v)dv < \infty$. Let a family of densities $\{F_N\}_{N \in \mathbb{N}}$, be defined with F_N as in(8) with $\sigma^{(N)}$ now being the normalized surface measure on $\mathbb{S}^{N-1}(\sqrt{N})$. Then $\{F_N\}_{N \in \mathbb{N}}$ is f -chaotic.*

Sznitman [13] constructed chaotic families of measures using a somewhat different approach: Let h_1, \dots, h_N be i.i.d random variables with law $\mu(dh) = f(h)dh$, where $h \in \mathbb{R}^d$. Assume that f is differentiable and satisfies the following condition

$$\int_{\mathbb{R}^d} (f(h) + |\nabla f(h)|)e^{\lambda|h|}dh < \infty, \tag{13}$$

where $\lambda \in \mathbb{R}$. Then, the conditional distribution μ^N of (h_1, \dots, h_N) subject to the constraint $\frac{h_1 + \dots + h_N}{N} = a$ is Υ -chaotic, where

$$\Upsilon = \frac{1}{Z_{\lambda_a}} e^{\lambda_a \cdot h} \mu(dh), \tag{14}$$

with λ_a determined by the equation $\int h d\Upsilon(h) = a$ and Z_λ is a normalisation constant. Note that, the choice of λ for which condition (13) is satisfied depends on a . Within the framework of this paper we think of the random variable $h \in \mathbb{R}$ as the energy of a particle, in this case $h = \phi(v) \geq 0$.

The organization of this paper is as follows. In section 2 we define the *micro-canonical* measure on $\Omega^{N-1}(\sqrt{N})$ and show that the uniform density on $\Omega^{N-1}(\sqrt{N})$ with respect to this measure is chaotic. In Section 3, we introduce a modified Kac model with particles having the energy $\phi(v)$ and the corresponding master equation. We also show that this master equation propagates chaos and obtain the limiting equation. In Section 4, we discuss how to generalize the results in Section 2 to the case $v \in \mathbb{R}^d$, $d > 1$, in which case the momentum is also conserved. An Appendix is also included to introduce the methods that are used in the paper.

2. Chaotic measures. This section is devoted to showing that the family of uniform probability densities on $\Omega^{N-1}(\sqrt{N})$ with respect to the microcanonical measure is chaotic. By Example 1.2, another approach would be show that the uniform density on $\Omega^{N-1}(\sqrt{N})$ with respect to surface measure is chaotic. However, since

we consider a particle system where the total energy is conserved it is natural to use the microcanonical measure on $\Omega^{N-1}(\sqrt{N})$. We show that the family of uniform probability densities on $\Omega^{N-1}(\sqrt{N})$ with respect to the microcanonical measure is $Ce^{-z_0\phi(v)}$ -chaotic where $z_0 > 0$ is the solution to a specific equation and C is a normalisation constant.

The microcanonical measure on $\Omega^{N-1}(\sqrt{N})$ is defined as follows: Let

$$H(v_1, \dots, v_N) = \sum_{i=1}^N \phi(v_i). \quad (15)$$

Definition 2.1. Provided that $|\nabla H| \neq 0$, the microcanonical measure, $\eta^{(E)}$, on $\Omega^{N-1}(\sqrt{E})$, is defined by

$$\eta^{(E)} = \frac{\sigma_\Omega}{|\nabla H|}, \quad (16)$$

where σ_Ω is the surface measure induced by the Euclidean measure in \mathbb{R}^N on $\Omega^{N-1}(\sqrt{E})$.

The microcanonical measure arises naturally in physics from the assumption of equal probability, meaning essentially that all microscopic particle configurations corresponding to the same energy are equally probable, see *e.g.* [11] or [16].

A more geometrical approach is given by the coarea formula, as explained *e.g.* in [6]: If $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is Lipschitz continuous and $n \geq m$, then for each measurable set $A \subset \mathbb{R}^n$,

$$\int_A J\Phi(x) dx = \int_{\mathbb{R}^m} \mathcal{H}^{n-m}(A \cap \Phi^{-1}(y)) dy, \quad (17)$$

where $J\Phi$ is the Jacobian of Φ , as defined in [6], and \mathcal{H}^{n-m} denotes the $n - m$ -dimensional Hausdorff measure. In our setting this can be rephrased as (see Theorem 3.13 in [6]): if $H : \mathbb{R}^n \rightarrow \mathbb{R}$ is Lipschitz continuous and $\text{ess inf} |\nabla H| > 0$, and $g : \mathbb{R} \rightarrow \mathbb{R}$ is integrable, then for almost all E ,

$$\frac{d}{dE} \int_{\mathbb{R}^n} g(x) \chi_{\{H(x) > E\}} dx = \int_{\{H=E\}} \frac{g}{|\nabla H|} d\mathcal{H}^{n-1}. \quad (18)$$

where χ is the indicator function of a set. Because $H(v_1, \dots, v_N) = \phi(v_1) + \dots + \phi(v_N)$ with ϕ convex and differentiable, the set $\{H = E\}$ is also convex and regular. Then

$$\begin{aligned} & \int_{\mathbb{R}^N} g(v_1, \dots, v_N) \delta(H(v_1, \dots, v_N) - E) dv_1 \dots dv_N \\ &= \int_{\mathbb{R}} \int_{\Omega^{N-1}(\sqrt{y})} g \delta(y - E) \frac{d\sigma_\Omega}{|\nabla H|} dy = \int_{\Omega^{N-1}(\sqrt{E})} g \frac{d\sigma_\Omega}{|\nabla H|}. \end{aligned}$$

This shows that

$$\delta(H(v_1, \dots, v_N) - E) = \frac{\sigma_\Omega}{|\nabla H|}.$$

Another concept that is relevant in this context is that of disintegration of a measure, which is a measure theoretic approach to almost the same problem, and a means of approaching conditional probabilities, see *e.g.* [5]. In our setting, the disintegration theorem states that given a measure $d\mu$ on \mathbb{R}^N there are a measure ν on \mathbb{R}^+ and a family of measures μ_E on the level surfaces $\{(v_1, \dots, v_N) \mid \sum \phi(v_j) = E\}$, such that

$$\int_{\mathbb{R}^N} f(y) d\mu(y) = \int_0^\infty \int_{\{\sum \phi(v_j) = E\}} f(y) d\mu_E(y) d\nu(E), \quad (19)$$

Remark 1. On $\mathbb{S}^{N-1}(\sqrt{E})$, we have that $|\nabla H| = 2\sqrt{E}$. This implies that the microcanonical measure is up to a constant factor equal to the surface measure on $\mathbb{S}^{N-1}(\sqrt{E})$.

Using the equality $H(v_1, \dots, v_N) = E$, we can express, at least locally, the variable v_N as function of v_1, \dots, v_{N-1} :

$$v_N = U(v_1, \dots, v_{N-1}).$$

By this parametrization, the surface $\Omega^{N-1}(\sqrt{E})$ can be represented as the graph of $U : \mathbb{R}^{N-1} \rightarrow \mathbb{R}$. The surface measure σ_Ω on $\Omega^{N-1}(\sqrt{E})$ is now given by

$$d\sigma_\Omega = \sqrt{1 + |\nabla U|^2} dv_1 \dots dv_{N-1}.$$

By the implicit function theorem it follows that

$$\frac{\partial U}{\partial v_k} = -\frac{\frac{\partial H}{\partial v_k}}{\frac{\partial H}{\partial v_N}} \quad k = 1, \dots, N - 1.$$

Thus

$$\frac{d\sigma_\Omega}{|\nabla H|} = \frac{1}{\left| \frac{\partial H}{\partial v_N} \right|} dv_1 \dots dv_{N-1}.$$

To carry out integration on $\Omega^{N-1}(\sqrt{E})$ with respect to the microcanonical measure $\eta^{(E)}$ we use the last equality:

$$\begin{aligned} \int_{\Omega^{N-1}(\sqrt{E})} g(v_1, \dots, v_N) d\eta^{(E)} \\ = \sum_{\epsilon=+,-} \int_{\sum_{i=1}^{N-1} \phi(v_i) \leq E} g(v_1, \dots, \epsilon v_N) \frac{1}{\left| \frac{\partial H}{\partial v_N} \right|_\epsilon} dv_1 \dots dv_{N-1}, \end{aligned}$$

where

$$v_N = \phi^{-1} \left(E - \sum_{i=1}^{N-1} \phi(v_i) \right),$$

and $\phi^{-1}(v)$ is the inverse of $\phi(v)$, $v \geq 0$. Moreover,

$$\left| \frac{\partial H}{\partial v_N} \right|_\epsilon = \left| \frac{\partial H}{\partial v_N} \left(v_1, \dots, v_{N-1}, \epsilon \phi^{-1} \left(E - \sum_{i=1}^{N-1} \phi(v_i) \right) \right) \right|.$$

The uniform density $F(v_1, \dots, v_N)$ on $\Omega^{N-1}(\sqrt{N})$ with respect to the microcanonical measure $\eta^{(N)}$ is given by

$$F_N(v_1, \dots, v_N) = \frac{1}{Z_\phi(\sqrt{N})}, \tag{20}$$

where

$$Z_\phi(\sqrt{E}) = \int_{\Omega^{N-1}(\sqrt{E})} d\eta(E).$$

To show that the uniform density on $\Omega^{N-1}(\sqrt{N})$ with respect to the microcanonical measure $\eta^{(N)}$ is $Ce^{-z_0\phi(v)}$ -chaotic we follow Kac [9], and start by determining the asymptotic behaviour of

$$Z_\phi(\sqrt{E}) = \sum_{\epsilon=+,-} \int_{\sum_{i=1}^{N-1} \phi(v_i) \leq E} \frac{1}{\left| \frac{\partial H}{\partial v_N} \right|_\epsilon} dv_1 \dots dv_{N-1}$$

with $E = N$ for large N . Since

$$\left| \frac{\partial H}{\partial v_N} \right|_\epsilon = |\phi'(v_N)| \quad \text{and} \quad v_N = \pm \phi^{-1} \left(E - \sum_{i=1}^{N-1} \phi(v_i) \right),$$

we have

$$Z_\phi(\sqrt{E}) = 2 \int_{\sum_{i=1}^{N-1} \phi(v_i) \leq E} \frac{1}{\left| \phi'(\phi^{-1}(E - \sum_{i=1}^{N-1} \phi(v_i))) \right|} dv_1 \dots dv_{N-1}.$$

To write $Z_\phi(\sqrt{E})$ as an integral over the sphere $\mathbb{S}^{N-1}(\sqrt{E})$, we make the change of variables $y_i^2 = \phi(v_i)$ with respect to sign of v_i , $i = 1, \dots, N - 1$. This leads to

$$Z_\phi(\sqrt{E}) = 2^N \int_{\sum_{i=1}^{N-1} y_i^2 \leq E} \frac{1}{\left| \phi'(\phi^{-1}(E - \sum_{i=1}^{N-1} y_i^2)) \right|} \prod_{i=1}^{N-1} \frac{|y_i|}{|\phi'(\phi^{-1}(y_i^2))|} dy_1 \dots dy_{N-1}. \tag{21}$$

Let

$$f(y) := \frac{|y|}{|\phi'(\phi^{-1}(y^2))|}. \tag{22}$$

The integrand in (21) is almost a product of N copies of $f(y)$. Multiply and divide the integrand by $|y_N| = \sqrt{E - \sum_{i=1}^{N-1} y_i^2}$. Recall the following formula for integration over a sphere

$$\begin{aligned} \int_{\mathbb{S}^{N-1}(E)} g(y_1, \dots, y_N) d\sigma^{(E^2)} \\ = \sum_{\epsilon = +, -} \int_{\sum_{i=1}^{N-1} y_i^2 \leq E^2} g(y_1, \dots, \epsilon y_N) \frac{E dy_1 \dots dy_{N-1}}{\sqrt{E^2 - \sum_{i=1}^{N-1} y_i^2}}. \end{aligned}$$

We now get

$$Z_\phi(\sqrt{E}) = \frac{2^{N-1}}{\sqrt{E}} \int_{\mathbb{S}^{N-1}(\sqrt{E})} \prod_{i=1}^N f(y_i) d\sigma^{(E)}. \tag{23}$$

Having Z_ϕ given by (23) is convenient in sense that, in [9], Kac determined the asymptotic behaviour of $Z_\phi(\sqrt{N})$ for large N by using the *saddle point* method (see e.g. [8]). For completeness, we present each step of the result with rigorous justification with $f(y)$ given by (22). A short description of the saddle point method is given in the Appendix.

We start by computing the Laplace transform of $E \mapsto Z_\phi(\sqrt{E})$. The Laplace transform of $Z_\phi(\sqrt{E})$ is defined provided that $Z_\phi(\sqrt{E})$ grows at most exponentially. Since the behaviour of $Z_\phi(\sqrt{E})$ depends on the function $f(y)$ defined by (22) we assume that $\phi(y)$ is such that

$$f(y) \leq K e^{by^2}, \tag{24}$$

for some $K \geq 0$ and $b > 0$. This condition ensures that $Z_\phi(\sqrt{E})$ grows at most exponentially.

Taking the Laplace transform of $Z_\phi(\sqrt{E})$, making the change of variable $r = \sqrt{E}$, we have, for $w \in \mathbb{C}$ where $\Re(w) > b$

$$\int_0^\infty e^{-wE} Z_\phi(\sqrt{E}) dE = 2 \int_0^\infty e^{-wr^2} r Z_\phi(r) dr.$$

Using (23), the last equality equals

$$2^N \left(\int_{-\infty}^{\infty} e^{-wy^2} f(y) dy \right)^N.$$

From condition (24) and since $\Re(w) > b$ it follows that

$$\Phi(w) := \int_{-\infty}^{\infty} e^{-wy^2} f(y) dy \tag{25}$$

is an analytic function of w for $\Re(w) > b$. By applying the inverse of the Laplace transform, we get

$$Z_\phi(\sqrt{E}) = \frac{2^N}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{zE} \left(\int_{-\infty}^{\infty} e^{-zy^2} f(y) dy \right)^N dz,$$

where $\gamma > b$ is such that the line $\gamma = \Re(z)$ lies in the half-plane where $\Phi(z)$ is analytic. Replacing E with N , we get

$$Z_\phi(\sqrt{N}) = \frac{2^{N-1}}{\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \left(e^z \int_{-\infty}^{\infty} e^{-zy^2} f(y) dy \right)^N dz.$$

By comparing with the saddle point integral (52) in the Appendix, we set

$$q(z) = 1 \quad \text{and} \quad S(z) = z + \log \Phi(z), \tag{26}$$

which is well defined because $\Phi(z)$ is nonzero on the line $\gamma = \Re(z)$. We now have

$$Z_\phi(\sqrt{N}) = \frac{2^{N-1}}{\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{NS(z)} dz. \tag{27}$$

The asymptotic behaviour of $Z_\phi(\sqrt{N})$ for large N is determined by the saddle points of $S(z)$. The next lemma concerns the saddle points of $S(z)$.

Lemma 2.2. *Assume that there exists a $\gamma > 0$ such that*

$$\int_{-\infty}^{\infty} e^{-\gamma y^2} f(y) dy < \infty. \tag{28}$$

Moreover, assume that

$$\int_{-\infty}^{\infty} (1 - y^2) f(y) dy < 0, \tag{29}$$

and

$$\int_{|y| \leq 1} (1 - y^2) f(y) dy > 0. \tag{30}$$

Let $S(z)$ be given by (26) with $\Phi(z)$ by (25). For $\Re(z) \geq \gamma$, the function $S(z)$ is analytic and there exists a unique saddle point z_0 to $S(z)$ such that z_0 is real, $z_0 \geq \gamma$ and $S''(z_0) > 0$. Moreover

$$Z_\phi(\sqrt{N}) \sim \frac{2^{N-1} e^{Nz_0}}{\sqrt{NS''(z_0)}} \left(\int_{-\infty}^{\infty} e^{-z_0 y^2} f(y) dy \right)^N. \tag{31}$$

Proof. Note that, for $z = \xi + i\eta$, for all ξ

$$\arg \max_{\eta} |e^{\xi+i\eta} \Phi(\xi + i\eta)| = \{0\}.$$

Hence, we only need to find saddle points on the real line.

Claim 1. *The function $S(z)$ has a unique saddle point z_0 where $z_0 \geq \gamma$.*

Proof. For $\Re(z) \geq \gamma$, we have

$$S'(z) = 1 - \frac{\int_{-\infty}^{\infty} y^2 e^{-zy^2} f(y) dy}{\int_{-\infty}^{\infty} e^{-zy^2} f(y) dy}. \quad (32)$$

For $z = \xi + i0$ where $\xi \geq \gamma$, the equation $S'(\xi) = 0$ is equivalent to

$$\int_{-\infty}^{\infty} e^{-\xi y^2} (1 - y^2) f(y) dy = 0.$$

Multiplying the last equality by e^{ξ} , we see that, $S'(\xi) = 0$ is equivalent to

$$\int_{-\infty}^{\infty} e^{-\xi(y^2-1)} (1 - y^2) f(y) dy = 0.$$

Let

$$A(\xi) = \int_{-\infty}^{\infty} e^{-\xi(y^2-1)} (1 - y^2) f(y) dy.$$

Since

$$A'(\xi) = \int_{-\infty}^{\infty} e^{-\xi(y^2-1)} (1 - y^2)^2 f(y) dy > 0,$$

it follows that $A(\xi)$ is an increasing function of ξ . Moreover, by (29) we have

$$A(0) < 0.$$

Note that

$$\lim_{\xi \rightarrow \infty} \int_{|y| > 1} e^{-\xi(y^2-1)} (1 - y^2) f(y) dy = 0.$$

Hence

$$\begin{aligned} \int_{-\infty}^{\infty} e^{-\xi(y^2-1)} (1 - y^2) f(y) dy &\sim \int_{|y| \leq 1} e^{-\xi(y^2-1)} (1 - y^2) f(y) dy \\ &= \int_{|y| \leq 1} e^{\xi(1-y^2)} (1 - y^2) f(y) dy. \end{aligned}$$

The last integral goes to infinity by (30) as $\xi \rightarrow \infty$. Hence, there exists a unique $z_0 \geq \gamma$ such that $S'(z_0) = 0$. \square

Claim 2. *The second derivative of $S(z)$ at z_0 is positive.*

Proof. We have

$$S''(z) = \frac{\Phi''(z)}{\Phi(z)} - \frac{\Phi'(z)^2}{\Phi(z)^2}.$$

For $z = z_0$, using the Jensen inequality, we get

$$\frac{\Phi''(z_0)}{\Phi(z_0)} = \frac{\int_{-\infty}^{\infty} y^4 e^{-z_0 y^2} f(y) dy}{\int_{-\infty}^{\infty} e^{-z_0 y^2} f(y) dy} > \left(\frac{\int_{-\infty}^{\infty} y^2 e^{-z_0 y^2} f(y) dy}{\int_{-\infty}^{\infty} e^{-z_0 y^2} f(y) dy} \right)^2 = \frac{\Phi'(z_0)^2}{\Phi(z_0)^2}.$$

This proves the claim. \square

We now turn to the proof of (31). We can write

$$\Phi(z) = \int_{-\infty}^{\infty} e^{-zy^2} f(y) dy = \int_0^{\infty} e^{-zy^2} (f(y) + f(-y)) dy.$$

By a change of variables, the last integral equals

$$\int_0^{\infty} e^{-zy} \frac{f(\sqrt{y}) + f(-\sqrt{y})}{2\sqrt{y}} dy.$$

Let $z = \xi + i\eta$, with $\xi \geq \gamma$. The last integral can be written as

$$\int_{-\infty}^{\infty} e^{-\xi y} \frac{f(\sqrt{y}) + f(-\sqrt{y})}{2\sqrt{y}} \mathbb{1}_{y \geq 0} e^{-i\eta y} dy.$$

The last integral is the Fourier transform of the function $\tilde{\Phi}_\xi$ at the point η , where

$$\tilde{\Phi}_\xi(y) = e^{-\xi y} \frac{f(\sqrt{y}) + f(-\sqrt{y})}{2\sqrt{y}} \mathbb{1}_{y \geq 0}.$$

For $\xi = z_0$, it follows that $\tilde{\Phi}_\xi \in L^1(\mathbb{R})$, and $|\mathcal{F}(\tilde{\Phi}_\xi)(\eta)| < \mathcal{F}(\tilde{\Phi}_\xi)(0)$. Moreover, By the Riemann Lebesgue lemma, it follows that

$$|\mathcal{F}(\tilde{\Phi}_\xi)(\eta)| \rightarrow 0 \text{ when } |\eta| \rightarrow \infty.$$

Let C_{+T} and C_{-T} be the curves in the complex plane given $[\gamma + iT, z_0 + iT]$ and $[\gamma - iT, z_0 - iT]$, respectively.

We have

$$\begin{aligned} \left| \int_{C_{\pm T}} e^{NS(z)} dz \right| &= \left| \int_{\gamma}^{z_0} e^{N(\xi+iT)} \mathcal{F}(\tilde{\Phi}_\xi)(T)^N d\xi \right| \\ &\leq \int_{\gamma}^{z_0} e^{N\xi} |\mathcal{F}(\tilde{\Phi}_\xi)(T)|^N d\xi \\ &\leq |z_0 - \gamma| \max_{\xi \in [\gamma, z_0]} e^{N\xi} |\mathcal{F}(\tilde{\Phi}_\xi)(T)|^N \rightarrow 0, \text{ as } |T| \rightarrow \infty. \end{aligned}$$

By Cauchy’s theorem, we can deform the contour in (27) to a contour passing through the saddle point z_0 . Hence

$$Z_\phi(\sqrt{N}) = \frac{2^{N-1}}{\pi i} \int_{z_0-i\infty}^{z_0+i\infty} e^{NS(z)} dz.$$

By the saddle point method, we obtain

$$\begin{aligned} Z_\phi(\sqrt{N}) &= \frac{2^{N-1}}{\pi i} \sqrt{\frac{-2\pi}{NS''(z_0)}} \left(1 + \mathcal{O}\left(\frac{1}{N}\right) \right) e^{NS(z_0)} \\ &\sim \frac{2^N e^{Nz_0}}{\sqrt{NS''(z_0)}} \left(\int_{-\infty}^{\infty} e^{-z_0 y^2} f(y) dy \right)^N. \end{aligned} \tag{33}$$

This finishes the proof of the lemma. □

The main theorem of this section is:

Theorem 2.3. *Let $f(y)$ defined by (22) satisfy the conditions (24), (29) and (30). Then, the family of uniform densities on $\Omega^{N-1}(\sqrt{N})$ with respect to the micro-canonical measure is $Ce^{-z_0\phi(v)}$ –chaotic. The positive constant z_0 is the unique real solution to the following equation*

$$\int_{-\infty}^{\infty} (1 - \phi(v)) e^{-z_0\phi(v)} dv = 0,$$

and C is a normalisation constant given by

$$C = \frac{1}{\int_{-\infty}^{\infty} e^{-z_0\phi(v)} dv}.$$

Proof. Let φ be a bounded continuous function on \mathbb{R}^k . For each fixed $k \in \mathbb{N}$, a modification of Lemma 2.2 shows that, for large N ,

$$\begin{aligned} \sum_{\epsilon=+,-} \int_{\sum_{i=1}^{N-1} \phi(v_i) \leq E} \varphi(v_1, \dots, v_k) \frac{1}{|\frac{\partial H}{\partial v_N}|^\epsilon} dv_1 \dots dv_{N-1} \\ \sim \int_{\mathbb{R}^k} \varphi(\phi^{-1}(y_1^2), \dots, \phi^{-1}(y_k^2)) \prod_{i=1}^k e^{-z_0 y_i^2} f(y_i) dy_1 \dots dy_k \\ \times \frac{2^N e^{(N-k)z_0}}{\sqrt{N S''(z_0)}} \left(\int_{-\infty}^{\infty} e^{-z_0 y^2} f(y) dy \right)^{N-k}. \end{aligned}$$

Using Lemma 2.2 again and making the change of variable $y_i^2 = \phi(v_i)$, where $dv_i = f(y_i) dy_i$, $i = 1, \dots, k$ leads to

$$\begin{aligned} \lim_{N \rightarrow \infty} \frac{\int_{\Omega^{N-1}(\sqrt{N})} \varphi(v_1, \dots, v_k) d\eta^{(N)}}{Z_\phi(\sqrt{N})} \\ = \frac{\int_{\mathbb{R}^k} \varphi(\phi^{-1}(y_1^2), \dots, \phi^{-1}(y_k^2)) \prod_{i=1}^k e^{-z_0 y_i^2} f(y_i) dy_1 \dots dy_k}{\int_{\mathbb{R}^k} \prod_{i=1}^k e^{-z_0 y_i^2} f(y_i) dy_1 \dots dy_k} \\ = \frac{\int_{\mathbb{R}^k} \varphi(v_1, \dots, v_k) \prod_{i=1}^k e^{-z_0 \phi(v_i)} dv_1 \dots dv_k}{\int_{\mathbb{R}^k} \prod_{i=1}^k e^{-z_0 \phi(v_i)} dv_1 \dots dv_k}. \end{aligned}$$

This is what we wanted to prove. □

We now consider two examples where Theorem 2.3 applies :

Example 2.4. In the classical Kac model $\phi(v) = v^2$. We get that $f(y) = \frac{1}{2}$ and thus, the conditions (28), (29) and (30) are fulfilled. To determine z_0 we need solve the equation

$$\int_{-\infty}^{\infty} (1 - v^2) e^{-z_0 v^2} dv = 0.$$

Direct calculations show that $z_0 = \frac{1}{2}$ is the unique real solution. Therefore the uniform density on $(\Omega^{N-1}(\sqrt{N}) = \mathbb{S}^{N-1}(\sqrt{N}))$ with respect to the microcanonical measure is $\frac{e^{-\frac{1}{2}v^2}}{\sqrt{2\pi}}$ -chaotic. This has already been discussed in Example 1.2. Recall that, on $\mathbb{S}^{N-1}(\sqrt{N})$, the microcanonical measure up to a constant is equal to the surface measure

Example 2.5. For a relativistic energy function $\phi(v) = \sqrt{v^2 + 1} - 1$, it follows that

$$f(y) = \frac{(y^2 + 1)|y|}{\sqrt{(y^2 + 1)^2 - 1}}.$$

It is easy to check that the conditions (28), (29) and (30) are satisfied. To find z_0 we solve the equation

$$\int_{-\infty}^{\infty} (1 - (\sqrt{v^2 + 1} - 1)) e^{-z_0(\sqrt{v^2+1}-1)} dv = 0.$$

By using numerical integration, we get that $z_0 \approx 0.734641$. Hence, the family of uniform densities on $\Omega^{N-1}(\sqrt{N})$ with $\phi(v) = \sqrt{v^2 + 1} - 1$ in (11) is $Ce^{-z_0(\sqrt{v^2+1}-1)}$ -chaotic, where $C \approx 4.082$.

We end this section by a comparison between our result and the result of Sznitman. The setting there is slightly different, and as commented by Sznitman it is not directly applicable for the case $h = v^2$ or the more general situation here. To see the relation between two results, assume that an individual velocity is distributed according the Gibbs measure $e^{-\beta\phi(v)}/Z$, where Z is normalization constant and $\beta > 0$. Then the random variable $h = \phi(v)$ that is the energy of a particle, has a distribution $\mu(dh) = f(h)dh$, where

$$f(h) = \frac{e^{-\beta h}}{|\phi'(\phi^{-1}(h))|}. \tag{34}$$

To apply Sznitman’s result, we need to show that f satisfies condition (13). This is a strong integrability condition on f and depends on the choice of ϕ . Unfortunately, even for the most classical case, the one studied by Kac, where $h = v^2$, condition (13) is not fulfilled since $f'(h)e^{-\beta h}$ is not integrable at $h = 0$. It is a technical condition needed to control the Fourier transform of f , and it could probably be relaxed, but it means that Sznitman’s result cannot be directly applied to our case.

3. Particle dynamics and master equation for general energy functions.

In this section we follow [2] and [9] to introduce dynamics between particles having the energy given by the function ϕ and obtain the corresponding master equation. By similar arguments as in [9], we will see that the master equation propagates chaos.

To introduce dynamics between the particles, let the *master vector* $\mathbf{V} = (v_1, \dots, v_N) \in \Omega^{N-1}(\sqrt{N})$. The master vector \mathbf{V} makes a jump on $\Omega^{N-1}(\sqrt{N})$ according the following steps:

1. Pick a pair (i, j) , $i < j$ according to the uniform distribution

$$P_{ij} = \frac{2}{N(N-1)}.$$

2. The pair of velocities (v_i, v_j) satisfies

$$\phi(v_i) + \phi(v_j) = h, \quad h > 0.$$

Let

$$y_i = \text{sign}(v_i)\sqrt{\phi(v_i)} \quad \text{and} \quad y_j = \text{sign}(v_j)\sqrt{\phi(v_j)}.$$

Then (y_i, y_j) is a point on the circle, and as in the original Kac model, the collision may be performed there.

Pick an angle θ uniformly on $(0, 2\pi]$ and let

$$y'_i(\theta) = y_i \cos \theta + y_j \sin \theta \quad \text{and} \quad y'_j(\theta) = -y_i \sin \theta + y_j \cos \theta.$$

The pair (v_i, v_j) is transformed on $\Omega^1(\sqrt{h})$ to $(v'_i(\theta), v'_j(\theta))$ according to

$$\begin{aligned} v'_i(\theta) &= \text{sign}(y'_i(\theta))\phi^{-1}(y'_i(\theta)^2), \\ v'_j(\theta) &= \text{sign}(y'_j(\theta))\phi^{-1}(y'_j(\theta)^2), \end{aligned} \tag{35}$$

where

$$\text{sign}(v) = \begin{cases} 1, & \text{if } v \geq 0 \\ -1, & \text{if } v < 0. \end{cases}$$

Note that

$$\phi(v'_i(\theta)) + \phi(v'_j(\theta)) = \phi(v_i) + \phi(v_j).$$

3. Update the master vector \mathbf{V} and denote the new master vector by $T_{i,j}(\theta)\mathbf{V}$.

By step 2, it follows that $T_{i,j}(\theta)\mathbf{V} \in \Omega^{N-1}(\sqrt{N})$. Repeat step 1, 2 and 3.

This is only a generalization of the dynamics in [9] where $\phi(v) = v^2$.

The steps above describe a random walk on $\Omega^{N-1}(\sqrt{N})$. As in [2], we define its Markov transition operator Q_ϕ . If V_k is the state of the particles after the k -th step of the walk, for a continuous function φ on $\Omega^{N-1}(\sqrt{N})$, the operator Q_ϕ is defined by

$$Q_\phi \varphi(y) = \mathbb{E}[\varphi(V_{k+1}) | V_k = y].$$

Writing out the expectation above, we get

$$Q_\phi \varphi(\mathbf{V}) = \frac{2}{N(N-1)} \sum_{i < j} \int_0^{2\pi} \varphi(T_{i,j}(\theta)\mathbf{V}) \frac{d\theta}{2\pi}. \quad (36)$$

If F_k is probability density of V_k with respect to the microcanonical measure $\eta^{(N)}$ on $\Omega^{N-1}(\sqrt{N})$, we have

$$\begin{aligned} \int_{\Omega^{N-1}(\sqrt{N})} \varphi F_{k+1} d\eta^{(N)} &= \mathbb{E}[\varphi(V_{k+1})] = \mathbb{E}[\mathbb{E}[\varphi(V_{k+1}) | V_k]] \\ &= \int_{\Omega^{N-1}(\sqrt{N})} Q_\phi \varphi F_k d\eta^{(N)}. \end{aligned}$$

By definition, the microcanonical measure is invariant under the transformation $\mathbf{V} \rightarrow T_{i,j}(\theta)\mathbf{V}$. It follows that Q_ϕ is self adjoint and

$$F_{k+1} = Q_\phi F_k.$$

So far, the process defined above is discrete in time. To obtain a time continuous process, we let the master vector \mathbf{V} be a function of time, and the times between the jumps (collisions) exponentially distributed. In this way, if $F_N(\mathbf{V}, 0)$ is the probability distribution of the N particles on $\Omega^{N-1}(\sqrt{N})$ at time 0, the time evolution of $F_N(\mathbf{V}, t)$ is described by the following master equation

$$\frac{\partial F_N(\mathbf{V}, t)}{\partial t} = \mathcal{K}_\phi F_N(\mathbf{V}, t), \quad (37)$$

where

$$\mathcal{K}_\phi = N[Q_\phi - I] \quad (38)$$

and I is the identity operator. A more complete discussion of master equations of this kind may be found in [2].

Note that, for $\phi(v) = v^2$, the collision operator \mathcal{K}_ϕ is the same as the collision operator \mathcal{K} in the Kac model. We have propagation of chaos for the master equation (37):

Theorem 3.1. *Assume that the family of initial densities $\{F_N(\mathbf{V}, 0)\}_{N \in \mathbb{N}}$ on $\Omega^{N-1}(\sqrt{N})$ is $f(v, 0)$ -chaotic. Then, the family of densities $\{F_N(\mathbf{V}, t)\}_{N \in \mathbb{N}}$, where*

$F_N(\mathbf{V}, t)$ is the solution to (37) is $f(v, t)$ -chaotic. Moreover, the density $f(v, t)$ satisfies the following equation

$$\frac{\partial}{\partial t} f(v, t) = \int_{\mathbb{R}} \int_0^{2\pi} [f(v(\theta), t) f(w(\theta), t) - f(v, t) f(w, t)] \frac{d\theta}{2\pi} dw, \quad (39)$$

with $f(v, 0) = f_0(v)$ and $v(\theta), w(\theta)$ given by (35).

Proof. The proof follows by the same arguments as in [9]. For a more detailed proof where propagation of chaos is shown for more general master equations, we refer to [2]. \square

4. Chaotic measures in higher dimensions. In Section 2 we proved that the uniform density with respect to the microcanonical measure on $\Omega^{N-1}(\sqrt{N})$ is $Ce^{-z_0\phi(v)}$ -chaotic, $v \in \mathbb{R}$. The goal of this section is to generalize this to the case $v \in \mathbb{R}^2$ where now both the energy and momentum are conserved; the generalization to \mathbb{R}^d , $d > 2$ may be treated in the same way. The calculations here are formal. For $p \in \mathbb{R}^2$, define

$$\Gamma^N(\sqrt{E}, p) = \left\{ (v_1, \dots, v_N) \in \mathbb{R}^{2N} \mid \sum_{i=1}^N \phi(v_i) = 2E, \sum_{i=1}^N v_i = p \right\}. \quad (40)$$

We assume that E and p are chosen such that $\Gamma^N(\sqrt{E}, p)$ is non-empty. The classical case when $\phi(v) = |v|^2$ has been thoroughly investigated in [3].

For $p = (p_1, p_2)$ and $v_i = (v_{i1}, v_{i2})$, $i = 1, \dots, N$, a measure μ_{E,p_1,p_2} concentrated on $\Gamma^N(\sqrt{E}, p)$ is defined by

$$\mu_{E,p_1,p_2} = \delta(2E - \sum_{i=1}^N \phi(v_i)) \delta(p_1 - \sum_{i=1}^N v_{i1}) \delta(p_2 - \sum_{i=1}^N v_{i2}). \quad (41)$$

The product of the Dirac measures is well defined since the hyper surfaces defined by setting the arguments of the Dirac measures to zero are mutually transversal. Let

$$Z(E, p_1, p_2) = \int_{\mathbb{R}^{2N}} \delta(2E - \sum_{i=1}^N \phi(v_i)) \delta(p_1 - \sum_{i=1}^N v_{i1}) \delta(p_2 - \sum_{i=1}^N v_{i2}) dv_1 \dots dv_N. \quad (42)$$

As in Section 2, we need to determine the asymptotic behaviour of $Z(E, p_1, p_2)$ with $E = N$ for large N . We note that in the case of relativistic collisions, with $\phi(v) = \sqrt{|v|^2 + 1} - 1$, the measure μ_{E,p_1,p_2} is Lorentz invariant (see [14]).

In the sense of distributions, the δ function is the inverse Fourier transform of the function 1 and formally can be written as

$$\delta(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{ix\xi} d\xi. \quad (43)$$

For $z = (z_1, z_2, z_3)$, we can formally write

$$\begin{aligned} & Z(E, p_1, p_2) \\ &= \frac{1}{(2\pi)^3} \int_{\mathbb{R}^{2N}} \int_{\mathbb{R}^3} e^{i(2E - \sum_{i=1}^N \phi(v_i))z_3} e^{i(p_1 - \sum_{i=1}^N v_{i1})z_1} e^{i(p_2 - \sum_{i=1}^N v_{i2})z_2} dz_1 dz_2 dz_3 dV, \end{aligned}$$

where $dV = dv_1 \dots dv_N$. With $E = N$, the last equality is

$$Z(N, p_1, p_2) = \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} e^{ip_1 z_1 + ip_2 z_2} \left(e^{2iz_3} \int_{\mathbb{R}^2} e^{-i\phi(v_1)z_3 - iv_{11}z_1 - iv_{12}z_2} dv_{11} dv_{12} \right)^N dz_1 dz_2 dz_3.$$

Note that here p_1 and p_2 are assumed to be independent of N . A natural and straightforward variation is to replace p_1 by Np_1 and p_2 by Np_2 .

Let

$$\begin{aligned} q(z_1, z_2, z_3) &= e^{ip_1 z_1 + ip_2 z_2}, \\ S(z_1, z_2, z_3) &= 2iz_3 + \log \int_{\mathbb{R}^2} e^{-i\phi(v_1)z_3 - iv_{11}z_1 - iv_{12}z_2} dv_{11} dv_{12}, \end{aligned} \tag{44}$$

where $q : \mathbb{C}^3 \rightarrow \mathbb{R}$, $S : \mathbb{C}^3 \rightarrow \mathbb{R}$. With these notations, we can write

$$Z(N, p_1, p_2) = \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} q(z_1, z_2, z_3) e^{NS(z_1, z_2, z_3)} dz_1 dz_2 dz_3. \tag{45}$$

The right hand side of the last equality is a saddle point integral in dimension 3. The asymptotic behaviour of $Z(N, p_1, p_2)$ for large N is determined by the saddle points of $S(z_1, z_2, z_3)$, i.e., points $(\bar{z}_1, \bar{z}_2, \bar{z}_3)$ such that

$$\nabla S(\bar{z}_1, \bar{z}_2, \bar{z}_3) = 0. \tag{46}$$

Using (44), we need to solve the following system of equations:

$$2i - \frac{i \int_{\mathbb{R}^2} \phi(v_1) e^{-i\phi(v_1)\bar{z}_3 - iv_{11}\bar{z}_1 - iv_{12}\bar{z}_2} dv_{11} dv_{12}}{\int_{\mathbb{R}^2} e^{-i\phi(v_1)\bar{z}_3 - iv_{11}\bar{z}_1 - iv_{12}\bar{z}_2} dv_{11} dv_{12}} = 0, \tag{47}$$

$$- \frac{i \int_{\mathbb{R}^2} v_{11} e^{-i\phi(v_1)\bar{z}_3 - iv_{11}\bar{z}_1 - iv_{12}\bar{z}_2} dv_{11} dv_{12}}{\int_{\mathbb{R}^2} e^{-i\phi(v_1)\bar{z}_3 - iv_{11}\bar{z}_1 - iv_{12}\bar{z}_2} dv_{11} dv_{12}} = 0, \tag{48}$$

$$- \frac{i \int_{\mathbb{R}^2} v_{12} e^{-i\phi(v_1)\bar{z}_3 - iv_{11}\bar{z}_1 - iv_{12}\bar{z}_2} dv_{11} dv_{12}}{\int_{\mathbb{R}^2} e^{-i\phi(v_1)\bar{z}_3 - iv_{11}\bar{z}_1 - iv_{12}\bar{z}_2} dv_{11} dv_{12}} = 0. \tag{49}$$

Since ϕ is even, it follows that $\bar{z}_1, \bar{z}_2 = 0$ are the unique solutions to equations (48) and (49). We can now obtain \bar{z}_3 by solving the following equation:

$$\int_{\mathbb{R}^2} (2 - \phi(v_1)) e^{-i\phi(v_1)\bar{z}_3} dv_1 = 0. \tag{50}$$

Assuming that $(0, 0, \bar{z}_3)$ is the unique solution to (46), and that we can deform the integration domain in (45) to contain $(0, 0, \bar{z}_3)$, we find that

$$\begin{aligned} Z(N, p_1, p_2) &\sim \frac{1}{(2\pi)^3} \left(\frac{2\pi}{N} \right)^{3/2} \frac{1}{(\det S''((0, 0, \bar{z}_3)))^{1/2}} e^{NS(0, 0, \bar{z}_3)} q(0, 0, \bar{z}_3) \\ &= \frac{1}{(2\pi N)^{3/2}} \frac{1}{(\det S''((0, 0, \bar{z}_3)))^{1/2}} e^{2Ni\bar{z}_3} \left(\int_{\mathbb{R}^2} e^{-i\phi(v_1)\bar{z}_3} dv_1 \right)^N. \end{aligned} \tag{51}$$

By the discussion in Section 2, this procedure shows formally that the uniform distributions on $\Gamma^N(\sqrt{N}, p)$ with respect to the measure μ_{N, p_1, p_2} is $Ce^{-z_0\phi(v)}$ -chaotic, where $z_0 = i\bar{z}_3$ and \bar{z}_3 the unique solution to (50) and

$$C = \frac{1}{\int_{\mathbb{R}^2} e^{-z_0\phi(v)} dv}.$$

Sznitman’s method, referred to in Section 1 and the end of Section 2, is not restricted to the one-dimensional setting, but could formally be used also here:

with N spatial dimensions, we would have $d = N + 1$, and h in Equations (13) and (14) would be a generalized four-momentum, $h = (v_1, v_2, \dots, v_N, \phi(v_1, \dots, v_N))$. However, the same difficulty as in the one-dimensional case would appear here, and a modification of Sznitman's argument would be needed, in the same way.

Appendix. The saddle point method is used to determine the asymptotic behaviour of integrals depending on a parameter. For a detailed description we refer to [8]. Without proof we only state below the saddle point method which is concerned with this paper.

One-dimensional saddle point method

Let γ be a contour in the complex plane. Assume that q and S are analytic functions in a neighborhood of the contour γ . Consider the following integral

$$F(\lambda) = \int_{\gamma} q(z) e^{\lambda S(z)} dz. \quad (52)$$

A point $z_0 \in \mathbb{C}$ is called a simple *saddle point* of the function $S : \mathbb{C} \rightarrow \mathbb{C}$ if $S'(z_0) = 0$ and $S''(z_0) \neq 0$. Assume that $z_0 \in \gamma$ is the unique simple saddle point of S . Then as $\lambda \rightarrow \infty$

$$F(\lambda) = \sqrt{\frac{-2\pi}{\lambda S''(z_0)}} e^{\lambda S(z_0)} \left(q(z_0) + \mathcal{O}\left(\frac{1}{\lambda}\right) \right). \quad (53)$$

If there are more than one saddle point, F will be expressed as a sum over these points.

Many-dimensional saddle point method

Let γ be an N -dimensional smooth compact manifold. Consider the following integral

$$F(\lambda) = \int_{\gamma} q(z) e^{\lambda S(z)} dz. \quad (54)$$

where $z = (z_1, \dots, z_N) \in \mathbb{C}^N$ and the functions $q(z)$ and $S(z)$ are assumed to be analytic in a domain D containing γ . A point z_0 is called a simple saddle point of $S(z)$ if $\nabla S(z_0) = 0$ and $\det S''(z_0) \neq 0$. Assume that $z_0 \in \gamma$ is the unique simple saddle point of S . Then as $\lambda \rightarrow \infty$

$$F(\lambda) = \left(\frac{2\pi}{\lambda}\right)^{N/2} \frac{1}{(\det S''(z_0))^{1/2}} e^{\lambda S(z_0)} \left(q(z_0) + \mathcal{O}\left(\frac{1}{\lambda}\right) \right). \quad (55)$$

If there are more than one saddle point, F is expressed as a sum over these points.

Acknowledgments. The authors would like to thank one of the referees for several suggestions. D.M. acknowledges support by the Swedish Science Council. B.W. acknowledges support by the Swedish Science Council, the Knut and Alice Wallenberg foundation and the Swedish Foundation for Strategic Research.

REFERENCES

- [1] E. A. Carlen, M. C. Carvalho, J. Le Roux, M. Loss and C. Villani, [Entropy and chaos in the Kac model](#), *Kinet. Relat. Models*, **3** (2010), 85–122.
- [2] E. A. Carlen, P. Degond and B. Wennbrg, [Kinetic limits for pair-interaction driven master equations and biological swarm models](#), *Math. Models Methods Appl. Sci.*, **23** (2013), 1339–1376.
- [3] K. Carrapatoso, [Quantitative and qualitative Kac's chaos on the Boltzmann's sphere](#), *Ann. Inst. Henri Poincaré Probab. Stat.*, **51** (2015), 993–1039.

- [4] C. Cercignani and G. Medeiros Kremer, *The Relativistic Boltzmann Equation: Theory and Applications*, Birkhäuser Verlag, Basel, 2002.
- [5] J. T. Chang D. Pollard, *Conditioning as disintegration*, *Statist. Neerlandica*, **51** (1997), 287–317.
- [6] L. C. Evans and R. F. Gariepy, *Measure Theory and Fine Properties of Functions, Revised Edition*, CRC Press, 2015.
- [7] I. Gallagher, L. Saint-Raymond and B. Texier, *From Newton to Boltzmann: Hard Spheres and Short-Range Potentials*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2013.
- [8] R. V. Gamkrelidze, Integral representations and asymptotic methods, in *Encyclopaedia of Mathematical Sciences*, Springer-Verlag, 1989.
- [9] M. Kac, Foundations of kinetic theory, in *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability*, **3** (1956), 171–197.
- [10] O. E. Landford III, Time evolution of large classical systems, *Dynamical Systems, Theory and Applications*, **38** (1975), 1–111.
- [11] T. Lelièvre, M. Rousset and G. Stoltz, *Free Energy Computations*, Imperial College Press, London, 2010.
- [12] M. Pulvirenti, C. Saffirio and S. Simonella, *On the validity of the Boltzmann equation for short range potentials*, *Rev. Math. Phys.*, **26** (2014), 1450001, 64pp.
- [13] A.-S. Sznitman, *Topics in propagation of chaos*, in *École d'Été de Probabilités de Saint-Flour XIX—1989*, **1464** (1991), 165–251.
- [14] R. M. Strain, *Coordinates in the relativistic Boltzmann theory*, *Kinet. Relat. Models*, **4** (2011), 345–359.
- [15] R. M. Strain and S.-B. Yun, *Spatially homogeneous Boltzmann equation for relativistic particles*, *SIAM J. Math. Anal.*, **46** (2014), 917–938.
- [16] M. Toda, R. Kubo and N. Saitô, *Statistical Physics I, Equilibrium Statistical Mechanics*, 2nd edition, Springer Verlag, 1992.

Received July 2015; revised April 2016.

E-mail address: dawan.dara@gmail.com

E-mail address: wennberg@chalmers.se