

Utah State University

DigitalCommons@USU

---

Memorandum

US/IBP Desert Biome Digital Collection

---

1979

## Seed-Eating Birds

H. R. Pulliam

Follow this and additional works at: [https://digitalcommons.usu.edu/dbiome\\_memo](https://digitalcommons.usu.edu/dbiome_memo)



Part of the [Earth Sciences Commons](#), [Environmental Sciences Commons](#), and the [Life Sciences Commons](#)

---

### Recommended Citation

Pulliam, H.R. 1979. Seed-Eating Birds. U.S. International Biological Program, Desert Biome, Utah State University, Logan, Utah. Final Progress Reports, Process Studies, RM 77-16.

This Article is brought to you for free and open access by the US/IBP Desert Biome Digital Collection at DigitalCommons@USU. It has been accepted for inclusion in Memorandum by an authorized administrator of DigitalCommons@USU. For more information, please contact [digitalcommons@usu.edu](mailto:digitalcommons@usu.edu).



**FINAL REPORT**

**SEED-EATING BIRDS**

\*H. R. Pulliam  
Museum of Northern Arizona

(\*now at Northern Arizona University)

**US/IBP DESERT BIOME  
RESEARCH MEMORANDUM 77-16**

in

**FINAL PROGRESS REPORTS**  
Process Studies, pp. 91-93

Proposal No. 2.3.2.5

**Printed 1979**

The material contained herein does not constitute publication.  
It is subject to revision and reinterpretation. The author(s)  
requests that it not be cited without expressed permission.

Citation format: Author(s). 1979. Title.  
US/IBP Desert Biome Res. Memo. 77-16.  
Utah State Univ., Logan. 3 pp.

Utah State University is an equal opportunity/affirmative action  
employer. All educational programs are available to everyone  
regardless of race, color, religion, sex, age or national origin.

Ecology Center, Utah State University, Logan, Utah 84322

## ABSTRACT

The objective of this study was to understand the effects of seed availability on the diets of sparrows and their population density. Seed availability was estimated from random soil samples and birds were censused weekly; sparrows captured from sample sites were forced to regurgitate. Between January 19 and March 2 seed abundance decreased from  $150 \times 10^6$  seeds/ha to  $121 \times 10^6$  seeds/ha, while the number of seed species increased from 51 to 57. A decrease in abundance occurred in most species, but *Ipomea* spp. showed a  $10^7$  seeds/ha increase. The observed diets were not the diets expected if the seed species consumed were eaten in the proportions available. By considering sparrow density as a function of grass seed production, we can assume that the birds eat most of the seeds available only in years when grass seed production is very low.

## INTRODUCTION

The study of seed-eating birds at the Silverbell IBP site had two primary goals. These were: 1) to understand how the availability of seeds affects the diets of sparrows, and 2) to ascertain how seed availability might regulate the population density of sparrows. It was hoped that the data would also bear on the question of the overall impact of granivorous birds on the seed reserves in the soil.

The study was funded for only the last year of the IBP program. In my original proposal, I pointed out that a one-year study would not be adequate to answer these questions but that something could be learned by comparison with my similar work at The Research Ranch (TRR). As it turned out, the year of the study had unusually low seed production and sparrow populations were too low at Silverbell to obtain diet samples. I therefore continued as planned to census granivorous birds and measure seed availability at Silverbell but I shifted my diet studies to the grasslands at TRR where wintering sparrows were more abundant. As the original plan had been to study the diet of Brewer's Sparrow (*Spizella breweri*) at Silverbell, I studied instead the closely related Chipping Sparrow, *Spizella passerina*.

The data on sparrow abundances and seed availability from the IBP site is analyzed for the one study year. In order to get an approximate answer to question 2, I compared this data to the several years of similar data from my studies at TRR.

## METHODS

As detailed descriptions of the methods and results will appear in published reports, I shall only outline them here.

At the Silverbell site, six study plots were chosen, three each in the two major plant associations. Each plot was 500 x 50 m. Birds were censused weekly using the standard Emlen census technique from a transect along the main axis of each plot. Seed availability in the soil was estimated from soil samples (each 10 x 10 x 0.5 cm) sampled at random (stratified) from one habitat in each plant association. All seeds were separated from the soil, identified and counted in my laboratory. Thus, the primary data consist of seed availability and bird abundance through an annual cycle.

The sparrow diet study plot was a 400 x 70 m area in an oak woodland at TRR. Birds were mist-netted on the same site

where soil samples were collected for analysis of seed availability. Each bird was given 0.5 cc of the emetic tincture of digitalis to cause it to regurgitate its crop contents. On each soil sampling date 28 soil samples were collected at random from the entire study plot.

## RESULTS

On January 19, there were approximately  $150 \times 10^6$  seeds/ha of 51 species at TRR. Between that date and March 2 many seed species decreased in abundance (Table 1), though *Ipomea* spp. showed an increase of about  $10^7$  seeds/ha. On March 2, total seed abundance was approximately  $121 \times 10^6$  seeds/ha, representing 57 species. All species of seeds which had more than an average of one seed per soil sample (100 cm<sup>2</sup>) on either sample date are shown in Table 1 along with a few species which were less common but showed up in Chipping Sparrow diets.

Table 1. The common seed species from soil samples taken January 19 and March 2, 1975 plus a few rare seed species which occurred in sparrow diets

Species	January 19		March 2	
	Seeds/100 cm <sup>2</sup>	$\sigma^2$ /mean	Seeds/100 cm <sup>2</sup>	$\sigma^2$ /mean
<i>Amaranthus retroflexus</i>	3.57 ± 2.49	48.4	2.18 ± 1.15	17.1
<i>Aristida divaricata</i>	1.11 ± 0.40	4.0	1.50 ± 0.42	3.3
<i>A. humulosa</i>	3.89 ± 2.50	45.1	0.54 ± 0.27	3.9
<i>Bidens leptoccephala</i>	1.04 ± 0.60	9.8	5.82 ± 3.84	70.9
<i>Ruellia chondrosioides</i>	0.93 ± 0.61	11.4	0.07 ± 0.05	1.0
<i>B. curtipendula</i>	2.32 ± 0.56	3.8	2.04 ± 0.67	6.1
<i>B. gracilis</i>	0.14 ± 0.07	0.9	0.46 ± 0.36	7.7
<i>Cassia leptadenia</i>	18.00 ± 6.07	57.4	18.61 ± 4.03	24.4
<i>Convolvulus</i> sp.	0.00		1.57 ± 0.57	5.7
<i>Croton</i> sp.	0.2 ± 0.22	4.6	0.86 ± 0.62	12.4
<i>Desmodium batocaulon</i>	3.32 ± 1.32	14.7	0.68 ± 0.34	4.8
<i>Diodia teres</i>	0.36 ± 0.26	5.2	0.43 ± 0.43	12.0
<i>Eragrostis lehmannia</i>	19.32 ± 4.74	32.7	19.57 ± 6.38	58.3
<i>Eriochloa gracilis</i>	0.25 ± 0.22	5.2	0.00	
<i>Euphorbia hyssopifolia</i>	7.75 ± 1.90	13.1	0.79 ± 0.35	4.5
<i>Ipomea</i> spp.	4.71 ± 1.32	10.3	14.14 ± 2.28	10.3
<i>Leptochloa dubia</i>	0.00		0.82 ± 0.78	21.0
<i>Lycopus phleoides</i>	2.07 ± 1.85	46.4	0.00	
<i>Mollugo verticillata</i>	37.64 ± 15.02	168.0	17.79 ± 6.12	59.9
<i>Panicum obtusum</i>	0.89 ± 0.31	3.0	2.54 ± 0.83	7.7
<i>Paspalum</i> sp.	0.00		1.25 ± 0.46	4.7
<i>Polanisia</i> sp.	0.89 ± 0.30	2.9	2.18 ± 0.81	8.4
<i>Portulaca oleracea</i>	1.46 ± 0.36	2.5	1.07 ± 0.25	1.7
<i>P. retusa</i>	28.61 ± 9.24	83.5	13.36 ± 2.89	17.5
<i>Rumex</i> sp.	1.14 ± 0.39	3.7	0.71 ± 0.21	1.8
<i>Schkuhria waltiana</i>	1.68 ± 0.72	8.6	1.96 ± 1.11	17.5
<i>Sida procumbens</i>	0.54 ± 0.24	3.1	0.61 ± 0.27	3.3
<i>Trichachne californica</i>	0.04 ± 0.04	1.0	0.00	
Unidentified E#64	1.29 ± 0.59	7.7	1.82 ± 0.75	8.7
Unidentified I#4	0.00		1.54 ± 1.02	19.1
Total	143.25		114.91	
25 other species	6.32		6.03	
Total seeds/100 cm <sup>2</sup>	149.57		120.94	

The variance/mean ratios shown in Table 1 indicate that the seeds were very clumped. These ratios do not rule out the possibility that the seeds were randomly distributed over a portion of the study site but absent from the rest of the site. Therefore, I compared the distribution of seed abundances to that expected for a Poisson with added zeros and found a poor fit with means still larger than variances for all common species. Thus, most species exhibited clumped distributions and could be adequately fitted to a negative binomial distribution.

The observed diets of Chipping Sparrows are shown in Table 2. Table 3 compares the observed diets to that expected if the sparrows were eating the seed species observed in their diets but eating them in the proportions available. It is clear that even among these species the diet is not a random subset of what is available.

Figure 1 shows the sparrow densities at TRR and Silverbell as functions of the grass seed production. The circles are data from TRR taken in two habitats over the years 1972-1976. It can be seen that the birds eat most of the seeds available (i.e., food limitation) only in years when grass seed production is very low. I have only a crude estimate of grass seed production at Silverbell; my methods were designed to study seed availability, not seed production. The two data points indicate that the Silverbell sites fall in line with the TRR results. However, this would have to be confirmed by estimating seed production and sparrow abundance over a variety of years, particularly when seed production is high.

Figure 1. Relation between sparrow density and available food at Silverbell (triangle) and The Research Ranch sites (circles). See text for discussion of food limitation hypothesis.

Table 2. Observed diets of Chipping Sparrows foraging in an oak woodland habitat and seed weights

Species	Seed wt (mg)	January 10	March 1
		Average seeds/sample ± S.E.	Average seeds/sample ± S.E.
<i>Amaranthus retroflexus</i>	0.05	14.44 ± 4.35 (28.3%)	6.50 ± 1.63 (13.8%)
<i>Aristida</i> spp.	1.00	1.07 ± 0.61 (2.1%)	0.44 ± 0.32 (0.9%)
<i>Bouteloua chondrostoides</i>	1.38	0.33 ± 0.27 (0.6%)	0
<i>B. curtipendula</i>	0.63	0.63 ± 0.32 (1.2%)	0.88 ± 0.81 (1.9%)
<i>Croton</i> sp.	0.50	0	0.06 ± 0.06 (0.1%)
<i>Eragrostis lehmanniana</i>	0.11	22.22 ± 6.49 (43.5%)	12.00 ± 4.49 (25.4%)
<i>Leptochloa dubia</i>	1.18	0	4.06 ± 2.83 (8.6%)
<i>Lycurus phleoides</i>	0.20	0.15 ± 0.15 (0.3%)	0.06 ± 0.06 (0.1%)
<i>Mollugo verticillata</i>	0.06	0.37 ± 0.23 (0.7%)	1.94 ± 0.88 (4.1%)
<i>Panicum obtusum</i>	1.00	1.22 ± 0.54 (2.4%)	0.13 ± 0.13 (0.3%)
<i>Paspalum</i> sp.	1.18	0	1.56 ± 1.03 (3.3%)
<i>Portulaca</i> spp.	0.22	10.00 ± 2.48 (19.6%)	12.56 ± 5.59 (26.6%)
<i>Sohkukhia wislizeni</i>	1.20	0	0.06 ± 0.06 (0.1%)
<i>Talinum</i> sp.	1.30	0	1.81 ± 1.15 (3.8%)
<i>Trichachne californica</i>	0.37	0.15 ± 0.12 (0.3%)	0
Unknown E#64	1.33	0	4.88 ± 3.02 (10.3%)
3 unidentified species	--	0.48 ± 0.27 (0.9%)	0.25 ± 0.15 (0.5%)
Total seeds/sample		51.06	47.19
Total number samples		27	16




Table 3. Expected and observed frequencies of seeds included in the diet. Expected values are from the frequencies of the seed species in soil samples

Species	January 19 sample			March 2 sample		
	Percent available	Expected number in diet	Observed number in diet	Percent available	Expected number in diet	Observed number in diet
<i>Amaranthus retroflexus</i>	3.4	46.4	390	3.2	23.4	104
<i>Aristida</i> spp.	4.7	63.8	29	3.0	21.9	7
<i>Bouteloua chondrostoides</i>	0.9	12.1	9	0.1	0.8	0
<i>B. curtipendula</i>	2.2	30.2	17	3.0	21.9	14
<i>Croton</i> sp.	0.3	3.8	0	1.3	9.2	1
<i>Eragrostis lehmanniana</i>	18.4	251.2	600	29.0	209.8	192
<i>Leptochloa dubia</i>	0.0	0.0	0	1.2	8.8	65
<i>Lycurus phleoides</i>	2.0	27.6	4	0.0	0.0	1
<i>Mollugo verticillata</i>	35.8	489.5	10	26.4	190.7	31
<i>Panicum obtusum</i>	0.8	11.6	33	3.8	27.2	2
<i>Paspalum</i> sp.	0.0	0.0	0	1.9	13.4	25
<i>Portulaca</i> spp.	28.6	391.0	270	21.4	154.7	201
<i>Sohkukhia wislizeni</i>	1.6	21.8	0	2.9	21.0	1
<i>Trichachne californica</i>	0.0	0.5	4	0.0	0.0	0
Unknown E#64	1.2	16.8	0	2.7	19.5	78