

OCCURRENCE OF RARE CADDISFLY (TRICHOPTERA) SPECIES AT THE CATCHMENT AREA OF LAKE BALATON**I. SZIVÁK^{1*} – A. MÓRA²**¹University of Pécs, Department of General and Applied Ecology, H-7624 Pécs, Ifjúság útja 6, Hungary, *Corresponding author, e-mail: szivaki@gamma.ttk.pte.hu²Hungarian Academy of Sciences, Balaton Limnological Research Institute, H-8237 Tihany, Klebelsberg Kuno út 3, Hungary**RITKA TEGZESFAJOK ELŐFORDULÁSA A BALATON VÍZGYŰJTŐJÉN (TRICHOPTERA)****SZIVÁK ILDIKÓ¹ – MÓRA ARNOLD²**¹PTE TTK KTI Általános és Alkalmazott Ökológiai Tanszék, 7624 Pécs, Ifjúság útja 6.²MTA Balatoni Limnológiai Kutatóintézete, 8237 Tihany, Klebelsberg Kuno út 3.

ABSTRACT: Although the caddisfly fauna of previously undiscovered regions in Hungary have adequately been explored, there are many regions (e.g. Great Hungarian Plain) and water types (e.g. backwaters of rivers, springs) whose caddisfly fauna is poorly known so far. In addition, occurrence of new species can be recorded in relatively well known regions as well. Although caddisfly fauna of catchment area of Lake Balaton is relatively well explored, during the most recent intensive faunistical investigations several rare and endangered caddisfly species were found. In this paper a compilation of formerly published results and new larval data are given on the following rare caddisfly species in Hungary: *Synagapetus krawanyi*, *Hydropsyche siltalai*, *Apatania muliebris*, *Limnephilus elegans*, *Silo nigricornis*, *Adicella reducta*, *Beraea maurus*.

Key words: endangered species, springs, *Synagapetus krawanyi*, *Hydropsyche siltalai*, *Apatania muliebris*, *Limnephilus elegans*, *Silo nigricornis*, *Adicella reducta*, *Beraea maurus*

KIVONAT: Habár a korábban kevéssé ismert hazai régiók tegzesfaunája mára már kielégítően feltárt, még mindig vannak olyan területek (pl.: Nagy Alföld) és vízterületek (pl.: folyók holtágai, források), melyek tegzesfaunájáról igen kevés információval rendelkezünk. Emellett a viszonylag alaposan feltárt területeken is kerülnek elő új fajok. A Balaton vízgyűjtőjén a legújabb, intenzív faunisztkai munkáink során számos ritka és veszélyeztetett tegzesfaj új előfordulását sikerült kimutatni annak ellenére, hogy a terület tegzesfaunája jól ismertnek tekinthető. Dolgozatunkban átfogó irodalmi áttekintést adunk és új előfordulási adatokat közlünk az alábbi, hazánkban ritka tegzesfajokról: *Synagapetus krawanyi*,

Hydropsyche siltalai, *Apatania muliebris*, *Limnephilus elegans*, *Silo nigricornis*,
Adicella reducta, *Beraea maurus*.

Kulcsszavak: veszélyeztetett fajok, források, *Synagapetus krawanyi*,
Hydropsyche siltalai, *Apatania muliebris*, *Limnephilus elegans*, *Silo nigricornis*,
Adicella reducta, *Beraea maurus*

Introduction

Until the end of 1970's the examination of the Hungarian caddisfly (Trichoptera) fauna was insufficient, with about 150 species recorded from Hungary by that time (NÓGRÁDI and UHERKOVICH 2002a). Furthermore caddisfly fauna of most regions of the country were poorly known or totally unknown. At the beginning of the 1980's, intensive collecting works were started, resulted in that 211 species are known from Hungary up to now (NÓGRÁDI and UHERKOVICH 2002a). Caddisfly fauna of previously undiscovered regions have adequately been explored and new important data have been available on ecology and distribution of many Hungarian species (NÓGRÁDI and UHERKOVICH 2002a). However, there are many regions (e.g. Great Hungarian Plain) and water types (e.g. backwaters of rivers, springs) whose caddisfly fauna is poorly known so far. In addition, occurrence of new species can be recorded in relatively well known regions as well.

The majority of our knowledge on the Hungarian caddisfly fauna is based on collection of imagines (NÓGRÁDI and UHERKOVICH 2002a and see other references in the mentioned publication), and few faunistical investigations were carried out on the basis of larvae (e.g. MÓRA and CSABAI 2002a, 2002b; MÓRA et al. 2006). It was caused by the lack of useful and exact identification keys for larvae. Collection of imagines can be no representative for individual waters, because the specimens captured by light traps could come far from their larval habitats. Thus collection of larvae is important when the aim is to explore the caddisfly fauna of an individual water (and not only a certain area). Of course, the collection of larvae is not adequate to all aspects. Larvae of some species are not distinguishable by morphological characters, and fewer species can be found arising from the nature of the method (SCHMERA and KISS 2000). Despite of these deficiencies, investigations on larvae is most appropriate to explore the accurate distribution of a certain species.

Caddisfly fauna of catchment area of Lake Balaton is relatively well explored based on collecting adults (NÓGRÁDI and UHERKOVICH 1994) as well as larvae (KÁLMÁN et al. 2006; MÓRA et al. 2007, 2008). During the most recent intensive faunistical investigations several rare and endangered caddisfly species were found in this region. Distributional data for *Hydropsyche siltalai* and *Limnephilus elegans* were already published (MÓRA et al. 2007, 2008), but due to the rarity of these species we also emphasize their occurrence in this work.

Materials and methods

Caddisfly larvae were captured by pond-net and manual selection during faunistical investigations between 2006 and 2008. Furthermore, quantitative samplings were carried out according to the AQEM protocol (AQEM CONSORTIUM

2002; HERING et al. 2004) in 2007 and 2008. Keys by WALLACE et al. (1990) and WARINGER and GRAF (1997) were used for identification of larvae. Distributional and ecological informations are given for all species on the basis of NÓGRÁDI and UHERKOVICH (2002a) and GRAF et al. (2002).

In Table 1 names of sampling sites are given with their administrative units in brackets, the accurate geographical co-ordinates (WGS-84) and the 10×10 km UTM-grid codes. In cases of some geographical terms we left the original Hungarian form for the localities being more identifiable: árok = valley, erdészet = forestry, forrás = spring, gátőrház = dike watchman's house, hegység = mount/hill, kert = garden, kút = spring, part(ja) = bank, patak = stream, víz = stream, völgy = valley.

All known literature data are given with the localities and the publications where they were mentioned. New records are given as follows: localities (with administration unit), date of sampling, total number of individuals and abbreviations of collectors' names in alphabetical order: IK – Katalin IHÁSZ, MA – Arnold MÓRA, SzA – András SZIVÁK, Szi – Ildikó SZIVÁK, except in cases of *Hydropsyche siltalai* and *Limnephilus elegans*.

Table 1. Sampling sites for new records at the catchment area of Lake Balaton with exact geographical co-ordinates (WGS-84) and 10×10 km UTM grid codes

Name of the sampling sites	Lat. (N)	Lon. (E)	UTM
1. Örvényesi-séd, Zádor-hegy (Pécsely)	46°58'14"	17°46'34"	YN 10
2. Széles-forrás (Monostorapáti)	46°56'06"	17°35'02"	XN 90
3. Tetves-patak (Védapuszta)	46°36'11"	17°48'22"	YM 16
4. Zádor-kút (Pécsely)	46°58'31"	17°46' 6"	YN 10
5. Zala (Csöde)	46°50'21"	16°32'30"	XM 18
6. Zala (Pókaszepetk)	46°55'21"	16°58'15"	XM 59
7. Zala (Zalaegerszeg)	46°51'02"	16°50'44"	XM 49
8. Zala (Zalalővő)	46°50'57"	16°37'36"	XM 18
9. Zala (Zalaszentgyörgy)	46°52'10"	16°42'43"	XM 29
10. Zala, Szentgrót (Zalaszentgrót)	46°56'33"	17°04'16"	XN 50
11. Zala, Zalakoppány (Zalaszentgrót)	46°53'15"	17°05'14"	XM 59

Results

Synagapetus krawanyi (Ulmer, 1938)

Literature data: Kisújbánya: Pásztor-forrás, Vár-völgy (NÓGRÁDI 1984, 1987b); Kőszeg: Hármas-patak (NÓGRÁDI and UHERKOVICH 1989); Magyaregregy: Egregyi-völgy (UHERKOVICH and NÓGRÁDI 2006); Óbánya (NÓGRÁDI 1987b; NÓGRÁDI and UHERKOVICH 1991); Pécs: Melegmányi-völgy, Nagy-Mély-völgy (UHERKOVICH and NÓGRÁDI 2006); Vékény: Iharos-kút, Vár-völgy (NÓGRÁDI 1984, 1987b; UHERKOVICH and NÓGRÁDI 2006); Velem: Hosszú-völgy, Szent-Vid-patak (NÓGRÁDI and UHERKOVICH 1989).

New record: Széles-forrás (Monostorapáti): 20/04/2008, 8I, IK-MA-Szi.

Synagapetus krawanyi has narrow distribution in Europe occurring in Eastern Alps and Southern Europe. Hungarian populations show southeastern Alpine connection with the most southeastern occurrence in Mecsek Mountains. Previously no data were known between the Alps (Kőszegi Mountains) and Mecsek Mountains (Fig. 1). Larvae live in xeno- and oligosaprofic springs and small brooklets. *S. krawanyi* is regarded as endangered caddisfly species in Hungary (NÓGRÁDI and UHERKOVICH 1999, 2002a).

***Hydropsyche siltalai* Döhler, 1963**

Literature data: Bükkösd: Szentdomján (UHERKOVICH and NÓGRÁDI 2006); Dobri: Kerka (UHERKOVICH 2004); Gyepükaján: Meleg-víz (UHERKOVICH and NÓGRÁDI 1999); Káptalanfa: Kígyós-patak (UHERKOVICH and NÓGRÁDI 1999); Kercaszomor: Kerca (UHERKOVICH 2004); Kőszeg: Chernel-kert, Keresztkút, Malom-árok (NÓGRÁDI 1988; NÓGRÁDI and UHERKOVICH 1992); Magyarföld: Kerka (UHERKOVICH 2004); Nagyvisnyó: Nagy-völgy (Kiss 1991; Kiss et al. 2002a; NÓGRÁDI et al. 1996); Órtilos: Dráva-part, (UHERKOVICH 2005); Szalafő (NÓGRÁDI and UHERKOVICH 1994); Szentpéterfölde: Erdészet, Szécsisziget: Kerka (NÓGRÁDI and UHERKOVICH 2002b); Velem (NÓGRÁDI 1988); Vízvár: Dráva-part (UHERKOVICH 2005).

New records: Zala [Pókaszepetk, Zalaegerszeg, Zalalövő, Zalaszentgyörgy, Szentgrót (Zalaszentgrót), Zalakoppány (Zalaszentgrót)] (for details see MÓRA et al. 2008)

Hydropsyche siltalai is widely distributed in Europe, but only sporadic data are known from Eastern and Central Europe. In Hungary, it occurs mostly in the Transdanubian region, where can locally be frequent (Fig. 2). Some records are known from Bükk Mountains as well. Larvae of *H. siltalai* live in oligo- and mesosaprobic streams. In Hungary, *H. siltalai* belongs to endangered species (NÓGRÁDI and UHERKOVICH 1999, 2002a). However, according to our recent investigations, *H. siltalai* seems to be a dominant *Hydropsyche* species in the middle reaches of Zala River (see MÓRA et al. 2008).

***Apatania muliebris* McLachlan, 1866**

Literature data: Kám: Jeli Arborétum (NÓGRÁDI 1994).

New records: Zádor-kút (Pécsely): 04/10/2006 – 19/10/2007, 150l, SzA-Szl.

Apatania muliebris is a parthenogenetic species due to only females are known. Several forms and subspecies were described for this morphologically various species, sometimes under synonymous species names (BARNARD and O'CONNOR 1987; SOLEM 1985). It is widely distributed in Northern Europe and the Alps, with the localities in Tyrol (Austria) being closest to Hungary. In Hungary, three females were previously captured at 'Hét-forrás' spring of the Jeli Arborétum, near Kám village (NÓGRÁDI 1994) (Fig. 3). SÁTORI (1939) mentioned *Apatania fimbriata* (Pictet, 1834) from a spring near Balatonhenye, but voucher specimens for this species are not available. However, this old data likely apply to *A. muliebris*. JUHÁSZ and KOVÁCS (1997) recorded *A. muliebris* from Jósva (Szinpetri, NE Hungary), but no voucher specimens are available for this record. Due to the conditions of the collecting sites, *A. muliebris* could unlikely occur in that stream. Larvae live in cool, unpolluted springs with permanent temperature and low concentrations of inorganic ions and organic matter. In early spring, larvae of this species were found in "Zádor-kút" spring (near Pécsely) in high abundance. *A. muliebris* is a postglacial relict, endangered species and protected by law in Hungary (NÓGRÁDI and UHERKOVICH 1999, 2002a).

***Limnephilus elegans* Curtis, 1834**

Literature data: Bükk-hegység: Disznóskút (Kiss 1978a, 1978b, 1979; Kiss et al. 2003); Györzámoly: Duna, Patkányosmajor, gátőrház (NÓGRÁDI 1994); Lipótfa (NÓGRÁDI 1988); Magyarszombatfa (UJHELYI 1981, NÓGRÁDI 1986).

New record: Tetves-patak (Vadépuszta) (see MÓRA et al. 2007).

Limnephilus elegans is a widespread species in Western and Northern Europe, with its most eastern occurrence in Hungary. It was firstly reported as larvae from Bükk Mountains, NE Hungary (Kiss 1978a), while adults were collected in the same year from a *Sphagnum* swamp at Magyarszombatfa, W Hungary (UJHELYI 1981) (Fig. 4).

Habitats of the species are vulnerable such *L. elegans* is endangered and protected by law in Hungary (NÓGRÁDI and UHERKOVICH 1999, 2002a).

***Silo nigricornis* (Pictet, 1834)**

Literature data: Felsőtárkány: Egeres-völgy, Toldi-kút, Vörös-kő-völgy, (Kiss 1984a, 1984b, 1991; NÓGRÁDI et al. 1996); Gyepükaján: Meleg-víz (UHERKOVICH and NÓGRÁDI 1999); Halászi: Mosoni-Duna partja (UHERKOVICH and NÓGRÁDI 2004); Lillafüred: Garadna-patak (SÁTORI 1938); Parádfürdő: Ilona-völgy (Kiss 1980, 1981); Púski: Zátönyi-Holt-Duna (NÓGRÁDI 2001); Szigetköz: Cikolasziget, gátórház (UHERKOVICH and NÓGRÁDI 2003); Vízvár: Dráva part (NÓGRÁDI 2001, NÓGRÁDI and UHERKOVICH 2001).

New record: Széles-forrás (Monostorapáti): 20/04/2008, 1l, IK-MA-Szl.

Silo nigricornis can be found all over the Palaearctic region, but sporadically distributed in Central Europe. It is a rare caddisfly species in Hungary (Fig. 5), but can locally be relatively frequent (UHERKOVICH and NÓGRÁDI 2003). Larvae of *S. nigricornis* inhabit hyporectal and rithral zones of streams. According to its rarity and vulnerable habitats, *S. nigricornis* is regarded as endangered species in Hungary (NÓGRÁDI and UHERKOVICH 1999, 2002a).

***Adicella reducta* (McLachlan, 1865)**

Literature data: Aszófő (UJHELYI 1971); Kercaszomor: Kerca (UHERKOVICH 2004).

New record: Zala (Csöde): 30/07/2008, 3l, AQEM.

Adicella reducta is widely distributed in Europe, but true rarity in Hungary due to only two adult males have been captured so far (UHERKOVICH 2004, UJHELYI 1971) (Fig. 6). *Adicella reducta* was also mentioned in some other publication (NÓGRÁDI 1989b, NÓGRÁDI and UHERKOVICH 1989, NÓGRÁDI 1995), but due to a revision, all of these specimens proved to be *Adicella filicornis* (Pictet, 1834) (see NÓGRÁDI and UHERKOVICH 2002a). *A. reducta* larvae live in oligo- and β-mesosaprobic, cobble covered mountain streams. According to its rarity, conservation status of the species has been unknown (NÓGRÁDI and UHERKOVICH 1999, 2002a), but the new larval record suggests that *A. reducta* is an endangered species in Hungary.

***Beraea maurus* (Curtis, 1834)**

Literature data: Bükk-hegység: Hárskút, Répáshutai-forráspata (Kiss 1979); Jósvafő: Nagy-Tohonya-forrás (OLÁH 1967); Kőszeg: Hármasháza-patak, Hét-forrás (NÓGRÁDI and UHERKOVICH 1989, 1992); Mátraháza (NÓGRÁDI 1989a); Szilvásvárad: Szalajka-völgy (Kiss 1991; Kiss et al 2002a, 2002b; NÓGRÁDI et al. 1996); Szőce (NÓGRÁDI 1987a, 1989b).

New records: Örvényesi-séd, Zádor-hegy (Pécsely): 04/10/2006 – 19/10/2007, 64l, SzA-Szl.

Beraea maurus is a widespread species in Europe, but rare in Hungary, especially in the Transdanubian region (Fig. 7). During our investigations, larvae were collected in relatively high abundance at upper reaches of Örvényesi-creek (near Pécsely), in late spring and early summer. Larvae of this species prefer xeno- and oligosaprobic waters in ecrenal and hyporectal regions of streams. *B. maurus* belongs to endangered species in Hungary (NÓGRÁDI and UHERKOVICH 1999, 2002a).




Fig. 1. Distribution of *Synagapetus krawanyi* (Ulmer, 1938) in Hungary (point: literature data, square: new data)




Fig. 2. Distribution of *Hydropsyche siltalai* Döhler, 1963 in Hungary (point: literature data, square: new data)




Fig. 3. Distribution of *Apatania muliebris* McLachlan, 1866 in Hungary (point: literature data, square: new data)




Fig. 4. Distribution of *Limnephilus elegans* Curtis, 1834 in Hungary (point: literature data, square: new data)




Fig. 5. Distribution of *Silo nigricornis* (Pictet, 1834) in Hungary (point: literature data, square: new data)




Fig. 6. Distribution of *Adicella reducta* (McLachlan, 1865) in Hungary (point: literature data, square: new data)




Fig. 7. Distribution of *Beraea maurus* (Curtis, 1834) in Hungary (point: literature data, square: new data)

Discussion

All examined species are sporadically distributed in Hungary. It is likely due to a narrow range of environmental factors are suitable for them (NÓGRÁDI and UHERKOVICH 2002a, GRAF et al. 2002). However, our knowledge on ecology of these species is still poor and needs further investigations. In all probability small and vulnerable populations of these species can only be found in Hungary, such all new records are essential information for conservation of these caddisflies. Four of the seven species were captured in and/or at very close to springs. The springs with their cool and clear water and low trophical status represent special habitats and are rather different from the other sections of streams (VANNOTE et al. 1980). Although springs are characterized by unique assemblages, as it is also suggested by the new records of the rare and specialized caddisfly species, but they are inadequately investigated in Hungary. Springs are vulnerable habitats, such further investigations on them are very important tasks for future hydrobiological studies.

Acknowledgements: We thank Dr. Júlia Török, Katalin Ihász and András Szivák for their comments and fieldwork. This study was financially supported partly by the Hungarian National Science Fund (OTKA K69033).

References

AQEM CONSORTIUM (2002): Manual for the application of the AQEM method. A comprehensive method to assess European streams using benthic

- macroinvertebrates, developed for the purpose of the Water Framework Directive. Version 1.0.
- BARNARD, P.C. – O'CONNOR, J.P. (1987): The populations of *Apatania muliebris* McLachlan in the British Isles (Trichoptera: Limnephilidae). – Entomologist's Gazette 38: 263–267.
- GRAF, W. – GRASSER, U. – WARINGER, J. (2002): Trichoptera (caddisflies). Part III. In: MOOG, O. (ed.): Fauna Aquatica Austriaca, Edition 2002. – Wasserwirtschaftskataster, Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft, Wien, 40 pp.
- HERING, D. – MOOG, O. – SANDIN, L. – VERDONSCHOT, P. F. M. (2004): Overview and application of the AQEM assessment system. – Hydrobiologia 516: 1–20.
- JUHÁSZ, P. – KOVÁCS, T. (1997): A szinpetri vízimalom létesítésével kapcsolatos zoológiai vizsgálatok eredményei (Hirudinoidea, Gastropoda, Bivalvia, Amphipoda, Odonata, Trichoptera, Pisces). – Manuscript, Research report, Gyöngyös.
- KÁLMÁN, A. – PÁLL-GERGELY, B. – CSER, B. – BODA, P. – CSABAI, Z. (2006): Makroszkópikus vízi gerinctelenek faunisztikai vizsgálata a Déli-Bakony és a Balaton-felvidék víztereiben. – Hidrológiai Közlöny 86: 161–164.
- Kiss, O. (1978a): A *Limnephilus elegans* Curtis (Trichoptera) előfordulása Magyarországon. – Folia entomologica hungarica 31/1: 227–228.
- Kiss, O. (1978b): A Bükk hegységi Disznóskút és Sebesvíz Trichoptera együtteseiről. – Acta Academiae Paedagogicae Agriensis, nova series 14: 493–507.
- Kiss, O. (1979): The Trichoptera of the Bükk Mountains. – Acta biologica debrecina 16: 45–55.
- Kiss, O. (1980): The Trichoptera in the Ilona Valley of the Mátra Mountains (Hungary). – Folia Historico-naturalia Musei Matraensis 6: 115–125.
- Kiss, O. (1981): Trichoptera in the Ilona stream of the Mátra Mountains, North Hungary. In: MORETTI, G.P. (ed.): Proceedings of the 3rd International Symposium on Trichoptera. – Series entomologica 20: 129–138.
- Kiss, O. (1984a): Fénycsapdával gyűjtött Trichopterák a Bükk hegységi Vöröskő-völgyből. – Acta Academiae Paedagogicae Agriensis, nova series 18: 709–717.
- Kiss, O. (1984b): Trichoptera in an intermittent rill of the Bükk Mountains, North Hungary. In: MORSE, J.C. (ed.): Proceedings of the 4th International Symposium on Trichoptera. – Dr. W. Junk, The Hague, pp. 191–195.
- Kiss, O. (1991): Trichoptera from a light trap in the Bükk Mountains, North Hungary, 1980–1988. In: TOMASZEWSKI, C. (ed.): Proceedings of the 6th International Symposium on Trichoptera. – Adam Mickiewicz University Press, Poznań, pp. 233–236.
- Kiss, O. – ANDRIKOVICS, S. – SZABÓ, T. – MOGYORÓSI, A. (2002a): Functional feeding groups of Trichoptera along streams typical of north Hungary. – Nova Supplementa Entomologica 15: 529–536.
- Kiss, O. – SCHMERA, D. – FEHÉR, I. (2003): Characteristics of caddis larvae assemblages from shallow lakes in the Bükk Mountains, North Hungary. – Hydrobiologia 506–509: 365–372.
- Kiss, O. – VILMI, L. – FEHÉR, I. (2002b): The water quality state of the Szalajka Stream, Bükk Mts., Hungary. – Braueria 29: 31–34.
- MÓRA, A. – BARNUCZ, E. – BODA, P. – CSABAI, Z. – CSER, B. – DEÁK, Cs. – PAPP, L. (2007): A Balaton környéki kisvízfolyások makroszkópikus gerinctelen faunája. – Acta biologica debrecina, Supplementum oecologica hungarica 16: 105–167.
- MÓRA, A. – BODA, P. – CSABAI, Z. – CSER, B. – DEÁK, Cs. – HORNYÁK, A. – JAKAB, T. – KÁLMÁN, Z. – KECSŐ, K. – KOVÁCS, T.Z. – PAPP, L. – POLYÁK, L. – SOÓS, N. (2008):

- A Zala és befolyói makroszkopikus gerinctelen faunája. – *Acta biologica debrecina, Supplementum oecologica hungarica* 18: 123–180.
- MÓRA, A. – CSABAI, Z. (2002a): Lárvaadatok az Aggtelek–Rudabányai-hegyvidék és a Putnoki-dombság tegzesfaunájához (Trichoptera). – *Folia Historico-naturalia Musei Matraensis* 26: 245–251.
- MÓRA, A. – CSABAI, Z. (2002b): Lárvaadatok a Dél-Alföld tegzesfaunájához (Trichoptera). – *Folia Historico-naturalia Musei Matraensis* 26: 262–267.
- MÓRA, A. – JUHÁSZ, P. – KISS, B. – MÜLLER, Z. (2006): Faunistical results of the Trichoptera investigations carried out in the frames of the ecological survey of the surface waters of Hungary (ECOSURV) in 2005. – *Folia Historico-naturalia Musei Matraensis* 30: 359–367.
- NÓGRÁDI, S. (1984): Six caddisfly species new in the Hungarian fauna (Trichoptera). – *Folia entomologica hungarica* 45/1: 159–165.
- NÓGRÁDI, S. (1986): New data to the caddisfly fauna of Hungary (Trichoptera). – *Folia entomologica hungarica* 47/1–2: 135–140.
- NÓGRÁDI, S. (1987a): Néhány adat Szőce tegzesfaunájához (Trichoptera). – *Praenorica Folia Historico-naturalia* 2: 113–118.
- NÓGRÁDI, S. (1987b): Caddisflies of the Eastern Mecsek Mountains, Hungary (Trichoptera), I. – *A Janus Pannonius Múzeum Évkönyve* 30–31: 11–22.
- NÓGRÁDI, S. (1988): New data to the caddisfly (Trichoptera) fauna of Hungary, II. – *Folia entomologica hungarica* 49: 205–210.
- NÓGRÁDI, S. (1989a): Locality data of the Trichoptera collection originating from the Carpathian Basin in the Hungarian Natural History Museum. – *Folia entomologica hungarica* 50: 147–156.
- NÓGRÁDI, S. (1989b): The Trichoptera fauna of Szőce and its environs (West Hungary). – *A Janus Pannonius Múzeum Évkönyve* 33: 27–33.
- NÓGRÁDI, S. (1994): New data to the caddisfly (Trichoptera) fauna of Hungary, III. – *Folia entomologica hungarica* 55: 271–280.
- NÓGRÁDI, S. (1995): Hungarian locality data of Ujhelyi's Trichoptera collection in the Hungarian Natural History Museum. – *Folia entomologica hungarica* 56: 119–131.
- NÓGRÁDI, S. (2001): Further data to the caddisflies (Trichoptera) of Hungary. – *Folia Historico-naturalia Musei Matraensis* 25: 83–90.
- NÓGRÁDI, S. – KISS, O. – UHERKOVICH, Á. (1996): The Trichoptera fauna of the Bükk National Park. In: MAHUNKA, S. (ed.): *The fauna of the Bükk National Park*. – Hungarian Natural History Museum, Budapest, pp. 397–409.
- NÓGRÁDI, S. – UHERKOVICH, Á. (1989): On the caddisfly fauna of the Kőszeg Mountains, Hungary (Trichoptera). – *A Janus Pannonius Múzeum Évkönyve* 33: 35–42.
- NÓGRÁDI, S. – UHERKOVICH, Á. (1991): Caddisflies of the Eastern Mecsek Mountains, Hungary (Trichoptera), II. – *A Janus Pannonius Múzeum Évkönyve* 35: 19–25.
- NÓGRÁDI, S. – UHERKOVICH, Á. (1992): Újabb vizsgálatok a Kőszegi-hegység Trichoptera faunáján. – *Savaria, a Vas megyei Múzeumok Értesítője, Pars historico-naturalis* 20/2: 149–157.
- NÓGRÁDI, S. – UHERKOVICH, Á. (1994): The Trichoptera fauna of the lake Balaton and its catchment area (Hungary). – *A Janus Pannonius Múzeum Évkönyve* 38: 27–45.
- NÓGRÁDI, S. – UHERKOVICH, Á. (1999): Protected and threatened caddisflies (Trichoptera) of Hungary. In: MALICKY, H. – CHANTARAMONGKOL, P. (eds.): *Proceedings of the 9th International Symposium on Trichoptera*. – Faculty of Science, Chiang Mai University, Chiang Mai, pp. 291–297.
- NÓGRÁDI, S. – UHERKOVICH, Á. (2001): Somogy megye tegzeseinek (Trichoptera) jegyzéke. – *Natura Somogiensis* 1: 295–301.

- NÓGRÁDI, S. – UHERKOVICH, Á. (2002a): Magyarország tegzesei (Trichoptera). – Dunántúli Dolgozatok, Természettudományi Sorozat 11: 1–386.
- NÓGRÁDI, S. – UHERKOVICH, S. (2002b): On the caddisflies (Trichoptera) from the catchment area of the rivers Mura and Kerka, Southwest Hungary. – Somogyi Múzeumok Közleményei 15: 129–144.
- OLÁH, J. (1967): Untersuchungen über die Trichopteren eines Bachsystems der Karpaten (Neue Erklärung des Mosaikmuster – „mosaic pattern“ – Prinzip in Flußwasser-Tiergemeinschaften). – Acta biologica debrecina 5: 71–91.
- SÁTORI, J. (1938): Adatok a Bükk-hegység rovarfaunájának ismeretéhez. – Állattani Közlemények 35: 51–61.
- SÁTORI, J. (1939): Insekten-faunistische Notizen aus dem Transdanubium im Mai und Juni 1938. – Fragmenta faunistica hungarica 2/2: 31–32.
- SCHMERA, D. – KISS, O. (2000): Mintavételezésből adódó eltérések tegzesek (Trichoptera) vizsgálata esetében. – Hidrológiai Közlöny 80: 383–384.
- SOLEM, J.O. (1985): Norwegian *Apatania* Kolenati (Trichoptera: Limnephilidae): identification of larvae and aspects of their biology in a high-altitude zone. – Entomologica scandinavica 16: 161–174.
- UHERKOVICH, Á. (2004): Further studies on the caddisflies (Trichoptera) of the water system of rivers Mura and Kerka, Southwest Hungary. – Folia Historico-naturalia Musei Matraensis 28: 187–194.
- UHERKOVICH, Á. (2005): Further faunistic results of the caddisfly (Trichoptera) examinations of the Dráva region, South Hungary. – Folia Historico-naturalia Musei Matraensis 29: 165–168.
- UHERKOVICH, Á. – NÓGRÁDI, S. (1999): Caddisflies (Trichoptera) of artificial water courses in the Bakony Mountains, Central Hungary. – Braueria 26: 21–23.
- UHERKOVICH, Á. – NÓGRÁDI, S. (2003): Trichoptera of Szigetköz, upper Hungarian Danube Region (Northwest Hungary), II. Species composition and its changes in some water bodies. – Folia Historico-naturalia Musei Matraensis 27: 237–258.
- UHERKOVICH, Á. – NÓGRÁDI, S. (2004): Trichoptera of the Szigetköz, upper Hungarian Danube Region (Northwest Hungary), III. Species composition in Moson Danube, and its changes. – Folia Historico-naturalia Musei Matraensis 28: 171–186.
- UHERKOVICH, Á. – NÓGRÁDI, S. (2006): Caddisflies (Trichoptera) of the Mecsek Mountains, South Hungary. – Natura Somogyiensis 9: 289–304.
- UJHELYI, S. (1971): Adatok a Leptoceridae (Trichoptera) család fajainak magyarországi elterjedéséhez. – Folia entomologica hungarica 24: 119–137.
- UJHELYI, S. (1981): Adatok az Alpokalja szitakötő-, álkérész- és tegzesfaunájához. – Savaria, a Vas megyei Múzeumok Értesítője, Pars historico-naturalis 11–12: 57–65.
- VANNOTE, R. L. – MINSCHALL G. W. – CUMMINS K. W. – SEDELL J. R. – CUSHING C. E. (1980): The river continuum concept. – Canadian Journal of Fisheries and Aquatic Sciences 37: 130–137.
- WALLACE, I.D. – WALLACE, B. – PHILIPSON, G.N. (1990): A key to the case bearing caddis larvae of Britain and Ireland. – Scientific Publication's Freshwater Biological Association 51, The Ferry House, 237 pp.
- WARINGER, J. – GRAF, W. (1997): Atlas der österreichischen Köcherfliegenlarven: unter Einschluss der angrenzenden Gebiete. – Facultas-Universitätsverlag, Wien, 286 pp.