

Purdue University Purdue e-Pubs

Department of Comparative Pathobiology Faculty
Publications

Department of Comparative Pathobiology

9-28-2012

Mycoplasma haemocanis – the canine hemoplasma and its feline counterpart in the genomic era

Naila C. do Nascimento
Purdue University

Andrea P. Santos
Purdue University

Ana MS Guimaraes
Purdue University

Phillip J. SanMiguel
Purdue University

Joanne Messick
Purdue University, jmessic@purdue.edu

Follow this and additional works at: <http://docs.lib.purdue.edu/cbpbpubs>

 Part of the [Veterinary Pathology and Pathobiology Commons](#)

Recommended Citation

This is the publisher pdf of "Mycoplasma haemocanis – the canine hemoplasma and its feline counterpart in the genomic era Naila C do Nascimento, Andrea P Santos, Ana MS Guimaraes, Phillip J SanMiguel, Joanne B Messick *Veterinary Research* 2012, 43:66 (28 September 2012) " and is available at: <http://www.veterinaryresearch.org/content/43/1/66>

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

RESEARCH

Open Access

Mycoplasma haemocanis – the canine hemoplasma and its feline counterpart in the genomic era

Naíla C do Nascimento^{1*}, Andrea P Santos¹, Ana MS Guimaraes¹, Phillip J SanMiguel² and Joanne B Messick^{1*}

Abstract

Mycoplasma haemocanis is a hemotrophic mycoplasma (hemoplasma), blood pathogen that may cause acute disease in immunosuppressed or splenectomized dogs. The genome of the strain Illinois, isolated from blood of a naturally infected dog, has been entirely sequenced and annotated to gain a better understanding of the biology of *M. haemocanis*. Its single circular chromosome has 919 992 bp and a low G + C content (35%), representing a typical mycoplasmal genome. A gene-by-gene comparison against its feline counterpart, *M. haemofelis*, reveals a very similar composition and architecture with most of the genes having conserved synteny extending over their entire chromosomes and differing only by a small set of unique protein coding sequences. As in *M. haemofelis*, *M. haemocanis* metabolic pathways are reduced and apparently rely heavily on the nutrients afforded by its host environment. The presence of a major percentage of its genome dedicated to paralogous genes (63.7%) suggests that this bacterium might use antigenic variation as a mechanism to evade the host's immune system as also observed in *M. haemofelis* genome. Phylogenomic comparisons based on average nucleotide identity (ANI) and tetranucleotide signature suggest that these two pathogens are different species of mycoplasmas, with *M. haemocanis* infecting dogs and *M. haemofelis* infecting cats.

Introduction

Hemotrophic mycoplasmas (hemoplasmas) are uncultivable cell-wall less bacteria, formerly classified as *Haemobartonella* and *Eperythrozoon* species, that adhere to the surface of the erythrocytes of their vertebrate hosts. These bacteria form a new clade within the *Mycoplasma* genus (class Mollicutes) and are phylogenetically related to the pneumoniae group of the mycoplasmas [1-5].

Mycoplasma haemocanis [*Haemobartonella canis*] was first described in Germany in 1928 in a splenectomized dog [6]. The name *Bartonella canis* was proposed and remained until 1939 when Tyzzer and Weinman created the new genus *Haemobartonella* [7]. *M. haemocanis*, proposed species name since 2002 [5], is a pleomorphic bacterium with coccoid and ring forms that can be visualized in the host's peripheral blood smear either singly or in chains that can resemble a "violin-bow" form [8]. It may cause overt, hemolytic anemia in

immunosuppressed [5,9] or splenectomized dogs [5,10], and has a worldwide distribution with prevalence of infection varying from 0.5% to 40% [11-14].

Similarities with the feline hemoplasma *M. haemofelis* [*Haemobartonella felis*], together with the fact that hemoplasmas are not species-specific as previously thought [15,16], led some research groups to hypothesize that these two bacteria could be the same species infecting different hosts [17,18]. Moreover, there are some reports in the old literature stating that *M. haemocanis* could infect cats; however *M. haemofelis* did not cause infection in dogs [19-21]. In 1961, Dr Lumb published the manuscript "Canine haemobartonellosis and its feline counterpart", reporting cross-transmission experiments: it was shown that when blood from cats infected with *M. haemocanis* was injected into susceptible splenectomized dogs, organisms could be seen on their peripheral blood smears [8]. It was concluded based on this evidence that the feline might act as a reservoir for *M. haemocanis*. However, blood from dogs previously injected with *M. haemofelis* inoculated into susceptible cats failed to result in circulating

* Correspondence: ncannes@purdue.edu; jmessic@purdue.edu

¹Department of Veterinary Pathobiology, Purdue University, 725 Harrison Street, West Lafayette, IN 47907, USA

Full list of author information is available at the end of the article

organisms, leading to the conclusion that these two bacteria were different species [8,22]. Forty years later, the sequences of the 16S rRNA genes of these two bacteria were reported to have 99% identity [17] raising the same question again. In 2002, the sequences of the RNase P genes of these bacteria were reported having 94.3 to 95.5% identity [18]. While the results of the RNase P genes did not support the hypothesis that *M. haemocanis* and *M. haemofelis* were identical, this additional data was still considered insufficient to determine whether these organisms should be classified as different species, subspecies, or strains of the same species [18].

Recently, three species of hemoplasmas, including *M. haemofelis*, had their genomes completely sequenced and annotated [23-27]. The aim of this study was to sequence the whole genome of *M. haemocanis* in order to better understand its biology and to perform a complete genomic comparison with its counterpart, *M. haemofelis*.

Materials and methods

Bacterial strain and DNA isolation

M. haemocanis organisms were isolated from the blood of a naturally infected dog at peak of bacteremia [17]. Written informed consent was obtained from the client for publication of this report. Bacterial genomic DNA was extracted using Quick-gDNA MiniPrep kit according to the manufacturer's instructions (Zymo Research, Irvine, CA, USA).

M. haemocanis strain Illinois sequencing and assembly

Whole genome was sequenced from paired-end libraries (TruSeq DNA sample preparation kit, Illumina, San Diego, CA, USA) using 20% of an Illumina® v3 chemistry lane (HiScanSQ). Sequencing resulted in 15.7 million high-quality filtered read pairs with an average read length of 2×100 nucleotides and a > 3400 X genome equivalent coverage. Reads were assembled using ABySS-PE v1.2.7 utilizing 20% of the reads with "kmer" set to 95 bases [28]. Predicted scaffolds with significant BLAST matches to canine DNA were excluded and the remaining mycoplasma scaffolds were then organized based on the orientation predicted in the assembly and on the genome sequence of *M. haemofelis* strain Ohio2. A total of 13 gaps were identified and closed using conventional PCR followed by Sanger sequencing.

Genome annotation and analyses

First pass annotation was achieved using the NCBI annotation pipeline. Manual annotation/curation of each gene was performed using the annotation tool Manatee, provided by the Institute for Genome Sciences (IGS) at the University of Maryland, School of Medicine. Comparative analyses with other bacterial genomes were

performed based on genomic data deposited in the NCBI database (NCBI, Bethesda, MD, USA).

The assignment of paralogous gene families was performed using BLASTclust tool provided by Max-Planck Institute for Developmental Biology [29], with 70% covered length and 30% sequence identity thresholds. Subcellular localization and protein sorting signals were predicted for each unique protein coding sequence (CDS) of *M. haemocanis* and *M. haemofelis* using PSORTb v.3.0 [30,31]. Metabolic pathways were predicted based on the KEGG pathway database [32] and the study reported by Yus et al. [33]. Presence of lipoproteins was predicted by LipoP version 1.0 software [34]. In addition, the tandem repeats were identified using the Tandem repeats finder program [35]. Comparative analyses of the whole genome of *M. haemocanis* and *M. haemofelis* strain Ohio2 were performed using the same tools mentioned above and all the CDSs from both genomes were evaluated using BLASTp and/or BLASTn in order to obtain a complete detailed comparison. CDSs were assigned using BLASTp and considered unique to *M. haemocanis* or *M. haemofelis* when there were no matching sequences in the aligned sequences list with $\geq 90\%$ coverage and $\geq 30\%$ identity or $\geq 80\%$ coverage and $\geq 40\%$ identity to the query sequence. Extended similarity group method for automated protein function prediction (ESG software) [36] was applied for both sets of unique CDSs.

Species differentiation analyses

The average nucleotide identity (ANI; MUMmer algorithm) and tetranucleotide signature correlation index between genomes were calculated using JSpecies software as previously described [37]. In addition to the genome of *M. haemocanis* strain Illinois, the following genome sequences were used in the analyses: *M. haemofelis* strain Ohio2 (CP002808.1), *M. haemofelis* strain Langford (FR773153.2), *M. suis* strain Illinois (CP002525.1), and *M. suis* strain KI3806 (FQ790233.1). If two organisms had ANIm and tetranucleotide coefficients greater than 94% and 0.99, respectively, they were considered the same species [37].

Results

Mycoplasma haemocanis strain Illinois genome features

The complete singular circular chromosome of *M. haemocanis* strain Illinois has a size of 919 992 base pairs (bp) and G + C content of 35%; these genomic features are similar to other hemoplasmas species sequenced to date [23,24,26,27] and within the range reported for other members of the genus *Mycoplasma* (Table 1). As described for all sequenced mycoplasmas (24 species to date), *M. haemocanis* also uses the opal stop codon (UGA) for tryptophan. The 16S, 23S and 5S rRNA genes

Table 1 General genomic characteristics of *Mycoplasma haemocanis* strain Illinois compared to members of pneumoniae group of mycoplasmas

Characteristic	Pneumoniae Group							
	Hemoplasmas				<i>M. pneumoniae</i>	<i>M. gallisepticum</i>	<i>M. genitalium</i>	<i>M. penetrans</i>
	<i>M. haemocanis</i> strain Illinois	<i>M. haemofelis</i> strain Ohio2	' <i>Candidatus M. haemominutum</i> ' strain Birmingham 1	<i>M. suis</i> strain Illinois				
Genome size (base pairs)	919 992	1 155 937	513 880	742 431	816 394	1 012 800	580 076	1 358 633
% of G + C	35.3	38.8	35.5	31.1	40	31	31.7	25.7
Total of genes	1207	1584	582	883	733	817	524	1069
Coding sequences (CDSs)	1173	1549	547	844	689	763	475	1037
CDSs with predicted function	286 (24.3%)	299 (19.3%)	219 (40%)	293 (34.7%)	333 (48.3%)	469 (61.46%)	323 (68%)	585 (56.4%)
No. of tRNAs	31	31	32	32	37	32	36	29
No. of rRNAs								
16S	1	1	1	1	1	2	1	1
23S	1	1	1	1	1	2	1	1
5S	1	1	1	1	1	3	1	1
Genes in paralogous families	748 (63.76%)	1103 (71.2%)	134 (24.5%)	361 (42.8%)	132 (19.1%)	110 (14.4%)	25 (5.2%)	245 (23.6%)

Data was obtained from GenBank database using the following accession numbers: *M. haemocanis* strain Illinois (CP003199.1), *M. haemofelis* strain Ohio2 (CP002808), '*Candidatus Mycoplasma haemominutum*' strain Birmingham 1 (HE613254.1), *M. suis* strain Illinois (CP002525), *M. pneumoniae* (U00089), *M. gallisepticum* (AE015450), *M. genitalium* (L43967), *M. penetrans* (BA000026). Paralogous gene families were assigned using BLASTclust, with 70% coverage and 30% sequence identity thresholds.

Table 2 Comparison of the total number of protein coding sequences (CDSs) of *M. haemocanis* strain Illinois and *M. haemofelis* strain Ohio2 genomes classified by role according to TIGR microbial role categories

Role Category	Number of CDSs (%)	
	<i>M. haemocanis</i> str. Illinois	<i>M. haemofelis</i> str. Ohio2
Purines, pyrimidines, nucleosides, and nucleotides	33 (2.81%)	29 (1.85%)
Fatty acid and phospholipid metabolism	6 (0.51%)	6 (0.38%)
Biosynthesis of co-factors, prosthetic groups, and carriers	8 (0.68%)	7 (0.45%)
Central intermediary metabolism	1 (0.09%)	1 (0.06%)
Energy metabolism	25 (2.13%)	22 (1.41%)
Transport and binding proteins	35 (2.98%)	32 (2.04%)
DNA metabolism	44 (3.75%)	52 (3.32%)
Transcription	21 (1.79%)	18 (1.15%)
Protein synthesis	96 (8.18%)	97 (6.2%)
Protein fate	21 (1.79%)	19 (1.21%)
Regulatory functions	4 (0.34%)	3 (0.19%)
Signal transduction	3 (0.26%)	2 (0.13%)
Cell envelope	3 (0.26%)	7 (0.45%)
Cellular processes	14 (1.19%)	10 (0.64%)
Unknown functions	8 (0.68%)	8 (0.51%)
Hypothetical proteins	887 (75.62%)	1253 (80.06%)
Total*	1209	1566

*Some of the CDSs are in more than one category.

are represented as single copies and share the same operon. The manual genome annotation suggests the presence of 1173 CDSs and 31 tRNAs, covering all amino-acids. Putative functions of most of the CDSs are represented as hypothetical proteins (75.62%), which are mostly due to its large repertoire of paralogous genes (63.76%) (Additional file 1: Table S1). These and other genome features were compared with other hemoplasmas and mycoplasmas members of the pneumoniae group (Table 1). The total number of CDSs of *M. haemocanis* classified by role (according to TIGR microbial role categories) was compared to those found in the *M. haemofelis* genome (Table 2).


Metabolic pathways predictions suggest similar growth requirements for *M. haemocanis* and *M. haemofelis*

Prediction of *M. haemocanis* metabolic pathways based on the KEGG pathway database [32] and Yus's report [33] revealed that they are identical to those predicted for *M. haemofelis* [24]. As shown for *M. haemofelis*, metabolic pathways in *M. haemocanis* are reduced with many of the nutrients and metabolic precursors imported from the blood environment [24]. ATP and DNA/RNA biosynthesis depend on the transport from the environment of glucose and ribose/base derivatives, respectively. Imported bases include: hypoxanthine, adenine, guanine, uracil and cytidine 5'-monophosphate (CMP). Furthermore, amino acids, nicotinamide and any

vitamins required for growth must be acquired from blood environment.

Comparative analyses of *M. haemocanis* and *M. haemofelis* genomes

M. haemocanis genome was compared in its entirety to *M. haemofelis* strain Ohio2 (Figure 1): *M. haemofelis* has 376 CDSs more than *M. haemocanis*, however the majority of these CDSs are members of paralogous gene families also present in *M. haemocanis*; a set of only 67 CDSs was found to be different between these hemoplasmas. The canine hemoplasma possesses only 20 CDSs not identified in *M. haemofelis* genome, while 47 CDSs were unique to *M. haemofelis*. Most of these CDSs are hypothetical proteins, including one family of paralogous genes from *M. haemocanis* and four paralogous gene families from *M. haemofelis* (Additional file 2: Table S2). Predicted functions based on protein sequence similarity for these particular sets of CDSs were assigned using ESG software [36] (Additional file 2: Table S2). Analyses based on PSORTb parameters show that 35% of the unique CDSs of *M. haemocanis* are associated with the cytoplasmic membrane, while 17% of *M. haemofelis* CDSs are predicted to be associated with the membrane and 6.4% with extracellular (signal peptide detected) localization. For most of the unique CDSs, an unknown subcellular localization was predicted, corresponding to 60% and 51% in *M. haemocanis* and



M. haemofelis genomes, respectively (Table 3). Thirteen out of the 20 (65%) unique CDSs of *M. haemocanis* and 35 out of 47 (74.5%) of *M. haemofelis* have at least one internal helix predicted. The number of predicted helices is also shown in the Additional file 2: Table S2. Fifteen lipoproteins were predicted in *M. haemocanis* compared

with 17 for *M. haemofelis* genome, 13 of them are conserved between the two species. In addition, we identified 33 variable number tandem repeats (VNTRs) in the genome of *M. haemocanis* genome (Additional file 3: Table S3), while 61 were reported for *M. haemofelis* [24]. As in *M. haemofelis* genome, most of the VNTRs of

Table 3 Subcellular localization of the unique protein coding sequences (CDSs) of *M. haemocanis* strain Illinois and *M. haemofelis* strain Ohio2 genomes

Subcellular Localization ^a	Number of unique CDSs*					
	<i>M. haemocanis</i> strain Illinois			<i>M. haemofelis</i> strain Ohio2		
	CDSs in paralogous families	CDSs not in paralogous families	CDSs with predicted function	CDSs in paralogous families	CDSs not in paralogous families	CDSs with predicted function
Cytoplasmic membrane	3	3	1	0	8	0
Cytoplasmic	0	0	1	3	7	2
Extracellular	0	0	0	3	0	0
Unknown	5	6	1	18	5	1
Total	8	9	3	24	20	3

^a prediction using PSORTb version 3 software.

* Unique CDSs were assigned using BLASTp when there were no matching sequences with $\geq 90\%$ coverage and $\geq 30\%$ identity or $\geq 80\%$ coverage and $\geq 40\%$ identity to the query sequence.

Table 4 Average nucleotide identity* (ANI) and tetranucleotides signature correlation indexes (Tetra) of selected hemotrophic mycoplasmas

	<i>M. haemofelis</i> Ohio2		<i>M. haemofelis</i> Langford		<i>M. suis</i> illinois		<i>M. suis</i> KI3806	
	ANI	Tetra	ANI	Tetra	ANI	Tetra	ANI	Tetra
<i>M. haemocanis</i> Illinois	85.11	0.959	85.21	0.962	85.59	0.452	85.5	0.453
<i>M. haemofelis</i> Ohio2			97.3	0.999	85.41	0.365	85.3	0.372
<i>M. haemofelis</i> Langford					84.83	0.366	87.74	0.372
<i>M. suis</i> illinois							97.74	0.997

* = ANI was calculated using MUMmer algorithm in JSpecies software.

M. haemocanis were localized within intergenic regions of hypothetical proteins. Five VNTRs were identified within the Type I restriction system operon; the presence of VNTRs in this operon was also described for *M. haemofelis* [24]. Other *M. haemocanis* VNTRs were identified within CDSs for SecD protein, efflux ABC transporter permease protein, PtsG protein, enolase, PotD protein, and for some of the hypothetical proteins.

Only 3 CDSs with known function are exclusive to *M. haemocanis* genome when compared to *M. haemofelis* strain Ohio2; however phosphotransferase system glucose-specific IIBC component (MHC_04460) is only present in the genome of *M. haemofelis* strain Langford, while two ribosomal proteins (MHC_00995 and MHC_05355) are in neither of the feline hemoplasma strains (Additional file 2: Table S2). Another 3 CDSs with known function were identified only in the genome of *M. haemofelis*: two of these proteins are C-5 cytosine-specific DNA methylases (MHF_1273 and MHF_1319), and the other protein is a type II site-specific deoxyribonuclease (MHF_1274) (Additional file 2: Table S2). Small CDSs (corresponding to 30–100 amino acids) characterized as fragments of paralogous genes were excluded from these analyses since they presented a coverage and/or identity below the cutoff to be considered as a member of a paralogous gene family (70% coverage and 30% identity threshold).

Phylogenomic comparison of *M. haemocanis* to other hemoplasmas

ANI and tetranucleotide signature correlation indexes are shown in Table 4. As indicated, *M. haemocanis* had an ANI of approximately 85% in comparison to all other hemoplasma genomes, including *M. haemofelis*. This is below the cutoff value of 94% for species circumscription. The tetranucleotide correlation indexes of *M. haemocanis* with other genomes were also below the 0.99 cutoff limit, being approximately 0.95 for *M. haemofelis* strains and 0.45 for *M. suis* strains. Based on these analyses, *M. haemocanis* is indeed a distinct species infecting the dog.

As expected, strains of the same species (*M. suis* Illinois and KI3806; *M. haemofelis* Ohio2 and Langford1) showed

high ANI and tetranucleotide correlation indexes, which were above the proposed thresholds for species definition. In contrast, ANI and tetranucleotide correlation indexes between *M. suis* and *M. haemofelis* were approximately 85% and 0.37, respectively, correctly separating these organisms as two different species of mycoplasmas.

Discussion

The complete genome sequence and annotation of *M. haemocanis* extends our understanding of the biology of hemoplasmas and provides clues about the growth requirements for *in vitro* cultivation of these bacteria. Based on the metabolic pathway predictions and specific metabolic deficiencies, a more comprehensive medium can be designed [33]. To date, only three other species of hemoplasmas have been entirely sequenced [23–27]. The genome features of *M. haemocanis*, including its small size, low G + C content and use of UGA codon to encode tryptophan, are similar to those of other hemoplasmas and are typical of members of the genus *Mycoplasma*. It is believed that the reduced metabolic pathways of hemoplasmas are probably a consequence of the adaptation to the nutrient-rich blood environment [23,24]. The predicted metabolic pathways of *M. haemocanis* are very similar to those of *M. haemofelis* having orthologs for all the CDSs identified in the genome of this feline hemoplasma [24]; this is not surprising since both species are obligate red cell pathogens that reside in the blood of their hosts. As suggested for other hemoplasmas, it is likely that *M. haemocanis* takes advantage of the erythrocyte's metabolism, scavenging nutrients, which leads to diminished erythrocyte life-span and exacerbation of anemia during acute disease.

Additional primary virulence factors were not identified in the genome of *M. haemocanis*. The *o*-sialoglycoprotein endopeptidase, related to the cleavage of glycophorin A, is conserved among hemoplasmas; the superoxide dismutase (SOD), identified in *M. haemofelis* [24,26] is also present in *M. haemocanis*, but not found in any other sequenced mycoplasma. Although SOD may protect these bacteria from superoxide anion toxicity faced in the blood environment, it is unlikely that this enzyme plays a determinant role in the primary

pathogenicity associated with *M. haemofelis* infection or in the opportunistic infection caused by *M. haemocanis*.

As with other hemoplasmas, *M. haemocanis* contains an abundance of paralogous gene families (63.7% of all its CDSs) and the presence of strategically located tandem repeats. Although there is evidence supporting the role of paralog genes and the presence of tandem repeats in the development of antigenic diversity in *Mycoplasma* species [38,39], additional studies are needed to verify the ability of hemoplasmas to undergo antigenic variation. The presence of irregular cyclic episodes of bacteremia in splenectomized dogs reported following experimental infection with *M. haemocanis* [40], and the possibility that such cycles are due to phase variation is also an area of active investigation in our laboratory.

Comparison of the genomes of *M. haemocanis* and *M. haemofelis* revealed remarkable genetic similarities. Most of the coding and non-coding sequences were conserved and topography of genes within their chromosomes was similar. Even the paralogous gene families were conserved between the two species; the only exceptions were one family with 8 members in *M. haemocanis*, and four small families of *M. haemofelis* with 8, 5, 4 and 3 members, and two with 2 members. The major difference in the paralogous families is the number of duplicate genes inside each of the common families. Thus, as with other bacteria that cannot survive without their host, it appears that maintaining paralogous gene families to generate antigenic variants is a high priority for the hemoplasmas too [41]. On the other hand, CDSs that are unique to *M. haemocanis* or *M. haemofelis* might represent a set of proteins related to differences in virulence and/or related to host specificity. Most of these unique proteins are hypothetical. Although we attempted to improve the function prediction accuracy using the ESG software [36], most of the probabilities assigned were less than 50% and results remained inconclusive (Additional file 2: Table S2). Regarding the sub-cellular localization of the unique CDSs, it is important to mention that the PSORTb software only predicts cytoplasmic membrane localizations when 3 or more transmembrane helices are present within the sequence, otherwise unknown localization is returned. Therefore, these predictions based on strict criteria might have underestimated the potential for membrane localization of these CDSs.

CDSs with known function that are unique to *M. haemocanis* do not appear to have a significant impact on its pathogenicity since they code for an enzyme involved in sugar transport and for ribosomal proteins. On the other hand, *M. haemofelis* possesses a type II restriction enzyme and two C-5 cytosine-specific DNA methylases (C5 Mtase); the restriction endonuclease is located in the same operon as one of the C5 Mtase, indicating that

this operon is functional [42]. Moreover, this endonuclease/methyltransferase pair is not present in any of the other hemoplasmas and the restriction enzyme is absent in the strain Langford 1 of *M. haemofelis*. DNA methylation has been associated with virulence in other bacteria [43]; however, the function of these pair in *M. haemofelis* Ohio2 is unknown.

As mentioned previously, the hemoplasmas cannot be cultivated *in vitro*. This has resulted in a lack of detailed phenotypic and genotypic characterization, which has hampered our ability to correctly classify these organisms within the *Mycoplasmataceae* family. In addition, the 16S rRNA gene failed to provide sufficient resolution to separate *M. haemocanis* and *M. haemofelis* as different species of *Mycoplasma* [5,17]. To date, the genotypic evidence for species differentiation of these two hemoplasmas is solely based on phylogenetic studies using a 177 bp fragment of their RNase P genes [18,44]. Herein, we performed a phylogenomic comparison between *M. haemocanis* and strains of *M. haemofelis* to resolve this long lasting controversy. In recent years, the sequencing of entire genomes has allowed the *in silico* evaluation of genomic similarities between different organisms. ANI and tetranucleotide signatures have been used as surrogates to previous methods of species circumscription, such as 16S rRNA gene phylogeny and DNA-DNA hybridization [37]. With both ANI and tetranucleotide indexes below the proposed thresholds for species definition, our results show that the *M. haemocanis* strain Illinois and *M. haemofelis* (strains Langford and Ohio2) are different species of mycoplasmas infecting two distinct animal species. This conclusion is also supported by the transmission studies done more than 50 years ago [8].

Taken together our results suggest that, although sharing very similar genomes, *M. haemocanis* and *M. haemofelis* are different mycoplasmal species infecting dogs and cats, respectively. The set of unique proteins may be a target for vaccine development against these hemoplasmas, especially for the feline hemoplasmosis that can cause acute disease in immunocompetent hosts.

Nucleotide sequence accession number

The genome of *M. haemocanis* strain Illinois was deposited in GenBank under the accession number CP003199.1.

Additional files

Additional file 1: Table S1. List of paralogous gene families* in the genome of *Mycoplasma haemocanis* strain Illinois. Complete list of paralogous gene families in *Mycoplasma haemocanis* strain Illinois genome assigned by BLASTclust tool provided by Max-Planck Institute for Developmental Biology, with 70% covered length and 30% sequence identity thresholds. GenBank accession numbers are provided.

Additional file 2: Table S2. Analyses of the unique protein coding sequences (CDSs) of *Mycoplasma haemocanis* strain Illinois and *M. haemofelis* strain Ohio2. Analyses of all the unique CDSs of *M. haemocanis* strain Illinois and *M. haemofelis* strain Ohio2 were performed using ESG software, which predicts protein functions based on sequence similarity, and PSORTb version 3 software, which predicts subcellular localization. GenBank accession numbers of proteins corresponding to each CDS are also provided.

Additional file 3: Table S3. Tandem repeats* identified in the genome of *Mycoplasma haemocanis* strain Illinois. Complete list of variable tandem repeats, identified using Tandem repeats finder program, indicating their sequences and position in the genome of *M. haemocanis*.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Conceived and designed experiments: NCN, APS, AMMSG, JBM. Performed experiments: NCN, APS, AMMSG, JBM. Analyzed the data: NCN, APS, AMMSG, PJS, JBM. Wrote and reviewed the paper: NCN, APS, AMMSG, PJS, JBM. All authors read and approved the final manuscript.

Acknowledgments

We are extremely grateful to The Genomic Core Facility at Purdue University and to Samuel Martin for constructing the library, Ann Feil for running the HiScanSQ, and Richard Westerman for performing the ABySS assembly. We also thank the University of Maryland's Institute of Genomic Sciences, which provided us with automatic annotation data and the manual annotation tool Manatee. This work was kindly supported by the Morris Animal Foundation (grant D10FE-0042), and the Brazilian government through CNPq Fellowship Program for Postdoctoral Research.

Author details

¹Department of Veterinary Pathobiology, Purdue University, 725 Harrison Street, West Lafayette, IN 47907, USA. ²Purdue Genomics Core Facility, Purdue University, 170 S. University Street, West Lafayette, IN 47907, USA.

Received: 8 June 2012 Accepted: 30 August 2012

Published: 28 September 2012

References

- Rikihisa Y, Kawahara M, W, Kociba G, Fuerst P, Kawamori F, Suto C, Shibata S, Futohashi M: **Western immunoblot analysis of *Haemobartonella muris* and comparison of 16S rRNA gene sequences of *H. muris*, *H. felis*, and *Eperythrozoon suis*.** *J Clin Microbiol* 1997, **35**:823–829.
- Neimark H, Kocan KM: **The cell wall-less rickettsia *Eperythrozoon wenyonii* is a *Mycoplasma*.** *FEMS Microbiol Lett* 1997, **156**:287–291.
- Neimark H, Johansson KE, Rikihisa Y, Tully JG: **Proposal to transfer some members of the genera *Haemobartonella* and *Eperythrozoon* to the genus *Mycoplasma* with descriptions of '*Candidatus Mycoplasma haemofelis*', '*Candidatus Mycoplasma haemomuris*', '*Candidatus Mycoplasma haemosuis*' and '*Candidatus Mycoplasma wenyonii*'.** *Int J Syst Evol Microbiol* 2001, **51**:891–899.
- Neimark H, Johansson KE, Rikihisa Y, Tully JG: **Revision of hemotrophic *Mycoplasma* species names.** *Int J Syst Evol Microbiol* 2002, **52**:683.
- Messick JB, Walker PG, Raphael W, Berent LM, Shi X: ***Candidatus Mycoplasma haemodidelphidis* sp. nov., '*Candidatus Mycoplasma haemolamae*' sp. nov. and *Mycoplasma haemocanis* comb. nov., hemotrophic parasites from a naturally infected opossum (*Didelphis virginiana*), alpaca (*Lama pacos*) and dog (*Canis familiaris*): phylogenetic and secondary structural relatedness of their 16S rRNA genes to other mycoplasmas.** *Int J Syst Evol Microbiol* 2002, **52**:693–698.
- Kikuth W: **Über einen neuen anamieerreger, *Bartonella canis* nov. sp.** *Klin Wchnschr* 1928, **7**:1729–1730 (in German).
- Tyzzler EE, Weinman D: ***Haemobartonella* n.g. (*Bartonella olim pro parte*), *H. microti* n. sp. of the field vole, *Microtus pennsylvanicus*.** *Am J Hyg* 1939, **30**:141–157.
- Lumb WV: **Canine Haemobartonellosis and Its Feline Counterpart.** *Calif Vet* 1961, **14**:24–25.
- Bellamy JE, Macwilliams PS, Searcy GP: **Cold-agglutinin hemolytic anaemia and *Haemobartonella canis* infection in a dog.** *J Am Vet Med Assoc* 1978, **173**:397–401.
- Hulme-Moir KL, Barker EN, Stonelake A, Helps CR, Tasker S: **Use of real-time polymerase chain reaction to monitor antibiotic therapy in a dog with naturally acquired *Mycoplasma haemocanis* infection.** *J Vet Diag Invest* 2010, **22**:582–587.
- Biondo AW, Dos Santos AP, Guimaraes AM, Vieira RF, Vidotto O, Macieira Dde B, Almosny NR, Molento MB, Timenetsky J, de Moraes HA, González FH, Messick JB: **A review of the occurrence of hemoplasmas (hemotrophic mycoplasmas) in Brazil.** *Rev Bras Parasitol Vet* 2009, **18**:1–7.
- Novacco M, Meli ML, Gentilini F, Marsilio F, Ceci C, Pennisi MG, Lombardo G, Loret A, Santos L, Carrapiço T, Willi B, Wolf G, Lutz H, Hofmann-Lehmann R: **Prevalence and geographical distribution of canine hemotropic mycoplasma infections in Mediterranean countries and analysis of risk factors for infection.** *Vet Microbiol* 2010, **142**:276–284.
- Roura X, Peters IR, Altet L, Tabar MD, Barker EN, Planellas M, Helps CR, Francino O, Shaw SE, Tasker S: **Prevalence of hemotropic mycoplasmas in healthy and unhealthy cats and dogs in Spain.** *J Vet Diagn Invest* 2010, **22**:270–274.
- Tennant KV, Barker EN, Polizopoulou Z, Helps CR, Tasker S: **Real-time quantitative polymerase chain reaction detection of hemoplasmas in healthy and unhealthy dogs from Central Macedonia, Greece.** *J Small Anim Pract* 2011, **52**:645–649.
- Luria BJ, Levy JK, Lappin MR, Breitschwerdt EB, Legendre AM, Hernandez JA, Gorman SP, Lee IT: **Prevalence of infectious diseases in feral cats in Northern Florida.** *J Feline Med Surg* 2004, **6**:287–296.
- Neimark H, Hoff B, Ganter M: ***Mycoplasma ovis* comb. nov. (formerly *Eperythrozoon ovis*), an eperythrocyclic agent of haemolytic anaemia in sheep and goats.** *Int J Syst Evol Microbiol* 2004, **54**:365–371.
- Brinson JJ, Messick JB: **Use of a polymerase chain reaction assay for detection of *Haemobartonella canis* in a dog.** *J Am Vet Med Assoc* 2001, **218**:1943–1945.
- Birkenheuer AJ, Breitschwerdt EB, Alleman AR, Pitulle C: **Differentiation of *Haemobartonella canis* and *Mycoplasma haemofelis* on the basis of comparative analysis of gene sequences.** *Am J Vet Res* 2002, **63**:1385–1388.
- Regendanz P, Reichenow E: **Beitrag zur kenntnis von *Bartonella canis*.** *Arch Schiffs-u Tropen-Hyg* 1932, **36**:305–322.
- Flint JC: **Feline infectious anemia.** In *PhD thesis*. St. Paul, Minnesota, USA: University of Minnesota, School of Veterinary Medicine; 1955.
- Griesemer RA: **Bartonellosis.** *J Natl Cancer Inst* 1958, **20**:949–954.
- Lumb WV: **More information on haemobartonellosis in dogs.** *J Am Vet Med Assoc* 2001, **219**:732–733.
- Guimaraes AM, Santos AP, SanMiguel PJ, Walter T, Timenetsky J, Messick JB: **Complete genome sequence of *Mycoplasma suis* and insights into its biology and adaptation to an erythrocyte niche.** *PLoS One* 2011, **6**:e19574.
- Santos AP, Guimaraes AM, do Nascimento NC, SanMiguel PJ, Martin SW, Messick JB: **Genome of *Mycoplasma haemofelis*, unraveling its strategies for survival and persistence.** *Vet Res* 2011, **42**:102.
- Oehlerking J, Kube M, Felder KM, Matter D, Wittenbrink MM, Schwarzenbach S, Kramer MM, Hoelzle K, Hoelzle LE: **Complete genome sequence of the hemotrophic *Mycoplasma suis* strain KI3806.** *J Bacteriol* 2011, **193**:2369–2370.
- Barker EN, Darby AC, Helps CR, Peters IR, Heesom KJ, Arthur CJ, Crossett B, Hughes MA, Radford AD, Tasker S: **Molecular characterization of the uncultivable hemotropic bacterium *Mycoplasma haemofelis*.** *Vet Res* 2011, **42**:83.
- Barker EN, Darby AC, Helps CR, Peters IR, Hughes MA, Radford AD, Novacco M, Boretti FS, Hofmann-Lehmann R, Tasker S: **Genome sequence for "*Candidatus Mycoplasma haemominutum*," a low-pathogenicity hemoplasma species.** *J Bacteriol* 2012, **194**:905–906.
- Simpson JT, Wong K, Jackman SD, Schein JE, Jones SJ, Birol I: **ABySS: a parallel assembler for short read sequence data.** *Genome Res* 2009, **19**:1117–1123.
- Biegert A, Mayer C, Remmert M, Söding J, Lupas AN: **The MPI Bioinformatics Toolkit for protein sequence analysis.** *Nucleic Acids Res* 2006, **34**:W335–339.
- Nakai K, Horton P: **PSORT: a program for detecting sorting signals in proteins and predicting their subcellular localization.** *Trends Biochem Sci* 1999, **24**:34–36.

31. Yu NY, Wagner JR, Laird MR, Melli G, Rey S, Lo R, Dao P, Sahinalp SC, Ester M, Foster LJ, Brinkman FS: **PSORTb 3.0: improved protein subcellular localization prediction with refined localization subcategories and predictive capabilities for all prokaryotes.** *Bioinformatics* 2010, **26**:1608–1615.
32. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M: **KEGG: Kyoto Encyclopedia of Genes and Genomes.** *Nucleic Acids Res* 1999, **27**:29–34.
33. Yus E, Maier T, Michalodimitrakis K, van Noort V, Yamada T, Chen WH, Wodke JA, Güell M, Martínez S, Bourgeois R, Kühner S, Raineri E, Letunic I, Kalinina OV, Rode M, Herrmann R, Gutiérrez-Gallego R, Russell RB, Gavin AC, Bork P, Serrano L: **Impact of genome reduction on bacterial metabolism and its regulation.** *Science* 2009, **326**:1263–1268.
34. Rahman O, Cummings SP, Harrington DJ, Sutcliffe IC: **Methods for the bioinformatic identification of bacterial lipoproteins encoded in the genomes of Gram-positive bacteria.** *World J Microbiol Biotech* 2008, **24**:2377–2382.
35. Benson G: **Tandem repeats finder: a program to analyze DNA sequences.** *Nucleic Acid Res* 1999, **27**:573–580.
36. Chitale M, Hawkins T, Park C, Kihara D: **ESG: extended similarity group method for automated protein function prediction.** *Bioinformatics* 2009, **25**:1739–1745.
37. Richter M, Rosselló-Móra R: **Shifting the genomic gold standard for the prokaryotic species definition.** *Proc Natl Acad Sci U S A* 2009, **106**:19126–19131.
38. van Belkum A, Scherer S, van Alphen L, Verbrugh H: **Short-sequence DNA repeats in prokaryotic genomes.** *Microbiol Mol Biol Rev* 1998, **62**:275–293.
39. Citti C, Nouvel LX, Baranowski E: **Phase and antigenic variation in mycoplasmas.** *Future Microbiol* 2010, **5**:1073–1085.
40. Lumb WW: **A study of canine haemobartonellosis.** In *PhD thesis*. St. Paul, Minnesota, USA: University of Minnesota, School of Veterinary Medicine; 1957.
41. Palmer GH: **The highest priority: what microbial genomes are telling us about immunity.** *Vet Immunol Immunopathol* 2002, **85**:1–8.
42. Bickle TA, Krüger DH: **Biology of DNA restriction.** *Microbiol Rev* 1993, **57**:434–450.
43. Donahue JP, Peek RM: ***Helicobacter pylori*: Physiology and Genetics.** In *Restriction and Modification Systems*. Edited by Mobley HLT, Mendz GL, Hazell SL. Washington (DC): ASM Press; 2001. Chapter 24.
44. Tasker S, Helps CR, Day MJ, Harbour DA, Shaw SE, Harrus S, Baneth G, Lobetti RG, Malik R, Beauvais JP, Belford CR, Gruffydd-Jones TJ: **Phylogenetic analysis of hemoplasma species: an international study.** *J Clin Microbiol* 2003, **41**:3877–3880.

doi:10.1186/1297-9716-43-66

Cite this article as: do Nascimento *et al.*: *Mycoplasma haemocanis* – the canine hemoplasma and its feline counterpart in the genomic era. *Veterinary Research* 2012 **43**:66.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

