



PRIFYSGOL  
**BANGOR**  
UNIVERSITY

## Cross-laboratory comparison of fluorimetric microplate and colorimetric bench-scale soil enzyme assays

Dick, Richard; Dick, Linda; Deng, Shiping ; Li, Xiufen ; Kandeler, Ellen ; Poll, Christian; Freeman, Christopher; Jones, Timothy; Weintraube, Michael N. ; Saxena, Jyotisna

### Soil Biology and Biochemistry

Published: 01/06/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Dick, R., Dick, L., Deng, S., Li, X., Kandeler, E., Poll, C., Freeman, C., Jones, T., Weintraube, M. N., & Saxena, J. (2018). Cross-laboratory comparison of fluorimetric microplate and colorimetric bench-scale soil enzyme assays. *Soil Biology and Biochemistry*, 121, 240-248.

#### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Manuscript Details

<b>Manuscript number</b>	SBB_2016_49_R1
<b>Title</b>	Cross-laboratory Comparison of Fluorimetric Microplate and Colorimetric Bench-scale Soil Enzyme Assays
<b>Article type</b>	Research Paper

### Abstract

There has been growing interest in fluorescence-based microplate methods to measure enzyme activities due to the sensitivity of fluorimetric detection and the potential for simultaneous and rapid assaying of multiple enzyme activities in the same soil suspension. However, micro-scale methods could introduce considerable operator error such as: 1) the requirement to put soil samples into a suspension; 2) the very small amounts of soil placed in each microplate well; 3) pipetting error because  $\mu\text{L}$  volumes are required; and 4) the need for standard curve calibration with every sample to account for quenching. For valid data comparison and interpretation, there is clearly a need to have a strict and agreed-upon enzyme assay protocol to standardize the microplate-based method. Therefore, the objectives were to: 1) determine the reproducibility and comparability of the standard p-nitrophenol bench-scale and 4-methylumbelliferone microplate enzyme assays measured by five laboratories for  $\beta$ -glucosidase (EC 3.2.1.21) and acid phosphomonoesterase (EC 3.1.3.2) on the same soil samples; and 2) determine the degree and the sources of variability associated with the assays within and among the laboratories. The results showed that overall variability was highest for replication on the microplate ( $n=4$ ), whereas suspension replication had low CVs. This suggests an important source of variation is from pipetting not variability from soil suspensions. A major effort was made to control for methodological differences by using air-dried soils (therefore more stable over varying storage periods) and operator consistency for each task across the labs (e.g. preheated reagents, microplate reader sensitivity set to the highest standard, readings taken within an hour of reaction termination, and controls for substrate autohydrolysis). As a result, the differences among labs were much smaller than differences due to soil type for the microplate method, indicating operator error can be minimized by following the same strict protocol. At the molar level, enzyme activity rates measured across the five labs were not the same between bench and MUF microplate methods (although they were within an order of magnitude), but were quite similar in terms of ranking of soil management treatments and soil types (Table 2). Correlations between bench and microplate assays were strong for both enzymes, although slightly stronger for acid phosphomonoesterase ( $r = 0.93$ ) than  $\beta$ -glucosidase ( $r = 0.81$ ). Additionally, for both acid phosphomonoesterase and  $\beta$ -glucosidase, correlation  $r$  values were mostly similar for MUF microplate and PNP bench method correlation with EL-FAME biomarkers, suggesting both methods were measuring activity originating from the same microbial groups. We conclude that different labs using the same MUF microplate protocol tested, gives reasonably similar absolute activity values, variability, and ranking of treatments (highest to lowest). We propose that the MUF microplate method described in this study be considered as a standard protocol for assaying soil enzyme activities, providing that the buffer pH for the incubation be adjusted to the optimal pH according to the enzyme of interest.

<b>Keywords</b>	cross-lab study; soil enzyme methods; microplate; 4-methylumbelliferone; p-nitrophenol
<b>Taxonomy</b>	Soil Microbial Ecology, Enzymatic Methods
<b>Manuscript region of origin</b>	North America
<b>Corresponding Author</b>	Richard Dick
<b>Order of Authors</b>	Richard Dick, Linda K. Dick, Shiping Deng, Xiufen Li, Ellen Kandeler, Christian Poll, Chris Freeman, Timothy Graham Jones, Michael Weintraub, Kawthar Esseili, Jyotisna Saxena
<b>Suggested reviewers</b>	Tom Speir, Denis Angers, David Coleman

## Submission Files Included in this PDF

### File Name [File Type]

Cover letter SBB Cross lab study.doc [Cover Letter]

Response Cross Lab Enzyme Study Reveiw Comments 4Oct17 (2).doc [Response to Reviewers]


Highlights.doc [Highlights]

Cross lab Enzyme Method Study Post Review Final 16Oct17.docx [Manuscript File]

Figures.pdf [Figure]

Procedures MUF Cross Lab Study 24Sep17 Final After Review.doc [e-Component]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.



**School of Environment and  
Natural Resources**

2021 Coffey Road  
Columbus OH  
43210-1085

Phone 614-292-  
2265 FAX 614-  
292-74

29 September, 2016

Dear Editor:

We are submitting the manuscript entitled “Cross-laboratory Comparison of Fluorimetric Microplate and Colorimetric Bench-scale Soil Enzyme Assays” that has the following authors: Richard P. Dick, Linda K. Dick, Shiping Deng, Xiufen Li, Ellen Kandeler, Christian Poll, Christopher Freeman, Timothy Graham Jones, Michael N. Weintraub, Kawthar A. Esseili, and Jyotisna Saxena.

This is original research and it was presented at the International Conference – Enzymes in the Environment: Activity, Ecology, and Applications, 24-27 July, 2016, Bangor, Wales, UK.

Thank you

A handwritten signature in cursive script that reads "Richard P. Dick".

Professor

## Cross Lab Enzyme Study

### Soil Biology and Biochemistry

#### **Reviewer 1**

1) I was enthusiastic about reading this manuscript and hoped to gain insights into variability and potential sources of error with the microplate approach. Unfortunately, rather than present a rationale and evidence-based argument that the method presented is the one to use, the manuscript contains many assertive statements (for example, that a strict method is needed), without good arguments for the case. The need for a strict, standardized method is repeatedly mentioned, but the experimental testing presented is at times vague on details, or variable between laboratories.

The comparisons performed are also somewhat bewildering: if part of the rationale was to compare small scale microplate methods to larger scale “bench top” methods, why use different artificial substrates for those comparisons (microplate MUF to benchtop pNP)? It’s very possible that microbial enzymes could show different affinities for different substrate types, and that this could vary between samples. It seems that it would have been easy to run both MUF and pNP assays in both microplate and benchtop form.

#### *RESPONSE:*

1a) pNP microplate method is less sensitive and troublesome. According to Deng et al. (2013), in the presence of soil suspension, as little as 50 pmol of 4-methylumbelliferone, compared to 16.28 nmol of pNP, can be detected in each assay. Furthermore, soil particles in the wells interferes with absorbance reading, lead to considerable background readings and cause high variability in pNP microplate method. Fluorescence measurements are known to be significantly less susceptible to the effects of turbidity compared to absorption based detection (Deng et al. 2011). This suggests that MUF microplate is more desirable.

1b) MUF bench method is not needed because the pNP bench method offers sufficient sensitivity for its detection, adequate stability for the convenience in routine laboratory analysis, and relatively low cost to be feasible (MUF at bench scale is very expensive) and adaptable for most research programs.

1c) Thus the effort in this study was focused on MUF microplate method, which is now widely used but yet with limited understanding on its reliability and sources of variability. We compared microplate method to the bench method which, for both enzyme assays in our study, has been well vetted. So the goal here was very practical in moving towards standardizing a MUF microplate procedure that if followed would allow cross study comparisons and meta-analysis of soil enzyme activity data. We used as much information as possible from previous research on microplate methods and implemented a protocol that was guided by the groundwork of research on the bench scale method.

1d) The employed bench method for the enzyme assays are vetted and standardized methods. The bench methods are based on sound chemistry/biochemistry principles that measures potential activity of enzymes in soils. We understand that it is very possible that an enzyme could show different affinities for different substrate types, and that this could vary between samples. In theory, the detected enzyme activity, however, should be comparable regardless of method of detection, providing that the methods are based on sound science and have been systematically evaluated. Our working hypothesis was that if both methods under different operators had statistically acceptable levels of variability, comparable absolute values (on molar basis), and consistent ranking of tested soil types, then we would conclude that the microplate method offers sufficient reproducibility as a reliable method for valid evaluation of soil enzyme activity. For meaningful comparison, variability and treatment ranking between methods are important, even if absolute values do not match up. This rationale is stated in the Introduction in the original submission (starting on L66).

2) Similarly, the number of replicates differs between the two methodological approaches. Why not standardize this?

*RESPONSE:* We assume that analytical replicates were implied in this comment. The required number of replicates in an assay is closely related to variability and reproducibility of an assay protocol. It is, therefore, method dependent. For the bench assays, duplicate assays were sufficient to offer statistically acceptable variability. These methods have been developed and systematically evaluated in M.A. Tabatabai's program, which have shown high precision (often <5% coefficient of variance) with the vetted protocols. For the microplate method, a larger number of analytical replicates are required because higher variability was expected due to the high sensitivity of detection, quenching of fluorescence by the soil suspension, and small amount of sample employed in each assay (Freeman et al., 1995; Deng et al., 2013; Deng et al., 2017). Unfortunately, most MUF-based microplate assays in the literature did not include assay replicates of soil suspensions as discussed in Deng et al. (2017). The replicates of up to 16 microplate wells in each assay were originated from the same soil suspension, which would account for variability introduced by pipetting. Deng et al. (J. Microbiol. Meth. 133:32–34. 2013) showed that 4 replicated wells were sufficient to address variability introduced by pipetting. The number of assay replications used in this study was based on previous research reports.

3) The FAME assays also seem out of place. They're not addressed in the abstract and seem more like an afterthought or an extra set of data that was just added and not really examined thoroughly.

*RESPONSE:* The focus of this study was on methods comparison. However, FAME data provided additional data validation and more in-depth data interpretation and understanding. The reason for the FAME assays was stated in the original submission (now L321-24).

“To give some insight into the relationship between the two enzyme activities assayed by bench and microplate methods and microbial community profiles, fatty acid methyl esters (EL-FAMES) were used as biomarkers for major functional microbial groups. The biomass of each microbial group was correlated with enzyme activities (Table 7).”

We revised the abstract to reflect the effort.

4) Finally, I'd also question the need and timing of a manuscript like this. Microplate assays are hardly new – they've been around for >20 years – so presenting them as a new approach that needs to be standardized compared to “traditional” methods seems somewhat dated.

*RESPONSE:* This is an excellent question and we agree that this research could have been done years ago. The microplate assay protocols were guided by the groundwork on the bench scale method; but were not systematically evaluated as done for the bench assays (Tabatabai, 1994). Over time, challenges have emerged. This study was motivated to address unaddressed issues facing the assay, for validate data comparison and meaningful interpretation. In research, reliability of a method is of paramount importance and particularly prudent when a method is widely being used, but not standardized; regardless if the method is relatively new or old.

Specific points by line

5) 8-9 Is this statement really true? Is it necessary to have a strict and agreed-upon protocol for valid data comparison and interpretation? Maybe for some comparisons between studies, but otherwise no. And of course the authors would like it to be their “strict and agreed-upon protocol” – although they don't present any comparisons to different protocols, just their own.

*RESPONSE:* We are surprised by this comment that agreed-upon protocols are necessary because for any research endeavor this is fundamental science. A method does not belong to anyone, but is one that is widely accepted by the scientific community. The method must be based on sound science, ideally easy to perform, and reproducible. The latter requires systematic evaluations of the assay protocol, which for enzyme activity assays is the “Tabatabai protocol” (some referred to them as “classic methods”). The microplate method in our study is based on the evolution across these method development papers - Marx et al., 2001; DeForest, J.L., 2009; German et al., 2011; Deng et al., 2011, 2013; Dick et al., 2013.

For a more detailed discussion of systematic method evaluation, please see: Dick, W.A. 2011. Development of a Soil Enzyme Reaction Assay. R.P. Dick (ed.) *Methods of Soil Enzymology*, SSSA Book Series. No. 9.. Soil Science Society of America, Madison, WI USA. doi:10.2136/sssabookser9.c14

6) 13-14 Sources/causes of variability aren't really addressed. Some speculation on the cause of variability but none were actually tested.

*RESPONSE:* The paper is fully focused on sources of variability from the field down to the lab analytical replication – as shown in Tables 3-5 with standard errors shown in Table 2.

7) 25 Don't cite a Table in an abstract

*RESPONSE:* Deleted

8) 22, 26-27 References to the importance of following a strict, standardized protocol, but this wasn't actually tested. Would allowing each lab to have conducted assays according to their own protocols have led to, for example, the same ranking of soils?

*RESPONSE:* We did not test strict protocols – we all agreed to strictly follow the same protocols for the 2 methods compared in the study – here by strict we mean that we agreed all follow the exact same protocol.

9) 83 Spell out THAM on first use

*RESPONSE:* We respectfully disagree – this is a widely used and recognized abbreviation in the discipline. We recognize that in biology/molecular biology it is often referred to as “Tris”).

10) 90-92 Is a quench standard curve necessary? Quench controls (MUF with sample) can also be used.

*RESPONSE:* Yes it is necessary

11) 98-102 The larger mass of soil used in benchtop assays is likely as big a factor (if not more) as pipetting error for more variability in microplate assays.

*RESPONSE:* Based on principles of analytical chemistry, the larger mass of soil used in bench assays would result in lowering variability and analytic errors from assay to assay.

12) 101-102 Why would variability in protocols exacerbate these methodological issues? Not clear

*RESPONSE:* We agree this was confusing as this was referring to lab to lab comparability and as such this statement was deleted.

13) 104-105 Again, why a strict and agreed-upon protocol? Who needs to agree (the limited number of authors?).

*RESPONSE:* Please see response to question #5 above.

14) 130 How was soil homogenized?



*RESPONSE:* L133 – Rewritten to clarify.

15) 131 (and elsewhere) I'm not convinced that air drying is a valid method for preservation. This allows all sorts of microbial processes to take place and in my experience is not a substitute for running fresh assays, regardless of the protocol used (MUF vs. pNP, microplate vs. bench). This would not be an acceptable method for other microbial assays (e.g. community composition) and it's bewildering why we think it is for a physiological measure. Why not at least freeze dry? Even if acceptable, no details on air drying (temperature, humidity, time) are presented.

*RESPONSE:* This was done to stabilize the sample and reduce variability because soil had to be shipped long distances and leaving them field moist would undoubtedly resulted in more variability. This was confirmed by the obtained results which showed that even though the time between soil sampling/processing and analysis, varied considerably across labs – the results were quite consistent. We agree that field fresh samples is needed when the study objective is focused on microbial analyses related to viable cells. However, decades of research have demonstrated that activities of soil enzymes are originated mostly from cell-free stabilized extracellular enzymes (Tabatabai, 1994). Enzyme activities in soil indicate a biochemical property of a soil, not necessarily correlated with microbial activities. Based on evaluation of 20+ soils under various conditions, Frankenberger and Dick (1983. SSSAJ 47:945-951) found that enzyme activities and microbial activities in soil were significantly correlated only when sugar was added to the soil prior to the test. Furthermore, for the enzymes used in this study – research data have shown that the activity for a given enzyme may go down some with air drying, the relative ranking of treatments or across soil types stays the same (e.g. phosphatase - Eivazi and Tabatabai, 1977; beta-glucosidase - Eivazi & Tabatabai, SBB 22:891; Bandick and Dick, 1999).

16) 167-168 “within two weeks of receiving...” is vague and variable. Why not follow a “strict” timeline? And the data from one laboratory that took “within four weeks” (presumably really 2-4 weeks) should be excluded.

*RESPONSE:* It was the best we could do given that 2 of the participating labs were in Europe. Again despite this variation in time to analysis, the results were quite consistent as expected.

17) 172-173 Why even include Lab 5 if they weren't doing the whole process? This, and the prior comment, just highlight how poorly standardized this experiment was.

*RESPONSE:* This lab was not set up to do the bench method, but has extensive experience in the microplate method. Including lab 5 strengthened data interpretation to further confirm or refute the study hypothesis that operator error was minimal in using the microplate method and gave 5 instead of 4 reps to determine microplate variability.

18) 178 Why 37C? What is the rationale for conducting assays of environmental enzymes at human body temperature? Combine a non-realistic temperature with a non-realistic pH and we're getting quite far removed from environmental conditions.

*RESPONSE:* See response to #15. An enzyme is very different than a viable cell and as such optimizing for pH and temperature to measure activity as an index of the isoenzymes present in the soil is fundamental biochemistry. In this study, potential enzyme activities in soil were measured. The study objective was not to evaluate “in situ” enzyme activities in soil. The choice of temperature and pH for the assay followed those used in the well accepted bench methods (i.e. Tabatabai, 1994). Yes, human body temperature was selected to be used. Enzyme activities increases with increasing temperature with a Q10 value often exceeding 2. The assaying temperature of 37°C allows sensitive detection of most soil enzymes, but does not raise concern on enzyme denaturation which occurs around 60-65 °C in soil. An enzyme assay is an index of the amount of isoenzymes that is in the soil sample that can perform a given reaction – for sensitivity and comparability, it is best to conduct the assay under optimized conditions. As soon as non-optimized conditions are not used, activity goes down and no longer is the assay truly able to be an index for the total amount of isoenzymes that are present. This could very well obscure treatment effects or even show non-significant differences when if they had been done under optimal conditions there would have been differences.

19) 180-181 Is using a stir bar and plate really “homogenizing” rather than just mixing?

*RESPONSE:* This is a matter of semantics - According to Webster:  
Homogenize: “to make uniform in structure or composition throughout”  
Mix: “to combine or blend into one mass”

20) 184-185 What are the substrates dissolved in?

*RESPONSE:* Water

21) 186 “several times” is vague; covered with what?

*RESPONSE:* 2-3 times, revised.

22) 191-193 Need more information on models of microplate readers and sensitivity settings. This is a major source of variability and it's glossed over.

*RESPONSE:* We disagree because if the sensitivity (gain) settings are based on the highest calibration standard (which all labs did), the results are normalized. All labs got comparable results would confirm this.

23) 194-198 This section on controls is vague. Does not adding substrate initially mean that a 0 M substrate solution was used (to keep volumes standardized) or that nothing was added (so volumes of controls were less than that of reactions)?

*RESPONSE:* Yes, the volume in the controls were less than that of the assayed sample during incubation. The substrate was added to the control following incubation and termination of the reaction. Detailed description can be found in the paper and/or the supplementary information.

24) 199-200 Again, varying reagents is yet another source of variation that should have been standardized for a “strict” protocol. These could have been shipped along with the soil samples.

*RESPONSE:* All substrates were the exact same compounds. Moreover, controls employed in the assay should have accounted for any potential variations in substrate auto-hydrolysis from variations in source and/or storage.

25) 211 Why 37 C?

*RESPONSE:* See above response to #18.

26) 216-218 As with the MUF assays, it’s not clear if an equal volume of 0 M substrate was used for controls or no additional volume.

*RESPONSE:* See response to #23. Controls and sample assays had the same volumes when taking the measurements.

27) 218 What instrument was used for absorbance measurements?

*RESPONSE:* Spectrophotometer

28) 227-228 Why not a 2-way ANOVA to allow for the detection of lab x sample interactions? Shouldn’t the follow-up tests (multiple pairwise comparisons) need something like the Bonferroni correction?

*RESPONSE:* 2-Way ANOVA was done and added to the M & M. We ran the Bonferoni test and got the same outcome.

29) 243-248 Seems off topic

*RESPONSE:* This puts the methods’ results in the context of the field treatments which in the next paragraph these soil properties are linked with the enzyme methods.

30) 277-278 As mentioned earlier, it would be reasonable to expect different affinities of different enzymes for pNP or MUF substrates

*RESPONSE:* We agree that this could be a factor. We added this on L291: “Although there is greater affinity by MUF over PNP (as shown by Km values presented by Marx et al, 2001 and Deng et al., 2013) which could change the activity rate, it should be a

consistent effect. A more likely factor contributing to variability detected by the MUF method is quenching of the emitted fluorescence by soil particles.”

31) 294-295 Speculation. No evidence is presented that different times between sampling had no effect

*RESPONSE:* We have recently shown that for B-glucosidase air dried soil kept at room temperature, 4 C or -20 C was stable on the order of months. This was added on L310.

32) 298-299 Didn't alter “ranking” is not the same as didn't alter activity

*RESPONSE:* We agree and this statement does not contradict that.

33) 300-302 Vague. Nothing in this study validates the use of air-dried samples

*RESPONSE:* These statements are not validating air drying over field moisture nor advocating one over the other - just making a comment enzyme assays can be run on air-dried samples. See REPSONE 39.

34) 317-328 The emphasis on FAME here seems out of order (presenting info in Table 7 ahead of Tables 3 and 4)

*RESPONSE:* This section was moved to the Bench vs. Microplate section (L321).

35) 360-363 Why not vary the mass of soil to see if this actually is a major factor?

*RESPONSE:* This was discussed based on principles in analytical chemistry. For the PNP bench method, sample size has been well tested and established. Reducing soil mass to the same level employed in the microplate methods is not possible. First, the enzyme activity would not be detectable because PNP is not as sensitive as MUF. Second, this would require development of a new assay protocol, which is beyond the scope of this paper. On the other hand, increasing soil mass in the MUF microplate method to that of the PNP bench method is not possible as well. In the microplate assay methods, soil suspensions are used in the assay. Pipetting mud into microplate wells is challenging. This change would also require systematic evaluation of the protocol, which again is beyond scope of the study. Moreover, studies of the MUF microplate method where substrate concentration curves were developed - the standard MUF substrate concentration is in excess - meaning the enzymes were saturated. This indicates the 1 g of soil put in to suspension when distributed to plate wells was not in excess to cause unsaturated substrate conditions.

36) 414-417 While my own assays do tend to use a standard pH, the argument that varying pH based on sample pH prevents direct comparisons isn't a good one. One could just as easily argue that not adjusting pH to the in situ pH of the soil prevents valid comparisons of actual real world activity, and at best gives just an estimate of potential

activity (under standardized conditions of pH and temperature that may have no real world application).

*RESPONSE:* This would be objective dependent. The objective of this study was to determine potential enzyme activities in soil. See responses to #18 above. Moreover, it is challenging to determine actual “in situ” soil enzyme activity in the laboratory, as the soil would be removed from the field and processed, and the assay conditions are totally different from the in situ environment.

37) 421-442 I don't follow why a 1 hour lag between reaction termination and reading was even necessary. Terminate the reaction, read the plate: 2 minute lag time, max.

*RESPONSE:* This is just to indicate the amount of time available in case for what ever reason there is variability in the time between terminating the reaction and reading fluorescence for a given operator. Deng et al. (2013) showed that there was very little changes in fluorescence within 3 hrs following termination with THAM (Fig. 1, Deng et al., 2013).

38) 426-428 The major problem here is that only one method is presented or tested, but then is proposed to be the standard method. No alternative methods were evaluated.

*RESPONSE:* Many methods in soil analyses have components that are operationally defined in order to allow results to be comparable from study to study. The microplate method we are proposing is based in part on the vetted bench method for the 2 enzymes in the study (Tabatabai, 1994) and on microplate studies (Freeman et al., 1995; Marx et al., 2001; Drouilon and Merckx, 2005; Deng et al, 2013, Dick et al., 2013). For an enzyme assay the criteria is straight forward: substrate is in excess, co-factors present (when needed), long enough incubation for adequate product detection, high enough temperature to minimize incubation time, adequate enzyme concentration (amount of soil), and optimum pH. So developing a soil enzyme assay is not done in comparison to alternative methods but rather establishing agreed upon operationally defined components and optimization.

## **Reviewer 2**

39) This is a well-written, interesting article that highlights potential sources of variation in assaying for soil enzymes using colorimetric and fluorimetric approaches. My major concern is that while the recommended protocols will likely reduce the variability among labs, will they still capture the biologically relative variation found within field conditions? Using air-dry and incubating at unnaturally warm temperatures seems like a biological filter that would reduce the natural variation found within a soil sample. While this approach has been used for decades, it doesn't necessary mean it is the best approach at all research questions. For example, drying soil will significantly reduce the activity of phosphatase (Sparling et al., 1986, SBB), although results can be mixed (Zornoza et al., 2006, SBB).

*RESPONSE:* Decades of research have demonstrated that activities of soil enzymes are originated mostly from cell-free stabilized extracellular enzymes (Tabatabai, 1994). Enzyme activities in soil indicate biochemical property of a soil, not necessarily correlated with microbial activities. Based on evaluation of 20+ soils under various conditions, Frankenberger and Dick (1983. SSSAJ 47:945-951) found that enzyme activities and microbial activities in soil were significantly correlated only when sugar was added to the soil prior to incubation and testing. Field-moist fresh soil samples are needed when the study objective is focused on microbial analyses related to viable cells. We are not advocating for or against air drying. Air-drying was done to stabilize the sample and reduce variability because soil had to be shipped long distances and leaving them field moist would undoubtedly resulted in more variability. Yes, air-drying would result in lower enzyme activity, but should not affect relative comparison of the samples. This was confirmed by the obtained results which showed that even though the time between soil sampling/processing and analysis, varied considerably across labs – the results were quite consistent across labs. Furthermore, for the enzymes used in this study – research data have shown that the activity for a given enzyme may go down some with air drying, but the relative ranking of treatments or across soil types stays the same (e.g. phosphatase - Eivazi and Tabatabai, 1977; beta-glucosidase - Eivazi & Tabatabai, SBB 22:891; Bandick and Dick, 1999)

40) This is the crux of these types of enzyme papers: Do you standardize to ensure repeatability among labs or do we optimize to best represent in-situ conditions? There is no ‘right’ answer to this question, but there clearly are researcher firmly in each camp. Regardless, I think this article should provide the potential down-side of strict standardization. For example, is a 1 hour incubation time at 37°C useful for all soils (e.g. Arctic/cold soils or ones with very low activity)? Likewise, consider in your recommendations that authors make it obvious in their methods section that they should report if methods were either optimized to standardized among lab or to optimize in-situ conditions.

*RESPONSE:* This is a reasonable question. Repeatability is important for any method. We think you meant optimized potential enzyme activity vs. in-situ enzyme activity. Method selection is objective dependent. If one were incubating and measuring viable microbial properties in situ conditions makes sense but an enzyme activity assay is straight biochemistry. In this study, potential enzyme activities were of interest and were measured. Some of the fundamentals are discussed in the response to #43. The length of incubation time should not affect the determined enzyme activity because activities are often expressed as the amount of product released per unit time by unit of soil, providing that substrate is not limited during the incubation period. Incubation temperature, on the other hand, is an important factor affecting the measured enzyme activity. Enzyme activities increases with increasing temperature with a Q10 value often exceeding 2. Therefore, enzyme activities are comparable only when quantified at the same incubation temperature. For Arctic/cold soils or ones with very low activity, one can simply extend incubation time for the activities to be detectable (extending incubation time should not alter outcome of the measurement – activities determined). The commonly used assaying

temperature of 37°C allows sensitive detection of most soil enzymes, but does not raise concern on enzyme denaturation which occurs around 60-65 °C in soil.

41) I think the whole issue of pipette seems odd and only seems an issue unless manual pipette were used. It's unclear in this paper if the authors used a manual or an electronic pipette.

*RESPONSE:* We used electronic pipettes – but putting uL levels of a soil suspension is still of concern because a typical value of the soil in each well was only 0.83 mg. Clogging and air bubble problems are much more likely when working at uL suspensions compared to larger volumes of solutions.

Specific Comments:

42) Line 29 Where in this study is the evidence for this statement? This seems at odds with the idea of standardization, unless there is an internationally recognized accepted resource on pH optimal for these enzymes. If so, provide a reference.

*RESPONSE:* The recognized pH optima is based the extensive work of M. A. Tabatabai (Tabatabai, 1994). However, if there has not been pH curves developed for a given enzyme assay, then this needs to be determined before the assay is used as a research tool. We have acknowledged this and added text to this effect in L438-442 and L457 in the manuscript.

43) 104 I don't agree entirely with this statement and it seems too draconian. We can still have valid comparison and interpretation using a proper approach. While some information would be lost, using enzymes stoichiometry will allow researcher to compare even if their methods are different that likely influence the absolute activity.

*RESPONSE:* We do not view this as draconian and stand by this statement. The concern is that the further the procedure is from optimal temperature or pH, the lower the activity which could mask treatment effects. Examples of this can be seen in some enzyme activity methods papers where on the extremes of the pH curve – soil types that are very different at the optimum pH then converge to levels that are very similar at very high or low pH. So in effect the results are confounded by less than optimal conditions (e.g. if one is comparing soils where some have a natural pH near the optimum and others are not close to the optimum and then the pH is not buffered – those soils not at the optimum are going to be lower automatically which confounds the goal of enzyme assays as an index of the total amount of enzyme present). Furthermore, soil is known to be heterogeneous. Micro-habitats in soil could have markedly different pH values than the bulk soil, making it challenging to truly measure “in-situ” soil enzyme activities.

44) 259 What is meant by ‘relative ranking’? This is not described in the methods section and it seems a considerable part of the comparison is based on this ranking approach. Please add some justification on how ranking is an appropriate approach to

compare the two methods. Maybe a comparison of the ratio between these enzymes may be a better approach in comparison.

*RESPONSE:* Relative ranking referred to relative enzyme activities among the tested soils. Consistent ranking by different methods validates their use in detecting treatment effects (e.g. soil type or various soil management treatments).

45) 262 The approach for this analysis was not described in the data analysis section. Please provide the details.

*RESPONSE:* This has been added to the M & M.

46)317 I like this approach of compared enzyme activities on something standard, like microbial biomass or soil C. The authors clearly show that microbial biomass is strongly correlated with enzyme activity.

*RESPONSE:* Thank you.

47) 342 Is this a mistake? Sandy soils are the easiest to homogenize and dispense – Do you mean to stay in suspension?

*RESPONSE:* Rewritten to indicate that to keep in solution is a major concern.

48) 385 Please remove the extra ‘,’

*RESPONSE:* Done

49) Tables 3-5 I don't find this tables very useful, I think they would be better placed in supplementary content.

*RESPONSE:* We respectfully disagree as these tables along with Table 2 are the heart of the paper. Tables 3-5 show the levels of variability at each stage for the bench and microplate methods. We acknowledge that they are complex, struggled how best to present this but in the end found this to be the most appropriate to support objectives of the study.



Bench and microplate enzyme activities were similar and highly correlated;

Following strict enzyme protocols across labs yielded similar activities;

Microplate method had highest variability for plate pipetting and not soil suspension;

MUF microplate and bench methods across labs gave similar ranking of treatments;

A standard microplate enzyme method is recommended to enable cross-study comparisons;

## **Cross-laboratory Comparison of Fluorimetric Microplate and Colorimetric Bench-scale Soil**

### **Enzyme Assays**

Richard P. Dick<sup>1\*</sup>, Linda K. Dick<sup>1</sup>; Shiping Deng<sup>2</sup>; Xiufen Li<sup>2</sup>, Ellen Kandeler<sup>3</sup>; Christian Poll<sup>3</sup>;

Christopher Freeman<sup>4</sup>; Timothy Graham Jones<sup>4</sup>; Michael N. Weintraub<sup>5</sup>; Kawthar A. Esseili<sup>5</sup>; and

Jyotisna Saxena<sup>1</sup>

<sup>1</sup> School of Environment and Natural Resources, The Ohio State University, Columbus OH, 43210, USA; <sup>2</sup> Department of Plant and Soil Sciences, Oklahoma State University, Stillwater, OK 74078, USA; <sup>3</sup>Institute of Soil Science and Land Evaluation, University of Hohenheim, Stuttgart D-70593, Germany; <sup>4</sup>School of Biological Sciences, Bangor University, Bangor, United Kingdom; <sup>5</sup>Department of Environmental Sciences, University of Toledo, Toledo, OH, 43606, USA.

\*Corresponding author:

. Phone: 614-247-7605, Fax: 614-247-6901; Email: dick.78@osu.edu

1 **Abstract**

2 There has been growing interest in fluorescence-based microplate methods to measure enzyme  
3 activities due to the sensitivity of fluorimetric detection and the potential for simultaneous and rapid  
4 assaying of multiple enzyme activities in the same soil suspension. However, micro-scale methods  
5 could introduce considerable operator error such as: 1) the requirement to put soil samples into a  
6 suspension; 2) the very small amounts of soil placed in each microplate well; 3) pipetting error  
7 because  $\mu\text{L}$  volumes are required; and 4) the need for standard curve calibration with every sample  
8 to account for quenching. For valid data comparison and interpretation, there is clearly a need to  
9 have a strict and agreed-upon enzyme assay protocol to standardize the microplate-based method.  
10 Therefore, the objectives were to: 1) determine the reproducibility and comparability of the standard  
11 *p*-nitrophenol bench-scale and 4-methylumbelliferone microplate enzyme assays measured by five  
12 laboratories for  $\beta$ -glucosidase (EC 3.2.1.21) and acid phosphomonoesterase (EC 3.1.3.2) on the  
13 same soil samples; and 2) determine the degree and the sources of variability associated with the  
14 assays within and among the laboratories. The results showed that overall variability was highest for  
15 replication on the microplate ( $n=4$ ), whereas suspension replication had low CVs. This suggests an  
16 important source of variation is from pipetting not variability from soil suspensions. A major effort  
17 was made to control for methodological differences by using air-dried soils (therefore more stable  
18 over varying storage periods) and operator consistency for each task across the labs (e.g. preheated  
19 reagents, microplate reader sensitivity set to the highest standard, readings taken within an hour of  
20 reaction termination, and controls for substrate autohydrolysis). As a result, the differences among  
21 labs were much smaller than differences due to soil type for the microplate method, indicating  
22 operator error can be minimized by following the same strict protocol. At the molar level, enzyme  
23 activity rates measured across the five labs were not the same between bench and MUF microplate

24 methods (although they were within an order of magnitude), but were quite similar in terms of  
25 ranking of soil management treatments and soil types (Table 2). Correlations between bench and  
26 microplate assays were strong for both enzymes, although slightly stronger for acid  
27 phosphomonoesterase ( $r = 0.93$ ) than  $\beta$ -glucosidase ( $r = 0.81$ ). Additionally, for both acid  
28 phosphomonoesterase and  $\beta$ -glucosidase, correlation  $r$  values were mostly similar for MUF  
29 microplate and PNP bench method correlation with EL-FAME biomarkers, suggesting both  
30 methods were measuring activity originating from the same microbial groups. We conclude that  
31 different labs using the same MUF microplate protocol tested, gives reasonably similar absolute  
32 activity values, variability, and ranking of treatments (highest to lowest). We propose that the MUF  
33 microplate method described in this study be considered as a standard protocol for assaying soil  
34 enzyme activities, providing that the buffer pH for the incubation be adjusted to the optimal pH  
35 according to the enzyme of interest.

36

37

38

39

40

41

42

43

44 Keywords: cross-lab study; soil enzyme methods; microplate; 4-methylumbeliferone; *p*-nitrophenol

45

46

47

48

49

50

51

## 52 **Introduction**

53           Soil enzyme assays have traditionally been performed at the bench scale using spectroscopic  
54 methods. There has been growing interest in fluorescence-based microplate methods to measure  
55 enzyme activities because fluorimetric detection is sensitive, and the microplate format has potential  
56 for simultaneous and rapid assaying of multiple enzyme activities from the same soil suspension. In  
57 contrast, bench-scale methodology has lower throughput and sensitivity. Microplate-format enzyme  
58 protocols have already been incorporated in many research laboratories; however, the lack of a  
59 standardized method for performing the assays and the potential for methodological problems  
60 (German et al. 2011) limits their usefulness for cross-study data comparisons and meaningful data  
61 interpretation. Several validation steps are recommended prior to standardization and widespread  
62 adaptation of new microbiological methods (AOAC, 2006; FEM, 2009; Green, 1996). Preliminary  
63 validation involves establishing performance characteristics such as specificity, sensitivity,  
64 reproducibility, and accuracy based on comparative testing with a reference method. Additionally, a  
65 collaborative study may be done to compare the assay's performance across laboratories.

66           Evaluative studies of various fluorimetric microplate assays using 4-methylumbelliferone  
67 (MUF) based substrates have been done in several laboratories (Deng et al., 2013; Drouillon and  
68 Merckx, 2005; Marx et al., 2001; Pritsch et al., 2004; Trap et al., 2012), and most were done in the  
69 same laboratories that developed the methods. However, the resulting data have not consistently had  
70 the same outcomes between the fluorescent microplate and the colorimetric *p*-nitrophenol (PNP)

71 bench-scale methods (Dick et al., 2013; Marx et al., 2001). Therefore, further cross-comparison  
72 between these methods is required to determine the sources of variation.

73         Methodological discrepancies may account for some differences in assay performance.  
74 Notable in the above studies were the different pHs under which the assays were performed. In  
75 some, the buffer pH was adjusted to the pH of the soil (Drouillon and Merckx, 2005; German et al.,  
76 2011; Trap et al., 2012), and in others, to the optimal pH for the enzyme (DeForest, 2009; Deng et  
77 al., 2013; Dick et al., 2013; Marx et al., 2001).

78         It has long been established that MUF fluorescence intensity is pH dependent, with the  
79 highest fluorescence signal around pH 10. Moreover, when NaOH is used to increase MUF  
80 fluorescence at the end of a fluorimetric enzyme assay, MUF fluorescence signal decreases over  
81 time after NaOH addition, making it challenging for obtaining quantitative fluorescence readings  
82 and for valid data comparison (DeForest, 2009; Drouillon and Merckx, 2005). German et al. (2011)  
83 found NaOH addition to be a significant source of variation and recommended against the NaOH  
84 addition for samples at pH 4.5 and higher, as they had no difficulty detecting the accumulation of  
85 MUF over time at  $\text{pH} \geq 4.5$ , even without increasing the sensitivity setting on their microplate  
86 fluorimeter. THAM pH 10-12 is the optimal solution to add after the enzyme assay because the  
87 fluorescence intensity highest due to the high pH and the signal remains stable up to three hours  
88 because of the THAM buffer (Deng et al., 2013).

89         One advantage of fluorimetric over colorimetric microplate methods is that fluorescence,  
90 unlike absorbance, is not increased by the presence of soil particles (Deng et al., 2013). However,  
91 fluorescence chemistry presents its own complexities because of quenching effects and the potential  
92 for chemical hydrolysis; both influence reproducibility. The quenching effect, due primarily to the  
93 presence of soil particles and dissolved organic matter in the assay mixtures, has been shown to

94 vary temporally and spatially (Freeman et al., 1995), and thus requires a calibration curve be  
95 developed for every sample assayed.

96 Thus, microplate enzyme assay methods introduce several sources of variation that have the  
97 potential to substantially affect assay reproducibility, and require careful execution of certain steps  
98 by the operator to minimize the variation. In part, this is due to the very small amounts of soil that  
99 must be used (0.83  $\mu\text{g}$  in a 250  $\mu\text{l}$  reaction volume in the current study) to minimize the quenching  
100 effect and to accommodate the small-volume microplate wells. Furthermore, dispensing  $\mu\text{L}$   
101 volumes of soil/buffer slurry with a pipette results in wide variations in the amount of soil added to  
102 each assay well. Because of this, the analytical error intrinsic to microplate enzyme assays is  
103 expected to be considerably larger than that of the conventional bench method. To minimize error,  
104 more replication is required compared to traditional bench-scale methods.

105 Thus, there are legitimate concerns about reproducibility and whether different labs can  
106 obtain comparable data from of the same samples. For valid data comparison and interpretation,  
107 there is clearly a need to have a strict and agreed-upon enzyme assay protocol to standardize the  
108 microplate-based method where different operators who follow these procedures obtain the same  
109 results. This is not case for the MUF microplate method as there has been considerable variation in  
110 the protocols used in the literature. Therefore, a cross-lab study was done on an optimized MUF  
111 procedure that has evolved from a number of labs (Marx et al., 2001; DeForest, J.L., 2009; German  
112 et al., 2011; Deng et al., 2011, 2013; Dick et al., 2013)

113 Since PNP bench-scale enzymes assays are widely used, largely vetted for standardization,  
114 and accepted, the objectives of this study were to determine the reproducibility by different  
115 laboratories for assaying activity of  $\beta$ -glucosidase (EC 3.2.1.21) and acid phosphomonoesterase (EC  
116 3.1.3.2) in the same soil samples using MUF microplate methods in comparison with standard PNP

117 bench-scale assays; and to determine the degree and the sources of variability associated with these  
118 two assays within and among five laboratories.

119

## 120 **Materials and Methods**

### 121 **Soils, sampling and storage**

122 The four soils selected were: 1) a sandy soil from the Ottokee fine sand series (sandy, mixed,  
123 mesic Aquic Udipsamments) under soybean (*Glycine max*) near Napoleon, Ohio (OTB); 2) a soil  
124 with higher clay content from the Miamian silt loam series (fine, mixed, active, mesic Oxyaquic  
125 Hapludalfs) under mixed grasses at the Ohio State University Waterman Farm, Columbus, Ohio  
126 (MMN); 3) a Jory silty clay loam (fine, mixed, active, mesic Xeric Palehumults) under Christmas  
127 tree (*Pseudotsuga menziesii* at time of sampling) management (32 yrs) and vegetation-free except  
128 for the trees, near Corvallis, Oregon (JMN); and 4) a Jory silty clay loam under > 90 yrs unmanaged  
129 Douglas Fir (*Pseudotsuga menziessi*) forest (JOG). The two Jory soil sites are side by side.

130 Approximately 3 kg of soil were collected at each sampling site along a transect at three  
131 spatially separated points (field replications) approximately 50 m apart. At each sampling site a 2 m  
132 diameter area was sampled by taking about 30 0-15 cm depth cores with a probe (2.54 cm dia.).  
133 Field moist soils were passed through a 2 mm sieve, which resulted in a thoroughly homogenized  
134 sample. A large portion was air-dried; while a small portion was left field-moist. Air drying was  
135 done by spreading soil on butcher block paper, spread at about 0.5 cm thick for 24 hours. The air-  
136 dried soils were separated into 500 g samples and placed in sealed zip lock bags for shipment on ice  
137 in Styrofoam containers to the collaborating laboratories. The field level replication was maintained  
138 throughout the research, with each laboratory receiving three separate replicates of each soil, for a  
139 total of 12 soil samples. The field-moist samples were either stored at 4°C for pH and soil texture



140 analyses, or stored at -20°C for Ester-Linked Fatty Acid Methyl Ester (EL FAME) analysis as an  
141 index microbial community composition.

## 142 **Soil chemical and biological properties**

143 Total C and N contents were measured by dry combustion (950°C) with a Vario Max CN  
144 Analyzer (Elementar; Hanau, Germany). Particle size distribution was determined by the pipette  
145 method (Kilmer and Alexander, 1949), and soil pH was determined using a 1:1 soil/water (v/v) ratio.  
146 Soil moisture content was measured by drying samples for 24 h at 105° C. Chemical and physical  
147 properties of the four soils used in the study are shown in Table 1.

148 The EL-FAME procedure was followed as described by Schutter and Dick (2000). In brief,  
149 soil samples (3 g wet weight) were extracted with 0.2 M KOH in methanol, then incubated for 1 h at  
150 37 °C with periodic vortexing followed by solution neutralization of the pH with 1.0 M acetic acid.  
151 Extracted FAMES were partitioned into an organic phase with hexane and centrifuged for 20 min at  
152 500 x g to with and then the separated hexane phase with EL-FAMES was dried under a stream of  
153 N<sub>2</sub>. FAMES were resuspended in a known volume of hexane. FAME composition in the extracts  
154 were determined using GC (Agilent 6890, Agilent Inc., Wilmington, DE) equipped with a 25–m HP  
155 Ultra-2 column (internal diameter, 0.2 mm; film thickness, 0.33 µm) and a flame ionization detector.  
156 The temperature program ramped from 170 to 280 °C at 4 °C per min, with 5 min at 280 °C  
157 between samples to clean the column. Individual fatty acids were identified relative to several  
158 standards: 37 FAMES mixture (FAME 37 47885-4; Supelco, Inc), 24 bacterial FAMES mixture (P-  
159 BAME 24 47080-U; Supelco, Inc.), and MIDI standards (Microbial ID, Inc.). Quantification of  
160 FAMES was accomplished by using varying concentrations of tridecanoic FAME (Supelco, Inc.)  
161 and allowed peak areas to be converted to a molar basis (correlations with enzyme activities were  
162 done on EL-FAMES on a per nmol basis). Fatty acids comprising less than 0.5 % of the total

163 relative abundance were not included in the data analysis. Sums of the fatty acids indicative of  
164 Gram-positive bacteria (GM+), Gram-negative bacteria (GM-), actinomycetes (actinos), plus the  
165 five additional fatty acids listed are used as a measure of total bacterial biomass (Frostegard and  
166 Bååth, 1996)

## 167 **Enzyme assays**

168 Two microplate enzyme assays using fluorescent methylumbelliferyl substrates, and two  
169 bench-scale assays using chromogenic *p*-nitrophenyl substrates were compared across five  
170 laboratories, each using identical protocols to measure  $\beta$ -glucosidase (EC 3.2.1.21) and acid  
171 phosphomonoesterase (EC 3.1.3.2) activities in each of the 12 soil samples. The enzyme assays  
172 were completed within two weeks of receiving the samples, with the exception of the bench-scale  
173 assay in one laboratory that was completed within four weeks.

174 The five laboratories are located in three countries across two continents. All participating  
175 laboratories had experience using fluorescence-based microplate methods. Labs 1 and 2 had  
176 experience with the protocol used in the current study, which was developed in Lab 2. Lab 5 did not  
177 have experience with the PNP bench-scale assay, and did not perform it in this study.

## 178 MUF Microplate Enzyme Assay

179 The MUF assay was done as described by Deng et al. (2011) with minor modifications.  
180 Microplate layouts in the assay are shown in Figure 1, and were followed by all participating  
181 laboratories. See supplemental section for detailed description of the method.

182 All reagents were brought to incubation temperature (37 °C) prior to beginning the assay.  
183 Two replicate soil suspensions were prepared for each sample by placing 1 g of soil into a 150 mL  
184 beaker with 120 mL of deionized H<sub>2</sub>O (dH<sub>2</sub>O). The soil was homogenized for 30 min using a 3.75  
185 cm magnetic stir bar and a stir plate set to 600 rpm. Soil suspension (100  $\mu$ L) was transferred with

186 continuous stirring to microplate wells containing 50  $\mu$ L of modified universal buffer (MUB, pH 6),  
187 using a multichannel pipette with four wide orifice tips, resulting in four microplate replicates from  
188 each suspension for each assay. Methylumbelliferyl- $\beta$ -D-glucoside or methylumbelliferyl-  
189 phosphate substrate dissolved in water (2 mM; 50  $\mu$ L) was subsequently added to each well.  
190 Following mixing by pipetting up and down 2 to 3 times, the plates were then covered, placed in a  
191 shallow water bath and incubated at 37° C for 1 h. Upon completing incubation, 50  $\mu$ L of Tris  
192 (hydroxymethyl) aminomethane (THAM) pH 10 was added to each well to bring the reaction  
193 mixtures to a pH that optimizes MUF fluorescence (pH 10 to pH 11) (Deng et al., 2013). Relative  
194 fluorescence was measured at 360 nm excitation and 460 nm emission. The volume in each reaction  
195 well during incubation was 200  $\mu$ L, and the total volume after reaction termination was 250  $\mu$ L. The  
196 labs did not use the same model microplate readers; however sensitivity (gain) settings based on the  
197 highest calibration standard were used to normalize results.

198 In addition to the four assay sample replicates, four control replicates were prepared for each  
199 suspension in the same way, except substrate was added after incubation and THAM addition. Four  
200 additional controls for autohydrolysis (AH) were set up one time only for each substrate, in which  
201 dH<sub>2</sub>O was used in the place of soil suspension, and substrate was added to two of the AH controls  
202 before incubation, and to two AH controls after incubation and THAM addition. Reagents were not  
203 all purchased from the same manufacturer because of limited availability and accessibility of  
204 laboratories involved in the study. However, controls employed in the assay accounted for substrate  
205 hydrolysis from substrate source and/or storage.

206 A soil-specific calibration curve was developed for each sample at the same time the assay  
207 was performed, using the same procedure and soil suspensions, except 50  $\mu$ L of each MUF standard  
208 was used in place of MUF substrate. Microplate wells of the prepared standards contained 0, 250,

209 500, 1000, 1500 or 2500 total pmol MUF. An average slope was obtained for each soil from two  
210 sets of standards (one from each replicate soil suspension), and was used to calculate enzyme  
211 activity ( $\mu\text{mol g}^{-1}$  dry soil  $\text{h}^{-1}$ ).

#### 212 PNP Bench-scale Enzyme Assay

213 The bench-scale method was performed as described previously (Tabatabai, 1994; Tabatabai  
214 and Bremner, 1969), but without the use of toluene (Drouillon and Merckx, 2005; Vuorinen and  
215 Saharinen, 1996). All reagents were brought to 37° C prior to performing the assay. Two assay  
216 replicates were performed for each soil by weighing 1 g of soil into a 50 mL Erlenmeyer flask,  
217 adding 4 mL of MUB (pH 6.0) and 1 mL of 0.05 M *p*-nitrophenyl- $\beta$ -D-glucoside or *p*-nitrophenyl  
218 phosphate, and then stoppering and swirling the flask to mix. A control was prepared for each  
219 sample following the same procedure, but with substrate added after the reaction was terminated;  
220 Flasks were incubated at 37° C for 1 h. Following incubation, 4 mL of THAM pH 12 and 1 mL of  
221 0.5 M  $\text{CaCl}_2$  were added, swirled to mix, and the suspension was passed through a Whatman #2  
222 filter. Filtrate absorbance was measured at 405 nm on spectrophotometer. Absorbance values from  
223 both controls were subtracted from those obtained for samples prior to calculating the amount of *p*-  
224 nitrophenol released during the incubation using a standard calibration curve, which was developed  
225 using the same procedure as for samples and with standards containing 0, 100, 200, 300, 400 or 500  
226 nmol *p*-nitrophenol in MUB. Dilutions of the sample filtrates were made with a 1:1 mixture of  
227 MUB pH 6.0 and 0.1 M THAM pH 12 when color intensity exceeded that of the highest *p*-  
228 nitrophenol standard solution.

#### 229 **Data analysis**

230 The SAS Univariate procedure (SAS version 9.3) was used to evaluate distribution of the  
231 data. Significant differences among soils and labs were determined using one-way analysis of

232 variance (ANOVA) and interactions between soil and lab factors were evaluated by 2-way ANOVA.  
233 Individual contrasts were subsequently analyzed using Tukeys HSD ( $p \leq 0.05$ ) post-hoc test. Simple  
234 correlation procedure (PROC CORR, SAS, 1999) was used to examine relationships between  
235 enzyme activity between the MUF microplate and bench methods.

236 To identify the source of variation between the methods and the degree of variation among  
237 the labs, coefficients of variation (CV) were calculated as standard deviation divided by the mean,  
238 and multiplied by 100 to convert to percentages. For the microplate method, the variability within  
239 suspensions was determined by calculating CVs using mean assay values, while those accounting  
240 for field replications were calculated using averaged activities from the two suspensions. Similarly,  
241 CVs accounting for the spatial and field scale of variability for the bench method were calculated  
242 using means from the replicate suspensions.

243

## 244 **Results and Discussion**

245 The selected soils were quite different in physical and chemical properties (Table 1). The pH values  
246 ranged from 5.2 to 7.3. Organic C varied widely, with over twofold more in the unmanaged Jory  
247 soil than the cultivated Ottokee soil. The higher organic C levels are likely due to the higher clay  
248 content of Jory and Miamian soils. However, in the case of the managed Jory soil, long-term  
249 Christmas tree cultivation has led to significant reduction in organic C content (Table 1). At this site,  
250 Christmas trees have been planted and harvested at about 5-year intervals and were sprayed with  
251 herbicides. Additionally, the Christmas tree operation keeps the inter-row area vegetation free. As a  
252 result, organic matter input to the soil was greatly reduced, and regular disturbance from equipment  
253 and tree harvesting has likely reduced soil organic matter in the managed Jory soil compared to the  
254 unmanaged forest Jory soil.

255 Both enzyme methods detected differences in activities between soil types or within a soil  
256 type (Jory soil) under different management practices (Table 2). Across both methods and for both  
257 enzymes, the unmanaged Jory soil consistently had the highest activities. For the most part, enzyme  
258 activities were highly correlated with clay content (Allison and Jastrow, 2006). Acid  
259 phosphomonoesterase activity was lowest in the sandy soil, Ottokee, which was expected as this  
260 soil had significantly lower levels of clay, organic C and microbial biomass (EL-FAME) than the  
261 other soils. However,  $\beta$ -glucosidase activity was lowest in the Miamian soil even though it has  
262 significantly higher clay content than the Ottokee soil.

### 263 **Bench vs. Microplate**

264 While there were clearly differences in enzyme activities detected by PNP bench-scale vs.  
265 MUF microplate methods, the relative ranking of the soils was basically the same for both methods,  
266 especially for acid phosphomonoesterase activity (Table 2). Correlations between bench and  
267 microplate assays were strong for both enzymes, although slightly stronger for acid  
268 phosphomonoesterase ( $r = 0.93$ ) than  $\beta$ -glucosidase ( $r = 0.81$ ) (data not shown).

269 Averaging the field replicates across all labs, the PNP bench-scale method consistently gave  
270 higher activities than those measured with the MUF microplate method (Table 2). For Jory managed,  
271 Jory unmanaged, Miamian, and Ottokee soils, mean activities of acid phosphomonoesterase by  
272 MUF microplate method were 73, 71, 67 and 28%, respectively, of those detected using the PNP  
273 bench method; while mean  $\beta$ -glucosidase activities determined by the MUF microplate method  
274 were 88, 93, 81 and 75%, respectively, of those measured by the PNP bench method.

275 Higher enzymes activities detected by PNP bench over MUF microplate methods are consistent  
276 with the results of Marx et al. (2001) and Trap et al., (2012). However, in the case of Marx et al.  
277 (2001) this difference between the methods could be due to use of different incubation temperatures

278 (microplate assays at 30 vs. 37°C for bench method). Drouillon and Merckx (2005) for  
279 phosphomonoesterase activity on a wide range of soils, found the PNP method was significantly  
280 lower than the MUF microplate method – nor were the activities of the two methods correlated.  
281 However, their assays were run at the soil pH. Variations in pH of these soils could have affected  
282 the comparability of phosphatase activities detected by the 2 methods. In contrast, some studies did  
283 not find significant differences between the two methods for N-actyl- $\beta$ -D-glucosaminidase (Popova  
284 and Deng, (2010) and for  $\beta$ -glucosidase (Dick et al., 2013) in diverse soil types. Deng et al. (2013)  
285 found that N-actyl- $\beta$ -D--glucosaminidase,  $\beta$ -glucosidase, and acid phosphomonoesterase activities  
286 in 16 soils were often significantly different (mostly with PNP being greater than MUF) but were in  
287 the same order of magnitude. As with this study, they found that activities determined by the two  
288 methods were significantly correlated.

289         The differences between the PNP and MUF-based methods in this study were not consistent  
290 across soil types and enzymes. There are likely a number of factors for differences between the  
291 methods that mechanistically cannot be isolated. Although there is greater affinity by MUF over  
292 PNP (Km values presented by Marx et al, 2001; Deng et al., 2013) which could change the activity  
293 rate, it should be a consistent effect. A more likely factor contributing to variability detected by the  
294 MUF method is quenching of the emitted fluorescence by soil particles. The degree of quenching  
295 varies temporally and spatially both within soil replicates as a function of uneven particle  
296 distribution resulting from pipetting variations, and between soil replicates as a function of  
297 differences in SOM content (Freeman et al., 1995). In theory, quenching of MUF in soil can be  
298 accounted for by having a standard calibration curve for each soil, but this is confounded by  
299 variations in particle density inherent to pipetting soil slurries. The resulting higher analytical

300 variability of the MUF microplate method (Tables 3 to 6) could explain some of the inconsistencies  
301 between the MUF microplate and PNP bench methods.

302         Although there were differences between the two methods, the differences were within an  
303 order of magnitude and did not alter the ranking of soil types between the two methods as discussed  
304 earlier. Furthermore, activities for the same enzyme in the same soil were highly correlated,  
305 suggesting that the same pool or a large portion of overlapping pools of enzymes were detected by  
306 the different approaches. This finding is supported by studies of Marx et al. (2001) and Dick et al.  
307 (2013) who reported similar  $V_{\max}$  for PNP- and MUF-based substrates in the detection of  $\beta$ -  
308 glucosidase and acid phosphomonoesterase activities using the microplate format assays.

309         Because the soils were air-dried, the time between sampling and analysis likely did not  
310 affect enzyme activities detected by either assay method. We did a preliminary study showing  $\beta$ -  
311 glucosidase on 2 diverse soils where the activity on air dried soil was constant p to 6 months  
312 (personal communication R Dick). Enzyme analyses were conducted from 3 to 6 weeks after soils  
313 were sampled, sieved, homogenized, and air-dried. The results were much closer for replicates of  
314 the same soil sample than between soil type providing evidence that the storage time that varied  
315 across labs was not a factor (meaning air dried samples were stable over time). This is in line with  
316 Bandick and Dick (1999) who showed that air drying soils for a range of enzyme activities  
317 (including acid phosphomonoesterase and  $\beta$ -glucosidase activity) did not alter the ranking of soil  
318 samples compared to analyses on fresh, field moist samples. The ability to use air-dried soil is an  
319 advantage over most other microbial properties that require rapid analysis on fresh soils.

320 Commercial labs often prefer the use of dried soils to accommodate for varying shipping times.

321         To give some insight into the relationship between the two enzyme activities assayed by  
322 bench and microplate methods and microbial community profiles, fatty acid methyl esters (EL-



323 FAMES) were used as biomarkers for major functional microbial groups. The biomass of each  
324 microbial group was correlated with enzyme activities (Table 3). For both acid  
325 phosphomonoesterase and  $\beta$ -glucosidase,  $r$  values were mostly similar for MUF microplate and  
326 PNP bench method correlation with EL-FAME biomarkers. Interestingly, the arbuscular  
327 mycorrhizal fungal marker was poorly correlated to  $\beta$ -glucosidase activity, but significantly  
328 correlated to acid phosphomonoesterase activity by both assay methods. The results support the  
329 conclusion that both enzyme methods detected activities from the same pools of enzymes in soil.  
330 Furthermore, these results suggest that there is a relationship between enzyme activities and the size  
331 of various microbial groups, and differences in the potential of these groups to produce  $\beta$ -  
332 glucosidase and acid phosphomonoesterase.

### 333 **Laboratory Comparisons**

334 Differences among labs were observed, and some were statistically significant (Table 2). A  
335 2-way ANOVA of the lab by soil type factors had a significant interaction ( $P < 0.001$ ). For both  
336 microplate and bench assays, there were slightly fewer significant differences among labs for  $\beta$ -  
337 glucosidase activity than were observed for acid phosphomonoesterase activity. Labs 3 and 4 tended  
338 to report higher enzyme activities compared to Labs 1 and 2 (both for MUF and PNP assays),  
339 whereas Lab 5 tended to report the lowest activities (MUF assay only). However, these differences  
340 were not always significant. Although there were differences between labs for both methods within  
341 a soil type, the mean values and ranking of the soils were very similar among the labs. Thus,  
342 differences among labs were small compared to differences among soils, and are more likely a  
343 reflection of low variance within each lab which enabled significant ( $P < 0.05$ ) detection of small  
344 differences among labs. However, it should be pointed out that analysis of variance can only  
345 determine differences between treatments and does not determine whether treatments are the same

346 on a probability basis. Thus, although there were at times small differences between labs for either  
347 enzyme method under evaluation, each lab's results would largely lead to the same conclusion in  
348 ranking the soil samples.

#### 349 **Levels of Variability Between and Within Assay Methods**

350 Coefficients of variation (CVs) were calculated for each lab and soil by enzyme assay for  
351 the MUF microplate method at three levels: microplate (four replicates); soil suspension (two  
352 replicates); and field (three replicates) (Tables 4 and 5). Average CVs across all soil samples and  
353 labs for the microplate method were 14% and 18 % for acid phosphomonoesterase and  $\beta$ -  
354 glucosidase activities, respectively. At the suspension level, average CVs were 12 and 11 %,   
355 respectively.

356 Coefficients of variation among the four replicate microplate wells ranged from 2 to 118 %  
357 (Tables 4 and 5). The highest CVs tended to occur in the sandy (Ottokee) soil at all levels  
358 replication (microplate and suspension). At the suspension level for the MUF microplate method,  
359 CVs ranged from 0 to 48% (Tables 4 and 5). Eleven of the 15 CVs in the sandy soil were greater  
360 than 10%, comparing to other soils where 2 to 5 of the 15 CVs were greater than 10%. However, it  
361 should be pointed out that the CVs would naturally be high simply because of the low number of  
362 replications (n=2). Sandy soils are inherently difficult to homogenize and dispense. Although the  
363 observed difference is not unexpected, this may affect precision when using microplate methods.  
364 These results suggest that greater suspension and microplate replication are needed to reduce the  
365 variability for sandy soils. Similar, but somewhat lower variability was also observed in the PNP  
366 bench method.

367 Pipetting error is another potential source of variability for the MUF assay due to the  
368 difficulty of evenly dispensing  $\mu$ L-scale volumes of soil suspensions and the possibility of soil

369 particles clogging the pipette tips and going unnoticed by the operator. Other sources of error  
370 associated with reagents (e.g. sources and reagent make up) and experimental conditions (e.g.  
371 managing temperature) would likely cause a systemic error or variation that is constant across all  
372 samples within a lab (thus contributing to differences among labs but not within a lab). Average  
373 CVs by lab across all soil samples ranged from 14 % (lab 1) to 23 % (lab 5) for  $\beta$ -glucosidase  
374 activities (Table 4) and 13 % (lab 2) to 17 % (lab 5) for acid phosphomonoesterase activities (Table  
375 5).

376 For the PNP bench method, the source of variation equivalent to MUF microplate  
377 suspension is assay replication (n=2 for bench method) (Tables 6 and 7), which overall had lower  
378 CVs than the MUF microplate suspensions (Tables 6 and 7). Averaging across all soil samples and  
379 labs for the PNP bench method, the CVs ranged from 4 % (lab 1) to 5 % (lab 3) and 3 % (lab 2) to  
380 6 % (lab 3) for acid phosphomonoesterase and  $\beta$ -glucosidase activities, respectively. An advantage  
381 of the bench method is that one gram of soil is used in each assay compared to the 0.83  $\mu$ g in each  
382 microplate well. This could be a major factor contributing to the substantially lower CVs of the  
383 bench over the microplate methods at the analytical level of replication. Alternatively, the  
384 microplate method allows for very small samples to be analyzed for studies at small spatial scales.

385 From a practical perspective, it is easier to increase the replicates on the microplate than it is  
386 to increase suspension replication, especially considering the 30 minutes required for  
387 homogenization of each suspension. Many laboratories include as many as 16 replicate wells per  
388 suspension (e.g. following the protocol of Saiya-Cork et al. 2002). However, replications of the  
389 same soil suspension account mainly for pipetting error, and do not account for errors originating  
390 from soil heterogeneity and/or treatments under evaluation. Results from this study suggest that  
391 activities obtained from two suspensions were consistent. This is in agreement with Dick et al

392 (2013), who did an in-depth analysis of the reproducibility of the suspension step in the MUF  
393 microplate method.

394 At the field level, CVs ranged from 1% to 90% among field replicates for the MUF assay,  
395 and 2% to 60% for the PNP assay, also with higher CVs in the sandy soil. Again, the bench-scale  
396 method resulted in CVs that were generally lower than those for the MUF microplate method.

397 A factorial analysis of mean activity for each assay across all four soils for each lab, and  
398 across all five labs for each soil showed significant differences based on both lab and soil (Table 2).  
399 Means of field replicate across all labs for each soil showed some significant differences, except in  
400 the sandy (Ottokee) soil, which may have been the result of the greater variability discussed above.

#### 401 **Standardization of the MUF Microplate Method**

402 The MUF microplate method has a number of protocol conditions or nuances that could  
403 make it difficult to obtain reproducible results by different operators and/or lab equipment. These  
404 include the need to fully suspend soils in solution, maintain buffer pH (Lakowicz, 1983) and a  
405 constant temperature throughout the incubation (Lakowicz, 1983), accurately pipette  $\mu\text{L}$  volumes,,  
406 and carefully account for quenching. An alternative protocol as outlined by Marx et al (2001) is not  
407 to add base and then to measure the linear increases in fluorescence. This omits the no-substrate  
408 control and potentially produces more accurate quantification of the enzymatic reaction rates based  
409 on multiple data points rather than one single data point. However, a calibration curve is needed  
410 under identical assay conditions. This method is also considerably less sensitive. As shown in Deng  
411 et al. (2013), the relative MUF fluorescence signal at pH 10 could be greater than 35000 times of  
412 those at pH 6. However, German et al. (2011) found that fluorescence signal was detectable in soils  
413 with  $\text{pH} \geq 4.5$ .

414 Methods used to prepare soil suspensions are rarely reported in the literature. Creating soil  
415 suspensions could be a major source of variation between labs, as type of homogenizer, size of  
416 glassware and stir bars, and differences in mixing speed and time could all affect the outcome. For  
417 this study, all labs followed the same soil suspension protocol (container, stir bar size, and stirring  
418 time and speed). This is a protocol point that should be considered as a standard inclusion for  
419 describing the MUF microplate method in the literature. Indeed, the basic soil suspension protocol  
420 followed in this study is based on the investigation of various factors that might affect soil  
421 suspension reproducibility as reported by Dick et al. (2013). They found using a stir bar provided  
422 reproducible results across 6 suspension replications for both a loamy soil and a sandy soil for the  
423 MUF microplate method.

424 Quenching was accounted for in the current study by using a standard curve for each soil.  
425 Variation in the amount of minerals and organic matter in each well affect the degree of quenching  
426 and therefore variability of the results. Temperature control is important as well because MUF  
427 florescence is very sensitive to temperature (Guilbault, 1990). We found that it is important to have  
428 all reagents at the incubation temperature prior to starting the substrate-soil incubation because it  
429 could take nearly 30 minutes of the incubation time to allow reaction mixtures to reach the protocol  
430 temperature of 37 °C.

431 It is also important to note that we used the optimal pH to maximize the activity of each  
432 enzyme (based on previous reports in the literature). Some researchers have argued that the  
433 incubation solution should be left unbuffered or adjusted to pH of the assayed soil (Drouillon and  
434 Merckx, 2005; German et al. 2011), with the goal to obtain in situ enzyme activities in the field.  
435 However, doing so prevents direct comparisons across studies and among different soils, which  
436 were goals of this study. Furthermore, if there are no treatment effects, it cannot be concluded that

437 there were no differences because it could be that it was simply due to using the non-optimized  
438 conditions of the assay. To determine what the optimal pH is for a given enzyme assay the reader is  
439 referred to Dick (2011) which is a comprehensive compilation of vetted methods, largely done by  
440 Tabatabai and co-workers, where each method has a recommended optimal pH. However, to be  
441 absolutely certain of the optimal pH for specific soils, pH curves should be developed before  
442 proceeding with investigations that use particular enzyme assays.

443 In this study, THAM buffer at pH 10 was used instead of NaOH to terminate the reaction.  
444 Deng et al. (2013) reported that fluorescence of MUF is not stable in the presence of NaOH and that  
445 THAM at pH 10-11 is optimal for maintaining fluorescence. We imposed a one-hour maximum for  
446 taking the readings after THAM addition (effectively terminating the reaction). This is because  
447 fluorescing expends energy, and once initiated fluoresce decreases over time. Hence, we  
448 recommend that this be a standard description in MUF microplate methods, and that authors report  
449 on how quickly the fluorescence was read.

450 We propose that the detailed MUF microplate method described in the Supplemental  
451 Section be the standard protocol except that the buffer pH for the incubation be adjusted to the  
452 optimal level according to the enzyme of interest. In brief we are suggesting that the following  
453 operationally defined protocols be adopted: 37<sup>0</sup>C incubation temperature, pH 10 THAM buffer to  
454 terminate the enzymatic reaction, 1 hour incubation, controlling for quenching in standard curve and  
455 autohydrolysis, pre-warming incubation reagents, and method of developing soil suspension. These  
456 can be adopted because the results of MUF microplate method to that of PNP bench method were  
457 quite similar in absolute values and in ranking of the soil treatments. It should be noted that the  
458 optimum pH for the enzyme assays where this is known, is largely based on soils from temperate  
459 regions (Tabatabai, 1994). As such further investigations are warranted on soils from other regions,

460 as variation across soils for enzyme sources and stabilization, and/or soil chemistry could affect  
461 optimum assay pH for a given soil.

## 462 **Conclusions**

463         The MUF microplate and bench methods each have advantages and disadvantages. The  
464 MUF microplate method in theory should be a high throughput method. From our experience this is  
465 the case if multiple enzymes are assayed together. However, if the assay is for one enzyme with  
466 multiple soil samples it is less clear that the microplate method takes less time because of the need  
467 to set up soil suspensions which is time consuming and takes considerable bench space. One  
468 additional advantage of the MUF microplate assay is the ability to do the assay on small amounts of  
469 soils (but only if the soil:solution ratio is kept the same as when 1 g of soil is put into solution).

470         To enable cross-literature comparisons and meta-analyses, it is important to employ enzyme  
471 assay protocols that minimize the potential for operator biases and generate reproducible results  
472 across labs. The MUF microplate method increases the number of points at which the operator can  
473 influence outcomes. These include homogenization and suspension of soil, retrieval and dispensing  
474 of  $\mu\text{L}$  volumes of soil slurries, and reading on a fluorimeter or spectrophotometer (DeForest, 2009;  
475 German et al., 2011; Trap et al., 2012). Of these, the most likely source of variation from an  
476 operator perspective is pipetting. Indeed, in this study overall variability was highest for replication  
477 on the microplate ( $n=4$ ), and the suspension replication actually had lower CVs with  $n$  only  
478 equaling 2. In comparison, the bench-scale method had significantly lower CVs for analytical and  
479 field replicates than the MUF microplate method. Most likely this is due the much larger soil  
480 amount that is used at the bench scale.

481         It has become increasingly apparent that MUF microplate protocols require very tight  
482 standardization and greater replication in order to yield results that are comparable either within or

483 across laboratories. In the current study, effort was made to control for methodological differences  
484 by using air-dried soils (therefore more stable over varying storage periods) and identical assay  
485 protocols and method for soil homogenization. Reagents were preheated and incubated in a water  
486 bath. Microplate reader sensitivity was set to the highest standard, readings were taken within an  
487 hour of base addition, and controls were included for substrate autohydrolysis.

488           When all labs adhered to a strict and consistent protocol, the differences among lab results  
489 were much smaller than differences due to soil type. In this study, the results for both bench and  
490 MUF microplate method from different labs were quite similar in terms of ranking of soil  
491 management treatments and soil types (Table 2). This suggests that different labs can come to the  
492 same conclusions on the same set of soil samples using a standardized MUF microplate protocol.  
493



## REFERENCES

- 494  
495  
496 Allison, S.D., Jastrow, J.D., 2006. Activities of extracellular enzymes in physically isolated  
497 fractions of restored grassland soils. *Soil Biol. Biochem.* 38, 3245-3256.
- 498 Association of Official Agricultural Chemists, 2006. AOAC Task Force recommends best practices  
499 for validation of microbiological methods.
- 500 Bandick, A.K., Dick, R.P., 1999. Field management effects on soil enzyme activities. *Soil Biol.*  
501 *Biochem.* 31, 1471-1479.
- 502 DeForest, J.L., 2009. The influence of time, storage temperature, and substrate age on potential soil  
503 enzyme activity in acidic forest soils using MUB-linked substrates and l-DOPA. *Soil Biol.*  
504 *Biochem.* 41, 1180-1186.
- 505 Deng, S., Kang, H., Freeman, C., 2011. Microplate fluorimetric assay of soil enzymes in: Dick, R.P.  
506 (Ed.), *Methods of Soil enzymology*. Soil Science Society of America, Madison, WI, pp.  
507 311-318.
- 508 Deng, S.P., Popova, I.E., Dick, L., Dick, R., 2013. Bench scale and microplate format assay of soil  
509 enzyme activities using spectroscopic and fluorometric approaches. *Appl. Soil Ecol.* 64, 84-  
510 90.
- 511 Drouillon, M., Merckx, R., 2005. Performance of para-nitrophenyl phosphate and 4-  
512 methylumbelliferyl phosphate as substrate analogues for phosphomonoesterase in soils with  
513 different organic matter content. *Soil Biol. Biochem.* 37, 1527-1534.
- 514 Dick, L.K., Jia, G., Deng, S., Dick, R.P., 2013. Evaluation of Microplate and Bench-scale  $\beta$ -  
515 glucosidase Assays for Reproducibility, Comparability, Kinetics, and Homogenization  
516 Methods in Two Soils. *Biol. Fertility Soils* In Press.
- 517 Eivazi, F., Tabatabaai, M.A., 1988. Glucosidases and galactosidase in soils. *Soil Biol. Biochem.* 20,  
518 601-606
- 519 Drouillon, M., Merckx, R., 2005. Performance of para-nitrophenyl phosphate and 4-  
520 methylumbelliferyl phosphate as substrate analogues for phosphomonoesterase in soils with  
521 different organic matter content. *Soil Biol. Biochem.* 37, 1527-1534.
- 522 Fem Microbiology Action Team, 2009. Method validation of U.S. Environmental Protection  
523 Agency microbiological methods of analysis. Fem Document Number 2009-01. October 7,  
524 2009.
- 525 Freeman, C., Liska, G., Ostle, N.J., Jones, S.E., Lock, M.A., 1995. The use of fluorogenic substrates  
526 for measuring enzyme-activity in peatlands. *Plant and Soil* 175, 147-152.
- 527 Frostegard, A., and E. Bååth. 1996. The use of phospholipid fatty acid analysis to estimate bacterial  
528 and fungal biomass in soil. *Biology and Fertility of Soils* 22:59-65.

- 529 German, D.P., Weibtraub, M.N., Grandy, A.S., Lauber, C.L., Rinkes, Z.L., Allison, S.D., 2011.  
530 Optimization of hydrolytic and oxidative enzyme methods for ecosystem studies. *Soil Biol.*  
531 *Biochem.* 43, 1387-1397.
- 532 Green, J.M., 1996. A practical guide to analytical method validation. *Anal. Chem.* 68, 305A-309A.
- 533 Guilbault, G.G., 1990. *Practical Fluorescence*, Second Edition ed. Marcel Dekker, Inc., New York.
- 534 Lakowicz, J.R., 1983. *Principles of Fluorescence Spectroscopy*. Plenum Press, New York.
- 535 Marx, M.C., Wood, M., Jarvis, S.C., 2001. A microplate fluorimetric assay for the study of enzyme  
536 diversity in soils. *Soil Biol. Biochem.* 33, 1633-1640.
- 537 Popova, I.E., Deng, S., 2010. A high-throughput microplate assay for simultaneous colorimetric  
538 quantification of multiple enzyme activities in soil. *Appl. Soil Ecol.* 45, 315-318.
- 539 Pritsch, K., Raidl, S., Marksteiner, E., Blaschke, H., Agerer, R., Schloter, M., Hartmann, A., 2004.  
540 A rapid and highly sensitive method for measuring enzyme activities in single mycorrhizal  
541 tips using 4-methylumbelliferone-labelled fluorogenic substrates in a microplate system. *J.*  
542 *Microbiol. Methods* 58, 233-241.
- 543 SAS Institute Inc., 1999. *SAS User's Guide: Statistics*. SAS Institute Inc, Cary, NC.
- 544 Schutter, M.E., and R.P. Dick. 2000. Extraction methods of fatty acids for microbial community  
545 structure analysis. *Soil Sci. Soc. Am. J.* 64:1659-1668.
- 546 Tabatabai, M.A., 1994. Soil enzymes, in: Weaver, R.W., Angle, J.S., Bottomley, P.S. (Eds.),  
547 *Methods of Soil Analysis part 2. Microbiological and Biochemical Properties*. Soil Science  
548 Society of America, Madison, WI, pp. 775-833.
- 549 Tabatabai, M.A., Bremner, J.M., 1969. Use of p-nitrophenyl phosphomonoesterase for assay of soil  
550 phosphomonoesterase activity. *Soil Biol. Biochem.* 1, 301-307.
- 551 Trap, J., Riah, W., Akpa-Vinceslas, M., Bailleul, C., Laval, K., Trinsoutrot-Gattin, I., 2012.  
552 Improved effectiveness and efficiency in measuring soil enzymes as universal soil quality  
553 indicators using microplate fluorimetry. *Soil Biol. Biochem.* 45, 98-101.
- 554 Vuorinen, A.H., Saharinen, M.H., 1996. Effects of soil organic matter extracted from soil on acid  
555 phosphomonoesterase. *Soil Biol. Biochem.* 28, 1477-1481.  
556  
557  
558

559  
 560  
 561  
 562  
 563  
 564  
 565  
 566  
 567  
 568  
 569  
 570  
 571  
 572  
 573  
 574  
 575  
 576  
 577  
 578  
 579  
 580  
 581  
 582  
 583  
 584  
 585  
 586  
 587  
 588  
 589

Table 1. Description and properties of soils

Soil	Vegetation	pH	Total C	Total N	Sand	Silt	Clay
			-----g kg <sup>-1</sup> -----			-----%-----	
Jory silty clay loam (fine, mixed, active mesic Xeric palehumult)	Christmas trees (managed)	5.2	27.9	2.2	46	38	16
Jory silty clay loam (fine, mixed, active mesic Xeric palehumult)	Douglas Fir (unmanaged)	6.0	36.8	2.7	52	34	14
Miamian silt loam (fine, mixed, active, mesic Oxyaquic Hapludalfs)	Mixed grasses (unmanaged)	5.8	27.9	2.0	38	40	22
Ottokee fine sand (mixed, mesic Aquic Udipsamnents)	Row cropped (managed)	7.3	17.6	ND†	90	8	2

†Not detectable.

Table 2. Soil enzyme activities by bench or microplate methods across labs

	Jory Managed	Jory Unmanaged	Miamian Unmanaged	Ottokee Row-Crop	
-----μmol g <sup>-1</sup> h <sup>-1</sup> -----					
Lab	<b><u>PNP bench-scale acid phosphomonoesterase</u></b>				□
1	1.93b <sup>†</sup> (0.179) <sup>††</sup>	3.90b (0.092)	3.18bc (0.343)	0.49a (0.219)	2.38B <sup>‡</sup>
2	1.76b (0.156)	3.30c (0.277)	2.76c (0.221)	0.42a (0.218)	2.06C
3	2.31a (0.143)	4.38a (0.187)	3.95a (0.468)	0.55a (0.280)	2.80A
4	2.07ab (0.071)	4.55a (0.168)	3.87b (0.490)	0.81a (0.297)	2.83A
5	NA	NA	NA	NA	NA
□	2.02C <sup>‡‡</sup>	4.03A	3.44B	0.57D	
	<b><u>MUF microplate acid phosphomonoesterase</u></b>				
1	1.40b (0.076)	2.75b (0.233)	1.97a (0.438)	0.19a (0.065)	1.58 BC
2	1.50b (0.081)	2.94b (0.191)	2.23a (0.442)	0.16a (0.071)	1.71 B
3	1.88a (0.085)	2.68b (0.225)	2.38a(0.264)	0.18a (0.044)	1.78 AB
4	1.24b (0.325)	3.85a (0.369)	2.53a (0.304)	0.14a (0.078)	1.94 A
5	1.34b (0.087)	1.99c (0.312)	2.40a (0.483)	0.14a (0.129)	1.47 C
□	1.47C	2.84A	2.30B	0.16D	
	<b><u>PNP bench-scale β-glucosidase</u></b>				
1	0.54b (0.015)	1.14a (0.225)	0.55a (0.038)	0.68a (0.065)	0.73B
2	0.50b (0.045)	1.08b (0.170)	0.55a (0.021)	0.63a (0.215)	0.69B
3	0.72a (0.036)	1.60a (0.210)	0.65a (0.389)	0.89a (0.235)	0.97A
4	0.51b (0.031)	1.08b (0.175)	0.74a (0.110)	0.72a (0.172)	0.76B
5	NA	NA	NA	NA	NA
□	0.57C	1.23A	0.62BC	0.73B	
	<b><u>MUF microplate β-glucosidase</u></b>				
1	0.48bc (0.006)	1.28a (0.279)	0.47ab (0.064)	0.63a (0.258)	0.72A
2	0.46bc (0.038)	1.18a (0.232)	0.50ab (0.090)	0.62a (0.193)	0.69A
3	0.75a (0.115)	1.33a (0.321)	0.55a (0.085)	0.57a (0.210)	0.80A
4	0.53b (0.206)	1.39a (0.185)	0.60a (0.035)	0.56a (0.194)	0.77A
5	0.28c (0.029)	0.53b (0.066)	0.39b (0.053)	0.39a (0.142)	0.40B
□	0.50B	1.14A	0.50B	0.55B	

<sup>†</sup>Values in a column within the same assay and soil type followed by the same lower case letter are not significantly different at  $p = 0.05$ .

<sup>††</sup> Values in parentheses are standard deviations.

<sup>‡</sup>Values in the mean column within an assay followed by the same upper case letter are not significantly different at  $p = 0.05$

<sup>‡‡</sup>Values in the mean row within an enzyme method followed by the same upper case letter are not significantly different at  $p = 0.05$ .

Table 3. Pearson's correlation coefficients (r values) for linear correlation between enzyme activities and biomass of microbial groups based on ester linked fatty acids (EL FAME) (n=24).

<b>Microbial Functional Group</b>	<b>Acid Phosphomonoesterase</b>		<b>β-Glucosidase</b>	
	<b>MUF Microplate</b>	<b>PNP Bench-scale</b>	<b>MUF Microplate</b>	<b>PNP Bench-scale</b>
<b>Arbuscular mycorrhizal fungi</b>	0.64***	0.69***	0.19 <sup>NS</sup>	0.12 <sup>NS</sup>
<b>Actinobacteria</b>	0.78***	0.75***	0.76***	0.70***
<b>Gram-negative bacteria</b>	0.80***	0.81***	0.89***	0.85***
<b>Gram-positive bacteria</b>	0.74***	0.71***	0.80***	0.75***
<b>Total bacteria</b>	0.77***	0.76***	0.84***	0.79***
<b>Total fungi</b>	0.81***	0.83***	0.83***	0.76***

Table 4. Coefficients of variation (CV) at three levels of replication for  $\beta$ -glucosidase activity using the MUF microplate method.

Replication Source	Jory Managed						Jory Unmanaged						Miamian Unmanaged						Ottokee Row Crop											
	1		2		3		1		2		3		1		2		3		1		2		3							
	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2						
Field Suspension	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2						
<b>Lab</b>	<b><u>Variation of microplate analytical replications (n=4) within a soil suspension</u></b>																													
1	8	8	4	7	17	4	4	6	6	8	11	11	19	17	7	11	11	9	15	15	10	4	13	48	48	22	26	26	24	36
2	17	14	15	14	6	17	7	8	8	5	10	10	9	10	22	18	18	4	21	21	9	14	12	20	20	38	118	118	19	13
3	9	22	13	36	23	22	12	7	7	10	20	20	18	9	17	70	70	11	15	15	20	39	36	69	69	22	26	26	22	21
4	21	16	6	6	14	42	18	9	9	13	6	6	24	38	17	14	14	9	9	9	9	10	6	11	11	11	30	30	10	9
5	22	30	23	4	17	10	24	7	7	18	18	18	10	14	19	7	7	5	22	22	10	35	36	16	16	8	16	16	38	22
	<b><u>Variation of soil suspensions (n=2)</u></b>																													
1	6	15	4	2	9	9	3	19	1	13	0	0																		
2	9	2	16	4	10	7	10	14	2	3	15	9																		
3	13	40	3	2	4	28	1	49	13	32	17	21																		
4	4	11	13	14	7	1	5	9	4	5	9	7																		
5	3	23	3	13	1	26	2	12	18	1	20	20																		
	<b><u>Variation of field replications (n=3)</u></b>																													
1		1				22																								
2		8				20																								
3		15				24																								
4		39				13																								
5		10				12																								

Table 5. Coefficients of variation (CV) at two levels of replication for acid phosphatase activity using the bench method (Lab #4 did not replicate the bench method, and Lab #5 did not perform the bench-scale assay).

Lab	Jory Managed			Jory Unmanaged			Miami Unmanaged			Ottokee Row Crop		
	1	2	3	1	2	3	1	2	3	1	2	3
<b>Variation of analytical replications (n=2)</b>												
1	3	4	2	2	2	2	2	3	0	13	2	8
2	1	1	0	3	4	16	2	2	4	3	3	8
3	1	2	7	7	7	6	2	1	1	0	13	14
<b>Variation of field replications (n=3)</b>												
1		9			2			11			45	
2		9			8			8			52	
3		6			4			12			51	
4		3			4			13			37	

Table 6. Coefficients of variation (CV) at two levels of replication for  $\beta$ -glucosidase activity using the bench method (Lab #4 did not replicate the bench method, and Lab #5 did not perform the bench-scale assay).

Lab	Jory Managed			Jory Unmanaged			Miamian			Ottokee		
	1	2	3	1	2	3	1	2	3	1	2	3
<b>Variation of analytical replications (n=2)</b>												
1	3	1	3	5	0	1	7	3	5	1	7	5
2	1	0	7	1	0	7	3	1	12	1	3	0
3	0	0	2	7	2	19	9	3	10	2	12	4
<b>Variation of field replications (n=3)</b>												
1		3			20			7			34	
2		9			16			4			34	
3		5			13			60			26	
4		6			16			15			24	






Figure 1. Microplate assay layouts. The bottom plate shows the layout for samples (light circles) and controls (dark circles) for one soil and two substrates, including three field replicates, two suspension

replicates, with four microplate well replicates each (adapted from Deng et al., 2011). The top two plates consist of standards with increasing MUF concentration and autohydrolysis controls.

Table 1. Description and properties of soils

Soil	Vegetation	pH	Total C	Total N	Sand	Silt	Clay
			-----g kg <sup>-1</sup> -----			-----%-----	
Jory silty clay loam (fine, mixed, active mesic Xeric palehumult)	Christmas trees (managed)	5.2	27.9	2.2	46	38	16
Jory silty clay loam (fine, mixed, active mesic Xeric palehumult)	Douglas Fir (unmanaged)	6.0	36.8	2.7	52	34	14
Miamian silt loam (fine, mixed, active, mesic Oxyaquic Hapludalfs)	Mixed grasses (unmanaged)	5.8	27.9	2.0	38	40	22
Ottokee fine sand (mixed, mesic Aquic Udipsamments)	Row cropped (managed)	7.3	17.6	ND†	90	8	2

†Not detectable.

Table 2. Soil enzyme activities by bench or microplate methods across labs

	<b>Jory Managed</b>	<b>Jory Unmanaged</b>	<b>Miamian Unmanaged</b>	<b>Ottokee Row-Crop</b>	
----- $\mu\text{mol g}^{-1} \text{h}^{-1}$ -----					
Lab	<b><u>PNP bench-scale acid phosphomonoesterase</u></b>				$\bar{x}$
1	1.93b <sup>†</sup> (0.179) <sup>††</sup>	3.90b (0.092)	3.18bc (0.343)	0.49a (0.219)	2.38B <sup>‡</sup>
2	1.76b (0.156)	3.30c (0.277)	2.76c (0.221)	0.42a (0.218)	2.06C
3	2.31a (0.143)	4.38a (0.187)	3.95a (0.468)	0.55a (0.280)	2.80A
4	2.07ab (0.071)	4.55a (0.168)	3.87b (0.490)	0.81a (0.297)	2.83A
5	NA	NA	NA	NA	NA
$\bar{x}$	2.02C <sup>‡‡</sup>	4.03A	3.44B	0.57D	
	<b><u>MUF microplate acid phosphomonoesterase</u></b>				
1	1.40b (0.076)	2.75b (0.233)	1.97a (0.438)	0.19a (0.065)	1.58 BC
2	1.50b (0.081)	2.94b (0.191)	2.23a (0.442)	0.16a (0.071)	1.71 B
3	1.88a (0.085)	2.68b (0.225)	2.38a (0.264)	0.18a (0.044)	1.78 AB
4	1.24b (0.325)	3.85a (0.369)	2.53a (0.304)	0.14a (0.078)	1.94 A
5	1.34b (0.087)	1.99c (0.312)	2.40a (0.483)	0.14a (0.129)	1.47 C
$\bar{x}$	1.47C	2.84A	2.30B	0.16D	
	<b><u>PNP bench-scale <math>\beta</math>-glucosidase</u></b>				
1	0.54b (0.015)	1.14a (0.225)	0.55a (0.038)	0.68a (0.065)	0.73B
2	0.50b (0.045)	1.08b (0.170)	0.55a (0.021)	0.63a (0.215)	0.69B
3	0.72a (0.036)	1.60a (0.210)	0.65a (0.389)	0.89a (0.235)	0.97A
4	0.51b (0.031)	1.08b (0.175)	0.74a (0.110)	0.72a (0.172)	0.76B
5	NA	NA	NA	NA	NA
$\bar{x}$	0.57C	1.23A	0.62BC	0.73B	
	<b><u>MUF microplate <math>\beta</math>-glucosidase</u></b>				
1	0.48bc (0.006)	1.28a (0.279)	0.47ab (0.064)	0.63a (0.258)	0.72A
2	0.46bc (0.038)	1.18a (0.232)	0.50ab (0.090)	0.62a (0.193)	0.69A
3	0.75a (0.115)	1.33a (0.321)	0.55a (0.085)	0.57a (0.210)	0.80A
4	0.53b (0.206)	1.39a (0.185)	0.60a (0.035)	0.56a (0.194)	0.77A
5	0.28c (0.029)	0.53b (0.066)	0.39b (0.053)	0.39a (0.142)	0.40B
$\bar{x}$	0.50B	1.14A	0.50B	0.55B	

<sup>†</sup> Values in a column within the same assay and soil type followed by the same lower case letter are not significantly different at  $p = 0.05$ .

<sup>††</sup> Values in parentheses are standard deviations.

<sup>‡</sup> Values in the mean column within an assay followed by the same upper case letter are not significantly different at  $p = 0.05$

<sup>‡‡</sup> Values in the mean row within an enzyme method followed by the same upper case letter are not significantly different at  $p = 0.05$ .

Table 3. Sources of variation (CV) for acid phosphatase activity using the MUF microplate method

Lab	Jory Managed				Jory Unmanaged				Miamian Unmanaged				Ottokee Row Crop											
	Field replication				Field replication				Field replication				Field replication											
	Srep <sup>†</sup>		Srep		Srep		Srep		Srep		Srep		Srep		Srep									
	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2								
<b>Four microplate replications within each of two soil suspensions</b>																								
1	4	3	7	3	3	13	5	2	3	5	4	4	4	7	10	4	6	5	9	12	110	48	22	106
2	7	13	13	14	9	12	6	16	12	18	8	11	13	5	6	2	4	10	7	4	50	39	32	4
3	12	2	11	36	3	40	5	9	11	4	6	13	8	9	8	9	4	11	24	27	23	22	21	17
4	12	3	16	11	11	4	9	3	6	31	14	7	2	4	2	5	5	31	8	4	25	32	14	7
5	18	13	5	5	13	20	13	9	41	12	14	26	7	12	3	9	9	6	11	24	16	59	13	63
<b>Two soil suspensions within each of three field replications</b>																								
1	1		5		15		5		6		10		0		2		1		8		18		33	
2	6		5		8		3		11		2		9		0		6		6		14		38	
3	2		12		1		6		10		8		8		48		27		47		5		39	
4	3		27		5		0		9		2		1		0		2		16		22		16	
5	7		10		24		29		9		21		10		6		2		27		47		0	
<b>Three field replications within each soil</b>																								
1			5						9						22								34	
2			5						7						20								43	
3			5						8						11								24	
4			26						10						12								56	
5			7						16						20								90	

†Suspension replication.

Table 4. Sources of variation (CV) for  $\beta$ -glucosidase activity using the MUF microplate method.

Lab	Jory Managed						Jory Unmanaged						Miami Unmanaged						Ottokee Row Crop					
	Field replication						Field replication						Field replication						Field replication					
	SRep†		SRep		SRep		SRep		SRep		SRep		SRep		SRep		SRep		SRep		SRep		SRep	
	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2	1	2
<b>Microplate assay replications (n=4) within a soil suspension</b>																								
1	8	8	4	7	17	4	4	6	8	11	19	17	7	11	9	15	10	4	13	48	22	26	24	36
2	17	14	15	14	6	17	7	8	5	10	9	10	22	18	4	21	9	14	12	20	38	118	19	13
3	9	22	13	36	23	22	12	7	10	20	18	9	17	70	11	15	20	39	36	69	22	26	22	21
4	21	16	6	6	14	42	18	9	13	6	24	38	17	14	9	9	9	10	6	11	11	30	10	9
5	22	30	23	4	17	10	24	7	18	18	10	14	19	7	5	22	10	35	36	16	8	16	38	22
<b>Soil suspension replications (n=2)</b>																								
1	6		15		4		2		9		9		3		19		1		13		0		0	
2	9		2		16		4		10		7		10		14		2		3		15		9	
3	13		40		3		2		4		28		1		49		13		32		17		21	
4	4		11		13		14		7		1		5		9		4		5		9		7	
5	3		23		3		13		1		26		2		12		18		1		20		20	
<b>Field replications (n=3)</b>																								
1			1						22						14						41			
2			8						20						18						31			
3			15						24						15						37			
4			39						13						6						35			
5			10						12						14						37			

† Suspension replication

Table 5. Sources of variation (CV) for acid phosphatase activity using the bench method (Lab #4 did not replicate the bench method, and Lab #5 did not perform the bench-scale assay).

Lab	Jory Managed			Jory Unmanaged			Miamian			Ottokee			Mean CV
	1	2	3	1	2	3	1	2	3	1	2	3	
-----%													
<b>Assay replications (n=2)</b>													
1	3	4	2	2	2	2	2	3	0	13	2	8	3
2	1	1	0	3	4	16	2	2	4	3	3	8	4
3	1	2	7	7	7	6	2	1	1	0	13	14	5
<b>Field replications (n=3)</b>													
1		9			2			11			45		17
2		9			8			8			52		19
3		6			4			12			51		18
4		3			4			13			37		14

Table 6. Sources of variation (CV) for  $\beta$ -glucosidase activity using the bench method (Lab #4 did not replicate the bench method, #5 did not perform the bench-scale assay).

Lab	Jory Managed			Jory Unmanaged			Miamian			Ottokee			Mean CV
	1	2	3	1	2	3	1	2	3	1	2	3	
-----%													
<b>Assay replications (n=2)</b>													
1	3	1	3	5	0	1	7	3	5	1	7	5	3
2	1	0	7	1	0	7	3	1	12	1	3	0	3
3	0	0	2	7	2	19	9	3	10	2	12	4	6
<b>Field replications (n=3)</b>													
1		3			20			7			34		16
2		9			16			4			34		16
3		5			13			60			26		26
4		6			16			15			24		15

Table 7. Pearson's correlation coefficients (r values) for linear correlation between enzyme activities and biomass of microbial groups based on ester linked fatty acids (EL FAME) (n=24).

Microbial Functional Group	Acid Phosphomonoesterase		$\beta$ -Glucosidase	
	MUF Microplate	PNP Bench-scale	MUF Microplate	PNP Bench-scale
Arbuscular mycorrhizal fungi	0.64***	0.69***	0.19 <sup>NS</sup>	0.12 <sup>NS</sup>
Actinobacteria	0.78***	0.75***	0.76***	0.70***
Gram-negative bacteria	0.80***	0.81***	0.89***	0.85***
Gram-positive bacteria	0.74***	0.71***	0.80***	0.75***
Total bacteria	0.77***	0.76***	0.84***	0.79***
Total fungi	0.81***	0.83***	0.83***	0.76***

\*\*\* P <0.001

<sup>NS</sup>Not significant at P<0.05






Figure 1. Microplate assay layouts. The bottom plate shows the layout for samples (light circles) and controls (dark circles) for one soil and two substrates, including three field replicates, two suspension replicates, with four microplate well replicates each (adapted from Deng et al., 2011). The top two plates consist of standards with increasing MUF concentration and autohydrolysis controls.

# MICROPLATE FLUORIMETRIC ASSAY OF SOIL ENZYMES PROCEDURES

## INTRODUCTION

Use of the microplate format in soil enzyme assays offers the advantage of simultaneous analysis of multiple enzymes using a small quantity of soil. A microplate reader can simultaneously, measure many samples (e.g., in 96 wells) for absorbance or fluorescence in microliter volumes, which allows researchers to substantially reduce reagent costs and possibly assay time over conventional bench-scale assays.

Enzyme activity assays using the microplate fluorescence methodology are gaining greater interest, in part, owing to the high sensitivity of fluorescence detection and significantly less susceptibility to turbidity interference, compared to absorption based detection (Deng et al., 2013). A major concern for fluorescence-based methods is the significant quenching of fluorescence of compounds such as 4-methylumbelliferyl (MUF) in soil, which varies temporally and spatially (Freeman et al., 1995). Fluorescence is also affected by pH and temperature (Lakowicz, 1983). The relative fluorescence signals of MUF are highest between pH 10 and 11, which could be >35,000-fold of those detected at pH 6.0 (Deng et al., 2013). As temperature increases, fluorescence decreases due to an increase in molecular motion that results in more frequent molecular collisions and subsequent loss of energy (Guilbault, 1990). Consequently, quantification of MUF in soil requires a calibration curve for each soil. For quantitative detection, it is also important to treat standards, blanks, and samples in exactly the same manner, prepare all solutions using the same reagents and preparation techniques, and measure at the same temperature after the same amount of time. The additional precautions and calibration steps add considerable labor and expense to the assay as well as additional sources of error.

Using the assay protocols described in Deng et al. (2011) that evolved from earlier studies (Drouillon and Merckx, 2005; Marx et al., 2001), several recent studies have shown comparable results between MUF-microplate and  $\rho$ -nitrophenol-bench scale assay approaches (Deng et al., 2013; Dick et al., 2013). Although uniformity of diluted soil suspensions is a potential concern for the MUF microplate method, Dick et al. (2013) found that taking samples from the top, middle, or bottom of the suspension had no significant effect ( $p = 0.49$ ) on variability or detecting treatment effects on enzyme activity. Furthermore, sonication of the suspension prior to sampling did not improve reproducibility of the MUF microplate method.

In recent years there have been a growing number of published papers attempting to mimic in situ pH or temperature. This may be appropriate depending on the goals of the research but such results need to be interpreted with caution. First because these less than optimal conditions can reduce the activity; this could obscure treatment effects or even change the ranking of treatment effects on soil enzyme activities. Secondly, other aspects of the assay such as saturating the enzyme with substrate (orders of magnitude, use of co-factors, and that soil is in slurry solution – all makes for very different conditions compared to in situ condition. The method described here is based on optimal conditions.

## PRINCIPLES

The assay described below is based on detection of MUF released by enzymatic hydrolysis of specific substrates when incubated with soil at the optimal pH of the assayed

enzyme. Following incubation for a defined time at the desired temperature, MUF is quantified upon addition of THAM (0.1 M, pH 10) as reported by Deng et al. (2013). Concentrations of MUF are calculated using a calibration curve and enzyme activities are expressed as mmole or  $\mu\text{mole}$  MUF released  $\text{kg}^{-1}$  soil  $\text{h}^{-1}$ .

## ENZYME ASSAYS PROTOCOL

### Apparatus

- 1-L beaker
- 1-L volumetric flasks
- 100-mL volumetric flasks
- 150-mL beakers
- Stir plate capable of regulating to 600 rpm
- 3.75-cm magnetic stir bar
- Multichannel pipette (0-250  $\mu\text{L}$ )
- Black solid polystyrene microplates, 96-well (CLS3915; Costar microplates; Corning, Inc., Lowell, MA).
- Incubator (37°C)
- Fluorescence microplate reader (360-nm excitation; 460-nm emission)

### Reagents (Table 1)

- Modified universal buffer (MUB) stock solution: Dissolve 12.1 g tris (hydroxymethyl) aminomethane (THAM), 11.6 g maleic acid, 14.0 g citric acid, and 6.3 g boric acid in 800 mL 0.5 M NaOH. Adjust to 1 L with 0.5 M NaOH and store at 4° C.
- MUB working solution (pH 6.0): Place 200 mL MUB stock solution in 1-L beaker containing magnetic stir bar. Place on stir plate and titrate the pH with HCl while stirring. Adjust the volume to 1 L with DI water.
- Methylumbelliferyl substrates (2 mM): Weigh 0.068 g methylumbelliferyl- $\beta$ -D-glucoside (MUF-G; Sigma Aldrich M3633) or 0.052 g methylumbelliferyl-phosphate (MUF-P; Sigma Aldrich M8883) into a 100-mL volumetric flask and adjust the volume to 100 mL with DI water. Prepare the solutions daily or aliquot and store for no more than two weeks at -20°C. Thaw one aliquot on the day of use and discard leftovers.
- THAM (0.1 M, pH 10): Dissolve 12.1 g Tris (hydroxymethyl) aminomethane (THAM) (MW 121.14) in 700 mL DI water. Adjust pH to 10.0 with 0.1 M NaOH, and then adjust volume to 1 L with DI water. Store in plastic container at room temperature.
- Methylumbelliferone (MUF) stock solution (100  $\mu\text{M}$ ): Dissolve 0.0202 g 4-methylumbelliferone sodium salt (98%; M1508; Sigma Aldrich) in 700 mL DI water, then adjust volume to 1 L. Store in the dark at 4° C for no more than 2 weeks or in aliquots at -20°C for no more than one month for this study. Thaw one aliquot on the day of use and discard leftovers.
- MUF working standards: 0, 5, 10, 20, 30, and 50  $\mu\text{M}$  MUF standards are prepared by diluting 0, 5, 10, 20, 30, or 50 mL of the MUF stock solution (100  $\mu\text{M}$ ) to 100 mL with DI water in 100 mL volumetric flasks. Store in the dark at 4° C for no more than 2 weeks.

**Procedure** (See Figure 1 for plate layout)

1. Warm reagents in the incubator at assay temperature for ~30 min prior to initiating the assay. Normally 37°C is used unless otherwise recommended in the original method development paper for a given assay.
2. Two replicate soil suspensions are prepared for each soil sample by weighing 1 g of soil into a 150 mL beaker, and adding 120 mL of DI water. The soil is homogenized for 30 min. using a 3.75 cm stir bar on a stir plate at 600 rpm.
3. A multichannel pipette with four tips is used, with continuous stirring, to load 100 µL of suspension into each microplate well containing 50 µL MUB pH 6.0. The same stir plate and speed should be used to carry out this step.
4. Add 50 µL of MUF substrate to each well, mix by pipetting up and down several times, and incubate at 37°C for 1 h.
  - a. The plates should be incubated in a shallow water bath in a pan that can be covered during incubation (can use aluminum foil). Make sure the water has reached 37°C prior to placing sample plates for incubation.
  - b. Substrate should be added to a plate as rapidly as possible but still with care taken in pipetting.
  - c. Make sure each plate is incubated for exactly 1 h.
5. Following incubation, add 50 µL of 0.1 M THAM (pH 10) to each well (in the same order as the substrate solution was added to keep the incubation time relatively consistent among samples) to terminate the reaction and increase fluorescence signal for its detection at 360 nm excitation and 460 nm emission. The relative fluorescence signal is stable under the stated conditions for several hours, with no detectable differences observed when readings were taken within three hour following the addition of THAM (Deng et al. 2013).
6. Four controls are performed in the same way except substrate is added after incubation. Controls are soil-specific.
7. Controls for autohydrolysis are prepared as for sample except DI water replaces soil suspension and substrate is added either before incubation or after the reaction is terminated by alkalization. These control will only need to be done once for each substrate.
8. Standards for soil-specific calibration curves are developed at the same time samples are assayed using the same soil suspension and procedure, except MUF standards are used in the place of MUF substrates. Briefly, 50 µL of each MUF working standard solution are placed into microplate wells that each contains 50 µL of MUB, followed by the addition of 100 µL of soil suspension. After incubation, 50 µL of THAM (pH 10) is added to each well as done for enzyme assays described above. The total volume in each well is 250 µL and contains MUF standards of 0, 250, 500, 1000, 1500, or 2500 pmol. Average data obtained from both soil suspensions are used to develop a calibration curve for the tested soil.
9. Set microplate reader sensitivity for sample plate according to the highest standard on the standard plate (Table 2).
10. Calculations (see Deng et al. 2011 for more details):
  - The average reading from the zero MUF standard should be subtracted from all other standard readings
  - The intercept of the calibration curve should be forced through zero.

- The average autohydrolysis is calculated by subtracting the average RFU of the autohydrolysis wells incubated with substrate added after reaction termination/alkalization from the average of the wells with substrate added before incubation.

$$\text{Corrected fluorescence } (F_{\text{corrected}}) = (F_{\text{sample}} - F_{\text{avg control}} - F_{\text{avg autohydrolysis}})$$

In each sample assay well,

$$\text{pmol MUF released} = \frac{F_{\text{corrected}}}{\text{Slope of the MUF calibration curve}}$$

Enzyme activity in soil,

$$\text{nmole MUF g}^{-1} \text{ soil h}^{-1} (\equiv \mu\text{mole MUF kg}^{-1} \text{ soil h}^{-1})$$

$$= \frac{\text{pmol MUF released}}{100 \mu\text{l soil suspension}} \times \frac{1000 \mu\text{l}}{1 \text{ mL}} \times \frac{120 \text{ mL}}{1 \text{ g soil}} \times \frac{1 \text{ nmol}}{1000 \text{ pmol}} \times \frac{1}{1 \text{ h}}$$

### COMMENTS

Pipetting is the most critical procedure for causing error in the MUF microplate method because of the small volumes that are used. In particular, retrieving sample volumes from the soil suspension is a critical pipetting step. To obtain accurate and reproducible data, it is important to check the pipette tips to ensure that the tips are tight and the multichannel pipette is calibrated and working properly. Also, caution should be exercised during pipetting to avoid trapping of air bubbles.

A fundamentally intractable issue with microplate assays is the variation in fluorescence over time of MUF standard's. Our experience has been that this is particularly acute with long term storage and reuse of MUF stock where the fluorescence is not stable over time.

We emphasize that the optimum pH be used and the optimal value chosen be based on pH curves found in the literature for a given enzyme. The most comprehensive compilation of soil enzyme assays is in Dick (2011) where optimal pHs are reported for methods that have been vetted. If this has not been done for an enzyme of interest, preliminary pH curve investigation should be done to determine the pH optima.

The above protocol is standardizing the operationally defined protocols (37 C incubation temperature, pH 10 THAM buffer to kill reaction, 1 hour incubation, controlling for quenching in the standard curve and autohydrolysis, fluorescence reading within 1 hour of stopping the enzyme reaction, pre-warming incubation reagents, and method of developing soil suspension) to enable cross paper and meta analyses. Since pH optima currently is based on PNP bench scale methods which did not use fluorescing substrates, further studies may show a different pH optima with MUF microplate for a given enzyme.

If an alternative methodology is used to attempt to mimic some aspect of in situ conditions, data should also be generated using the standardized approach to allow cross paper comparison and enable firm conclusions and interpretation of the data.

## REFERENCES

- Deng S., Kang H., Freeman, C., 2011. Microplate fluorimetric assay of soil enzymes. In: Dick, R.P. (Ed.), *Methods of Soil Enzymology*, Vol 9. Soil Science Society of America, Madison, WI, pp. 311-318.
- Deng, S., Popova, I.E., Dick, L., Dick, R., 2013. Bench scale and microplate format assay of soil enzyme activities using spectroscopic and fluorometric approaches. *Applied Soil Ecology* 64, 84-90.
- Dick, R.P. (ed). 2011. *Methods of Soil Enzymology*. Soil Science Society of America, Madison Wisconsin.
- Dick, L.K., Jia, G., Deng, S., Dick, R.P., 2013. Evaluation of microplate and bench-scale  $\beta$ -glucosidase assays for reproducibility, comparability, kinetics, and homogenization methods in two soils. *Biology and Fertility of Soils* 49, 1227-1236.
- Freeman, C., Liska, G.P, Ostle, N.H., Jones, S.E., Lock, M.A., 1995. The use of fluorogenic substrates for measuring enzyme activity in peatlands. *Plant Soil* 175, 147-152.
- Guilbault, G.G., 1990. *Practical Fluorescence*, Second Edition, Marcel Dekker, Inc., New York.
- Lakowicz, J.R., 1983. *Principles of Fluorescence Spectroscopy*. Plenum Press, New York.




Figure 1. Plate layouts for assaying activities of two enzymes in four soils. The plates above represent requirements for one soil and two substrates, including three field replications, two soil suspension replications and four analytical replications. The top two plates consist of standards with increasing MUF concentration and autohydrolysis controls. The bottom plates show layout for samples (light circles) and controls (dark circles).

**Table 1. Suggested source of substrates**

Sigma Catalog #	Substrate/Standard
M3633	4-Methylumbelliferyl $\beta$ -D-glucopyranoside
M8883	4-Methylumbelliferyl phosphate
M1508	4-Methylumbelliferone sodium salt
N7006	4-Nitrophenyl $\beta$ -D-glucopyranoside
P4744	Phosphatase substrate
1048	4-Nitrophenol
T6066	Tris base (THAM: MW 121.14)

**Table 2. Microplate reader and gain settings**

Microplate Reader (brand/model)	Does Reader have Auto-Sensitivity Setting?	Determination of Gain Setting (sensitivity)
BioTek Synergy H1	Yes	Set auto-sensitivity to highest well of calibration plate, then apply same value to sample plates