

Are Rises in Electro-Magnetic Field in the Human Environment, interacting with Multiple Environmental Pollutions, the Tripping Point for Increases in Neurological Deaths in the Western World?

#Professor Colin Pritchard, Dr. Anne Silk, Faculty of Health & Social Sciences, Bournemouth University and Dr. Lars Hansen, Dept Psychiatry, Hon Senior Lecturer, University of Southampton.

Corresponding author. Professor Colin Pritchard, Research Professor, Faculty of Health & Social Sciences, Bournemouth University, Royal London House, Christchurch Rd; Bournemouth BH13LT UK. cpritchard@bournemouth.ac.uk 07825248004.

Abstract

Whilst humans evolved in the earth's Electro-Magnetic-Field (EMF) and sun-light, both being essential to life but too much sun and we burn. What happens if background EMF rise to critical levels, coinciding with increasing environmental pollutants?

Two of the authors can look back over 50 clinical years and appreciate the profound changes in human morbidity across a range of disparate conditions - autoimmune diseases, asthma, earlier cancer incidence and reduced male sperm counts. In particular have been increased autism, dyslexia, Attention Deficit Hyperactivity Disorder and neurological diseases, such as Amyotrophic Lateral Sclerosis, Multiple Sclerosis, Parkinson's Disease, Early Onset Dementia, Multiple System Atrophy and Progressive Supranuclear Palsy. What might have caused these change- whilst genetic factors are taken as given, multiple environmental pollutants are associated with neurological disease although the mechanisms are unclear. The pace of increased neurological deaths far exceeds any Gompertzian explanation that because people are living longer they are more likely to develop more age-related problems such as neurological disease.

Using WHO global mortality categories of Neurological Disease Deaths (NDD) and

Alzheimer's and Dementia deaths (Alz), updated June 2018, together they constitute Total Neurological Mortality (TNM), to calculate mortality rates per million for people aged 55-74 and for the over-75's in twenty-one Western countries.

Recent increases in American people aged over-75's rose 49% from 1989 to 2015 but US neurological deaths increased five-fold. In 1989 based on Age-Standardised-Deaths-Rates America USA was 17th at 324pm but rising to 539pm became second highest.

Different environmental / occupational factors have been found to be associated with neuro-degenerative diseases, including background EMF. We briefly explore how levels of EMF interact upon the human body, which can be described as a natural antennae and provide new evidence that builds upon earlier research to propose the following hypothesis.

Based upon recent and new evidence we hypothesise that a major contribution for the relative sudden upsurge in neurological morbidity in the Western world (1989-2015), is because of increased background EMF that has become the tipping point-impacting upon any genetic predisposition, increasing multiple-interactive pollutants, such as rises in petro-chemicals, hormone disrupting chemicals, industrial, agricultural and domestic chemicals. The unprecedented neurological death rates, all within just twenty-five years, demand a re-examination of long-term EMF safety related to the increasing background EMF on human health. We do not wish to 'stop the modern world', only make it safer.

Key Words: neurological aetiology Electro-Magnetic-Fields environmental epigenetic

Acknowledgements: We confirm that we have no vested or conflict of interest in this study. Nor was there any external funding for the project.

Text (4,630).

Introduction:

The Hypothesis.

It is an evolutionary fact that humans evolved within the Earth's constant Electro-Magnetic-Field (EMF) and the sun's radiation which are essential to life. If there is too much sun, we burn but what if there is too much EMF?

We hypothesise that over the last 25 years increases in background Electro-Magnetic-Fields within the human environment, impacting upon the extant and rising multiple interactive-environmental pollutants is the Tipping Point for the accelerating increases in neurological deaths in the Western world ¹⁻¹³. For example, impacting on rising problems of air pollution, on what has been described as the neuro-inflammation hypothesis ^{14,15}, whilst increased background of EMF are increasingly linked to both neurodegenerative disease and leukaemia, both associated with oxidative stress ¹⁶.

To make the case we need to look back at changing environmental pollutants and think across disciplines to consider apparently unconnected changes in human morbidity. Two of the authors have the advantage of being able to look back over 50 clinical years and can appreciate the profound changes that have occurred. What is often difficult to grasp is how recent are changes in the world environment. There has been an explosive development in a wide range of chemicals that have been employed by human in the relentless pursuit of mastering our environment.

The digital world that now surrounds us with new levels of Electro-Magnetic Fields [EMFs] and it is the magnetism that might be the problem. Many environmentally linked diseases often need 20 or more years to develop ^{17,18}, so as the digital world is still a relatively a new phenomenon, the possible twenty years needed for pathological effects of raised EMF is perhaps beginning to emerge?

The first major wake-up call of the impact of environmental pollution upon human morbidity was Rachel Carson's 1968 'Silent Spring'. She highlighted the damage to wild-life by environmental degradation and alerted us to the fact that human beings are not exempt or immune to environmental pollutants. The slow response to emerging scientific alarm bells was seen in the reluctant acceptance of the problem

with asbestos, which has led to today a growing epidemic of people entering their sixties with asbestos related diseases and with the previously assumed neutral organophosphates¹⁷⁻¹⁹. The recent Volkswagen scandal who hid the extent of diesel particulates shows we cannot always trust even the most prestigious companies to acknowledge inadvertent negative impacts on human health of economic activity. We will ignore notable changes in human morbidity such as rises in autoimmune disease, allergies, falling sperm counts but focus upon conditions which can be said to be broadly neurologically related.

Attention Deficit Hyperactivity Disorder (ADHD) which was virtually unheard of until the 1980's. It might well be an early soft-neurological sign but reflects undoubted change. It was described as 'minimal cerebral dysfunction' and from 1963 onwards it took 20 years to have a 100 papers on the medical data base 'medline'. Re-defined as ADHD in 1987, from January to September 2018 there were 700 papers, from across the world.

Autism, rare before the 1970's, initially being described as a primarily psychological disorder in children of 'refrigerator parents'. Now widely recognised and every Western country has an Autistic Society, with an estimated 700,000 British people on the autistic spectrum with a projected 20million people world- wide²⁰. Whilst this may be to do with a greater willing to make the diagnosis, does this really explain the enormous rise in numbers?

Another 'ambiguous' condition is Chronic Fatigue Syndrome (CFS) or Myalgic Encephalomyelitis (ME) and is still debatable whether it is a 'functional' rather than neurological disorder. Two other new but controversial diagnostic categories are Electromagnetic Hypersensitivity (EHS) and Multiple Chemical Sensitivity (MCS). The first paper on medline for EHS was in 2005 but in the last three years there were 223. In regard to MCS the first paper was in 1975 and only a second study by 1988 but in 2018 there were eight such papers.

If ADHD, Autism-Asperger spectrum, Dyslexia and ME / CFS, all initially ascribed to children but now diagnosed in adults, are thought of as basically neurological disorders this raises the question are there environmental influences related to these increased 'neurological' conditions. Plus rises in the established conditions such as Motor Neurone and Parkinson's Diseases etc ¹⁻¹⁶?

It is hypothesised that the impact of Electro-Magnetic-Fields, *interacting and impacting on top of genetic predisposition and extant and growing multiple environmental changes*, is the tipping point that triggers these unprecedented accelerating rises.

However, it has to be acknowledged that there are some inconsistent findings regarding EMF, as some have found no EMF link with neuro-pathology, or that EMFG has been therapeutic ²¹⁻²⁷. Alternative explanations, such as the Gompertzian hypothesis ^{28, 29} and improved diagnosis ³⁰ are sometimes used to suggest the rises in neurological morbidity are essentially an artefact.

The major artefact argument is the Gompertzian hypothesis that explains the raised incidence is because people are now living longer to develop aged-related disease, so it is argued that the increases mainly reflect improved longevity ^{28, 29}. This hypothesis will be evidentially refuted in the course of this study.

Clinical Studies of Incidence of Neurological Disorders: We briefly explore human clinical level studies reporting rises in neurological diseases, especially Parkinson's Disease and Amyotrophic Lateral Sclerosis (ALS), designated Motor Neurone Disease in Europe. We have decided to exclude all animal studies, despite many results indicating possible aetiological EMF connections but there is always the question of transferability to humans. As will be shown, there are sufficient human clinical studies to support the EMF tipping point hypothesis.

A series of clinical type studies highlighted rises in a range of neurological disease such as ALS, could be described as epidemic ¹⁻¹⁶, mainly in Western countries but reports from other continents are increasing ⁷⁻¹⁰ and rises in Parkinson's Disease

adds to the growing burden upon services and families ^{4, 13}. Another notable feature over the past decade is the rise in what might be considered the rarer neurological conditions such as Multiple System Atrophy and Progressive Supranuclear Palsy ³¹⁻³⁵.

Nonetheless, it cannot be denied that having proportionally more elderly people might well be a minor factor, but when the extent of the changes, based on the latest WHO data, updated June 2018, this explanation fails to account for the increases ³⁶. A major challenge to the Gompertzian explanation is starkly challenged by evidence of the explosive rise in early onset dementias reported in the Western world and other continents ³⁷⁻⁴⁵. Furthermore, in practice terms the Gompertzian explanation is at practical odds with important but different type of evidence found in the need for the British Parkinson's Society to establish a Young Persons section for their services, with people under-40's and a new British charity, Young Dementia UK that has many clients under 50 years old! It is when these conditions are juxtaposed together then the possible connections between the psychical and socio-psychological environments can be seen.

New Comparative International Study: Before considering the inter-related multiple causes of neurological conditions, and, addressing the central tenant that EMF might be the tipping point for the rises in neurological morbidity, we provide the very latest population-based analysis of neurological deaths. This builds upon our earlier epidemiological studies ¹¹⁻¹³ but extrapolates further from the recently updated WHO data ³⁶.

Table [1] shows the two WHO global neurological categories Nervous Disease Deaths (NDD) and Alzheimer & other Dementia (Alz) deaths in Age-Standardised-Death-Rates (ASDR) – in effect controlled for total population between baseline years 1989-912 to 2013-15, which combined gives a Total Neurological Mortality (TNM) rate per million (pm) of population ³⁶.

The highest TNM was Finland at 911pm, the USA 539pm, the Netherlands and the UK 424pm. Over the period 1989 to 2015 this represents rises of 161%, 196%, 128% and 67% respectively. The lowest countries were Japan at 103pm, Greece 104pm, Austria 181pm and Portugal 229pm yet these rates are also equivalent to rises of 56%, 44%, 46% and 129% respectively, with an overall average rise for the 21 countries of 90% over the period, all controlled for population.

A perusal of the table shows that in the 1989-91 period seventeen countries had higher NDD rates than Alz but by 2013-15 only eight nations' NDD rates were higher than Alz although *both* death rates increased over the period.

A further challenge to the Gompertzian explanation²⁸⁻²⁹, is shown in Table [2], which compares the increases in population of men and women aged 75+ juxtaposed against rises in the Elderly (75+) TNM rates from which to calculate a population and Total Neurological (TNM) time ratio.

The highest rate was Finland at 7,204pm but by 2013-15 had risen to 31,246pm, which is 4.33 ratio of change, compared to only 66% rise in the 75+ population, yielding an odds ratio of 1:2.61. Followed by the USA at 18056pm, with a five-fold 75+ TNM increase compared to a 49% rise in population yields a ratio of change of 1:3.59.

Other notable population to TNM ratios were Canada 1:4.89, Denmark 1:4.21, Sweden 1:3.40, Portugal 1: 3.25, Norway 1:2.97, Netherlands 1:2.77 and the UK 1:2.49 and six other countries had ratios greater than 1:1.50, only Belgium at 1:0.88 had greater population rises than TNM, and though Greece and Japan more than doubled their TNM over the period they had the next smallest population to TNM ratios, 1:1.10 and 1:1.33 respectively.

Insert Table [2]

Though the Gompertzian explanation of these increases might have made some contribution but not at this rate or in such a relatively short time. Moreover, in 1989-

91 the USA was 14th highest of the 21 countries but by 2013-15 were second highest in the Western world!

An indication of the speed of change in neurological deaths in the 55-74 year age band, which is *below the life-expectancy* of all the countries is given in Table [3], ranked by biggest increase in just the last 10 years.

Thirteen countries had rises of 20% and more over the decade, the biggest rises were in Greece 68%, Japan 50%, albeit starting from a low base and the Netherlands at 56%, with ten countries having rises of >30%, including the UK at 45% and the USA at 32%, though there were slight falls in Belgium and France. In the first international comparison of changing patterns of neurological deaths which focused upon the 55-74 age band ¹¹, when compared to later studies ^{12,13}, it is clear these rates are increasing and accelerating as well as involving more countries as the years have unfolded.

Insert Table [3]

To provide a policy and practice perspective, as rates are statistics but numbers are actual people. The numbers of people in Britain and America dying from a neurological disorder in the age-band 55-74 will illustrate the speed of change.

Between 1989 until 2015 the numbers of neurological deaths of people aged 55-74 in Britain went from 3,518 to 5,177 an effective 47% increase over the period. The *numbers* of Total Neurological Mortality in the UK rose from 27,419 to 90,535, equivalent to a 330% rise.

In the USA, the 55-74 age band numbers grew from 16,608 to 76,244, more than a four-fold increase. As to total numbers America went from 68,071 to 409,217, a six-fold increase. It is questionable whether neurological services have increased in anything like these proportions.

What there can be little doubt about is that there an epidemic rise in neurological deaths, outstripping any demographic or Gompertzian explanation, or due to neurological disease diagnosis becoming more 'fashionable' but surely not at the extent of these accelerating changes.

Therefore our hypothesis asks might the changed pattern of EMF in the human environment be the trigger / tipping point impacting upon of any genetic predisposition and multiple inter-acting environmental influences?

Some Aetiological Associations: This raises the question of what might be the cause of such changes? Evidence comes from studies that point towards multiple-interactive- environmental factors. A number of occupational groups have been associated with raised ALS ⁴⁶⁻⁵². With similar associations in regard to occupations and exposure to a range of chemicals and solvents in regard to Parkinson's Disease ⁵³⁻⁵⁶. Also there have been indications of background Electro-Magnetic-Fields been associated with various neurological conditions ^{14,54-59}. Although, as mentioned previously, some studies have shown that short term exposure to low level EMF can help improve memory in Alzheimer, and can assist healing of wounds, fractures and improve stroke patients ²¹⁻²⁷.

Epigenetics: As with many pathologies there are degrees of under-lying genetic predisposition/ vulnerability awaiting the environmental trigger, seen in the epigenetics of ALS with its familial and sporadic forms, interacting with genetic predisposition influenced by environment ⁶⁰⁻⁶⁶.

Some research is quite specific about broad environmental triggers interacting with a person's genetic background, with such examples related to solvents, Electro-Magnetic-Field (EMF) and a range of 'natural products' ^{18, 48, 52,66-69}. Bearing in mind the increase in early onset dementias, at unprecedented levels, the epigenetic influence is confirmed by a series of studies that show a degree of familial inheritance ^{38, 40,42, 68,69}. Finland has long had the highest rate of neurological pathology. With some studies showing a degree of familial origins, however, the

extent of the recent changes supports the finding that increased rates of neurological morbidity are more sporadic than familial in origin ^{4, 5, 38, 40-43, 54, 60-69}. Moreover, it should be remembered that the USA neurological death rate rose from being 14th highest to become second highest in the Western world.

A key paper comes from Finland that explored early onset dementia and interactive epigenetic factors, whose rate of increase far exceeds any possible inherent genetic or Gompertzian influences but indicates that the logical explanation of the environment in which rises in background EMF appears to have played a significant and perhaps tipping point to fuel the neurological epidemic ^{42,43, 70}.

EMF: Impacting Upon Human Neurology: The key question is how does EMF affect the brain and human neurology? After all the brain is essentially an electro-biochemical organ and electrical impulses operate the nervous system, so in one sense, the EMFs are unlikely to be 'neutral' on human neurology over time ¹⁴.

Moreover, human's evolved in the earth's magnetic field, whose flow can be seen by any compass, whilst sunlight is essential for life, too much damages or kills.

Whereas the increased background Electro-Magnetic-Field on people over the past two decades, with the remarkable increases in digital devices, raises the question are we beginning to see the negative health impact, especially over the last decade, similar to the delayed development of environmental based disorder with a twenty year lead-in, such as asbestosis and impact on the previously assumed neutral organophosphates ^{17- 19}.

The EMF comes from a range of social as well as domestic objects such as radar power with pulsed MHz and GHz signals, some for microseconds can be very high, whilst at a low domestic level wiring creates a EMF with harmonic spikes, which by taking a compass close to electric sockets produces observable but noticed magnetic field. Some authorities showed that an accumulating dose of GSM 900-MHz RE-EMF might have devastating effects on neural stem cells proliferation and neurogenesis, leading to differentiation and apoptosis ^{71- 90}.

From a recent study in Military Medical Research , the strength of the various EMF objects, with which most people come into frequent contact are as follows ⁸².

Domestic: - FM Radio Antennas 80-800 MHz.

Mobile Phones 453-1980MHz.

Microwave ovens 915-2450MHz.

Cordless phones 1800-1900MHz.

Bluetooth devices 2450MHz.

Previously hairdryers produced seventeen time the ambient EMF but modern hairdryers have now been adapted to emit around 50-60Mhz.

Locality:- Mobile Phone base station 463-2170MHz

Surgical equipment 2450MHz.

Wireless Local Radio 2450-5000MHz.

Radar 30-30,000MHz (300 GHz).

Twenty-five years ago apart from FM radio antennas and domestic TV's and radio the sources of EMFs were not yet in general use! With such a major increase in background EMF can we really be sure of its neutrality? When illustrating changes to students and asking males to point to where they keep their mobile phones, mainly in their trouser pockets, means their gonads are being bathed in between 450-1900 Mhz, a totally new phenomena on human evolution but space precludes the possibility related to falls in male sperm counts ²². Whilst, an early study showed that even at a low frequency such as 50-Hz this could influence sleep patterns ⁷¹, but crucially the EMF interacts at cell level and at chromosomal level ⁷²⁻⁸¹.

Burke described in detail the magnetic phenomena on human neurology cellular and EMF interaction with the body ⁸¹. Whilst, we have generally eschewed studies of the impact of EMF on animals, we need to acknowledge that even at very low frequencies on simple cell development, such as sea urchins, the impact make a dramatic effect on new cell development ⁸².

There are very large numbers and types of charged metallic particles in our bodies and brains. The magnetic fields and unpaired electrons that result from charged particle movements are part of the chemical bonds holding all matter together. These traverse a variety of paths and some will spin. These two fundamentally different types of particle movements are *diamagnetic*, those with axes orientation patterns are called *paramagnetic*, *ferromagnetic*, *antiferromagnetic*, *ferromagnetic*, *antiferromagnetic* or *helical*, depending on which pattern of spin-axis orientation is stable under given environmental and chemical circumstances.

Spinning electrons have much larger Magnetic Moments than do spinning nuclei. This is because the small mass of an electron makes it spin much faster than any nuclei. Thus high levels of Reactive Oxygen Species can be generated in the body. Hence the increased EMF in the home such as radios, electric plugs, microwave and conductive ovens, TV set computers, as EMF, especially the magnetic component, interacts with the intricate electrical activities of the central nervous system as synapses which are the core structure that involves the electro-chemical signals that direct the body⁸¹. Chronic or over exposure to EMF has been found to alter chromosomes and DNA and gene expression^{73, 74, 78, 82}. It can also disturb the mitochondrial ADP phosphorylation process - crucially stem cell functioning, and can create apoptosis of neural cells^{68, 72-76}. The EMF can also influence melatonin^{71, 79} and in particular create oxidative stress⁸³⁻⁸⁶ which is linked to shrinkage of dendritic spines in the hippocampus⁸⁷⁻⁸⁸. All these features appear to be precursors of neurological pathology.

Whilst other environmental influences might well be interacting with the EMF, such as nanoparticles from Petro-chemical pollutants, that can also result in increased in oxidative stress, which is associated with the development of neuro-degenerative disease^{34, 51-59, 72-93}

Hence these relatively new EMF effects extenuate the extant multiple environmental pollutants, albeit at low, and individually, at officially 'safe' levels!

Whilst not forgetting that EMF helps in some conditions low levels of EMF, especially over time, exposure to EMF have also been found to be linked with memory problems ^{57,91-96}. This apparent inconsistency is highlighted as traumatic EMF, in the sense that electric shocks and head injury have also been associated with neurodegenerative diseases ⁹⁷⁻¹⁰⁰. On balance it would appear the pathology occurs with high intensity but short duration and low intensity but long and frequent intensity ¹⁰⁰⁻¹⁰⁴ (the Bunsen-Roscoe effect), which might account for the apparent inconsistency of EMF being therapeutic ²⁴⁻²⁷, not unlike radiotherapy for cancer ⁸⁵⁻¹⁰⁵. A key paper on how high gradient EMF affects cell life found that even relatively small EMF can crucially change the cell membrane and disturb its biological function leading to cell death ¹⁰⁶. One study looked at the preventive role of resveratrol in reducing oxidative stress in a four-year controlled study of long-term occupational exposure to EMF, to find after four years the index group had substantially lower cellular damage than controls ¹⁰⁵. As there are similar new studies from China and Russia does this imply that the safety of workers in the EMF industries is a matter of growing interest?

Electro-MAGNETIC-Fields Tipping Point: The studies that initially led to our hypothesis was by Hallberg who analysed Swedish national mortality data on Alzheimer's from 1948 until 2014 ⁵⁹. The graph on increased Alzheimer's from 1948 to 1988 rose at an angle of 5 degrees but from 1990 to 2014 the curve rose to a 75 degree angle. The researchers linked these changes to increases in background EMF. The second was a small cluster study examining Motor Neurone Disease in a Southern English village to find an eight times the county rate of MND and four times the UK national rate ⁹⁹. The methodological problems of cluster studies were readily acknowledged but what was striking was the local geography of the village which was aligned with a busy airport, compounded all the environmental health problems (chemical, solvent, EMF) associated with airports ^{32,46,48,51,53,55,85, 91,108}. On either side of the runway were two large downs. On the West was the airport's radar equipment

and on the East were the county commercial mobile masts. The local geography meant that the flight take-off was low for five miles to the village which had an excess of MND patients ¹⁰³.

There are a range of studies identifying neurological pathology to links with solvents ⁶⁷, chemicals ²⁻⁴; electronics and IT ^{73-85,106} and occupations that include the former ^{46-52,72, 97,102}. Thus this airport, with the relative high density of Petro-chemical, nanoparticles and electro-radiation, brought together in one site key interactive environmental pollutants ¹⁰³. Whilst other studies linking neurological disorders to airport environments and risks to airline pilot's health ^{32, 106,107}.

Taken together it can be seen that there are strong indications of the increased EMF impacting upon human health, both neurological, as well as with some cancers ^{1,10, 72, 105, 111, 112}.

However the aetiological issues of neurological disease is complex as quite apart from the genetic-environmental, some studies have found a viral link with neurological disease ¹¹³, another with diabetes ¹¹⁴, alcohol being associated with Alzheimer's ¹¹⁵ and the notable rises in allergies over the past two decades ¹¹⁶, whilst an earlier study explored EMF on male reproductive patterns which are known to be falling in many Western developed countries ²².

One very important feature is the role of any under-lying genetic predisposition, which might explain why oxidative stress is associated both with the cancers neurological diseases. This complicates pinning down the precise interactive mechanisms of people living in the same environmental situation but develop different conditions or no obvious pathologies, possibly depending on genetic predispositions ^{1-3, 37, 46, 48, 49, 52, 54, 57, 60, 64, 65, 72, 91, 103, 112}.

It would be conclusive if we could accurately measure EMF changes in the human environment but all we can do is remind readers of the massive intrusion into our world of EMF devices which really only began in the 1990's.

Discussion of the Cautionary Principle:

There is little doubt that there has been a marked increase in episodic neurological disease which seems to be a response changing interactive multiple environmental pollution, impacting upon under-lying genetic factors. The question is has the case been made that the EMF might be a key tipping point for this hitherto relatively ignored neurological epidemic?

The answer paradoxically, might be found in how the various national and official authorities have sought greater awareness of the dangers of EMF that are so helpful and convenient to our lives, ranging from mobile phones, PC's to induction cookers⁸¹. Early official regulations on the safety of EMF in the USA in 2002 were mildly speculative and cautionary¹¹⁷. Later in 2006 the Committee on 'Man and Radiation' was acknowledging that there could be possible health problems¹¹⁸ whilst in 2016 the EU called for greater surveillance and investigation of related health problems of EMF, crucially it acknowledged that that there were emerging health problems¹¹⁹. Indeed urging an cautionary approach from early to late life and the need for importance of radiation studies of workers^{105, 107, 120-123}.

From Rachel Carson in 1968 alerting us to the simple fact that humans are affected by the environment in which they live which is the very driving force of evolution. Recent studies related to ionising radiation related to workers in the nuclear industry whilst focusing upon cancers reasserted the precautionary principle found in the use of the linear-no-threshold (LNT) dose-response model. This points to the risk being related to received doses but acknowledge there are no threshold doses below which there is no risk. This might well be true for background EMF impacting upon long term human health and development^{105, 115, 116, 122-126}.

To an extent this paper challenges one of the most exciting human developments, the emergence of the universal digital age therefore it may arouse the wrath of possible vested interests. We reiterate, we do not want to stop the modern digital age but make it safer, as the long term effects of being closely surrounded by new

compounded levels electro-magnetism are simply unknown but there are symptomatic danger signals.

This study provides sufficient indicative evidence that we need to proceed, guided by the cautionary principle, to re-consider health and safety guidelines in relation to compounded increases in EMF and human health, especially neurological. At the time of writing this has been given greater urgency as charities with Progressive Supranuclear Palsy patients are seeking help under the age of forty, which reflects recent major studies of a cohort of MSA patients mean onset of 37 years^{31-33, 100.}

Such young people with a dementing neurodegenerative disease would have been unknown forty years ago and barely credible thirty years ago. Neurological disease, as with other conditions has a combination of interactive causes but with human neurology, which is essentially an electro-biochemical system, the EMF is likely to play a significant role.

Despite the size of the literary review we returned to Medline for the latest papers on the topic and despite the continuing inconsistencies. There are current papers showing negative associations and one suggesting a nocebo effect, with is the reverse of the placebo effect, self-suggestion that something is wrong¹²⁷ but the further need for caution has not been ruled out. There is beginning to be research across the Western world showing a growing sense of urgency related to EMF impact upon child and adolescent developing brains¹²⁸⁻¹³⁰. Whilst the pre-cautionary measured introduction the use of Resveratrol which appears to reduce oxidative stress and is claimed to be offer some protection against adverse effects of EMF!¹⁰⁵ The key is to think outside the box and considered what might EMF do to the other existing pollutants in the human environment and the number of years it takes to develop neurodegenerative diseases, as with other classic environmental induced pathology^{17,18.}

In closing, we take a backward look over just 25 clinical years to consider the rises in the incidence of Autism and Asperger syndrome, Chronic Fatigue Syndrome, ADHD,

Early Onset Dementias, rises in the WHO diagnostic categories Nervous Disease Deaths and Alzheimer's and Other Dementia deaths across the Western world, all alongside intensive increases in background environmental pollutants and the now ubiquitous EMF. Part of the problem is that these influences have been slow and insidious and the new digital world, so full of 'good things', perhaps we may need to begin to realise that possibly we have opened Pandora's Box. There is a need to re-think the modern world's impact upon human health and its epigenetic development, almost matching as it does climate change, which for so long has relatively been denied.

It is stressed that we do not want to 'stop the modern' EMF world, only make it safer. Finally, we close as we began. It is an evolutionary fact that humans evolved within the Earth's electro-magnetic-field and the sun's radiation. If we have too much sun we burn so are we beginning to get too much EMF that stirs up the soup of interactive multiple-pollutants that impact upon human health?

References

1. Chang PA, Wu YJ. Motor neurone disease and neuro-toxic substances: a possible link? *Chem Biol Interact.* 2009;180:127-130.
2. Cho JW, Jeon BS, Jeong D, et al. Association between Parkinsonism and participation in agriculture in Korea. *J Clin Neurol.* 2008;4(1):23–28.
3. Fang F, Valdimarsdóttir U, Bellocco R, Ronnevi LO, Sparén P, et al. (2009) Amyotrophic lateral sclerosis in Sweden, 1991-2005. *Arch Neurol.* 66: 515-519.
4. Tobin K, Gilthorpe MS, Rooney J, et al. Age- period- cohort analysis of trends in amyotrophic lateral sclerosis incidence. *J Neurol.* 2016;263:1919-1926.
5. Alonso A, Logroscino G, Jick SS, Hernán MA (2009) Incidence and lifetime risk of motor neuron disease in the United Kingdom: a population-based study. *Eur J Neurol.* 16: 745-751.
6. Gordon PH, Artaud F, Aouba A, et al. Changing mortality for motor neuron disease in France (1968-2007):an age-period cohort analysis. *Eur J Epidemiol.* 2011;26(9):729–737.

7. Moura MC, Casulari LA, Carvalho Garbi MR. Ethnic and demographic incidence of amyotrophic lateral sclerosis (ALS) in Brazil: A population based study. *Amyo Lat Scler Front Degen*. 2016;17(3-4):275–281.
8. Shahrizaila N, Sobue G, Kuwabara S, et al. Amyotrophic lateral sclerosis and motor neuron syndromes in Asia. *J Neurol Neurosurg Psychiatry*. 2016;87:821-830.
9. Zhang L, Cao B, Zou Y, et al. Causes of death in Chinese patients with multiple system atrophy. *Aging Dis*. 2018;9(1):102–108.
10. Pakpoor J, Goldacre M. Neuroepidemiology: The increasing burden of mortality from neurological diseases. *Nat Rev Neurol*. 2017. 13:518-519.
11. Pritchard C, Mayers A, Baldwin D. Changing patterns of neurological mortality in the 10 major developed countries--1979-2010. *Public Health*. 2013. 127: 357-368.
12. Pritchard C, Rosenorn-Lanng E. Neurological deaths of American adults (55–74) and the over 75's by sex compared with 20 Western countries 1989–2010: Cause for concern. *Surg Neurol Int*. 2015;6:123.
13. Pritchard C, Rosenorn-Lanng E, Silk A, Hansen L. Controlled population- based comparative study of USA and international adult [55- 74] neurological deaths 1989-2014. *Acta Neurol Scand*. 2017. 136; 698-707.
14. Consales C, Merla C, Marino C, Bennassi B. Electromagnetic Fields, Oxidative Stress and Neurodegeneration. *Int J Cell Biol*. 2012; 16. Doi:10.1155/2012/683897.
15. Chen CY, Hung HJ, Chang KH, et al. Long-term exposure to air pollution and the incidence of Parkinson's disease: A nested case-control study. *PLoS One*. 2017;12(8):e0182834.
16. Jayaraj RL, Rodriguez EA, Wang Y, Blocl ML. Outdoor Ambient Air Pollution and Neurodegenerative Diseases: The neuro-inflammation Hypothesis. *Curr Environ Health Reports*. 4:166-179.
17. Furuya S, Chimed-Ochir O, Takahashi K, David A, Takala J. Global Asbestos Disaster. *Int J Environ Res Public Health*. 2018 May 16;15(5). pii: E1000. doi: 0.3390/ijerph15051000.
18. Kerger BD. Longevity and pleural mesothelioma: age-period-cohort analysis of incidence data from the Surveillance, Epidemiology, and End Results (SEER) Program, 1973-2013. *BMC Res Notes*. 2018 May 23;11(1):337. doi: 10.1186/s13104-018-3436-0.
19. Ross SM, McManus IC, Harrison V, Mason O (2013) Neurobehavioral problems following low-level exposure to organophosphate pesticides: a systematic and meta-analytic review. *Crit Rev Toxicol*. 43: 21-44. VIP
20. Autistic Society (2018) www.austistic.society.org
21. Guerriero F, Ricevuti G. Extremely low frequency electromagnetic fields stimulation modulates autoimmunity and immune responses: A possible immune-

modulatory therapeutic effect in neurodegenerative diseases. *Neural Regen Res.* 2016. 11:1888-1895.

22. Altun G, Deniz ÖG, Yurt KK, Davis D, Kaplan S. Effects of mobile phone exposure on metabolomics in the male and female reproductive systems. *Environ Res.* 2018 Nov;167:700-707.

23. Villarini M, Gambelunghe A, Giustarini D, Ambrosini MV, Fatigoni C, Rossi R, Dominici L, et al. No evidence of DNA damage by co-exposure to extremely low frequency magnetic fields and aluminium on neuroblastoma cell lines. *Mutat Res.* 2017 Nov;823:11-21.

24. Zubko O, Gould RL, Gay HC, Cox HJ, Coulson MC, Howard RJ. Effects of Electromagnetic field emitted by GSM phones on working memories: a meta-analysis. *Int J Geriatr Psychiatry.* 2017. 32: 125-135.

25. Saliev T, Mustapova Z, Kulsharova G, Bulanin D, Mikhalovsky S. Therapeutic potential of EMF for tissue engineering and wound healing. *Cell Prolif.* 2014. 47:485-493.

26. Chen Z, Liu H, Wang H, Wu C, Feng H, Han J. Effects of low-frequency rotary magnetic fields on advanced gastric cancer: Survival and Palliation of symptoms. *J Cancer Res.*(2018) 14: 815-819.

27. Cichoń N, Rzeźnicka P, Bijak M, Miller E, Miller S, Saluk J. Extremely low frequency electromagnetic field reduces oxidative stress during the rehabilitation of post-acute stroke patients. *Adv Clin Exp Med.* 2018 Jul 19. doi: 10.17219/acem/73699.;

28. Easton DM. Gompertzian growth and decay: a powerful descriptive tool for neuroscience. *Physiol Behav.* 2005;86:407-414.

29. Riggs JE, Schochet SS Jr (1992) Rising mortality due to Parkinson's disease and amyotrophic lateral sclerosis: a manifestation of the competitive nature of human mortality. *J Clin Epidemiol.* 45: 1007-1012.

30. Goldacre MJ, Duncan M, Griffith M, Turner MR (2010) Trends in death certification for multiple sclerosis, motor neuron disease, Parkinson's disease and epilepsy in English populations 1979-2006. *J Neurol.* 257: 706-715.

31. Sulena, Gupta D, Sharma AK, et al. Clinical profile of dysphagia in patients with Parkinson's disease, progressive supranuclear. *J Assoc Physicians India.* 2017;65:32-37.

32. Vanacore N. Epidemiological evidence on multiple system atrophy. *J Neural Transm.* 2005;112(12):1605-1612.

33. Bjornsdottir A1, Gudmundsson G, Blondal H, et al. Incidence and prevalence of multiple system atrophy: a nationwide study in Iceland. *J Neurol Neurosurg Psychiatr.* 2013. 84:136-140.

34. Chrysostome V, Tison F, Yekhle F, et al. Epidemiology of multiple system atrophy: a prevalence and pilot risk factor study in Aquitaine, France. *Neuroepidemiology.* 2004. 23:201-208.

35. Yoo HS, Chung SJ, Kim SJ, Oh JS, Kim JS, Ye BS, Sohn YH, Lee PH. The role of 18F-FP-CIT PET in differentiation of progressive supranuclear palsy and frontotemporal dementia in the early stage. *Eur J Nucl Med Mol Imaging*. 2018. 45:1585-1595.
36. World Health Organization. World Statistical Annual 11: 126-139. Geneva, Switzerland: World Health Organization. www.who.int/whosis/mort/table1.process.cfm. 1980-2015 (2018).
37. Sabatelli M, Madia F, Conte A, Luigetti M, Zollino M.) Natural history of young-adult amyotrophic lateral sclerosis. *Neurology*. 2008. 71: 876-881.
38. Nilsson C, Lundqvist LM, Nilsson K, Santillio A, Vestberg S. Age- related incidence and family history in frontotemporal dementia: data from the Swedish Dementia Registry. *PLoS One*. 2014;9:90-94.
39. Panegyres PK, Chen HY. Early- onset Alzheimer's disease: a global cross-sectional analysis. *Eur J Neurol*. 2014;21:1149-1154.
40. Cuyvers E, van der Zee J, Bettern K. Genetic variability in SQSTM1 and risk of early- onset Alzheimer dementia: a European early- onset dementia consortium study. *Neurobiol Ageing*. 2015;36:15-22.
41. Li J, Wu L, Tang Y, Zhou A, Wang F, Xing Y, Jia J. Differentiation of neuropsychological features between posterior cortical atrophy and early onset Alzheimer's disease. *BMC Neurol*. 2018 May 10;18(1):65. doi: 10.1186/s12883-018-1068-6.
42. Perrone F, Cacace R, Van Mossevelde S, Van den Bossche T, De Deyn PP, Cras P, Engelborghs S, van der Zee J, Van Broeckhoven C. Genetic screening in early-onset dementia patients with unclear phenotype: relevance for clinical diagnosis. *Neurobiol Aging*. 2018 Sep;69:292.e7-292.e14. doi: 10.1016/j.neurobiolaging.2018.04.015. Epub 2018 May 9.
43. Rantalainen V, Lahti J, Henriksson M, Kajantie E, Eriksson JG, Räikkönen K. Cognitive ability in young adulthood predicts risk of early-onset dementia in Finnish men. *Neurology*. 2018 Jul 10;91(2):e171-e179. doi: 10.1212/WNL.0000000000005757. Epub 2018 Jun 6.
44. Sanchez AM, Scharovsky D, Romano LM, et al., et al. Incidence of early- onset dementia in Mar del Plata. *Neurologia*. 2015;30: 77-82.
45. Batla A, De Pablo-Fernandez E, Erro R, Reich M, Calandra-Buonaura G, Barbosa P, Balint B, Ling H, Islam S, Cortelli P, Volkmann J, Quinn N, Holton JL, Warner TT, Bhatia KP. Young-onset multiple system atrophy: Clinical and pathological features. *Mov Disord*. 2018.33:1099-1107.
46. Beard JD, Steege AL, Ju J. Mortality from amyotrophic lateral sclerosis and Parkinson's disease among different occupation groups - United States, 1985-2011. *Morb Mortal Wkly Rep*. 2017;66(22):718-722.
47. Bergman BP, Mackay DF, Pell JP. Motor neurone disease and military service: evidence from the Scottish Veterans Health Study. *Occup Environ Med*. 2015;72:877-879.

48. Parlett LE, Bowman JD, van Wijngaarden E. Evaluation of occupational exposure to magnetic fields and motor neuron disease mortality in a population-based cohort. *J Occup Environ Med*. 2011 53: 1447-1451.
49. Sutedja NA, Fischer K, Veldink JH, van der Heijden GJ, Kromhout H. What we truly know about occupation as a risk factor for ALS: a critical and systematic review. *Amyotroph Lateral Scler*. 2009.10: 295-301.
50. Stampfer MJ. Welding occupations and mortality from Parkinson's disease and other neurodegenerative diseases among United States men, 1985-1999. *J Occup Environ Hyg*. 2009. 6:267-272.
51. Callaghan B, Feldman D, Gruis K, Feldman E (2011) The association of exposure to lead, mercury and selenium and the development of amyotrophic lateral sclerosis and the epigenetic implications. *Neurodegener Dis* 8: 1-8.
52. Fischer H, Kheifets L, Huss A, et al. Occupational exposure to electric shocks and magnetic fields and amyotrophic lateral sclerosis in Sweden. *Epidemiology*. 2015.26:137-146.
53. Chen H, Kwong J, Copes R, Tu K, Villeneuve PJ, van Donkenlarn A. Living near major roads and the incidence of dementia, Parkinson's disease and multiple sclerosis. *Lancet*. 2017. 389:718-726.
54. Goldman SM, Quinlan PJ, Ross GW, et al. Solvent exposures and Parkinson disease risk in twins. *Ann Neurol*. 2012.71:776-784.
55. Gore AC. Neuro- endocrine targets of endocrine disruptors. *Hormones*. 2010. 9: 16-27.
56. Han YH, Lee JH, Kang BM. Topographical differences of brain iron deposition between progressive supranuclear palsy and parkinsonian variant multiple system atrophy. *J Neurol Sci*. 2013. 325:29-35.
57. Begum N, Wang B, Mori M, Vares G (2012) Does ionizing radiation influence Alzheimer's disease risk? *J Radiat Res*. 53: 815-822.
58. Zhang Y, Zhang Y, Yu H, Yang Y, Li W, Qian Z. Theta-gamma coupling in hippocampus during working memory deficits induced by low frequency electromagnetic field exposure. *Physiol Behav*.
59. Hallberg O. A trend modal Alzheimer's disease. *ADMET*. 2015;3:281-286.
60. Xu Y, Liu X, Shen J, Tian W, Fang R, Li B, Ma J, Cao L, Chen S, Li G, Tang H. The Whole Exome Sequencing Clarifies the Genotype- Phenotype Correlations in Patients with Early-Onset Dementia. *Aging Dis*. 2018.9 696-705.
61. Bastos AF, Orsini M, Machado D, Mello MP, Nader S, et al. (2011) Amyotrophic lateral sclerosis: one or multiple causes? *Neurol Int*. 2011 Jun;3(1):e4. doi: 10.4081/ni.2011.e4.
62. Al-Chalabi A, Lewis CM (2011) Modelling the effects of penetrance and family size on rates of sporadic and familial disease. *Hum Hered*. 71: 281-288.

63. Little J, Barakat-Haddad C, Martino R, Pringsheim T, Tremlett H, McKay KA, van Lieshout P, Walsh SJ, Gomes J, Krewski D. Genetic variation associated with the occurrence and progression of neurological disorders. *Neurotoxicology*. 2017. 61:243-264.
64. Lunke S, El-Osta A. The emerging role of epigenetic modifications and chromatin remodelling in spinal muscular atrophy. *J Neurochem*. 2009. 109: 15571569.
65. Sanyal J, Chakraborty DP, Sarkar B, et al. Environmental and familial risk factors of Parkinson's disease: case-control study. *Can J Neurol Sci*. 2010.37.:637–642.
66. Gangisetty O, Murugan S. Epigenetic modifications in neurological disease: natural products as epigenetic modulators- a treatment strategy. *Adv Neurobiol*. 2016;12:1-25.
67. Goldman SM, Quinlan PJ, Ross GW, et al. Solvent exposures and Parkinson disease risk in twins. *Ann Neurol*. 2012.71:776–784.
68. Dai MH, Zheng H, Zeng LD, Zhang Y. The genes associated with early-onset Alzheimer's disease. *Oncotarget*. 2017. 15:15132-15143.
69. Blank M. Protein and DNA reactions stimulated by electromagnetic fields. *Electromagn Biol Med*. 2008. 27: 3-23.
70. Pasanen P, Myllykangas L, Pöyhönen M, Kiviharju A, Siitonen M, Hardy J, Bras J, Paetau A, Tienari PJ, Guerreiro R, Verkkoniemi-Ahola A. Genetics of dementia in a Finnish cohort. *Eur J Hum Genet*. 2018 Jun;26(6):827-837.
71. Akerstedt T, Arnetz B, Ficca G, Paulsson LE, Kallner A. A 50-Hz electromagnetic field impairs sleep. *J Sleep Res*. 1999.8:77-81.
72. Balamuralikrishnan B, Balachandar V, Kumar SS, Stalin N, Vasha P. Evaluation of chromosomal alteration in electrical workers occupationally exposed to low frequency of electro-magnetic-fields in Coimbatore population, India. *Asian Pac J Cancer Prev*. 2012. 13:2961-2966.
73. Lasalvia M, Scrima R, Perna G, Piccoli C, Capitano N, Biagi PF, Schiavulli L. Exposure to 1.8 GHz electromagnetic fields affects morphology, DNA-related Raman spectra and mitochondrial functions in human lympho-monocytes. *PLoS One*. 2018 Feb 20;13(2):e0192894. doi: 10.1371/journal.pone.0192894.
74. Buchachenko AL, Kuznetsov DA, Berdinskii VL. New mechanisms of biological effects of electromagnetic fields. *Biophysics*. 2006. 51: 489-496.
75. Buchachenko AL, . Spin biochemistry: magnetic ²⁴Mg-²⁵Mg-²⁶Mg isotope effect in mitochondrial ADP phosphorylation. *Cell Biochem Biophys*. 2005. 43:243-51.
76. Davarpanah Jazi S, Modolo J, Baker C, Villard S, Legros A. Effects of A 60 Hz Magnetic Field of Up to 50 milliTesla on Human Tremor and EEG: A Pilot Study. *Int J Environ Res Public Health*. 2017 Nov 24;14(12). pii: E1446. doi: 10.3390/ijerph14121446.

77. Eghlidospouyhr M, Ghanbari A, Mortazavi SMJ, Azari H., Effects of radiofrequency exposure emitted from a GSM mobile phone on proliferation, differentiation and apoptosis of neural stem cells. *Anal Cell Biol.* 2017. 50: 115-123.
78. Halgamuge MN (2013) Pineal melatonin level disruption in humans due to electromagnetic fields and ICNIRP limits. *Radiat Prot Dosimetry* 154: 405-416.
79. Kesari KK¹, Siddiqui MH, Meena R, Verma HN, Kumar S. Cell phone radiation exposure on brain and associated biological systems. *Indian J Exp Biol.* 2013. 51:187-200.
80. Burke, H (1986) *Handbook of Magnetic Phenomena*. New York, Van Nostrand.
81. Zhi, WJ, Wang LF, Hu, XJ. Recent advances in the effect of microwave radiation on brains. *Military Medical Research* 2017. 12: 4-29.
82. Ravera S, Falugi C, Calzia, D; Pepe, IM, Panfoli, I. First Cell Cycles of Sea Urchin *Paracentrotus lividus* Are Dramatically Impaired by Exposure to Extremely Low-Frequency Electromagnetic Field . *Biology of Reproduction*, 2006, 75: 948–953
83. Kimsa-Dudek M, Synowiec-Wojtarowicz A, Derewniuk M, Gawron S, Paul-Samojedny M. Impact of fluoride and a static magnetic field on the gene expression that is associated with the antioxidant defence system of human fibroblasts. *Chem Biol Interact.* 2018. 287:13-19.
84. Reale M, Kamal MA, Patruno A, Costantini E, D'Angelo C, Pesce M, Greig NH. Neuronal cellular responses to extremely low frequency electromagnetic field exposure: implications regarding oxidative stress and neurodegeneration. *PLoS One.* 2014 Aug 15;9(8):e104973. doi: 10.1371/journal.pone.0104973. eCollection 2014.
85. Migliore L, Coppedè F. Environmental-induced oxidative stress in neurodegenerative disorders and aging. *Mutat Res.* 2009. 31:73-84.
86. Kivrak EG, Yurt KK, Kaplan AA, Alkan I, Altun G. Effects of electromagnetic fields exposure on the antioxidant defence system. *J Microsc Ultrastruct.* 2017. 5:167-176.
87. Zablotskii V, Polyakova T, Dejneka A. Cells in Non-Uniform Magnetic World: How Cells Respond to High-Gradient Magnetic Fields. *Bioessays.* 2018 Jun 25:e1800017. doi: 10.1002/bies.201800017.
88. Zhi WJ, Peng RY, Li HJ, Zou Y, Yao BW, Wang CZ. Microwave radiation leading to shrinkage of dendritic spines in hippocampal neurons mediated by SNK-SPAR pathway. *Brain Res.* 2018. 1679: 134-143.
89. Deniz OG, Kaplan S, Selçuk MB, Terzi M, Altun G, Yurt KK, Aslan K, Davis D. Effects of short and long term electromagnetic fields exposure on the human hippocampus. *J Microsc Ultrastruct.* 2017 Oct-Dec;5(4):191-197.
90. Ross CL, Pettenati MJ, Procita J, Cathey L, George SK, Almeida-Porada G. Evaluation of Cytotoxic and Genotoxic Effects of Extremely Low-frequency Electromagnetic Field on Mesenchymal Stromal Cells. *Glob Adv Health Med.* 2018 May 18;7:2164956118777472. doi: 10.1177/2164956118777472. eCollection 2018.

91. Teschke K, Marion SA, Tsui JK, et al. Parkinson's disease and occupation: differences in associations by case identification method suggest referral bias. *Am J Ind Med*. 2014. 57:163–171.
99. Purdey M. Elevated levels of ferrimagnetic metals in food chains supporting the Guam cluster of neurodegeneration: do metal nucleated crystal contaminants [corrected] evoke magnetic fields that initiate the progressive pathogenesis of neurodegeneration. *Med Hypotheses*. 2004. 63:793-809.
91. Johnansen C (2004) Electromagnetic fields and health effects – epidemiological studies of cancer, diseases of the central nervous system and arrhythmia-related heart disease. *Scand J Work Environ Health*. Supplement 1: 1-80.
92. Hug K, Rösli M, Rapp R. Magnetic field exposure and neurodegenerative diseases--recent epidemiological studies. *Soz Präventivmed*. 2006. 51:210-20.
93. Hasan GM, Skeihk A, Karim S, Hague A, Kamal MA. Effect of electromagnetic radiations on neurodegenerative disease – technological revolution as a curse in disguise. *CNS Neurol Disord Drug Targets*. 2014. 13:1406-1412.
94. Ghadamgahi M¹, Monazzam MR², Hosseini M³. Memory loss risk assessment for the students nearby high-voltage power lines-a case study. *Environ Monit Assess*. 2016 Jun;188(6):355. doi: 10.1007/s10661-016-5358-4. Epub 2016 May 18.
95. Foerster M, Thielens A, Joseph W, Eeftens M, Rösli M. A Prospective Cohort Study of Adolescents' Memory Performance and Individual Brain Dose of Microwave Radiation from Wireless Communication. *Environ Health Perspect*. 2018 Jul 23;126(7):077007. doi: 10.1289/EHP2427. eCollection 2018 Jul.
96. Kheifets L, Repacholi M, Saunders R, van Deventer E. The sensitivity of children to electromagnetic field. *Pediatrics*. 2005.116:303-313.
97. Zhou H, Chen G, Chen C, Yu Y, Xu Z (2012) Association between extremely low-frequency electromagnetic fields occupations and amyotrophic lateral sclerosis: a meta-analysis. *PLoS One* 7: e48354.
98. van der Mark M, Vermeulen R, Nijssen PC, et al. Extremely low-frequency magnetic field exposure, electrical shocks and risk of Parkinson's disease. *Int Arch Occup Environ Health*. 2015. 88:227–234.
99. Campbell RB, Dini DA. *Occupational Injuries from Electrical Shock and Arc Flash Events*. New York: Springer; 2016.
100. Pritchard C, Silk A. Patient's occupation, electric & head trauma in a cohort of 88 multiple system atrophy patients compared with the general population: a hypothesis stimulating pilot study. *J Neurol Stroke*. 2018. 8:178–183.
101. Huss A, Vermeulen R, Bowman JD, et al. Electric shocks at work in Europe: development of a job exposure matrix. *Occup Environ Med*. 2013;70(4):261–267.
102. Huss A, Spoerri A, Egger M. Swiss National Cohort. Occupational exposure to magnetic fields and electric shocks and risk of ALS: the Swiss National Cohort. *Amyo Latl Scler Front Degener*. 2015.16: 80–85.

103. Pritchard C, Silk A. A Case-Study Survey of an Eight-year Cluster of Motor Neurone Disease (MND) Referrals in a Rural English Village: Exploring Possible Aetiological Influences in a Hypothesis Stimulating Study. *J Neurol Disord* . 2014. 2: 1-5.
104. Huss A, Peters S, Vermeulen R. Occupational exposure to extremely low-frequency magnetic fields and the risk of ALS: A systematic review and meta-analysis. *Bioelectromagnetics*. 2018. 39:156-163.
105. Zhang D, Zhang Y, Zhu B, Zhang H, Ye Sun C. Resveratrol may reverse the effects of long-term occupational exposure to EMF on workers of a power plant. 2017. *Oncotarget*. 8:4797-47506.
106. Clem PA, Navathe PD, Drane MA. Identifying Pilots with Parkinson's Disease. *Aerosp Med Hum Perform*. 2016.87:545-549.
107. Arva P, Wagstaff AS. Medical disqualification of 275 commercial pilots: changing patterns over 20 years. *Aviat Space Environ Med*. 2004. 75:791-4.
108. Jagathesan T, O'Brien MD. The Aeromedical Implications of Parkinson's Disease. *Aerosp Med Hum Perform*. 2015. 86: 1046-1051.
109. Capozzella A¹, Sacco C¹, Chighine A¹, Loreti B¹, Scala B¹, Casale T¹, Sinibaldi F¹, Tomei G², Giubilati R¹, Tomei F¹, Rosati MV¹. Work related aetiology of amyotrophic lateral sclerosis (ALS): a meta-analysis. *Ann Ig*. 2014. 26:456-72.
110. Hobson P, Gallagher J, Meara J. Cross-sectional survey of Parkinson's disease and parkinsonism in a rural area of the United Kingdom. *Mov Disord*. 2005.20:995-998.
111. Wakeford R (2018) The growing importance of radiation worker studies. *Brit J Cancer*. 119:527-529.
112. Haylock RGE, Gillies M, Hunter N, Zhang, W , Phillipson M. Cancer mortality and incidence following external occupational radiation exposure: An update of the 3rd analysis of the UK Registry for Radiation Workers. *Brit J Cancer*. 2018. 199: 530-44.
113. Pakpoor J, Noyce A, Goldacre R, Selkikhova M, Mullin S, Schrag A, Lees A, Goldacre M. Viral hepatitis and Parkinson disease: A national record-linkage study. *Neurology*. 2017. 88:1630-1633.
114. De Pablo-Fernandez E, Goldacre R, Pakpoor J, Noyce AJ, Warner TT. Association between diabetes and subsequent Parkinson disease: A record-linkage cohort study. *Neurology*. 2018 Jul 10;91(2):e139-e142. doi: 10.1212/WNL.0000000000005771..
115. Schwarzingler M, Pollock BG, Hasan OSM, Dufouil C, Rehm J; QalyDays Study Group. Contribution of alcohol use disorders to the burden of dementia in France 2008-13: a nationwide retrospective cohort study. *Lancet Public Health*. 2018 Mar;3(3):e124-e132. doi: 10.1016/S2468-2667(18)30022-7. Epub 2018 Feb 21.
116. Sarchet P. The Allergy Epidemic. *New Scientist*. 2018.3190: 28-35.

117. United States Department of Labor (2002). "Controlling Electric Hazards, Revision. Washington DC.
118. Committee on Man and Radiation (COMAR). COMAR technical information statement: expert reviews on potential health effects of radiofrequency electromagnetic fields and comments on the bioinitiative report. *Health Phys.* 2009. 97:348-56.
119. EUROPAEM EMF Guideline 2016 for the prevention, diagnosis and treatment of EMF-related health problems and illnesses. *Bioelectromagnetics*.2017. 38:386-396.
120. Trempe CL, Lewis TJ. It's Never Too Early or Too Late-End the Epidemic of Alzheimer's by Preventing or Reversing Causation From Pre-birth to Death. *Front Aging Neurosci.* 2018 Jul 12;10:205. doi: 10.3389/fnagi.2018.00205. eCollection.
121. Carpenter DO. Human disease resulting from exposure to electromagnetic fields. *Rev Environ Health.* 2013. 28: 159-172.
122. Christ A, Guldimann R, Bühlmann B, Zefferer M, Bakker JF, et al. (2012) Exposure of the human body to professional and domestic induction cooktops compared to the basic restrictions. *Bioelectromagnetics* 33: 695-705.
123. Fernández C, de Salles AA, Sears ME, Morris RD, Davis DL. Absorption of wireless radiation in the child versus adult brain and eye from cell phone conversation or virtual reality. *Environ Res.* 2018 Nov;167:694-699.
124. Birks LE, Struchen B, Eeftens M, van Wel L, Huss A, Gajšek P, Kheifets L, Gallastegi M, Dalmau-Bueno A, Estarlich M, Fernandez MF, Meder IK, Ferrero A, Jiménez-Zabala A, Torrent M, Vrijkotte TGM, Cardis et al. Spatial and temporal variability of personal environmental exposure to radio frequency electromagnetic fields in children in Europe. *Environ Int.* 2018 Aug;117:204-214. doi: 10.1016/j.envint.
125. Bhatt CR, Redmayne M, Billah B, Abramson MJ, Benke G. Radiofrequency-electromagnetic field exposures in kindergarten children. *J Expo Sci Environ Epidemiol.* 2017 Sep;27(5):497-504. doi: 10.1038/jes.2016.55.
126. Pedersen C, Poulsen AH, Rod NH, et al. Occupational exposure to extremely low-frequency magnetic fields and risk for central nervous system disease: an update of a Danish cohort study among utility workers. *Int Arch Occup Environ Health.* 2017. 90:619–628.
127. Eltiti S, Wallace D, Russo R, Fox E. Symptom Presentation in Idiopathic Environmental Intolerance With Attribution to Electromagnetic Fields: Evidence for a Nocebo Effect Based on Data Re-Analyzed From Two Previous Provocation Studies. *Front Psychol.* 2018 Aug 28;9:1563. doi: 10.3389/fpsyg.2018.01563. eCollection 2018.
128. Zeleke BM, Brzozek C, Bhatt CR, Abramson MJ, Croft RJ, Freudenstein F, Wiedemann P, Benke G. Personal Exposure to Radio Frequency Electromagnetic Fields among Australian Adults. *Int J Environ Res Public Health.* 2018 Oct 12;15(10). pii: E2234. doi: 10.3390/ijerph15102234.

129. Guxens M, Vermeulen R, Steenkamer I, Beekhuizen J, Vrijkotte TGM, Kromhout H, Huss A. Radiofrequency electromagnetic fields, screen time, and emotional and behavioural problems in 5-year-old children. *Int J Hyg Environ Health*. 2018 Oct 9. pii: S1438-4639(18)30502-9. doi: 10.1016/j.ijheh.2018.09.006. [Epub ahead of print]

130. Yamaguchi-Sekino S, Sekino M, Ueno S. Biological effects of electromagnetic fields and recently updated safety guidelines for strong static magnetic fields. *Magn Reson Med Sci*. 2011;10(1):1-10.