



University of Dundee

Lignin engineering to improve saccharification and digestibility in grasses

Halpin, Claire

Published in:
Current Opinion in Biotechnology

DOI:
[10.1016/j.copbio.2019.02.013](https://doi.org/10.1016/j.copbio.2019.02.013)

Publication date:
2019

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Halpin, C. (2019). Lignin engineering to improve saccharification and digestibility in grasses. *Current Opinion in Biotechnology*, 56, 223-229. <https://doi.org/10.1016/j.copbio.2019.02.013>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Lignin engineering to improve saccharification and digestibility in grasses

Claire Halpin

Address:

Division of Plant Sciences

School of Life Sciences

University of Dundee at the JHI

Invergowrie

Dundee DD2 5DA

United Kingdom

E-mail: c.halpin@dundee.ac.uk

Abstract

The digestibility of plant biomass has a major influence on its value as a forage for livestock and as a feedstock for industrial biotechnology. For both processes, the concentration, structure, and composition of lignin influence the accessibility of wall carbohydrate polymers to microbes and digestive enzymes during biochemical decomposition. Although lignin engineering has been less tractable in monocots than in model dicots, a body of work is accumulating on the effects of manipulating lignin biosynthesis in energy grasses and cereal crops. In addition to conventional targets for lignin engineering, several novel features of grass lignin have recently become amenable to targeted manipulation through the identification of genes involved in their synthesis.

Introduction

Many feedstocks that might be used for future biorefinery applications come from grasses, be it biomass from perennial energy crops (switchgrass, miscanthus, sugarcane) or agricultural coproducts such as cereal straw from food crops. In these materials, lignin can represent 20% or more of the biomass, and will inhibit biomass digestion in several ways, mainly by preventing microbes and enzymes from gaining access to cellulose, but also by binding digestive enzymes or releasing inhibitory breakdown products. Physical and chemical pretreatments are therefore necessary to facilitate biomass digestion by removing some xylans and lignin to enable enzymes to gain access to the hydrophobic cellulose face [1] and release simple sugars (saccharification). An ideal pretreatment might remove as much lignin as possible while minimizing polysaccharide modification to retain a near-native microfibrillar structure that enzymes can attack [1]. Ruminant animals have solved the problem of releasing sugars from plant biomass in ways somewhat analogous to industrial biochemical processing, i.e., by effective physical pretreatment (chewing, and rechewing regurgitated solids), then slow microbial fermentation in the rumen and peptic/enzymatic digestion of non-fermented material. Although biomass digestion in the rumen, and in industry, may not always depend on the same factors, there is sufficient commonality for synergy between both camps in identifying targets to increase biomass digestibility.

Special features of lignin biosynthesis in grasses

Lignin biosynthesis and structure are discussed elsewhere in this issue [2,3]. Briefly, lignin is a polymer of predominantly three monolignols *p*-coumaryl alcohol, coniferyl alcohol, and sinapyl alcohol, giving rise to H, G and S units. Like dicots, grasses have an SG-type lignin but with significantly more H units, although H levels are still low. The enzymes of the phenylpropanoid pathway that produce these monolignols are obvious targets for lignin genetic engineering to improve biomass digestibility (Figure 1). This 'conventional' lignin pathway operates similarly in dicots and grasses with a few distinctions. A key intermediate, *p*-coumarate, derives partially from deamination of tyrosine in grasses, whereas it derives solely from deamination of phenylalanine followed by 4-hydroxylation in dicots. Some grass PAL enzymes also possess tyrosine ammonia-lyase activity and are bifunctional phenylalanine/tyrosine ammonia-lyases, PTALs [4,5]. In addition, CSE is crucial to lignin biosynthesis in several dicots and influences saccharification efficiency but only some grasses have an orthologous gene [6-8].

In addition to conventional components, grass lignins have several special features that offer targets for lignin engineering beyond those usually possible in dicots (Figure 1). These include the incorporation of the flavone tricetin as an additional lignin monomer [9,10], the *p*-coumaroylation of (mainly) S lignin units, and the incorporation of ferulates into lignin [11]. Genes controlling incorporation of these components are only now coming to light and present novel lignin engineering opportunities. Because of the clear differences in structure between grass and dicot lignins, the results of specific gene manipulations in dicots should not be extrapolated to possible digestibility benefits in grasses – this must be evaluated directly in grass species.

Conventional targets for reducing lignin content in grasses

It is absolutely clear that the amount of lignin in grass cell walls is generally negatively correlated with digestibility. For example, in maize inbreds, diverse miscanthus species and a variety of cereal crops, biomass digestibility in rumen fluid or glucose yields on saccharification were negatively correlated with Klason lignin [12-14]. To reduce lignin content, engineering strategies can be targeted at the early phenylpropanoid pathway to alter flux into all subsequent steps, i.e., manipulating *PTAL* and *4CL*. Manipulating *C4H* is unlikely to be effective as grasses can bypass a requirement for *C4H* using *PTAL* (Figure 1). When multiple *PAL* genes were suppressed by RNAi in brachypodium the result was a large 43% decrease in lignin and nearly 2-fold increase in sugars released after a limited-extent digestion [4]. Increases in S/G and H units were explained by potential differences in relative flux through those arms of the pathway. However, the plants showed developmental delays and increased susceptibility to some pathogens [4]. More specific suppression of *PTAL* would be instructive but, although such plants have been produced [5], digestibility data has not been reported.

Suppression of class I *4CLs* (i.e., *4CLs* potentially involved in monolignol biosynthesis) in switchgrass, rice, sugarcane and sorghum *bmr2* mutants leads to around 17-20% lignin reduction, usually with brown midrib-like phenotypes [15-18]. In cases in which lignin monomer ratios were determined, these generally showed equivalent reductions in G and S, although switchgrass showed differential effects on the three monolignols. Biomass yield was normal in most cases although rice plants were shorter and less fertile than controls [16]. Sugarcane had reduced height but produced more tillers so biomass was maintained in most lines, even in the field [17]. Saccharification efficiency after dilute acid pretreatment was 52-76% improved in *4CL*-suppressed switchgrass and field-grown sugarcane [15,17]. Thus it appears that *4CL* can be an effective target for moderate reduction of lignin content with significant benefits for digestibility even in field-grown plants.

Genes common to all three branches of the lignin pathway (i.e., *CCR* and *CAD*) might also be expected to reduce overall lignin content. This has proven to be the case although changes to monomeric ratios have also been seen. Digestibility improvements consequent on *CCR* deficiency (by RNAi or mutation) have been reported in maize and perennial ryegrass [19-22]. Lignin content was lowered by 9-37%, with all three monomers generally reduced (although results on H units were inconsistent between maize studies), and S/G ratio either not altered or increased. Both species showed no change to growth, even in field-grown plants, and digestibility in rumen fluid was 14-15% improved [19,20]. Glucose release after pretreatment and limited saccharification increased by up to 40% in maize. As with *CCR*, *CAD*-

suppression should affect all three monolignol branches. Mutants or transgenics for the major lignin biosynthetic CAD exist in many grasses and consistently show 6-26% reductions in lignin [23-30]. Impacts on G and S vary, possibly reflecting a contribution of other *CAD* genes, or differences between *CAD* knock-out mutants and knocked-down/suppressed plants. S units are generally the most reduced component, with G units reduced or proportionally increased in different cases, generally resulting in a drop in S/G [24-26] or no S/G change [23,27,29]. Sinapaldehyde [24,26] and sometimes coniferaldehyde [29] is present in the lignin which may have a more condensed structure. An enrichment in lignin free-phenolic end-groups makes the polymer more soluble in alkali. Both 44-46% increased sugar yields after limited saccharification [26], and increased ethanol yields [28] are reported after mild alkaline pretreatments. Digestibility in rumen fluid is also increased [24,29].

Conventional targets for altering lignin monomer ratios in grasses

Studies conflict on whether changes to lignin monomer ratios could have benefits for biomass digestibility. Differences between experiments in species studied, growth conditions, tissue harvested, lignin analysis method, pretreatment used, etc., likely influence results and make it difficult to draw conclusions. Illustrative experiments in which both lignin compositional data and digestibility are reported are discussed below to enable some evaluation.

The proportion of H units in lignin might be increased by suppressing enzymes that divert *p*-coumaroyl-CoA into other branches, i.e., HCT, C3H and CSE. To date, only the first two have been manipulated in grasses. Maize RNAi-suppressed in one of two *C3H* genes had the proportion of lignin H units increased from 2% to nearly 8% with little change in overall lignin levels; one line had improved biomass degradability but growth was affected [31]. Rice CRISPR/Cas9-mediated *C3H* knockouts arrested development at the 4-5 leaf stage, although less severely affected *C3H*-RNAi-suppressed plants had only slightly reduced height [32]. *C3H*-RNAi lines had reductions in stem total lignin, G, S and ferulate units, with increased levels of β -ether-linked H units, triclin and *p*-coumarate [32]. Most likely several of these changes contributed to the observed improved saccharification yields [32]. Nevertheless, the consistent appearance of severe phenotypes, both in grasses and in dicots, when *HCT* or *C3H* are suppressed, suggest that they are not robust targets for lignin engineering.

It has been suggested from dicot studies that S/G ratio influences digestibility. S/G might be specifically manipulated by targeting CCoAOMT, F5H (CAld5H) or COMT. Maize and rice have been produced with manipulated CCoAOMT or F5H, but digestibility was not reported [33,34]. By comparison, *COMT*, the gene underlying improved digestibility in *bm3* mutants, has been an obvious target for transgenic manipulation in biomass and forage grasses and cereal crops [e.g., 35-38] with several evaluations progressing to the field [38-42]. Together, these demonstrate changes in *COMT*-suppressed transgenics consistent with those in *bm3* plants, i.e., moderate reductions to lignin (6-14%), with more dramatic reductions in S units (20-46%) accompanied by a decrease in S/G. Some reports record reduced levels of ester-linked *p*-coumarate in lignin with normal levels of ferulate [37,40] and/or the appearance of 'signature' 5-OH-G units [35,37]. Growth is generally normal although field-grown sugarcane had normal biomass at 6% lignin reduction, but decreased biomass at 8-12% lignin reduction [40]. Digestibility for forage use was increased in

COMT-suppressed maize, ryegrass, and tall fescue by 9-11% [35,38,43]. Likewise, saccharification efficiency was increased in sugarcane and switchgrass under a range of conditions [40,41,44] and product yields on simultaneous saccharification and fermentation, or after fermentation with *Clostridium thermocellum* (capable of consolidated bioprocessing), were also increased [36]. Switchgrass transgenics had over 20% improvement in cellulose accessibility [44], likely a crucial factor in the improved digestibility. Similarly, *COMT* brown-midrib mutants are thought to be more digestible only because of their lower lignin content, the modified lignin being more inhibitory to biomass digestion [45].

Novel targets for lignin engineering in grasses

Given that tricetin is a lignin monomer and appears to function as a nucleation site for lignification in grasses [46,47], inhibiting such nucleation might reduce lignin content and increase digestibility. However, a maize mutant in *chalcone synthase C2* (acting at the entry point to the flavonoid pathway) with dramatically reduced tricetin had increased leaf lignin due to redirection of flux from the blocked flavonoid pathway into lignin, reducing leaf saccharification efficiency [48]. *COMT* is involved in later steps in tricetin biosynthesis and its suppression led to reduced tricetin [37,49,50]; conversely suppression of *C3H* in rice increased tricetin [32], yet both types of manipulation improved saccharification. A rice mutant in the *FNSII* gene specific to the flavonoid pathway and devoid of tricetin had enhanced saccharification efficiency, but also unexpectedly showed reduced lignin S/G, as well as incorporating the intermediate naringenin in place of tricetin [51]. Given the range of changes to lignin in these variously tricetin-modified plants, improved saccharification cannot be assigned to any specific change. Clearly, the interactions between flavonoid and lignin biosynthesis require more study before they can be manipulated in a predictable way.

The *p*-coumarate in grass lignin mainly acylates S units; it is attached to monolignols intracellularly by a BAHD *p*-coumaroyl-CoA:monolignol transferase or *PMT* [52-55]. Suppression and overexpression of *PMT* in brachypodium resulted in 10% decrease and 3-fold increase respectively in *p*-coumarate on lignin. *PMT*-overexpressing plants gave higher sugar yields on saccharification after certain pretreatments, although they also had reduced Klason lignin and changes to lignin monomer ratios [53]. Similarly, brachypodium *PMT* expressed in arabidopsis under the control of the *C4H* promoter produced levels of lignin coumaroylation comparable to those in grasses, decreased lignin content by 10-30%, and increased saccharification yields with and without pretreatment [54]. The greatest saccharification improvements were seen in an Arabidopsis *CCR*-mutant background in which lignin ferulate esters were also increased; these plants had a higher incidence of lignin free-phenolic end-groups with 10% decreased lignin and released about 50% more glucose on saccharification [54].

The presence of ferulates is another interesting phenomenon of grass cell walls. Quite high levels acylate arabinoxylans and produce important cross-links between arabinoxylan chains and between arabinoxylans and lignin [55]. The chemistry of such cross-links is reviewed elsewhere in this issue [56] and their presence in cell walls could clearly impact wall pore-size and the mobility of cell wall-digesting enzymes. Several *BAHD* genes have been implicated in the conjugation of ferulate to xylan's arabinosyl units as their downregulation reduces wall-bound ferulate in rice [57], brachypodium [58] and *Setaria* [59]. In the later case, a significant improvement in saccharification efficiency was also demonstrated. The changes in these plants

appear relatively specific to ester-linked ferulate reduction as lignin content was not altered, although there were increases in wall-esterified *p*-coumarate. It was recently discovered that ferulates also acylate monolignols, primarily sinapyl alcohol in grasses, and are incorporated into lignins to produce so-called 'zip-lignins' [60]. *BAHD* genes related to those known to conjugate ferulate to arabinoxylan are likely involved in conjugating ferulate to monolignols. A rice feruloyl-CoA monolignol transferase (FMT) has been identified and its overexpression in activation-tagged mutants or transgenics increases β -ether-bonded monolignol ferulates [60]. Such conjugates were also increased in maize CCR-deficient mutants in which the flux of feruloyl-CoA toward G and S lignin biosynthesis is blocked [21], adding yet another change to the lignin modifications described in CCR-deficient plants that may contribute to improved saccharification sugar yields.

Conclusions

Unravelling the influences on digestibility in grasses of different components and structures in lignin is difficult, as several features concomitantly change in most genetic interventions, illustrating inter-dependency between them. A constant feature in many improved-digestibility transgenic grasses and mutants is a moderate reduction in lignin. This has no, or minimal, impacts on biomass yield for several gene targets (*4CL*, *CCR*, *COMT*, *CAD*, *PMT*). Other lignin changes seen in several but not all improved-digestibility transgenics include reduced or unaltered S/G, an increase in free-phenolic end-groups, and possibly a decrease in esterified *p*-coumarate. These generalisations need to be viewed with caution given the small number of underpinning experiments, and the lack of digestibility data for several potential targets. Adverse effects on growth, if they occur, might be mitigated by careful screening of transgenic events and by introgressing into several cultivars to explore the most compatible background genetics. In addition, the advent of gene editing enables more precise and stable engineering of specific gene/enzyme activity levels to help avoid adverse effects. The several experiments and targets that have now progressed as far as field trials and still show beneficial effects of lignin engineering for digestibility, should encourage optimism for the future and the exploration of conventional and novel targets in increasingly sophisticated ways.

Acknowledgements

CH is a Royal Society Wolfson Research Merit Award holder and is also supported by BBSRC (grants BB/N023455/1; BB/P022499/1; BB/N013689/1) and Royal Society (grant CHG\R1\170091).

References

- [1] Ding S-Y Liu, YS, Zeng YN, Himmel MR, Baker JO, Bayer EA: **How does plant cell wall nanoscale architecture correlate with enzymatic digestibility.** *Science* 2012, **338**:1055-1060.
- [2] Vanholme R, De Meester B, Ralph J and Boerjan W. **Lignin biosynthesis and its integration into metabolism.** *Curr Opin Biotechnol* 2019, **this issue**.
- [3] Ralph J, Lapierre C, Boerjan W: **Lignin structure and its engineering.** *Curr Opin Biotechnol* 2019, **this issue**.

[4] Cass CL, Peraldi A, Dowd PF, Mottiar Y, Santoro N, Karlen SD, Bukhman YV, Foster CE, Thrower N, Bruno LC *et al.*: **Effects of *PHENYLALANINE AMMONIA LYASE (PAL)* knockdown on cell wall composition, biomass digestibility, and biotic and abiotic stress responses in *Brachypodium***. *J Exp Bot* 2015, **66**: 4317–4335.

[5] *Barros J, Serrani-Yarce JC, Chen F, Baxter D, Venables BJ, Dixon RA: **Role of bifunctional ammonia-lyase in grass cell wall biosynthesis**. *Nat Plants* 2016, **2**:16050

Demonstration, by downregulation in transgenic plants, that the brachypodium bifunctional ammonia-lyase (PTAL), provides nearly 50% of the total lignin deposited, with a preference for S-lignin and wall-bound *p*-coumarate biosynthesis. Isotope dilution experiments indicate that the L-phenylalanine and L-tyrosine pathways into lignin are distinct beyond the formation of *p*-coumarate, supporting the concept of metabolon organization during lignin synthesis.

[6] Vanholme R, Cesarino I, Rataj K, Xiao Y, Sundin L, Goeminne G, Kim H, Cross J, Morreel K, Araujo P, *et al.*: **Caffeoyl shikimate esterase (CSE) is an enzyme in the lignin biosynthetic pathway in *Arabidopsis***. *Science* 2013, **341**:1103-1106.

[7] Ha CM, Escamilla-Trevino L, Yarce JC, Kim H, Ralph J, Chen F, Dixon RA: **An essential role of caffeoyl shikimate esterase in monolignol biosynthesis in *Medicago truncatula***. *Plant J* 2016, **86**:363-375.

[8] Saleme MLS, Cesarino I, Vargas L, Kim H, Vanholme R, Goeminne G, Van Acker R, Fonseca FCA, Pallidis A, Voorend W, *et al.*: **Silencing *CAFFEOYL SHIKIMATE ESTERASE* affects lignification and Improves Saccharification in Poplar**. *Plant Physiol* 2017, **175**:1040-1057.

[9] del Río JC, Rencoret J, Prinsen P, Martínez ÁT, Ralph J, Gutiérrez A: **Structural characterization of wheat straw lignin as revealed by analytical pyrolysis, 2D-NMR, and reductive cleavage methods**. *J Agric Food Chem* 2012, **60**: 5922–5935.

[10] Lan W, Lu F, Regner M, Zhu Y, Rencoret J, Ralph SA, Zakai UI, Morreel K, Boerjan W, Ralph J: **Tricin, a flavonoid monomer in monocot lignification**. *Plant Physiol* 2015, **167**:1284–1295.

[11] Ralph J: **Hydroxycinnamates in lignification**. *Phytochem Rev* 2010, **9**:65-83.

- [12] Mechin V, Argillier O, Menanteau V, Barriere Y, Mila I, Pollet B, Lapierre C: **Relationship of cell wall composition to in vitro cell wall digestibility of maize inbred line stems.** *J Sci Food Agric* 2000, **80**:574-580.
- [13] van der Weijde T, Torres AF, Dolstra O, Dechesne A, Visser RGF, Trindade LM: **Impact of different lignin fractions on saccharification efficiency in diverse species of the bioenergy crop Miscanthus.** *Bioenerg Res* 2016, **9**:146-156.
- [14] Ostos Garrido FJ, Pistón F, Gómez LD, McQueen-Mason SJ: **Biomass recalcitrance in barley, wheat and triticale straw: Correlation of biomass quality with classic agronomical traits.** *PLoS ONE* 2018, **13**:e0205880.
- [15] Xu B, Escamilla-Trevino LL, Sathitsuksanoh N, Shen ZX, Shen H, Zhang YHP, Dixon RA, Zhao BY: **Silencing of 4-coumarate:coenzyme A ligase in switchgrass leads to reduced lignin content and improved fermentable sugar yields for biofuel production.** *New Phytol* 2011, **192**:611-625.
- [16] Gui JS, Shen JH, Li LG: **Functional characterization of evolutionarily divergent 4-coumarate: coenzyme A ligases in rice.** *Plant Physiol* 2011, **157**:574-586.
- [17] Jung JH, Kannan B, Dermawan H, Moxley GW, Altpeter F: **Precision breeding for RNAi suppression of a major 4-coumarate:coenzyme A ligase gene improves cell wall saccharification from field grown sugarcane.** *Plant Mol Biol* 2016, **92**:505-517.
- [18] Saballos A, Sattler SE, Sanchez E, Foster TP, Xin ZG, Kang C, Pedersen JF, Vermerris W: **Brown midrib2 (Bmr2) encodes the major 4-coumarate:coenzyme A ligase involved in lignin biosynthesis in sorghum (Sorghum bicolor (L.) Moench).** *Plant J* 2012, **70**:818-830.
- [19] Tamasloukht B, Lam MSJWQ, Martinez Y, Tozo K, Barbier, Jourda C, Jauneau A, Borderies G, Balzergue S, Renou JP, et al.: **Characterization of a cinnamoyl-CoA reductase 1 (CCR1) mutant in maize: effects on lignification, fibre development, and global gene expression.** *J Exp Bot* 2011, **62**:3837-3848.
- [20] Tu Y, Rochfort S, Liu ZQ, Ran YD, Griffith M, Badenhorst P, Louie GV, Bowman ME, Smith KF, Noel JP et al.: **Functional analyses of caffeic acid O-methyltransferase and cinnamoyl-CoA-reductase genes from perennial ryegrass (Lolium perenne).** *Plant Cell* 2010, **22**:3357-3373.
- [21] Smith RA, Cass CL, Mazaheri M, Sekhon RS, Heckwolf M, Kaeppler H, De Leon N, Mansfield SD, Kaeppler S, Sedbrook JC, et al.: **Suppression of**

CINNAMOYL-CoA REDUCTASE increases the level of monolignol ferulates incorporated into maize lignins. *Biotechnol Biofuel* 2017, **10**:109.

- [22] Park SH, Mei CS, Pauly M, Ong RG, Dale BE, Sabzikar R, Fotoh H, Nguyen T, Sticklen M: **Downregulation of maize cinnamoyl-coenzyme A reductase via RNA interference technology causes brown midrib and improves ammonia fiber expansion-pretreated conversion into fermentable sugars for biofuels.** *Crop Sci* 2012, **52**:2687–2701.
- [23] Halpin C, Holt K, Chojecki J, Oliver D, Chabbert B, Monties B, Edwards K, Barakate A, Foxon GA: **Brown-midrib maize (*bm1*) – a mutation affecting the cinnamyl alcohol dehydrogenase gene.** *Plant J* 1998, **14**:545-553.
- [24] Chen L, Auh CK, Dowling P, Bell J, Chen F, Hopkins A, Dixon RA, Wang ZY: **Improved forage digestibility of tall fescue (*Festuca arundinacea*) by transgenic down-regulation of cinnamyl alcohol dehydrogenase.** *Plant Biotechnol J* 2003, **1**:437-449
- [25] Trabucco GM, Matos DA, Lee SJ, Saathoff AJ, Priest HD, Mockler TC, Sarath G, Hazen SP: **Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid O-methyltransferase in *Brachypodium distachyon*.** *BMC Biotechnol* 2013, **13**:61.
- [26] d'Yvoire MB, Bouchabke-Coussa O, Voorend W, Antelme S, Cezard L, Legée F, Lebris P, Legay S, Whitehead C, McQueen-Mason SJ, *et al.*: **Disrupting the cinnamyl alcohol dehydrogenase 1 gene (*BdCAD1*) leads to altered lignification and improved saccharification in *Brachypodium distachyon*.** *Plant J* 2013, **73**:496-508.
- [27] Zhang KW, Qian Q, Huang ZJ, Wang YQ, Li M, Hong LL, Zeng DL, Gu MH, Chu CC, Cheng ZK: **GOLD HULL AND INTERNODE2 encodes a primarily multifunctional cinnamyl-alcohol dehydrogenase in rice.** *Plant Physiol* 2006, **140**:972-983.
- [28] Fornale S, Capellades M, Encina A, Wang K, Irar S, Lapierre C, Ruel K, Joseleau JP, Berenguer J, Puigdomenech P, *et al.*: **Altered lignin biosynthesis improves cellulosic bioethanol production in transgenic maize plants down-regulated for cinnamyl alcohol dehydrogenase.** *Mol Plant* 2012, **5**:817-830.
- [29] Barriere Y, Chavigneau H, Delaunay S, Courtial A, Bosio M, Lassagne H, Derory J, Lapierre C, Mechin V, Tatout C: **Different mutations in the *ZmCAD2* gene underlie the maize brown-midrib1 (*bm1*) phenotype with**

similar effects on lignin characteristics and have potential interest for bioenergy production. *Maydica* 2013, **58**:6-20.

- [30] Fu CX, Xiao XR, Xi YJ, Ge YX, Chen F, Bouton J, Dixon RA, Wang ZY: **Downregulation of cinnamyl alcohol dehydrogenase (CAD) leads to improved saccharification efficiency in switchgrass.** *Bioenergy Res* 2011, **4**:153-164.
- [31] Fornal S, Rencoret J, Garcia-Calvo L, Capellades M, Encina A, Santiago R, Rigau J, Gutierrez A, del Rio JC: **Cell wall modifications triggered by the down-regulation of Coumarate 3-hydroxylase-1 in maize.** *Plant Sci* 2015, **236**:272–282.
- [32] Takeda Y, Tobimatsu Y, Karlen SD, Koshiha T, Suzuki S, Yamamura M, Murakami S, Mukai M, Hattori T, Osakabe K, et al. **Downregulation of *p*-coumaroyl ester 3-hydroxylase in rice leads to altered cell wall structures and improves biomass saccharification.** *Plant J* 2018, **95**: 796-811.
- [33] Li XY, Chen WJ, Zhao Y, Xiang Y, Jiang HY, Zhu SW, Cheng BJ: **Downregulation of caffeoyl-CoA O-methyltransferase (CCoAOMT) by RNA interference leads to reduced lignin production in maize straw.** *Genet Mol Biol* 2013, **36**:540–546.
- [34] Takeda Y, Koshiha T, Tobimatsu Y, Suzuki S, Murakami S, Yamamura M, Rahman MM, Takano T, Hattori T, Sakamoto M *et al.*: **Regulation of CONIFERALDEHYDE 5-HYDROXYLASE expression to modulate cell wall lignin structure in rice.** *Planta* 2017, **246**:337-349.
- [35] Piquemal J, Chamayou S, Nadaud I, Beckert M, Barrière Y, Mila I, Lapierre C, Rigau J, Puigdomenech P, Jauneau A, *et al.*: **Down-regulation of caffeic acid O-methyltransferase in maize revisited using a transgenic approach.** *Plant Physiol* 2002, **130**:1675-1685.
- [36] Fu C, Mielenz JR, Xiao X, Ge Y, Hamilton CY, Rodriguez M Jr, Chen F, Foston M, Ragauskas A, Bouton J *et al.*: **Genetic manipulation of lignin reduces recalcitrance and improves ethanol production from switchgrass.** *Proc Natl Acad Sci* 2011, **108**:3803-3808.
- [37] Daly P, McClellan C, Maluk M, Oakey H, Lapierre C, Waugh R, Stephens J, Marshall D, Barakate A, Tsuji Y *et al.*: **RNAi-suppression of barley caffeic acid O-methyltransferase modifies lignin despite redundancy in the gene family.** *Plant Biotechnol J* 2018, <https://doi.org/10.1111/pbi.13001>.

- [38] Tu Y, Rochfort S, Liu Z, Ran Y, Griffith M, Badenhorst P, Louie GV, Bowman ME, Smith KF, Noel JP, *et al.*: **Functional analyses of caffeic acid O-methyltransferase and cinnamoyl-CoA-reductase genes from perennial ryegrass (*Lolium perenne*)**. *Plant Cell* 2010, **22**:3357-3373.
- [39] Pichon, M Deswartes, C Gerentes, D Guillaumie, S Lapierre, C Toppan, A Barriere, Y Goffner, D: **Variation in lignin and cell wall digestibility in caffeic acid O-methyltransferase down-regulated maize half-sib progenies in field experiments**. *Mol Breeding* 2006, **18**:253-261.
- [40] Jung JH, Vermerris W, Gallo M, Fedenko JR, Erickson JE, Altpeter F: **RNA interference suppression of lignin biosynthesis increases fermentable sugar yields for biofuel production from field-grown sugarcane**. *Plant Biotechnol J* 2013, **11**:709-716.
- [41] Baxter HL, Mazarei M, Labbe N, Kline LM, Cheng Q, Windham MT, Mann DG, Fu C, Ziebell A, Sykes RW *et al.*: **Two-year field analysis of reduced recalcitrance transgenic switchgrass**. *Plant Biotechnol J* 2014, **12**:914–924.
- [42] Kannan B, Jung JH, Moxley GW, Lee SM, Altpeter F: **TALEN-mediated targeted mutagenesis of more than 100 COMT copies/alleles in highly polyploid sugarcane improves saccharification efficiency without compromising biomass yield**. *Plant Biotechnol J* 2018, **16**:856-866.
- [43] Chen L, Auh CK, Dowling P, Bell J, Lehmann D, Wang ZY: **Transgenic down-regulation of caffeic acid O-methyltransferase (COMT) led to improved digestibility in tall fescue (*Festuca arundinacea*)**. *Funct Plant Biol* 2004, **31**:235-245.
- [44] Li M, Pu Y, Yoo CG, Gjersing E, Decker SR, Doeppke C, Shollenberger T, Tschaplinski TJ, Engle NL, Sykes RW *et al.*: **Study of traits and recalcitrance reduction of field-grown COMT down-regulated switchgrass**. *Biotechnol Biofuel* 2017, **10**:12.
- [45] Thorstensson EMG, Buxton DR, Cherney JH: **Apparent inhibition to digestion by lignin in normal and brown midrib stems**. *J Sci Food Agric* 1992, **59**:183-188.
- [46] Lan W, Lu F, Regner M, Zhu Y, Rencoret J, Ralph SA, Zakai UI, Morreel K, Boerjan W, Ralph J: **Tricin, a flavonoid monomer in monocot lignification**. *Plant Physiol* 2015, **167**:1284–1295.

- [47] *Lan W, Morreel K, Lu F, Rencoret J, del Río JC, Voorend W, Vermerris W, Boerjan WA, Ralph J: **Maize triclin-oligolignol metabolites and their implications for monocot lignification.** *Plant Physiol* 2016, **171**:810–820.

Identification of triclin-containing compounds in the metabolite pool of maize lignifying tissues helps establish triclin as an important monomer in grass lignification, and a nucleation site for polymer extension.

- [48] Eloy NB, Voorend W, Lan W, Saleme MDS, Cesarino I, Vanholme R, Smith RA, Goeminne G, Pallidis A, Morreel K et al.: **Silencing *CHALCONE SYNTHASE* in maize impedes the incorporation of triclin into lignin and increases lignin content.** *Plant Physiol* 2017, **173**:998–1016.
- [49] Eudes A, Dutta T, Deng K, Jacquet N, Sinha A, Benites VT, Baidoo EEK, Richel A, Sattler SE, Northen TR et al.: **SbCOMT (Bmr12) is involved in the biosynthesis of triclin-lignin in sorghum.** *PLoS One* 2017, **12**:e0178160.
- [50] Fornale S, Rencoret J, Garcia-Calvo L, Encina A, Rigau J, Gutierrez A, del Rio JC, Caparros-Ruiz D: **Changes in cell wall polymers and degradability in maize mutants lacking 3'- and 5'-O-methyltransferases involved in lignin biosynthesis.** *Plant Cell Physiol* 2017, **58**:240-255.
- [51] Lam PY, Tobimatsu Y, Takeda Y, Suzuki S, Yamamura M, Umezawa T, Lo C: **Disrupting *flavone synthase II* alters lignin and improves biomass digestibility.** *Plant Physiol* 2017, **174**:972–985.
- [52] Withers S, Lu FC, Kim H, Zhu YM, Ralph J, Wilkerson CG: **Identification of grass-specific enzyme that acylates monolignols with *p*-coumarate.** *J Biol Chem* 2012, **287**:8347–8355.
- [53] Petrik DL, Karlen SD, Cass CL, Padmakshan D, Lu FC, Liu S, Le Bris P, Antelme S, Santoro N, Wilkerson CG et al.: ***p*-Coumaroyl-CoA: monolignol transferase (PMT) acts specifically in the lignin biosynthetic pathway in *Brachypodium distachyon*.** *Plant J* 2014, **77**:713–726.
- [54] **Sibout R, Le Bris P, Legée F, Cezard L, Renault H, Lapierre C: **Structural redesigning Arabidopsis lignins into alkali-soluble lignins through the expression of *p*-coumaroyl-CoA: monolignol transferase PMT.** *Plant Physiol* 2016, **170**:1358–1366.

This paper demonstrates that a dicot, Arabidopsis, can be engineered to produce similar levels of *p*-coumaroylation on lignin to that found in grasses (8-9%), without any impact on plant development, by introducing either the BdPMT1 or BdPMT2

genes under the control of the Arabidopsis cinnamate-4-hydroxylase promoter. Lignin content was reduced and its structure was altered and saccharification efficiency was improved.

- [55] **Hatfield RD, Rancour DM, Marita JM: **Grass cell walls: a story of cross-linking**. *Front Plant Sci* 2017, **7**:2056.

Excellent recent review on the incorporation of ferulates and p-coumarates into grass secondary cell walls and the consequences for cell wall properties.

- [56] Terrett OM, Dupree P: **Covalent interactions between lignin and hemicelluloses in plant secondary cell walls**. *Curr Opin Biotechnol* 2019, **56**:97-104.

- [57] Piston F, Uauy C, Fu LH, Langston J, Labavitch J, Dubcovsky J: **Downregulation of four putative arabinoxylan feruloyl transferase genes from family PF02458 reduces ester-linked ferulate content in rice cell walls**. *Planta* 2010, **231**:677–691.

- [58] Buanafina MMD, Fescemyer HW, Sharma M, Shearer EA: **Functional testing of a PF02458 homologue of putative rice arabinoxylan feruloyl transferase genes in *Brachypodium distachyon***. *Planta* 2016, **243**:659–674.

- [59] **de Souza WR, Martins PK, Freeman J, Pellny TK, Michaelson LV, Sampaio BL, Vinecky F, Ribeiro AP, da Cunha BADB, Kobayashi AK et al.: **Suppression of a single BAHD gene in *Setaria viridis* causes large, stable decreases in cell wall feruloylation and increases biomass digestibility**. *New Phytol* 2018, **218**:81–93.


This paper provides the strongest evidence yet that feruloylation of arabinoxylan in grass cell walls is mediated by the BAHD01 gene and orthologues, and that it may be a suitable target for manipulating digestibility.


- [60] *Karlen SD, Peck ML, Zhang C, Smith RA, Padmakshan D, Helmich KE, Free HCA, Lee S, Smith BG, Lu F, et al.: **Monolignol ferulate conjugates are naturally incorporated into plant lignins**. *Sci Adv* 2016, **2**:1–9.

Demonstration that rice OsAT5 and other grass orthologues are likely BAHD feruloyl-coenzyme A monolignol transferases (FMTs) and a survey of the presence of monolignol ferulate in lignins among plant species.

Figure 1. Lignin biosynthesis in grasses. Most likely routes to intermediates are shown but do not preclude other routes. Only abbreviated enzyme names are used in the text. Red arrows denote pathways of specific importance in grasses. Blue enzyme/gene names denote membrane-bound P450 enzymes that may be in close proximity to each other.

PAL - phenylalanine ammonia lyase; PTAL - bifunctional phenylalanine:tyrosine ammonia lyase; C4H - cinnamate-4-hydroxylase; 4CL - 4-coumarate:CoA ligase; HCT - *p*-hydroxycinnamoyl-CoA:quininate/shikimate *p*-hydroxycinnamoyltransferase; C3H - *p*-coumarate 3-hydroxylase; CSE - caffeoyl shikimate esterase; CCoAOMT - caffeoyl-CoA O-methyltransferase; COMT - caffeate O-methyltransferase; CCR - cinnamoyl-CoA reductase; CAD - cinnamyl alcohol dehydrogenase; F5H - ferulate 5-hydroxylase; PAT - *p*-coumaroyl-CoA arabinoxylan transferase; PMT - *p*-coumaroyl-CoA monolignol transferase; FAT - feruloyl-CoA arabinoxylan transferase; FMT - feruloyl-CoA monolignol transferase; CHS - chalcone synthase; CHI - chalcone isomerase; FNS - flavone synthase; F3'H - naringenin 3-dioxygenase; C5'H - chrysoeriol 5-hydroxylase





Chew/Ferment/Digest

Sugar
Energy
Products

Chop/Pre-treat/Digest/Ferment

Biomass