



University of Dundee

X-ray spectral transitions of black holes from RXTE all-sky monitor

Gierliski, Marek; Newton, Jo

Published in:
Monthly Notices of the Royal Astronomical Society

DOI:
[10.1111/j.1365-2966.2006.10514.x](https://doi.org/10.1111/j.1365-2966.2006.10514.x)

Publication date:
2006

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
Gierliski, M., & Newton, J. (2006). X-ray spectral transitions of black holes from RXTE all-sky monitor. *Monthly Notices of the Royal Astronomical Society*, 370(2), 837-844. <https://doi.org/10.1111/j.1365-2966.2006.10514.x>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

X-ray spectral transitions of black holes from *RXTE* All-Sky Monitor

Marek Gierliński^{1,2*} and Jo Newton¹

¹*Department of Physics, University of Durham, South Road, Durham DH1 3LE*

²*Astronomical Observatory, Jagiellonian University, Orla 171, 30-244 Kraków, Poland*

Accepted 2006 April 30. Received 2006 April 4; in original form 2006 January 30

ABSTRACT

We have analysed X-ray outbursts from several Galactic black hole (GBH) transients, as seen by the All-Sky Monitor (ASM) on board *Rossi X-ray Timing Explorer (RXTE)*. We have used the best estimates of distance and black hole mass to find their luminosity (scaled to the Eddington limit), which allowed for direct comparison of many sources. We have found two distinct hard-to-soft state transitions in the initial part of the outburst. The distinction is made on the basis of the transition luminosity, its duration, the shape of the track in the hardness–luminosity diagram and evolution of the hardness ratio. The bright/slow transition occurs at ~ 30 per cent of Eddington (estimated bolometric) luminosity and takes $\gtrsim 30$ d, during which the source quickly reaches the intermediate/very high state and then proceeds to the soft state at much slower pace. The dark/slow transition is less luminous ($\lesssim 10$ per cent of Eddington), shorter ($\lesssim 15$ d) and the source does not slow its transition rate before reaching the soft state. We speculate that the distinction is due to irradiation and evaporation of the disc, which sustains the Comptonizing corona in the bright intermediate/very high state.

Key words: accretion, accretion discs – instabilities – X-rays: binaries.

1 INTRODUCTION

Black holes are remarkably simple objects, entirely characterized by their mass and spin, so we should expect similar appearance of their accretion flows. In Galactic binaries, black holes have comparable masses, so the accretion flow properties should be then determined by their spin and (instantaneous) accretion rate. It has been established recently that the accretion rate history plays an important role in shaping the accretion flow as well (e.g. van der Klis 2001; Maccarone & Coppi 2003). The inclination angle can also have some, but rather weak, effect on the observed properties, due to anisotropy of the disc or corona emission. Most of these effects can be easily accounted for: the mass by scaling the observed luminosity to Eddington (L_{Edd}) limit, the accretion history by tracing the entire outburst of a transient and the inclination angle effects are easy to understand within existing models of accretion. Perhaps the most challenging are the effects of black hole spin as they involve the properties of the accretion flow near and below the last stable orbit (e.g. Done & Gierliński 2006).

Galactic black holes (GBHs) are observed in several distinct X-ray spectral states (see Zdziarski & Gierliński 2004; McClintock & Remillard 2006, and references therein). This is best seen in X-ray transients undergoing occasional outbursts, when their accretion rate (and luminosity) can change by several orders of magnitude. With varying accretion rate, the X-ray spectrum can also change dramatically, passing through various states on time-scales

from days to months. These spectral transitions are probably the result of changes in the geometry of the accretion flow (for a review see Done & Gierliński 2004).

Typically, a transient begins its outburst in the low/hard (LH) state, where its spectrum can be roughly described by a hard (photon spectral index $\Gamma \sim 1.5\text{--}2$) power law with high-energy cut-off at ~ 100 keV (e.g. Wilson & Done 2001). The LH state can be seen at luminosities up to $\sim 0.2L_{\text{Edd}}$ during the rise of an outburst (Done & Gierliński 2003), but is less bright ($\lesssim 0.04L/L_{\text{Edd}}$) in the decaying part of the outburst (Maccarone 2003). A truncated disc, replaced by the optically thin, hot, Comptonizing inner flow can explain the hard state (e.g. Poutanen, Krolik & Ryde 1997; Esin et al. 2001), but other geometries have been also proposed (e.g. Beloborodov 1999; Young et al. 2001).

After the initial LH state, the X-ray spectrum usually undergoes transition into the high/soft (HS) state, via intermediate (IM) or very high (VH) state. The X-ray spectrum is then dominated by the disc emission of temperature of ~ 1 keV, accompanied by a soft ($\Gamma \sim 2\text{--}2.5$) power-law tail to higher energies. In the soft state, the cold disc probably extends down to or near the last stable orbit, and the hard component is produced by Comptonization in active regions or corona above the disc surface (e.g. Gierliński et al. 1999).

The IM/VH state is characterized by a soft X-ray spectrum dominated by a steep ($\Gamma \sim 2.5$) power law, extending to several hundred keV without apparent cut-off (Grove et al. 1998). Initially, it was recognized as the brightest state of a given object (Miyamoto et al. 1991), but HS states brighter than IM/VH state were seen (e.g. Zdziarski et al. 2001). IM/VH state has been observed in various sources at a wide range of luminosities of $\sim 0.01\text{--}1L_{\text{Edd}}$. Another

*E-mail: Marek.Gierlinski@durham.ac.uk

characteristic feature of IM/VH state is strong rapid aperiodic variability often accompanied by a distinct quasi-periodic oscillation.

The IM/VH state is typically related to the transitions between the hard and soft states (Rutledge et al. 1999). In many transients, it has been seen after the initial hard state, when the source reached high luminosity ($\gtrsim 0.1L_{\text{Edd}}$). This intermediate phase can last up to several dozen days before the source moves to the standard HS state. There are, however, instances of IM/VH state observed during ‘failed’ transitions or between two periods of the soft state (e.g. Homan et al. 2001; Kubota & Done 2004).

The X-ray spectral states and transitions of GBHs were analysed from the *Rossi X-ray Timing Explorer (RXTE)* spectral data by many authors (e.g. Sobczak et al. 2000; Homan et al. 2001; Done & Gierliński 2003; Fender, Belloni & Gallo 2004; Homan & Belloni 2005). A lot of work has been also done on the *RXTE* All-Sky Monitor (ASM) data (e.g. Grimm, Gilfanov & Sunyaev 2002; Zdziarski et al. 2002; Maccarone & Coppi 2003; Šimon 2004; Zdziarski et al. 2004; McClintock & Remillard 2006). However, few of these works attempted to discuss properties of spectral states in terms of absolute *luminosity*. In this paper, we present the first systematic study of black hole X-ray spectral states, based on hardness and luminosity derived from *RXTE* ASM data. We use the best known estimates of distance and mass of accreting black holes to derive their luminosity as a fraction of L_{Edd} . This allows us for direct comparison between different sources. We also trace *spectral evolution* during the outburst. Using these data, we show that there are two distinct types of transition between the hard and soft spectral states in the initial part of the outburst.

2 DATA REDUCTION

We have analysed publicly available light curves of eight GBH candidates from ASM (Levine et al. 1996) on board *RXTE* (Jahoda et al. 1996). These sources were selected because their data had statistics sufficient to build hardness–luminosity (H–L) diagrams. We used one-day averages in three ASM energy bands, roughly corresponding to 1.5–3, 3–5 and 5–12 keV. To allow for direct comparison between different sources, we converted the observed count rates into luminosities.

First, we calculated the energy flux in each ASM channel, adopting the method of Zdziarski et al. (2002), who built the ASM response matrix, comparing various pointed Cyg X-1 observations with ASM count rates. To check the validity of this method for the IM/VH spectral shape, which is the main topic of this paper, we have used the physical model fitted to the VH state *RXTE* spectrum of XTE J1550–564 (model HYB in table 4 in Gierliński & Done 2003). This model gave the 1.5–12 keV flux of $4.3 \times 10^{-8} \text{ erg s}^{-1} \text{ cm}^{-2}$, while the corresponding ASM data converted into flux resulted in $4.5 \times 10^{-8} \text{ erg s}^{-1} \text{ cm}^{-2}$, in excellent agreement with the pointed observation.

Then, we found the total 1.5–12 keV (isotropic) luminosity and expressed it as a fraction of Eddington luminosity [$L_{\text{Edd}} = 1.26 \times 10^{38} (M/M_{\odot}) \text{ erg s}^{-1}$] using the best known estimates of mass and distance, summarized in Table 1. We also calculated the hard flux (not count rate) hardness ratio (HR) of energy bands 5–12/3–5 keV. For clarity, we have discarded the data points with poor statistics (relative error on HR greater than 0.3) from H–L diagrams.

3 RESULTS

For our analysis, we have selected 10 outbursts of the transients listed in Table 1. In Figs 1 and 2, we present their light curves and

Table 1. The list of the sources used in this paper, together with assumed mass (M) and distance (D) estimates and their uncertainties. The numbers in square brackets refer to quantities that were not known in the time of writing of this paper and were assumed: mass of $10 M_{\odot}$ and distance of 8.5 kpc.

Source name	$M (M_{\odot})$	D (kpc)
4U 1543–47	9.4 (7.4–11.4) ^a	7.5 (7–8) ^a
XTE J1550–564	10 (9.7–11.6) ^b	5.3 (2.8–7.6) ^b
XTE J1650–500	4 ($\lesssim 7.3$) ^c	2.6 (1.9–3.3) ^d
GX 339–4	6 (2.5–10) ^e	8 (6.7–9.4) ^f
XTE J1739–278	[10]	8.5 (6–11) ^g
H1743–322	[10]	[8.5]
XTE J1859+226	[10]	7.6 (4.6–8) ^h
XTE J2012+381	[10]	[8.5]

^aPark et al. (2004); ^bOrosz et al. (2002); ^cOrosz et al. (2004); ^dHoman et al. (2006); ^eCowley et al. (2002); ^fZdziarski et al. (2004); ^gGreiner et al. (1996); ^hHynes et al. (2002).

H–L diagrams. In order to guide the eye and make comparison of various sources easier, we have copied a schematic shape of the H–L track of XTE J1550–564 1998 outbursts (shown in Fig. 1a) in all other H–L diagrams (the inverted Y-shaped curve).

Fig. 3 shows the evolution of the HR of these outbursts. The horizontal lines show positions of hard and soft branches. These lines were used to estimate the duration of the hard-to-soft transition, which was defined as the time between the last point on the hard branch and the first point on the soft branch. Certainly, this is—to some extent—arbitrary, since we have defined these branches by reading the diagrams. The soft branch (and hard branch, where possible) corresponds to HR which stabilized at a certain value. Where only one or a few points were available in the hard state, we used them as the hard state HR. The position of each branch varies slightly from source to source, but in most cases they are easy to identify from the diagrams. We will discuss possible drawbacks of this method in Section 4.

We have estimated the transition luminosity as L/L_{Edd} at hardness of HR = 1.25. We also found the fluence of each outburst, defined as the dimensionless L/L_{Edd} integrated over the entire outburst (so it has units of time). We summarize these results in Table 2.

These results suggest existence of two different types of hard-to-soft state transition. The bright/slow (BS) transition [Figs 1(a)–(d) and 3(a)–(d)] is characterized by the transition luminosity of $\sim 0.1L/L_{\text{Edd}}$ and duration of $\gtrsim 30$ d. In the H–L diagram, they trace a semicircular, anticlockwise track between the LH and HS branches. The best examples of this transition are shown by XTE J1550–564 (1998 outburst) and XTE J1859+226. The dark/fast (DF) transition [Figs 1(e)–(g) and 3(e)–(g)] is significantly dimmer ($\lesssim 0.05L/L_{\text{Edd}}$) and faster ($\lesssim 15$ d). In the H–L diagram, they move on an almost straight line between the LH and HS branches. The best examples of this transition are shown by XTE J1650–500 and XTE J2012+381. The last column in Table 2 shows the derived type of the transition. We discuss the details of this categorization in Sections 3.1 and 3.2.

3.1 Bright/slow transition

Fig. 1(a) shows the light curve and H–L diagram of the 1998 outburst of XTE J1550–564. Fig. 3(a) shows the evolution of its HR. This particular outburst had a very good *RXTE* coverage of pointed observations and its broad-band energy spectra were analysed in details (e.g. Sobczak et al. 2000; Gierliński & Done 2003; Kubota & Done 2004). It is known to cover all spectral states (e.g. Homan




Figure 1. Light curves and H–L diagrams of the outbursts of sources listed in Table 1, except for GX 339–4, which is shown in Fig. 2. The light curves are shown by light grey (cyan in colour) crosses. The thin black curve follows the transition from the hard to the soft state and joins the same data points in the light curve and H–L plot. The thick grey (green in colour) inverted Y-shaped line in the H–L diagrams illustrates the hard-to-soft transition of XTE J1550–564 and is used for comparison with other sources.

et al. 2001). In Figs 1(a) and 3(a), we connect the data points of the initial part (first 76 d) of the outburst. The outburst began in the LH state (Wilson & Done 2001) on the right-hand branch in the H–L diagram (at HR ~ 2.5). Then, as the luminosity increased, the spectrum softened fairly quickly and the source moved anticlockwise in the H–L diagram, reaching the top point corresponding to a bright flare in the light curve. This region in the H–L diagram corresponds to the IM/VH state (Gierliński & Done 2003; Kubota & Done 2004). Then, the luminosity decreased at much slower rate, and spectrum softened even more, while the source reached the HS (left-hand) branch in the H–L diagram at HR ~ 0.5 . The entire transition from leaving the LH branch until reaching the HS branch took 54 d.

Panels (a)–(d) of Figs 1 and 3 show the results from outbursts which we have categorized as the BS transition. They all begin in the LH branch of the H–L diagram, travel to the top ‘cusp’ at similar luminosity of $\sim 0.1L_{\text{Edd}}$ and then move on to the HS branch. XTE J1550–564 and XTE J1859+226 had a flare (brighter in the former

source) around the ‘cusp’. In terms of the HR evolution (Fig. 3), they seem to begin the outburst with a fairly fast transition to the IM/VH state at HR ~ 1.3 , after which the transition to the soft branch is significantly slower. This is illustrated in Fig. 4. The transition to the IM/VH state takes about 5 d, while the full transition from LH branch to HS branch takes more than ~ 30 d (see Table 2). These properties of the BS transition can be particularly clearly seen in XTE J1550–564 and XTE J1859+226.

GRS 1739–278 and H1743–322 have limited statistics, but show many resemblances to the XTE J1550–564 H–L track, transition luminosity and duration of the transition. Alas, the distance to H1743–322 is not known, so its luminosity was calculated for the assumed distance of 8.5 kpc and we cannot make any claims about its actual luminosity. Their HR evolution, shown in Figs 3(c) and (d), is similar to the two transitions shown in panels (a) and (b). Though errors on HR are substantially larger in GRS 1739–278 (Fig. 3c), its behaviour is entirely consistent with the BS transition. H1743–322 shows a complex HR curve with the hard-to-soft




Figure 2. Light curves and H–L diagrams of GX 339–4. Left-hand panels, (a) and (b), show two outburst assuming the mass and distance from Table 1, i.e. $6 M_{\odot}$ and 8 kpc, respectively. Right-hand panels, (c) and (d), show the same outbursts calculated for the mass of $10 M_{\odot}$ and distance of 6 kpc. The symbols and lines are the same as in Fig. 1.




Figure 3. Evolution of the HR during the initial part of the outburst for several sources. The black lines connect the same points as in Figs 1 and 2. The thick (green in colour) horizontal lines show our definition of the hard (upper line) and the soft (lower line) state for each source. The softest state of XTE J1550–564 achieved during the 2000 outburst (panel e) was harder than soft state in 1998 outburst (panel a), which is marked in panel (e) with a dashed line.

transition repeated (Fig. 3d) immediately after reaching the soft branch. The first transition lasts 25 d, the total transition $\gtrsim 80$ d.

The light curves of these outbursts can be complex in shape and generally long, extending over 200 d (with the exception of XTE J1859+226).

3.2 Dark/fast transition

Figs 1(e)–(g) and 3(e)–(g) show three outbursts with quite different behaviour. They do not follow the XTE J1550–564 H–L track but make a ‘shortcut’ between the LH and HS branches at a much lower luminosity of $\lesssim 0.05 L_{\text{Edd}}$. The transition time is much shorter, taking less than 16 d (Table 2). While the BS transients slow down their rate of transition after reaching the IM/VH state at $\text{HR} \sim 1.3$, the DF transients proceed to the soft branch at the same rate (see

Fig. 4). The entire outburst seems to be shorter than in BS state transients.

XTE J1650–500 makes the most clear example of this transition, moving between the LH and HS branches at the luminosity of only about $0.02 L_{\text{Edd}}$ in about 16 d. The 2000 outburst of XTE J1550–564 was very different from its 1998 outburst. The transition was much shorter (about 7 d) and occurred at much lower luminosity. However, while during the 1998 outburst the source reached $\text{HR} = 0.4$ in the soft state (marked by a dashed line in Fig. 3(e), the HR decreased only to 0.6 in 2000. Either it has not reached the ‘proper’ soft state at all, or the soft state was slightly different than in the 1998 outburst.

XTE J2012+381 is more problematic, as we do not know its distance, so the luminosity is calculated for the assumed distance of 8.5 kpc. However, its LH evolution is very similar to the two

Table 2. Transition luminosity (in Eddington luminosity), transition duration (in days) and outburst fluence (in days) for sources where transition was observed. The transition luminosity was read from the colour–luminosity diagram at $HR = 1.25$, corresponding to a ‘cusp’ in the XTE J1550–554 diagram. The luminosity range was given for the distance range in Table 1. The numbers in square brackets correspond to the assumed distance of 8.5 kpc. Superscripts in the first column denote different outbursts, referring to figure numbers. Asterisks mark the distance to GX 339–4 of 6 kpc and its mass of $10 M_{\odot}$, as opposed to the quantities listed in Table 1.

Source name	$L_{\text{trans}}/L_{\text{Edd}}$	t_{trans}	Fluence	Type
XTE J1550–564 ^{1a}	0.13 (0.04–0.28)	54	26	BS
XTE J1859+226	0.12 (0.05–0.14)	31	5.7	BS
XTE J1739–278	0.13 (0.06–0.21)	43	13	BS
H1743–322	[~0.17]	25 or ≥ 80	[17]	BS
XTE J1550–564 ^{1e}	0.03 (0.01–0.07)	7	1.2	DF
XTE J1650–500	0.02 (0.01–0.05)	16	1.3	DF
XTE J2012+381	[0.02]	4	[3.7]	DF
4U 1543–47	0.07 (0.06–0.08)	2	9	DF
GX 339–4 ^{2a,2c*}	0.15 (0.11–0.21) or 0.05*	9	49 or 16*	DF?
GX 339–4 ^{2b,2d*}	0.06 (0.04–0.08) or 0.02*	~8	21 or 7*	DF?




Figure 4. Comparison of the HR evolution of all transitions shown in Fig. 3 (the second transition of H1743–322, after MJD 52762, was omitted for clarity), zeroed at the beginning of the hard-to-soft transition. Solid curves represent BS transitions and dotted curves represent DF transitions. This figure illustrates the crucial difference between the two types of transitions. After the initial fast phase, the BS transition is much slower after reaching $HR \sim 1.3$, while DF transients continue to the soft branch at the same rate.

other DF outbursts, both in track shape and speed of movement. Therefore, we classify this source as a DF transient.

Figs 1(h) and 3(h) show the outburst of 4U 1543–47. It made a fast transition from the LH state to the unusually bright HS state, which might be due to its low inclination angle (see discussion in Section 4). However, the luminosity, duration of the transition and the evolution of HR allow us to classify this outburst as a DF transition.

Fig. 2 shows GX 339–4. In panel (a), we show the 2002/2003 outburst based on the most recent distance estimate of 8 kpc (Zdziarski et al. 2004). The light curve is long and complex, resembling BS transients. On the other hand, the shape of the H–L diagram, the HR evolution (Fig. 3i) and the fast transition time (9 d) strongly suggest the DF state transition. However, the transition luminosity $>0.1 L_{\text{Edd}}$ is inconsistent with DF transitions. Additionally, both LH and HS branches are unusually bright, not consistent with any other source in our sample, except perhaps 4U 1543–47 (Fig. 1h).

Another (2004/2005) GX 339–4 outburst, shown in Fig. 2(b), is less luminous and perhaps consistent with DF transient behaviour.

In order to explain the unusual 2002/2003 outburst of GX339–4, we consider the possibility whether it can be still consistent with other DF transients within uncertainties of its mass and distance. Hynes et al. (2004) gave a fairly robust lower limit on the kinematic distance of 6 kpc. Cowley et al. (2002) gave the upper limit on the mass of $10 M_{\odot}$. We use these two extremes to create a new plot, shown in Fig. 2(c) and (d). Clearly, both outbursts now seem to be consistent both in the track shape and transition luminosity with other DF transients. Though the entire light curve is more similar to other BS transients, the *transition itself* is short and marginally consistent with typical DF luminosity, so we tentatively classify GX 339–4 as a DF transient.

4 CAVEATS

In Section 3, we postulated existence of two distinct hard-to-soft spectral transitions in the initial part of the outburst of black hole transients. Our classification is based on the transition luminosity, duration, the shape of the track in the H–L diagram and the evolution of the HR. Plainly, there are several uncertainties here. First of all, the luminosity strongly depends on highly uncertain distance and (to less extent) black hole mass. In Table 2, we show the range of luminosities due to the distance uncertainty. The uncertainties are large indeed, and there is an overlap between the two categories around $\sim 0.05 L_{\text{Edd}}$.

Another uncertainty comes from the interstellar absorption, affecting the derived luminosity and HR. Fortunately, most of sources analysed in this paper have moderate absorption column of $N_{\text{H}} \lesssim 10^{22} \text{ cm}^{-2}$. XTE J1739–278 and H1743–322 have higher columns of about $2 \times 10^{22} \text{ cm}^{-2}$ (Greiner, Dennerl & Predehl 1996; Capitanio et al. 2005) which, depending on the shape of the spectrum, can diminish the 1.5–12 keV flux down to ~ 0.7 of the unabsorbed flux. The effect of this column on 3–12 keV hardness is negligible.

The luminosities quoted in this paper are calculated in the 1.5–12 keV energy band, but the bolometric luminosity is definitely larger. To estimate the bolometric correction in the IM/VH state, we used the same physical model of XTE J1550–564 as in Section 2. It gives the bolometric luminosity 2.3 times larger than the 1.5–12 keV luminosity. Since the IM/VH spectra are similar in shape, this seems to be a good estimate of the bolometric correction factor. Thus, we estimate that the bolometric luminosity is ~ 30 per cent of L_{Edd} during the BS transition and $\lesssim 10$ per cent of L_{Edd} during the DF transition (though 4U 1543–47 transition is brighter, ~ 16 per cent of L_{Edd} , which might be due to its lower inclination angle, see next paragraph). We have performed similar tests for hard-state (observation id. 30188-06-01-00) and soft-state (30191-01-43-00) data and found bolometric corrections of 2.8 and 2.5, respectively.

The inclination angle of the accretion disc can play a significant role in the disc-dominated HS state. The emission from optically thin Comptonization, which dominates the LH state, is probably more isotropic, though it might be mildly boosted at low angles if it originates from a jet/outflow. Inclinations of the sources in our sample range (where known) between 21° for 4U 1543–47 (Park et al. 2004) and 72° for XTE J1550–564 (Orosz et al. 2002). We expect the apparent disc in the former source to be brighter by factor $\cos 21^\circ / \cos 72^\circ \approx 3$ than the latter one. This might explain why the HS branch of 4U 1543–47 is about three times brighter than in XTE J1550–564. The relatively low inclination of $(40 \pm 20)^\circ$ (Cowley

et al. 2002) cannot account for the unusually high luminosity of GX 339–4, as it is also very bright on the LH branch.

Opportunely, all the caveats regarding distance, absorption and inclination angle can be dropped in the case of two different outbursts observed from *the same* source, XTE J1550–564. It displayed the BS transition in 1998 (Fig. 1a) and the DF transition in 2000 (Fig. 1e). Each outburst showed typical characteristics derived for each transition category, which significantly strengthens our result.

One possible uncertainty comes from the identification of the soft- and hard-state branches, which can subsequently affect measurements of the transition duration. Usually, the spectral state of an X-ray source is established on the basis of its spectral *and* variability properties. In this work, we define the hard and soft state branches on the sole basis of their HR, which might not necessary agree with the full spectral/timing definition. In particular, ASM data show that XTE J1650–500 reached the HS branch at HR = 0.35 around MJD 52176. We have calculated the same HR from available PCA data and found it to be in perfect agreement with the ASM data. However, after the transition, around MJD 52190 (see Fig. 4f), where the ASM data are sparse and of poor statistics, PCA shows a further decrease in the HR to about HR = 0.15. Rossi et al. (2004) classified the state at HR = 0.35 as the VH state, but they also noted that its properties were quite different from the IM/VH state during the transition. A similar situation is in the case of the 2000 outburst of XTE J1550–564, which never reached the soft state similar to that in 1998. It is then possible, that our HS branch derived from the ASM data contains softer IM/VH states, as defined from the spectral and variability properties. Thus, the transition time from the hard state to the *true* soft state can be longer than the transition between the ASM LH and HS branches, derived in this paper.

This, however, does not seem to affect our main result about the difference between the BS and DF transitions. The transition between the LH and HS branches, *as derived from ASM data*, is markedly faster and less luminous in DF transients. Even if our HS branch does not exactly correspond to the true soft state, there is still a *systematic* difference between those transitions, as illustrated in Fig. 4.

We would like to stress that, despite all the caveats discussed in this section, there are two criteria distinguishing between the DF and BS transitions, which are fairly robust. These are the duration of transition between LH and HS branches (or evolution of the HR) and the shape of the LH track. Though in some cases the limited statistics makes recognizing the track rather difficult, we find that when all the clues are taken together, there is a significant, systematic difference between the two categories of transitions. Therefore, we find our classification robust and well defined.

5 DISCUSSION AND CONCLUSIONS

We have studied light curves and H–L diagrams of several GBH transients. We have found two distinct categories of spectral transitions occurring after the initial hard state during the rise of the outburst and before the disc-dominated HS state. The BS transition reaches the estimated bolometric luminosity of $0.3 L_{\text{Edd}}$ or more. The initial transition from the LH state is fast, but it is significantly slowed down after reaching the bright IM/VH state. The entire transition to the HS state takes over 30 d. The LH and HS branches in the H–L track are connected by a roughly semicircular, counterclockwise track. The DF transition typically takes place at luminosities $\lesssim 0.1 L_{\text{Edd}}$, it is faster (less than ~ 15 d), does not slow down at the IM/VH state and makes a ‘shortcut’ between the LH and HS




Figure 5. Schematic diagram of two types (BS and DF) of hard-to-soft transitions. Left-hand panel: H–L diagram. Right-hand panel: evolution of the HR.

branches in the H–L diagram. The difference between these two categories is shown in Fig. 5.

The distinction between the two categories seems to be clear, and can be particularly well seen in Fig. 4. However, we only have four transitions in each category, so from purely statistical point of view the bimodal behaviour is not yet significant. Plainly, more data are necessary to firmly confirm our discovery.

The terms ‘very high’ and ‘intermediate’ state have been used in the literature (e.g. Miyamoto et al. 1991; Homan et al. 2001), and sometimes the distinction between the higher and lower luminosities had been made (e.g. Méndez & van der Klis 1997). Later, this distinction was abandoned, when it became clear that the IM/VH states can occur at various luminosities, often less than that in the disc-dominated HS state. Instead, the definition based on both spectral and timing properties was proposed (e.g. McClintock & Remillard 2006). The energy and power spectra of these states are similar, regardless of luminosity (e.g. Homan et al. 2001; Gierliński & Done 2003; Kalemci et al. 2003; Montanari, Frontera & Amati 2004). No systematic distinction between bright and dark IM/VH spectral states has been found so far. In this paper, we have shown that the hard-to-soft state transition can take place at two different luminosities, and that there is a systematic difference between the low- and high-luminosity transitions. If physical properties of the accretion flow are different between the DF and BS transitions, then we might expect some difference in the spectral state as well. If this were true, then perhaps the distinction between the IM and VH states might be valid, after all. Confirming this potential difference would require additional studies.

There are certain limitations to the BS/DF distinction we have proposed in this paper. The estimated BS transition bolometric luminosity of $\sim 0.3 L_{\text{Edd}}$, observed in the four sources shown in Figs 1(a)–(d), is not entirely typical, and can be easily surpassed. Some of the transients reached and exceeded the Eddington luminosity during their outbursts, e.g. V404 Cyg (e.g. Życki, Done & Smith 1999), Nova Mus (Życki et al. 1999) or V4641 Sgr (Revnivtsev et al. 2002). GRS 1915+105 is also known to be very bright (Done, Wardziński & Gierliński 2004), but it is a rather untypical transient (Truss & Done 2006). It is interesting to note that GX 339–4 made two transitions at significantly different luminosity, but with time-scales and HR evolution typical for DF transitions. Thus, the DF transition luminosity is not constant and the transition can occur at a range of luminosities below a certain critical value. The opposite, i.e. a range of luminosities above a critical limit, could be true for the BS transitions. It is difficult to ascertain using currently available data, whether these two critical luminosities are equal or not.

We have seen both types of transitions from the same source, XTE J1550–564. This obviously means that the outburst type

is not predestined by the properties of the binary, such as the disc size, the orbital period or mass/spin of the black hole. The distinguishing factor must be hidden in the accretion flow itself. The numerical model of the disc instability by Dubus, Hameury & Lasota (2001) shows that two types of outbursts can develop in the accretion disc, depending on the radius at which the hydrogen ionization instability is triggered. When the ignition radius is small (the ‘inside-out’ outburst) the propagation time of the inward heating front is short, while the outward front can easily stall in the regions of higher density, which leads to short, low-amplitude outbursts. When the ignition radius is large (the ‘outside-in’) outburst, the inward front progresses through regions of decreasing density and can heat the whole disc, leading to longer, brighter outbursts.

This, in principle, could explain the difference between bright and dark transitions/outbursts. However, we can see from the light curves that the length of the outburst is not simply correlated with its amplitude. Though generally BS outbursts have longer and more complex light curves than DF outbursts, there are a few exceptions. For example, XTE J1859+226 (Fig. 1b) and XTE J2012+381 (Fig. 1g), have light curves very similar in shape and duration, despite undergoing different types of LH–HS transition. GX 339–4, tentatively categorized as a DF transient, displays very long and complex light curves.

Perhaps a better quantity that could distinguish between ‘inside-out’ and ‘outside-in’ outbursts is the fluence, which is a rough estimate of the total amount of the accreted material. We have estimated the fluence of each outburst (Table 2). BS outbursts tend to have larger fluence, though the distinction is not unique. For example, the BS outburst of XTE J1748–288 has smaller fluence than the DF outburst of XTE J2012+381, though distances and BH masses of both sources are not known, so this estimate is very uncertain. And again, GX 339–4 does not easily fit into the picture with its large fluence, even assuming extreme black hole mass and distance, as in Section 4. Clearly, the radius at which the initial instability occurs can be an important factor, but does not entirely explain the observed properties of spectral transitions.

Thus, it seems that the total amount of accreted material in an outburst does not predestine the type of the LH–HS transition. The clue to the distinction between the DF and BS transitions might lay in the irradiation of the disc by the central source. Irradiation can lengthen the bright state of the disc (King & Ritter 1998). At the same time, it can increase evaporation rate (Róžańska & Czerny 2000; Dullemond & Spruit 2005; Meyer-Hofmeister, Liu & Meyer 2005), which in turn can help sustaining the bright corona providing the source of irradiation. An interesting feature of the observed outbursts is that the initial transition to the IM/VH state takes roughly the same time, irrespectively of the luminosity, and only after that transitions diversify into slow and fast, for high and low luminosities, respectively. We speculate, that the luminosity in the IM/VH state defines the type of the LH–HS transition. When it reaches $\gtrsim 0.3 L_{\text{Edd}}$, it triggers the self-sustaining bright corona, and prolongs the bright BS transition. The dimmer and much shorter DF transition does not reach the luminosity required to lengthen the bright state and sustain the corona, so it disappears and the source quickly moves on to the disc-dominated HS state.

ACKNOWLEDGMENTS

We thank the anonymous referee for their helpful comments and Chris Done for stimulating discussions. MG acknowledges support through a PPARC PDRF.

REFERENCES

- Beloborodov A. M., 1999, *ApJ*, 510, L123
 Capitanio F. et al., 2005, *ApJ*, 622, 503
 Cowley A. P., Schmidtke P. C., Hutchings J. B., Crampton D., 2002, *AJ*, 123, 1741
 Done C., Gierliński M., 2003, *MNRAS*, 342, 1041
 Done C., Gierliński M., 2004, *Prog. Theor. Phys. Suppl.*, 155, 9
 Done C., Gierliński M., 2006, *MNRAS*, 367, 659
 Done C., Wardziński G., Gierliński M., 2004, *MNRAS*, 349, 393
 Dubus G., Hameury J.-M., Lasota J.-P., 2001, *A&A*, 373, 251
 Dullemond C. P., Spruit H. C., 2005, *A&A*, 434, 415
 Esin A. A., McClintock J. E., Drake J. J., Garcia M. R., Haswell C. A., Hynes R. I., Munro M. P., 2001, *ApJ*, 555, 483
 Fender R. P., Belloni T. M., Gallo E., 2004, *MNRAS*, 355, 1105
 Gierliński M., Done C., 2003, *MNRAS*, 342, 1083
 Gierliński M., Zdziarski A. A., Poutanen J., Coppi P. S., Ebisawa K., Johnson W. N., 1999, *MNRAS*, 309, 496
 Greiner J., Dennerl K., Predehl P., 1996, *A&A*, 314, L21
 Grimm H.-J., Gilfanov M., Sunyaev R., 2002, *A&A*, 391, 923
 Grove J. E., Johnson W. N., Kroeger R. A., McNaron-Brown K., Skibo J. G., Philips B. F., 1998, *ApJ*, 500, 899
 Homan J., Belloni T., 2005, *Ap&SS*, 300, 107
 Homan J., Wijnands R., van der Klis M., Belloni T., van Paradijs J., Klein-Wolt M., Fender R., Méndez M., 2001, *ApJS*, 132, 377
 Homan J., Wijnands R., Kong A., Miller J. M., Rossi S., Belloni T., Lewin W. H. G., 2006, *MNRAS*, 366, 235
 Hynes R. I., Haswell C. A., Chaty S., Shrader C. R., Cui W., 2002, *MNRAS*, 331, 169
 Hynes R. I., Steeghs D., Casares J., Charles P. A., O’Brien K., 2004, *ApJ*, 609, 317
 Jahoda K., Swank J. H., Giles A. B., Stark M. J., Strohmayer T., Zhang W., Morgan E. H., 1996, *SPIE*, 2808, 59
 Kalemci E., Tomsick J. A., Rothschild R. E., Pottschmidt K., Corbel S., Wijnands R., Miller J. M., Kaaret P., 2003, *ApJ*, 586, 419
 King A. R., Ritter H., 1998, *MNRAS*, 293, L42
 Kubota A., Done C., 2004, *MNRAS*, 353, 980
 Levine A. M., Bradt H., Cui W., Jernigan J. G., Morgan E. H., Remillard R., Shirey R. E., Smith D. A., 1996, *ApJ*, 469, L33
 Maccarone T. J., 2003, *A&A*, 409, 697
 Maccarone T. J., Coppi P. S., 2003, *MNRAS*, 338, 189
 McClintock J. E., Remillard R., 2006, in Lewin W. H. G., van der Klis M., eds, *Compact Stellar X-ray Sources*. Cambridge Univ. Press, Cambridge
 Méndez M., van der Klis M., 1997, *ApJ*, 479, 926
 Meyer-Hofmeister E., Liu B. F., Meyer F., 2005, *A&A*, 432, 181
 Miyamoto S., Kimura K., Kitamoto S., Dotani T., Ebisawa K., 1991, *ApJ*, 383, 784
 Montanari E., Frontera F., Amati L., 2004, *Nucl. Phys. B*, 132, 412
 Orosz J. A. et al., 2002, *ApJ*, 568, 845
 Orosz J. A., McClintock J. E., Remillard R. A., Corbel S., 2004, *ApJ*, 616, 376
 Park S. Q. et al., 2004, *ApJ*, 610, 378
 Poutanen J., Krolik J. H., Ryde F., 1997, *MNRAS*, 292, L21
 Revnivtsev M., Gilfanov M., Churazov E., Sunyaev R., 2002, *A&A*, 391, 1013
 Rossi S., Homan J., Miller J. M., Belloni T., 2004, *Nucl. Phys. B*, 132, 416
 Róžańska A., Czerny B., 2000, *A&A*, 360, 1170
 Rutledge R. E. et al., 1999, *ApJS*, 124, 265
 Šimon V., 2004, *A&A*, 418, 617
 Sobczak G. J., McClintock J. E., Remillard R. A., Cui W., Levine A. M., Morgan E. H., Orosz J. A., Bailyn C. D., 2000, *ApJ*, 544, 993
 Truss M., Done C., 2006, *MNRAS*, 368, L25
 van der Klis M., 2001, *ApJ*, 561, 943
 Wilson C. D., Done C., 2001, *MNRAS*, 325, 167
 Young A. J., Fabian A. C., Ross R. R., Tanaka Y., 2001, *MNRAS*, 325, 1045

- Zdziarski A. A., Gierliński M., 2004, *Prog. Theor. Phys. Suppl.*, 155, 99
Zdziarski A. A., Grove J. E., Poutanen J., Rao A. R., Vadawale S. V., 2001, *ApJ*, 554, L45
Zdziarski A. A., Poutanen J., Pacias W. S., Wen L., 2002, *ApJ*, 578, 357

- Zdziarski A. A., Gierliński M., Mikołajewska J., Wardziński G., Smith D. M., Alan Harmon B., Kitamoto S., 2004, *MNRAS*, 351, 791
Życki P. T., Done C., Smith D. A., 1999, *MNRAS*, 309, 561

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.