

This is a repository copy of *Genetic modification of western wheatgrass (Pascopyrum smithii) for the phytoremediation of RDX and TNT*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/138740/>

Version: Accepted Version

---

**Article:**

Zhang, Long, Rylott, Elizabeth Lucy [orcid.org/0000-0002-1609-414X](https://orcid.org/0000-0002-1609-414X), Bruce, Neil Charles [orcid.org/0000-0003-0398-2997](https://orcid.org/0000-0003-0398-2997) et al. (1 more author) (2019) Genetic modification of western wheatgrass (*Pascopyrum smithii*) for the phytoremediation of RDX and TNT. *Planta*. pp. 1007-1015. ISSN 0032-0935

<https://doi.org/10.1007/s00425-018-3057-9>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

1 Genetic modification of western wheatgrass (*Pascopyrum smithii*) for the  
2 phytoremediation of RDX and TNT

3 Long Zhang,<sup>1</sup> Elizabeth L. Rylott,<sup>2</sup> Neil C. Bruce,<sup>2</sup> Stuart E. Strand<sup>1\*</sup>

4 <sup>1</sup>Department of Civil and Environmental Engineering, University of Washington, Box 355014, Seattle,  
5 WA 98195-5014, USA;

6 <sup>2</sup>Centre for Novel Agricultural Products, Department of Biology, University of York, Wentworth Way,  
7 York YO10 5DD, UK.

8

9

10

11 **Word count:** 3899, 4366 including captions

12

13 **Main Conclusion:** Transgenic western wheatgrass degrades the explosive RDX and detoxifies TNT

14 **ABSTRACT**

15 Contamination, from the explosives, hexahydro-1, 3, 5-trinitro-1, 3, 5-triazine (RDX) and 2, 4, 6-  
16 trinitrotoluene (TNT), especially on live-fire training ranges, threatens environmental and human health.  
17 Phytoremediation is an approach that could be used to clean-up explosive pollution, but it is hindered by  
18 inherently low *in planta* RDX degradation rates, and the high phytotoxicity of TNT. The bacterial genes,  
19 *xplA*, and *xplB*, confer the ability to degrade RDX in plants, and a bacterial nitroreductase gene *nfsI*,  
20 enhances the capacity of plants to withstand and detoxify TNT. While previous studies have used model  
21 plant species to demonstrate the efficacy of this technology, trials using plant species able to thrive in the  
22 challenging environments found on military training ranges are now urgently needed. Perennial western  
23 wheatgrass (*Pascopyrum smithii*) is a United States native species that is broadly distributed across North  
24 America, well-suited for phytoremediation, and used by the US military to re-vegetate military ranges.  
25 Here we present the first report of the genetic transformation of western wheatgrass. Plant lines  
26 transformed with *xplA*, *xplB* and *nfsI* removed significantly more RDX from hydroponic solutions and  
27 retained much lower, or undetectable, levels of RDX in their leaf tissues when compared to wild-type  
28 plants. Furthermore, these plants were also more resistant to TNT toxicity, and detoxified more TNT than  
29 wild-type plants. This is the first study to engineer a field-applicable grass species capable of both RDX  
30 degradation and TNT detoxification. Together, these findings present a promising biotechnological  
31 approach to sustainably contain, and remove, RDX and TNT from training range soil and prevent  
32 groundwater contamination.

33 **Keywords:** phytoremediation; RDX; TNT; western wheatgrass; transformation, monocot promoters;  
34 stacked genes.

## 35 INTRODUCTION

36 The explosives hexahydro-1, 3, 5-trinitro-1, 3, 5-triazine (RDX) and 2,4,6-trinitrotoluene (TNT) are used  
37 extensively by the US military. These compounds are damaging to the environment and human health:  
38 RDX targets the nervous system and can cause seizures in humans and animals (Ramasahayam et al.  
39 2017) while TNT is considered to be highly toxic and mutagenic (Bolt et al. 2006; Honeycutt et al. 1996;  
40 Travis et al. 2008). The US Environmental Protection Agency has classified both RDX and TNT as  
41 possible human carcinogens (USEPA, 2014), and during explosives manufacture, transportation, storage  
42 and disposal, these toxic pollutants are released into the environment. Additionally, live-fire training at  
43 military bases has resulted in the contamination of soils around targets with particulates of munitions  
44 compounds, which have leached RDX into the underlying groundwater. The total area of operational  
45 ranges in the United States contaminated with munitions constituents is estimated to be more than 16  
46 million acres (United States General Accounting, 2004), with clean-up of active ranges estimated by the  
47 US Department of Defense to cost between US\$16 billion and US\$165 billion (United States General  
48 Accounting, 2004). Chemical-based methods to remediate explosives, such as alkaline hydrolysis  
49 (Sviatenko et al. 2017), redox reactions using iron-bearing materials (Oh et al. 2016), and permeable  
50 reaction barriers (Ahmad et al. 2007) have proven too cost-intensive for the scale of the contaminated  
51 areas. Phytoremediation of RDX and TNT has been developed as a potentially more cost-efficient, and  
52 sustainable method to clean up contaminated soil.

53 Plants readily take up and translocate RDX from the soil to the aerial tissues (Vila et al. 2007; Brentner et  
54 al. 2010), with studies of poplar (*Populus deltoides* x *nigra* DN-34) showing that RDX can be reduced  
55 to hexahydro-1-nitroso-3,5-dinitro-1,3,5-triazine (MNX) and hexahydro-1,3-dinitroso-5-nitro-1,3,5-  
56 triazine (DNX) (Van Aken et al. 2004). However, despite high uptake rates, plants have relatively low  
57 rates of RDX degradation (Mezzari et al. 2004; Best et al. 1999; Winfield 2004) with RDX accumulated  
58 in living plant tissues returning to the environment following plant senescence (Yoon et al. 2006).

59 In contrast to RDX, TNT remains predominantly in root tissues with less than 25 % translocated to aerial  
60 tissues (Vila et al. 2007; Brentner et al. 2010). TNT is also more phytotoxic (Pavlostathis and Jackson  
61 2002; Hannink et al. 2007; Johnston et al. 2015) than RDX, and symptoms of toxicity such as stunted  
62 growth and chlorosis have been observed at concentrations comparable to those found on military training  
63 ranges (Pichtel 2012). Plants have a limited ability to detoxify TNT. Endogenous nitroreductases,  
64 including oxophytodienoate reductases (Beynon et al. 2009) reduce TNT to hydroxyl-amino and then  
65 amino-dinitrotoluenes (HADNTs and ADNTs). These transformed intermediates are then conjugated to  
66 hydrophilic molecules such as sugars (Gandia-Herrero et al. 2008) and glutathione (Gunning et al. 2014),  
67 then appear to be incorporated into cell wall materials (Rylott and Bruce 2009).

68 While plants are limited in their ability to remediate these explosives, bacteria able to degrade RDX and  
69 detoxify TNT have been isolated from explosives-contaminated soils, and the genes responsible  
70 characterized: a fused flavodoxin-cytochrome P450 XplA and associated flavodoxin reductase XplB from  
71 *Rhodococcus rhodococcus* strain 11Y confer aerobic biodegradation of RDX (Seth-Smith et al. 2002) and  
72 from *Enterobacter cloacae*, *nfsI* encodes a nitroreductase (NR), which transforms TNT, favoring the  
73 production of the 4-ADNT isomer (Hannink et al. 2001). Numerous species of bacteria containing *xplA*  
74 and *xplB* have since been isolated from RDX-contaminated locations around the world (Seth-Smith et al.  
75 2008; Rylott et al. 2011b; Andeer et al. 2009; Andeer et al. 2013; Bernstein et al. 2011; Thompson et al.  
76 2005; Binks et al. 1995; Coleman 1998; Adrian and Arnett 2004). Yet despite the presence of RDX-  
77 degrading bacteria, RDX contamination in these regions persists, suggesting that other limitations are  
78 preventing bacterial degradation of RDX.

79 To increase the ability of plants to degrade RDX, *xplA* and *xplB* have been transformed into a range of  
80 plant species. The co-pollutant TNT is phytotoxic, and also inhibits XplA activity (Jackson et al. 2007).  
81 To overcome these hindering factors, the TNT detoxifying nitroreductase has been expressed in tobacco  
82 (*Nicotiana tabaccum*; Hannink et al. 2001), the RDX-metabolizing *xplA* and *xplB* in *Arabidopsis*  
83 (*Arabidopsis thaliana*; Rylott et al. 2006); and all three genes have since been simultaneously expressed

84 in *Arabidopsis* (Rylott et al. 2011a), and creeping bentgrass (*Agrostis stolonifera*; Zhang et al. 2017a).  
85 These transgenic plants were all able to degrade RDX and resist high levels of TNT when compared to  
86 wild-type plants. In our recent report, *nfsI* transformed into the plastid genome of tobacco conferred  
87 enhanced resistance to TNT (Zhang et al. 2017b). However, *Arabidopsis*, tobacco, and creeping bentgrass  
88 are not well-adapted to thriving in the challenging environments found in military training ranges.  
89 Switchgrass (*Panicum virgatum*), is a United States native species, with additional attributes that make it  
90 suitable for phytoremediation of explosives on military training ranges, and recently plants have been  
91 transformed with *xplA*, *xplB* and *nsfI* and shown to remediate RDX (Zhang et al. 2017a). However, since  
92 the range of the lowland switchgrass species is limited, and planting a single species in the field is likely  
93 to result in vulnerability to pests, diseases and other environmental stresses; additional grass species are  
94 needed. Western wheatgrass (*Pascopyrum smithii*) is a long-lived, cool season, United States native  
95 species that is broadly distributed across North America, with characteristics that make it a desirable plant  
96 for the phytoremediation of RDX and TNT on live-fire training ranges. Its low maintenance requirements  
97 and vigorously spreading rhizomes give it the ability to recover quickly after damage caused by military  
98 vehicle traffic, explosive fire or the range fires that occur on training ranges (Palazzo et al. 2005).  
99 Furthermore, cultivars, which have improved performance over existing commercially available cultivars,  
100 have been developed by the US military specifically for the re-vegetation of ranges including western  
101 wheatgrass (Palazzo et al. 2005). However, until now wheatgrass species have proven recalcitrant to  
102 transformation. This study uses *xplA*, *xplB*, and *nfsI* to demonstrate an effective transformation protocol,  
103 and validate the ability of this species to remediate RDX and TNT.

## 104 **MATERIALS AND METHODS**

### 105 **Plasmid construction**

106 The binary vector pEDLZ2014 (Figure 1) was constructed using a protocol similar to that used to  
107 transform switchgrass, prcs2-NABNR (Zhang et al. 2017a) with the following modifications. The *Osact*  
108 sequence was amplified from pANIC 5A (Mann et al. 2012) and used to replace the 35S promoter in

109 pSAT4a (Tzfira et al. 2005) to produce pNSAT4a. The *xplA*, *xplB* genes (GenBank accession number  
110 AF449421 for both), and *nfsI* gene (M63808) were cloned by PCR from the vectors pMLBart-*xplA*,  
111 pART27-*xplB*, and pART27-*nfsI* (Rylott et al. 2011a) and inserted into pNSAT3a, pNSAT4a, and  
112 pNSAT6a, respectively, to produce pNSAT3a-*xplA*, pNSAT4a-*nfsI*, and pNSAT6a-*xplB* (Zhang et al.  
113 2017a). The hygromycin resistance gene, hygromycin B phosphotransferase (*hpt*), was cloned by PCR  
114 from pCambia1301 and inserted into pNSAT1a to produce pNSAT1a-*hpt*. The expression cassettes of  
115 *OsActin-hpt-35s* T, *ZmUbi-xplA-mas* T, *OsActin-nfsI-35S* T, and *PvUbi-xplB-rbc* T were sequentially  
116 inserted into the pRCS2 binary vector at these respective restriction sites *AscI-AscI*, *I-PpoI-I-PpoI*, *I-SceI-*  
117 *I-SceI*, *PI-PspI-PI-PspI* to produce pEDLZ2014. All the primers used in this research are listed in  
118 Supplement Table 1 and the sequences of all the elements in the vector were listed in Supplement  
119 Table 2.

## 120 **Culture condition for plant materials**

121 Western wheatgrass plants (including callus regeneration, RDX and TNT uptake experiments) were  
122 cultured at 25°C under 50  $\mu\text{E m}^{-2} \text{sec}^{-1}$  constant light with a 16 h day/8 h night photoperiod.

## 123 **Transformation of western wheatgrass**

124 The grain husks of western wheatgrass were removed and the seeds surface-sterilized in 10 % bleach for  
125 30 min, then rinsed four times with sterile water. Sterile seeds were cultured on callus induction (CI)  
126 medium (Murashige and Skoog's (MS) basic medium (Murashige and Skoog 1962) with 0.5 g/L proline,  
127 0.5 g/L N-Z amine, 2.5 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D), 0.1 mg/L 6-benzylaminopurine (6-  
128 BAP), 30 g/L sucrose, pH 5.8) in the dark for embryogenic callus induction. The embryogenic calli were  
129 sub-cultured onto CI medium at one-month intervals.

130 A biolistic particle system (Bio-Rad PDS-1000) was used for delivery of pEDLZ2014 with 7,584 kPa  
131 (1,100 psi) rupture disks, a microcarrier flight distance of 9 cm and a vacuum of 97 kPa (27 in) Hg, with  
132 all hardware and reagents produced by Bio-Rad. Microprojectile preparation was as described earlier

133 (Lutz et al. 2006). The embryogenic calli were incubated on osmotic medium (CI medium with 36.4 g/L  
134 sorbitol and 36.4 g/L mannitol) for 6 h prior to bombardment. After bombardment, the calli were keep in  
135 the dark overnight on osmotic medium, then transferred to CI medium for two days and then to CI  
136 medium with 75 mg/L hygromycin. After two months of growth, hygromycin-resistant calli were  
137 transferred to fresh selective CI medium for further selection and callus propagation. After three more  
138 months of growth, the hygromycin resistant calli were transferred to regeneration medium (MS medium  
139 with 1 mg/L 6-BAP, 0.1 mg/L 1-naphthaleneacetic acid (NAA), 30 g/L sucrose, pH 5.8) with 75 mg/L  
140 hygromycin. Regenerated plantlets were transferred to MS medium containing 50 mg/L hygromycin, then  
141 rooted plantlets transferred to shoot induction medium (MS medium with 2 mg/L 6-BAP, 0.2 mg/L NAA,  
142 30 g/L sucrose, pH 5.8) with 50 mg/L hygromycin for cluster shoots induction. After two months, new  
143 branches in cluster shoots were excised and transferred to fresh MS medium with hygromycin at 50 mg/L  
144 to promote growth and rooting. The rhizomes were cut into sections with a node on each fragment and  
145 cultured on MS medium for new plantlet development. One week later, new shoots and roots developed  
146 from nodes of rhizomes fragments. Healthy plants from the cluster shoots or rhizomes were selected and  
147 transferred to soil.

#### 148 **Transgene expression analysis**

149 DNA was extracted using DNeasy plant mini kits (Qiagen, Valencia, CA, USA), and PCR reactions  
150 performed using the primers listed in Table S1. For transcript analysis, mRNA was extracted from mature  
151 western wheatgrass leaf blades using RNeasy plant mini kits (Qiagen). One microgram of total RNA was  
152 treated with DNase I (Invitrogen) and used to synthesize cDNA with random primers (Invitrogen) and M-  
153 MLV reverse transcriptase (Invitrogen). Quantitative real-time PCR (qPCR) was performed using a  
154 SensiFast SYBR No-ROX kit (Bioline). Data from RT-qPCR were normalized to the western wheatgrass  
155 5.8S gene (GenBank: GU557082.1). Protein analysis used SDS-PAGE with 8 µg of crude protein per  
156 sample, and antibodies were used as reported previously, XplA, (Rylott et al. 2006); XplB, (Jackson et al.  
157 2007) and nitroreductase (Rylott et al. 2011a).



158 **RDX uptake studies**

159 Wild-type and transformed plants of comparable biomass and developmental status were selected and  
160 cultured in 5 mL ½ MS media with 30 mg/L RDX for 12 days. Levels of RDX in the medium were  
161 assayed at regular time intervals, with volumes of medium refilled back to 5 mL with tap water each time  
162 before sampling. After 12 days, the RDX concentrations in the plant tissues were determined.

163 **TNT uptake studies**

164 Mature leaf blades were cut into sections about 1.5 cm length and cultured in 20 mL water containing  
165 12.5, and 32.5 mg/L TNT. The starting biomass in each bottle was  $0.572 \pm 0.003$  g. Samples of medium  
166 were removed at regular intervals for analysis. After 66 hours, the concentrations of TNT and 4-ADNT in  
167 the plant tissues were determined.

168 **Extraction of RDX, TNT, and 4-ADNT from plant tissue**

169 Plant leaf tissue (100 mg) was freeze dried and ground to a powder using a Fast Prep 24 (MP  
170 Biomedicals, LLC. Solon, USA). RDX, TNT and ADNT were extracted from leaf materials using 1 mL  
171 methanol with shaking for 12 hours, then three repeat extractions with shaking for two hours. The  
172 combined extraction products were dried using an Eppendorf Vacufuge 5301, re-dissolved in 1.5 mL  
173 methanol, centrifuged at 13,000 for ten minutes and the supernatant (1000 µL) analyzed by HPLC.

174 **Determination of RDX, TNT and ADNT levels**

175 RDX, TNT and ADNT concentrations were analyzed using a modular Waters HPLC system consisting of  
176 a Waters 717 autosampler, two Waters 515 HPLC pumps, and a Waters 2996 photodiode array detector.  
177 A 4.6- by 250-mm Waters C18 column was used for separation under conditions similar to those  
178 described previously (Andeer et al. 2013), with concentration determined based on absorbance at 240 nm.  
179 Peak integrations and analyses were conducted using Millennium32 software (Waters, Milford, MA). The  
180 limit of detection of RDX by this method is 0.01 mg/L.

181 **Data analysis**

182 Data were analyzed for statistical significance using ANOVA in Microsoft Excel software (Microsoft  
183 Excel 2016 MSO). When ANOVA analysis gave a significant difference, Fishers Least Significant  
184 Difference (LSD) method was performed to compare the means. Groupings differing by statistical  
185 significance are labeled by letters in the figures.

186 **RESULTS**

187 **Creation of transgenic western wheatgrass**

188 Following results from the expression of *xplA*, *xplB*, and *nfsI* in switchgrass and creeping bentgrass, the  
189 vector used in our former report (Zhang et al. 2017a), was modified to improve transgene expression. The  
190 35S promoter driving expression of *nfsI* was replaced with the rice (*Oryza sativa*) actin promoter (*Osact*);  
191 and the switchgrass ubiquitin promoter (*Pvubi*) driving *xplA* expression was replaced with the maize (*Zea*  
192 *mays*) ubiquitin promoter (*ZmUbi*). The structure of the resulting vector, pEDLZ2014, is shown in  
193 Figure 1.

194 To successfully transform western wheatgrass, vigorously-growing embryogenic calli were needed.  
195 Figure 2a shows the white or light-yellow, friable embryogenic calli obtained after 1-2 months of growth.  
196 Following additional culturing, calli suitable for transformation were obtained (Figure 2b) and bombarded  
197 with pEDLZ2014 vector DNA. Figure 2c shows the appearance of the surviving calli two months after  
198 bombardment; after three to four months, green and healthy plantlets were obtained (Figure 2d). In total,  
199 10 plates of embryogenic calli were bombarded, with 98 calli showing hygromycin resistance and of  
200 these, 30 produced green and healthy plantlets (Figure 2e). To produce biomass for subsequent analysis,  
201 the plantlets were propagated on cluster shoot induction medium which initiated the formation of robust  
202 shoots and rhizomes as shown in Figure 2f. The rhizomes were cut into nodal sections, as shown in  
203 Figure 2g, and cultured to regenerate new plantlets. Figure 2h shows the appearance of the fully  
204 regenerated plants in soil.

205 To confirm the integration and expression of the transgenes in the hygromycin-resistant plantlets, PCR  
206 and qRT-PCR analysis were conducted on thirteen, independently-transformed lines. The results,  
207 presented in Figure 3a, show expression of *xplA*, *xplB* and *nfsI* in the transgenic lines, while the  
208 transgenes were not detectable in the wild-type, untransformed plants. Expression of the transgenes was  
209 highest in lines S35, N1, N9 and D22 and subsequent western blot analysis (Figure 3b) of these lines  
210 revealed the presence of 60, 45 and 24 kDa bands, which corresponded in size to the XplA, XplB and NR  
211 proteins, respectively.

### 212 **RDX uptake studies**

213 To determine the rate of RDX uptake, the plants were grown in liquid MS medium in open test tubes.  
214 Evaporation of RDX from the medium was minimal due to its low volatility (Xiong et al. 2009). Over the  
215 course of the experiment, all the transformed lines removed RDX from the medium faster than the wild-  
216 type plants (Figure 4a). After 3 days, the RDX concentration in the medium containing transformed plants  
217 was significantly lower than that of wild-type plants ( $p = 2 \times 10^{-6}$ ). After 6 days, the difference between  
218 wild-type and transformed lines further increased with lines N9 and D22 removing significantly more  
219 RDX than lines S35 and D45 ( $p = 0.0003$ ). After 12 days of culture, 65 % of the RDX had been removed  
220 by wild-type plants while 86 % and 87 % by line N9 and D22 respectively. After 12 days, wild-type RDX  
221 leaf tissue levels were found to be  $0.12 \pm 0.03$  mg/g RDX, in comparison, RDX was not detected in tissue  
222 from transformed lines tested, N9, D22, and S35, and RDX was at  $0.033 \pm 0.001$  mg/g in transformed line  
223 D45 (Figure 4b). To confirm that the lack of accumulation of RDX in transformed tissue was caused by  
224 degradation, rather than by dilution in growing plant tissues, leaf blades of wild-type and lines, N9 and  
225 D22 were cultured in water with 20 mg/L or 40 mg/L RDX. Figure 4c shows that while the concentration  
226 of RDX in the wild-type leaves increased from day 2 to day 8, no RDX was detected in the transformed  
227 line D22. Although RDX was detected in leaf tissue of transformed line N9, it was 28 % lower than that  
228 in wild-type ( $p = 0.004$ ) at day 2 when cultured in RDX solution at 20 mg/L, and 53% lower ( $p = 0.033$ )  
229 at day 2, with 40 RDX mg/L. In contrast to wild-type, the RDX concentration in line N9 decreased after 2

230 days of culture and could not be detected on day 8. This result confirmed that both N9 and D22 are able to  
231 degrade RDX.

### 232 **TNT detoxification studies**

233 Though the main target of this research was to engineer grasses for RDX degradation, the grasses also  
234 need to be able to resist the phytotoxicity of TNT since training ranges are contaminated by both RDX  
235 and TNT, and TNT can also inhibit XplA activity (Jackson et al. 2007). Thus the *xplA-xplB-nfsI*  
236 transformed lines N9, and D22, which had the highest RDX degradation abilities, were tested for their  
237 resistance to TNT conferred by NR activity. When leaf sections were cultured in water containing 15  
238 mg/L TNT, the transformed lines removed TNT more quickly than wild-type plants ( $p = 0.0019$  at 18 hr  
239 incubation). Line D22 removed TNT most rapidly, with TNT undetectable in the solution after 48 hr.  
240 After 66 hr, nearly all TNT had disappeared from solution from wild-type and transformed plants (Figure  
241 5a). Though wild-type western wheatgrass removed TNT from the solution, when compared to the  
242 transgenic lines after 66 hr, the wild-type leaf sections exhibited symptoms of toxicity, with overall  
243 yellowing of the leaf surfaces and darkening of the cut edges (Supplement Figure 1a). After 66 hours of  
244 culture in water containing 32 mg/L TNT, the wild-type plants were unable to remove any TNT, while the  
245 transformed lines removed nearly all the TNT ( $0.38 \pm 0.16$  mg/L remaining for N9 and  $0.51 \pm 0.35$  for  
246 D22; Figure 5b). Furthermore, the wild-type plants showed signs of toxicity, as indicated by a reduction  
247 in growth rate such that the biomass gain of wild-type was 24 % and 30 % of that of N9 and D22,  
248 respectively after 3 days (Supplement Figure 1b).

249 While in whole plants, TNT and transformation products are localized predominantly in the roots, when  
250 grown in liquid culture, leaf tissues are known to also contain these products (Hannink et al. 2007;  
251 Hannink et al. 2001). To verify that TNT was transformed in the tissues of the modified plants, TNT and  
252 ADNT levels, in leaf tissue were determined after 42 hours culture in TNT solution (Figure 5c). When  
253 cultured in 12.5 mg/L TNT solution, TNT was not detected in the tissues of either wild-type or  
254 transformed plants, while ADNT was detected in wild-type leaf tissue at a level lower than for N9 and

255 D22 ( $p = 0.0188$ ). When cultured in 25 mg/L TNT solution for 42 hours, the wild-type leaf tissue  
256 accumulated  $11.91 \pm 1.64$   $\mu\text{g/g}$  TNT, about 3 times of that in lines N9 and D22 ( $p = 0.0004$ ). At the same  
257 time, the ADNT concentration in the wild-type plants was  $6.01 \pm 0.92$   $\mu\text{g/g}$ , about 60 % of that in leaf  
258 tissue of transformed plants ( $p = 0.0012$ ). These results demonstrated that though wild-type plants have  
259 the ability to uptake and detoxify limited amounts of TNT, the *nfsI*-transformed western wheatgrass are  
260 able to detoxify significantly more TNT, resulting in increased tolerance, and subsequent ability to take  
261 up overall more TNT.

## 262 **DISCUSSION**

263 On live-fire training ranges, deposition of RDX and TNT onto soils around targets is ongoing and thus a  
264 continuing source of soil and groundwater contamination. Phytoremediation could be a relatively low-  
265 cost method to contain and clean-up these sites but many plant species grow poorly in RDX and TNT  
266 contaminated soil (Vila et al. 2007; Via and Zinnert 2016; Panz and Miksch 2012; Via et al. 2015) and are  
267 not able to degrade significant amounts of RDX (Best et al. 1999; Yoon et al. 2006; Best et al. 2006).

268 Studies in model plant species have demonstrated that genetically modifying plants with *xplA*, *xplB* and  
269 *nfsI* conferred the ability to degrade RDX and detoxify TNT (Hannink et al. 2007; Rylott et al. 2011a).

270 The subsequent development of genetically modified switchgrass and creeping bentgrass confirmed that  
271 this system for RDX degradation worked efficiently in monocot grasses (Zhang et al. 2017a). However,  
272 expression of the *nfsI* gene, under the control of the 35S promoter, in these lines was poor, and  
273 accordingly, the transformed switchgrass did not show enhanced resistance to TNT. Expression of the  
274 35S promoter has been reported to be lower in monocots (Jang et al. 2002; Christensen et al. 1992). In  
275 this research, the 35S promoter driving the expression of *nfsI* was replaced with the monocot *OsAct*  
276 promoter. The resulting western wheatgrass plants showed both enhanced RDX degradation and  
277 enhanced TNT detoxification.

278 Western wheatgrass has many desirable attributes that make it suitable for deployment on training ranges.  
279 It is native to the US, growing across a broad geographical location with includes military lands (USDA,  
280 2002). Furthermore, its robust leaves and rhizomes make it able to withstand stresses associated with  
281 military ranges such as fire, military vehicular traffic, and nutrient poor soils. Some of these attributes  
282 also make western wheatgrass valuable as a forage grass, so advances in its genetic modification are  
283 important in agriculture. Here we describe a method for reliably transforming western wheatgrass. This  
284 method could also be adapted to other forage grasses, such as slender wheatgrass (*Elymus trachycaulus*)  
285 and Siberian wheatgrass (*Agropyron fragile*). In our experience, a prerequisite for successful development  
286 of transformed callus is the selection of the friable white callus morphology (Figure 2b).

287 In conclusion, this is the first report on the genetic transformation of western wheatgrass. This protocol  
288 will contribute to wider plant biotechnology uses, here, we utilize it to show the efficacy of *xplA*, *xplB*,  
289 *nfsI*-expressing plants to degrade RDX and detoxify TNT. These species neatly complement transgenic  
290 switchgrass lines, enabling this technology to be used across a broader range of environmental, and  
291 geographic conditions for the containment and detoxification of explosives on military training ranges.

292

## 293 **Acknowledgements**

294 The authors declare no conflict of interest. This work was funded by US DoD SERDP ER-1498 and  
295 ESTCP ER-201436. We thank Mr. Ryan Routsong for his work of maintaining the HPLC system and  
296 advice for preparing HPLC samples.

297

## 298 **References**

299  
300 Adrian NR, Arnett CM (2004) Anaerobic biodegradation of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX)  
301 by *Acetobacterium malicum* strain HAAP-1 isolated from a methanogenic mixed culture. *Curr*  
302 *Microbiol* 48:332-340

303 Ahmad F, Schnitker SP, Newell CJ (2007) Remediation of RDX- and HMX-contaminated groundwater  
304 using organic mulch permeable reactive barriers. *J Contam Hydrol* 90:1-20

305 Andeer P, Stahl DA, Lillis L, Strand SE (2013) Identification of microbial populations assimilating  
306 nitrogen from RDX in munitions contaminated military training range soils by high sensitivity  
307 stable isotope probing. *Environ Sci Technol* 47:10356-10363

308 Andeer P, Stahl DA, Bruce NC, Strand SE (2009) Lateral transfer of genes for hexahydro-1,3,5-trinitro-  
309 1,3,5-triazine (RDX) degradation. *Appl Environ Microbiol* 75:3258-3262

310 Bernstein A, Adar E, Nejidat A, Ronen Z (2011) Isolation and characterization of RDX-degrading  
311 *Rhodococcus* species from a contaminated aquifer. *Biodegradation* 22:997-1005

312 Best EP, Geter KN, Tatem HE, Lane BK (2006) Effects, transfer, and fate of RDX from aged soil in  
313 plants and worms. *Chemosphere* 62:616-625

314 Best EP, Sprecher SL, Larson SL, Fredrickson HL, Bader DF (1999) Environmental behavior of  
315 explosives in groundwater from the Milan Army Ammunition Plant in aquatic and wetland plant  
316 treatments. Uptake and fate of TNT and RDX in plants. *Chemosphere* 39:2057-2072

317 Beynon ER, Symons ZC, Jackson RG, Lorenz A, Rylott EL, Bruce NC (2009) The role of  
318 oxophytodienoate reductases in the detoxification of the explosive 2,4,6-trinitrotoluene by  
319 *Arabidopsis*. *Plant Physiol* 151:253-261

320 Binks PR, Nicklin S, Bruce NC (1995) Degradation of hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX) by  
321 *Stenotrophomonas maltophilia* PB1. *Appl Environ Microbiol* 61:1318-1322

322 Bolt HM, Degen GH, Dorn SB, Plottner S, Harth V (2006) Genotoxicity and potential carcinogenicity of  
323 2,4,6-TNT trinitrotoluene: structural and toxicological considerations. *Rev Environ Health*  
324 21:217-228

325 Brentner LB, Mukherji ST, Walsh SA, Schnoor JL (2010) Localization of hexahydro-1,3,5-trinitro-1,3,5-  
326 triazine (RDX) and 2,4,6-trinitrotoluene (TNT) in poplar and switchgrass plants using phosphor  
327 imager autoradiography. *Environ Pollut* 158:470-475

328 Christensen AH, Sharrock RA, Quail PH (1992) Maize polyubiquitin genes: structure, thermal  
329 perturbation of expression and transcript splicing, and promoter activity following transfer to  
330 protoplasts by electroporation. *Plant Mol Biol* 18:675-689

331 Coleman NV, D.R. Nelson, and T. Duxbury (1998) Aerobic biodegradation of hexahydro-1,3,5-trinitro-  
332 1,3,5-triazine (RDX) as a nitrogen source by a *Rhodococcus sp.*, strain DN22. *Soil Biol Biochem*  
333 30:1159-1167

334 Gandia-Herrero F, Lorenz A, Larson T, Graham IA, Bowles DJ, Rylott EL, Bruce NC (2008)  
335 Detoxification of the explosive 2,4,6-trinitrotoluene in *Arabidopsis*: discovery of bifunctional O-  
336 and C-glucosyltransferases. *Plant J* 56:963-974

337 Gunning V, Tzafestas K, Sparrow H, Johnston EJ, Brentnall AS, Potts JR, Rylott EL, Bruce NC (2014)  
338 *Arabidopsis* glutathione transferases U24 and U25 exhibit a range of detoxification activities with  
339 the environmental pollutant and explosive, 2,4,6-trinitrotoluene. *Plant Physiol* 165:854-865

340 Hannink N, Rosser SJ, French CE, Basran A, Murray JA, Nicklin S, Bruce NC (2001)  
341 Phytodetoxification of TNT by transgenic plants expressing a bacterial nitroreductase. *Nature*  
342 *Biotechnol* 19:1168-1172

343 Hannink NK, Subramanian M, Rosser SJ, Basran A, Murray JA, Shanks JV, Bruce NC (2007) Enhanced  
344 transformation of tnt by tobacco plants expressing a bacterial nitroreductase. *Int J*  
345 *Phytoremediation* 9:385-401

346 Honeycutt ME, Jarvis AS, McFarland VA (1996) Cytotoxicity and mutagenicity of 2,4,6-trinitrotoluene  
347 and its metabolites. *Ecotoxicol Environ Saf* 35:282-287

348 Jackson RG, Rylott EL, Fournier D, Hawari J, Bruce NC (2007) Exploring the biochemical properties and  
349 remediation applications of the unusual explosive-degrading P450 system XplA/B. *Proc Natl*  
350 *Acad Sci U S A* 104:16822-16827

351 Jang IC, Choi WB, Lee KH, Song SI, Nahm BH, Kim JK (2002) High-level and ubiquitous expression of  
352 the rice cytochrome c gene *OsCc1* and its promoter activity in transgenic plants provides a useful  
353 promoter for transgenesis of monocots. *Plant Physiol* 129:1473-1481

354 Johnston EJ, Rylott EL, Beynon E, Lorenz A, Chechik V, Bruce NC (2015) Monodehydroascorbate  
355 reductase mediates TNT toxicity in plants. *Science* 349:1072-1075

356 Lutz KA, Svab Z, Maliga P (2006) Construction of marker-free transplastomic tobacco using the Cre-  
357 loxP site-specific recombination system. *Nat Protoc* 1:900-910

358 Mann DG, Lafayette PR, Abercrombie LL, King ZR, Mazarei M, Halter MC, Poovaiah CR, Baxter H,  
359 Shen H, Dixon RA, Parrott WA, Neal Stewart C, Jr. (2012) Gateway-compatible vectors for high-  
360 throughput gene functional analysis in switchgrass (*Panicum virgatum* L.) and other monocot  
361 species. *Plant Biotechnol J* 10:226-236

362 Mezzari MP, Van Aken B, Yoon JM, Just CL, Schnoor JL (2004) Mathematical modeling of RDX and  
363 HMX metabolism in poplar (*Populus deltoides* x *Populus nigra*, DN34) tissue culture. *Int J*  
364 *Phytoremediation* 6:323-345

365 Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco tissue  
366 cultures. *Physiol Plantarum* 15:473-497

367 Oh SY, Yoon HS, Jeong TY, Kim SD, Kim DW (2016) Reduction and persulfate oxidation of nitro  
368 explosives in contaminated soils using Fe-bearing materials. *Environ Sci Process Impacts* 18:863-  
369 871

370 Palazzo AJ, Jensen KB, Waldron BL, Cary TJ (2005) Effects of tank tracking on range grasses. *J*  
371 *Terramechanics* 42:177-191

372 Panz K, Miksch K (2012) Phytoremediation of explosives (TNT, RDX, HMX) by wild-type and  
373 transgenic plants. *J Environ Manage* 113:85-92

374 Pavlostathis SG, Jackson GH (2002) Biotransformation of 2,4,6-trinitrotoluene in a continuous-flow  
375 *Anabaena sp.* system. *Water Res* 36 (7):1699-1706

376 Pichtel J (2012) Distribution and fate of military explosives and propellants in soil: a review. *Appl*  
377 *Environ Soil Sci* 2012:33

378 Ramasahayam S, Jalgama S, Atwa SM, Salley JT, Thongdy M, Blaylock BL, Meyer SA (2017)  
379 Megakaryocyte expansion and macrophage infiltration in bone marrow of rats subchronically  
380 treated with MNX, N-nitroso environmental degradation product of munitions compound RDX  
381 (hexahydro-1,3,5-trinitro-1,3,5-triazine). *J Appl Toxicol* 37:913-921

382 Rylott E, Bruce N (2009) Plants disarm soil: engineering plants for the phytoremediation of explosives.  
383 *Trends Biotechnol* 27:73-81

384 Rylott EL, Budarina MV, Barker A, Lorenz A, Strand SE, Bruce NC (2011a) Engineering plants for the  
385 phytoremediation of RDX in the presence of the co-contaminating explosive TNT. *New Phytol*  
386 192:405-413

387 Rylott EL, Jackson RG, Edwards J, Womack GL, Seth-Smith HM, Rathbone DA, Strand SE, Bruce NC  
388 (2006) An explosive-degrading cytochrome P450 activity and its targeted application for the  
389 phytoremediation of RDX. *Nat Biotechnol* 24:216-219

390 Rylott EL, Jackson RG, Sabbadin F, Seth-Smith HM, Edwards J, Chong CS, Strand SE, Grogan G, Bruce  
391 NC (2011b) The explosive-degrading cytochrome P450 XplA: biochemistry, structural features  
392 and prospects for bioremediation. *Biochim Biophys Acta* 1814:230-236

393 Seth-Smith HM, Rosser SJ, Basran A, Travis ER, Dabbs ER, Nicklin S, Bruce NC (2002) Cloning,  
394 sequencing, and characterization of the hexahydro-1,3,5-Trinitro-1,3,5-triazine degradation gene  
395 cluster from *Rhodococcus rhodochrous*. *Appl Environ Microbiol* 68 (10):4764-4771

396 Seth-Smith HMB, Edwards J, Rosser SJ, Rathbone DA, Bruce NC (2008) The explosive-degrading  
397 cytochrome P450 system is highly conserved among strains of *Rhodococcus spp.* *Appl Environ*  
398 *Microbiol* 74:4550-4552

399 Sviatenko LK, Gorb L, Leszczynska D, Okovytyy SI, Shukla MK, Leszczynski J (2017) In silico kinetics  
400 of alkaline hydrolysis of 1,3,5-trinitro-1,3,5-triazinane (RDX): M06-2X investigation. *Environ*  
401 *Sci Process Impacts* 19:388-394

402 Thompson KT, Crocker FH, Fredrickson HL (2005) Mineralization of the cyclic nitramine explosive  
403 hexahydro-1,3,5-trinitro-1,3,5-triazine by *Gordonia* and *Williamsia* spp. *Appl Environ Microbiol*  
404 71:8265-8272



- 405 Travis ER, Bruce NC, Rosser SJ (2008) Short term exposure to elevated trinitrotoluene concentrations  
 406 induced structural and functional changes in the soil bacterial community. *Environ Pollut*  
 407 153 :432-439
- 408 Tzfira T, Tian GW, Lacroix B, Vyas S, Li J, Leitner-Dagan Y, Krichevsky A, Taylor T, Vainstein A,  
 409 Citovsky V (2005) pSAT vectors: a modular series of plasmids for autofluorescent protein  
 410 tagging and expression of multiple genes in plants. *Plant Mol Biol* 57:503-516
- 411 United States Department of Agriculture (2002) Plant fact sheet-western wheatgrass (*Pascopyrum*  
 412 *smithii*) [https://plants.usda.gov/factsheet/pdf/fs\\_pasm.pdf](https://plants.usda.gov/factsheet/pdf/fs_pasm.pdf)
- 413 United States Environmental Protection Agency (2014) Technical fact sheet –hexahydro-1,3,5-trinitro-  
 414 1,3,5-triazine (RDX) [https://www.epa.gov/sites/production/files/2014-](https://www.epa.gov/sites/production/files/2014-03/documents/ffrrofactsheet_contaminant_rdx_january2014_final.pdf)  
 415 03/documents/ffrrofactsheet\_contaminant\_rdx\_january2014\_final.pdf and technical fact sheet –  
 416 2,4,6-trinitrotoluene (TNT) [https://www.epa.gov/sites/production/files/2014-](https://www.epa.gov/sites/production/files/2014-03/documents/ffrrofactsheet_contaminant_tnt_january2014_final.pdf)  
 417 03/documents/ffrrofactsheet\_contaminant\_tnt\_january2014\_final.pdf
- 418 United States General Accounting Office (2004) Department of Defense operational ranges, more  
 419 reliable cleanup cost estimates and a proactive approach to identifying contamination are needed.  
 420 <http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA435939&Location=U2&doc=GetTRDoc.pdf>
- 421 Van Aken B, Yoon JM, Just CL, Schnoor JL (2004) Metabolism and mineralization of hexahydro-1,3,5-  
 422 trinitro-1,3,5-triazine inside poplar tissues (*Populus deltoides* x *nigra* DN-34). *Environ Sci*  
 423 *Technol* 38:4572-4579
- 424 Via SM, Zinnert JC (2016) Impacts of explosive compounds on vegetation: A need for community scale  
 425 investigations. *Environ Pollut* 208:495-505
- 426 Via SM, Zinnert JC, Young DR (2015) Differential effects of two explosive compounds on seed  
 427 germination and seedling morphology of a woody shrub, *Morella cerifera*. *Ecotoxicology*  
 428 24:194-201
- 429 Vila M, Lorber-Pascal S, Laurent F (2007) Fate of RDX and TNT in agronomic plants. *Environ Pollut*  
 430 148 :148-154
- 431 Winfield LE, Rodger, J. H., & D'surney, S.,J. (2004) The responses of selected terrestrial plants to short  
 432 (<12 days) and long term (2, 4 and 6 weeks) hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX)  
 433 exposure. part I: Growth and developmental effects. *Ecotoxicology* 13:335-347
- 434 Xiong RC, Fern JT, Keffer DJ, Fuentes-Cabrera M, Nicholson DM (2009) Molecular simulations of  
 435 adsorption and diffusion of RDX in IRMOF-1. *Mol Simulat* 35:910-919
- 436 Yoon JM, Van Aken B, Schnoor JL (2006) Leaching of contaminated leaves following uptake and  
 437 phytoremediation of RDX, HMX, and TNT by poplar. *Int J Phytoremediation* 8:81-94
- 438 Zhang L, Routsong R, Nguyen Q, Rylott EL, Bruce NC, Strand SE (2017a) Expression in grasses of  
 439 multiple transgenes for degradation of munitions compounds on live-fire training ranges. *Plant*  
 440 *Biotechnol J* 15:624-633
- 441 Zhang L, Rylott EL, Bruce NC, Strand SE (2017b) Phytodetoxification of TNT by transplastomic tobacco  
 442 (*Nicotiana tabacum*) expressing a bacterial nitroreductase. *Plant Mol Biol* 95:99-109

443

## 444 **Legends**

445 **Figure 1** Construction of vector pEDLZ2014 for transformation of western wheatgrass.

446 T-DNA region of the binary vector plasmid pEDLZ2014 used to transform western wheatgrass. The RDX  
 447 degradation gene *xplA*, flavodoxin reductase gene *xplB*, and TNT detoxifying nitroreductase gene *nfsI*  
 448 were constructed into versatile cloning vector pNSATs. Arrows show the direction of transcription.

449 Abbreviations: *Osact*, *Oryza sativa* actin promoter; *hpt*, selection marker gene, *hygromycin B*  
450 *phosphotransferase (hpt)*; 35s T, terminator of CaMV 35s gene; *Zmubi*, *Zea mays* ubiquitin promoter;  
451 *mas T*, terminator of *manopine synthase* gene of *Agrobacterium*; *Pvubi*, *Panicum virgatum* (switchgrass)  
452 ubiquitin promoter; *rbc T*, terminator of rubisco small subunit of tobacco; RB left border; RB right border  
453

454 **Figure 2** Tissue culture, transformation and propagation of western wheatgrass.

455 (a) Two month old calli induced from mature seeds. (b) Embryogenic calli that were used for  
456 bombardment with vector DNA. (c) Calli screened on Callus Induction medium with 75 mg/L  
457 hygromycin for two months after bombardment. (d) Hygromycin resistant plantlets from calli that were  
458 transferred to regeneration medium for 1-2 month. (e) Regenerated plantlets growing on MS medium with  
459 75 mg/L hygromycin. (f) PCR and qRT-PCR positive plants were transferred to cluster shoot induction  
460 medium to induce cluster shoots and rhizomes. (g) Shoot induced on rhizomes sections with nodules on MS  
461 medium. (h) Transgenic plants growing in soil.

462

463

464 **Figure 3** Molecular characterization of *xplA-xplB-nfsI* transformed western wheatgrass.

465 a) Transcript abundance measured using quantitative RT-PCR on plant lines transformed with *xplA*, *xplB*  
466 and *nfsI*. Values were normalized to the 5.8s rRNA gene and relative to expression level of the transformed  
467 line D22. Data are the means  $\pm$  SE, n = 3.

468 b) Western blot analysis on leaf blades of western wheatgrass lines expressing XplA, XplB and  
469 nitroreductase (NR) protein. WT, wild type; PC, positive control, *xplA-xplB-nfsI* transformed *Arabidopsis*  
470 (Rylott et al. 2011a).

471

472 **Figure 4** Uptake and degradation of RDX by *xplA-xplB-nfsI* transformed western wheatgrass. (a)  
473 Concentration of RDX in culture medium over the course of the experiment. (b) Concentration of RDX in  
474 plant tissue after 12 days culture. (c) Accumulation of RDX in leaf tissue during 8 days culture in 20 mg/L


475 and 40 mg/L RDX solution. Letters indicate that RDX concentration in tissue were significantly different  
476 ( $p < 0.05$ ) from other lines. Data are the means  $\pm$  SE,  $n = 3$ ; ND = None Detected; NPC, No Plant Control

477

478 **Figure 5** Uptake of TNT by *xplA-xplB-nfsI* transformed western wheatgrass. Plants were grown in water  
479 with TNT at (A) 15 mg/L, and (b) 32 mg/L. (C) The concentration of TNT and 4-ADNT in the leaf tissue  
480 after 42 hours culture in 12.5 and 25 mg/L TNT solution. The lowercase letters and uppercase letters  
481 indicate that TNT and 4-ADNT concentration in tissue were significantly different from other lines at  
482 0.05 or 0.01 level (ND = None Detected). Data are the means  $\pm$  SE,  $n = 3$ .

483

484



493 **Figure 1** Construction of vector pEDLZ2014 for transformation of western wheatgrass.

494 T-DNA region of the binary vector plasmid pEDLZ2014 used to transform western wheatgrass. The RDX  
495 degradation gene *xplA*, flavodoxin reductase gene *xplB*, and TNT detoxifying nitroreductase gene *nfsI*  
496 were constructed into versatile cloning vector pNSATs. Arrows show the direction of transcription.

497 Abbreviations: *Osact*, *Oryza sativa* actin promoter; *hpt*, selection marker gene, *hygromycin B*

498 *phosphotransferase (hpt)*; 35s T, terminator of CaMV 35s gene; *Zmubi*, *Zea mays* ubiquitin promoter;

499 *mas T*, terminator of *manopine synthase* gene of *Agrobacterium*; *Pvubi*, *Panicum virgatum* (switchgrass)  
500 ubiquitin promoter; *rbc T*, terminator of rubisco small subunit of tobacco; LB left border; RB right border

501

502

503

504

505

506


507

508

509

510

511



512

513

514 **Figure 2** Tissue culture, transformation and propagation of western wheatgrass.

515 (a) Two month old calli induced from mature seeds. (b) Embryogenic calli that were used for

516 bombardment with vector DNA. (c) Calli screened on Callus Induction medium with 75 mg/L

517 hygromycin for two months after bombardment. (d) Hygromycin resistant plantlets from calli that were


518 transferred to regeneration medium for 1-2 month. (e) Regenerated plantlets growing on MS medium with

519 75 mg/L hygromycin. (f) PCR and qRT-PCR positive plants were transferred to cluster shoot induction

520 medium to induce cluster shoots and rhizomes. (g) Shoot induced on rhizomes sections with nodules on MS

521 medium. (h) Transgenic plants growing in soil.


522



523

524

525



526

527

528

529 **Figure 3** Molecular characterization of *xplA-xplB-nfsI* transformed western wheatgrass.

530 **a)** Transcript abundance measured using quantitative RT- PCR on plant lines transformed with *xplA*, *xplB*  
 531 and *nfsI*. Values were normalized to the 5.8s rRNA gene and relative to expression level of the transformed  
 532 line D22. Data are the means  $\pm$  SE, n = 3.



533 **b)** Western blot analysis on leaf blades of western wheatgrass lines expressing XplA, XplB and  
 534 nitroreductase (NR) protein. WT, wild type; PC, positive control, *xplA-xplB-nfsI* transformed Arabidopsis  
 535 (Rylott et al. 2011a).

536

537


538

539



541





542

543

544 **Figure 4** Uptake and degradation of RDX by *xplA-xplB-nfsI* transformed western wheatgrass. **(a)**

545 Concentration of RDX in culture medium over the course of the experiment. **(b)** Concentration of RDX in

546 plant tissue after 12 days culture. **(c)** Accumulation of RDX in leaf tissue during 8 days culture in 20 mg/L


547 and 40 mg/L RDX solution. Letters indicate that RDX concentration in tissue were significantly different

548 ( $p < 0.05$ ) from other lines. Data are the means  $\pm$  SE,  $n = 3$ ; ND = None Detected; NPC, No Plant Control

549


550

551



552

553



554

555


556

557

558

559

560



561

562 **Figure 5** Uptake of TNT by *xplA-xplB-nfsI* transformed western wheatgrass. Plants were grown in water  
563 with TNT at (A) 15 mg/L, and (b) 32 mg/L. (C) The concentration of TNT and 4-ADNT in the leaf tissue  
564 after 42 hours culture in 12.5 and 25 mg/L TNT solution. The lowercase letters and uppercase letters  
565 indicate that TNT and 4-ADNT concentration in tissue were significantly different from other lines at  
566 0.05 or 0.01 level (ND = None Detected). Data are the means  $\pm$  SE, n = 3.

567

568

569