



Comparison Inequalities for Order Statistics of Gaussian Arrays

K. Dębicki, E. Hashorva, L. Ji and C. Ling

Mathematical Institute, University of Wrocław
pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland.
E-mail address: Krzysztof.Debicki@math.uni.wroc.pl

Department of Actuarial Science, University of Lausanne
UNIL-Dorigny, 1015 Lausanne, Switzerland.
E-mail address: Enkelejda.Hashorva@unil.ch

Institute for Information and Communication Technologies,
HEIG-VD, University of Applied Sciences of Western Switzerland
Route de Cheseaux 1, 1401 Yverdon-les-Bains, Switzerland
E-mail address: lanpeng.ji@unil.ch

School of Mathematics and Statistics, Southwest University
Beibei District, 400715 Chongqing, China.
E-mail address: lcx98@swu.edu.cn

Abstract. Normal comparison lemma and Slepian's inequality are essential tools for the analysis of extremes of Gaussian processes. In this paper we show that the Normal comparison lemma for Gaussian vectors can be extended to order statistics of Gaussian arrays. Our applications include the derivation of mixed Gumbel limit laws for the order statistics of stationary Gaussian processes and the investigation of lower tail behavior of order statistics of self-similar Gaussian processes.

1. Introduction

In the recent contributions of Dębicki et al. (2015a, 2014b, 2015b) order statistics of Gaussian and stationary processes are studied. Given a random process $\{X(t), t \geq 0\}$ with almost surely (a.s.) continuous trajectories, and $X_1, \dots, X_n, n \in \mathbb{N}$ independent copies of X we define $X_{r:n}(t)$ generated by X as the r th lower order statistics of $X_1(t), \dots, X_n(t)$ for any fixed $t \geq 0$, and thus $X_{1:n}(t) \leq \dots \leq$

Received by the editors February 15, 2016; accepted January 9, 2017.

2010 *Mathematics Subject Classification.* Primary 60G15; secondary 60G70.

Key words and phrases. Slepian's inequality; conjunction probability; Normal comparison inequality; order statistics process; mixed Gumbel limit theorem; lower tail probability; self-similar Gaussian process.

Research supported by SNSF grant (200021-140633/1, 200021-166274), NCN Grant No 2013/09/B/ST1/01778, NSFC11601439 and cstc2016jcyjA0036.

$X_{n:n}(t), t \geq 0$. The calculation of the so-called r -th conjunction probability

$$p_{r:n}(u) = \mathbb{P} \left\{ \sup_{t \in [0, T]} X_{r:n}(t) > u \right\} \quad (1.1)$$

for fixed r, T and large u is of both theoretical and applied interest; see e.g., [Alodat \(2011\)](#); [Alodat et al. \(2010\)](#); [Ling and Peng \(2016\)](#); [Worsley and Friston \(2000\)](#).

Order statistics processes play a crucial role in various statistical applications, for instance in models concerned with the analysis of the surface roughness during all machinery processes and functional magnetic resonance imaging (fMRI) data. Given the fact that $p_{r:n}(u)$ cannot be in general calculated explicitly, asymptotic expansions as $u \rightarrow \infty$ and the so-called Gumbel limit results (with $u = u_T \rightarrow \infty$ as $T \rightarrow \infty$) are derived in [Dębicki et al. \(2015a,b\)](#). Indeed, such limit theorems have been in the focus of many theoretical and applied contributions, see e.g., [Aue et al. \(2009\)](#); [Berman \(1982, 1992\)](#); [Piterbarg \(1996, 2004\)](#) and the recent contributions [Dębicki et al. \(2015c\)](#); [Jarušková \(2015\)](#). The crucial tool for establishing Gumbel limit theorems is the so-called Normal comparison lemma, which has been shown to be one of the most important tools in the study of Gaussian processes and random fields, see e.g., [Berman \(1982, 1992\)](#); [Leadbetter et al. \(1983\)](#); [Li and Shao \(2002, 2004\)](#); [Piterbarg \(1996\)](#). The lack of a comparison lemma for order statistics processes has already been noted in [Dębicki et al. \(2015a\)](#); therein some results are derived only for the minimum process.

In the simpler framework of two d -dimensional Gaussian distributions $\Phi_{\Sigma^{(1)}}$ and $\Phi_{\Sigma^{(0)}}$ with $N(0, 1)$ marginal distributions, the normal comparison inequality gives explicit bounds for the difference

$$\Delta(\mathbf{u}) := \Phi_{\Sigma^{(1)}}(\mathbf{u}) - \Phi_{\Sigma^{(0)}}(\mathbf{u}), \quad \forall \mathbf{u} = (u_1, \dots, u_d) \in \mathbb{R}^d$$

in terms of the covariance matrices $\Sigma^{(k)} = (\sigma_{ij}^{(k)})_{d \times d}, k = 0, 1$. The derivation of the bounds for $\Delta(\mathbf{u})$, by [Slepian \(1962\)](#), [Berman \(1964, 1992\)](#) and [Piterbarg \(1996, 2015\)](#) relies strongly on Plackett's partial differential equation; see [Plackett \(1954\)](#). The most elaborate version of the normal comparison inequality is due to [Li and Shao \(2002\)](#). Specifically, Theorem 2.1 therein shows that

$$\Delta(\mathbf{u}) \leq \frac{1}{2\pi} \sum_{1 \leq i < j \leq d} \left(\arcsin(\sigma_{ij}^{(1)}) - \arcsin(\sigma_{ij}^{(0)}) \right)_+ \exp \left(-\frac{u_i^2 + u_j^2}{2(1 + \rho_{ij})} \right), \quad \forall \mathbf{u} \in \mathbb{R}^d,$$

where $\rho_{ij} := \max(|\sigma_{ij}^{(0)}|, |\sigma_{ij}^{(1)}|)$ and $x_+ = \max(x, 0)$. Clearly, if $\sigma_{ij}^{(0)} \leq \sigma_{ij}^{(1)}, 1 \leq i, j \leq d$, then

$$\Phi_{\Sigma^{(0)}}(\mathbf{u}) \leq \Phi_{\Sigma^{(1)}}(\mathbf{u}),$$

which is the well-known Slepian's inequality derived in [Slepian \(1962\)](#). Based on the results of [Li and Shao \(2002\)](#), [Yan \(2009\)](#) showed that for W an $N(0, 1)$ random variable and $\mathbf{u} \in (0, \infty)^d$

$$1 \leq \frac{\Phi_{\Sigma^{(1)}}(\mathbf{u})}{\Phi_{\Sigma^{(0)}}(\mathbf{u})} \leq \exp \left(\frac{1}{\sqrt{2\pi}} \sum_{1 \leq i < j \leq d} \frac{e^{-\frac{(u_i + u_j)^2}{8}}}{\mathbb{E} \left\{ \left(W + \frac{u_i + u_j}{2} \right)_+ \right\}} \ln \left(\frac{\pi - 2 \arcsin(\sigma_{ij}^{(0)})}{\pi - 2 \arcsin(\sigma_{ij}^{(1)})} \right) \right)$$

provided that $0 \leq \sigma_{ij}^{(0)} \leq \sigma_{ij}^{(1)} \leq 1$. Recent extensions of the normal comparison inequalities are presented in [Chernozhukov et al. \(2015\)](#); [Dębicki et al. \(2015a\)](#); [Harper \(2013, 2017\)](#); [Hashorva and Weng \(2014\)](#); [Lu and Wang \(2014\)](#).

Our principal goal of this paper is the derivation of comparison inequalities for order statistics of Gaussian arrays, which are useful in several applications. In order to fix the notation, we denote by $\mathcal{X} = (X_{ij})_{d \times n}$ and $\mathcal{Y} = (Y_{ij})_{d \times n}$ two $d \times n$ random arrays with $N(0, 1)$ components and jointly Gaussian (hereafter referred to as standard Gaussian arrays), and let $\Sigma^{(1)} = (\sigma_{ij,lk}^{(1)})_{dn \times dn}$ and $\Sigma^{(0)} = (\sigma_{ij,lk}^{(0)})_{dn \times dn}$ be the covariance matrices of \mathcal{X} and \mathcal{Y} , respectively, with $\sigma_{ij,lk}^{(1)} := \mathbb{E}\{X_{ij}X_{lk}\}$ and $\sigma_{ij,lk}^{(0)} := \mathbb{E}\{Y_{ij}Y_{lk}\}$. Furthermore, define $\mathbf{X}_{(r)} = (X_{1(r)}, \dots, X_{d(r)})$, $1 \leq r \leq n$ to be the r th order statistics vector generated by \mathcal{X} as follows

$$X_{i(1)} = \min_{1 \leq j \leq n} X_{ij} \leq \dots \leq X_{i(r)} \leq \dots \leq \max_{1 \leq j \leq n} X_{ij} = X_{i(n)}, \quad 1 \leq i \leq d.$$

Similarly, we write $\mathbf{Y}_{(r)} = (Y_{1(r)}, \dots, Y_{d(r)})$ which is generated by \mathcal{Y} . Clearly, in the case of independent rows of Gaussian arrays, the study of $\mathbf{X}_{(r)}$ reduces to that of the component-wise order statistics $X'_{i(r)}$ s for Gaussian random vector, see [Chernozhukov et al. \(2015\)](#). Our principal results, stated in [Theorem 2.1](#) and [Theorem 2.4](#), derive bounds for the difference

$$\Delta_{(r)}(\mathbf{u}) := \mathbb{P}\{\mathbf{X}_{(r)} \leq \mathbf{u}\} - \mathbb{P}\{\mathbf{Y}_{(r)} \leq \mathbf{u}\}, \quad \mathbf{u} \in \mathbb{R}^d. \quad (1.2)$$

Two applications of those bounds are discussed in [Section 3](#), including the study of the mixed Gumbel limit theorems for order statistics of stationary Gaussian processes and the lower tail probability of order statistics of self-similar Gaussian processes.

We organize this paper as follows. In [Section 2](#) we display our main results. [Section 3](#) is devoted to the applications. The proofs are relegated to [Section 4](#) and [Appendix](#).

2. Main Results

This section is concerned with sharp bounds for $\Delta_{(r)}(\mathbf{u})$ defined in (1.2), which go in line with Li and Shao's normal comparison inequality (see [Li and Shao, 2002](#)). For notational simplicity we set below

$$Q_{ij,lk} = \left| \arcsin(\sigma_{ij,lk}^{(1)}) - \arcsin(\sigma_{ij,lk}^{(0)}) \right|, \quad Q_{ij,lk}^+ = (\arcsin(\sigma_{ij,lk}^{(1)}) - \arcsin(\sigma_{ij,lk}^{(0)}))_+.$$

Theorem 2.1. *If \mathcal{X} and \mathcal{Y} are two standard $d \times n$ Gaussian arrays, then for any $1 \leq r \leq n$ we have*

$$\begin{aligned} |\Delta_{(r)}(\mathbf{u})| \leq & \frac{1}{2\pi} \left(\sum_{\substack{1 \leq i \leq d \\ 1 \leq j < k \leq n}} Q_{ij,ik} \exp\left(-\frac{u_i^2}{1 + \rho_{ij,ik}}\right) \right. \\ & \left. + \sum_{\substack{1 \leq i < l \leq d \\ 1 \leq j, k \leq n}} Q_{ij,lk} \exp\left(-\frac{u_i^2 + u_l^2}{2(1 + \rho_{ij,lk})}\right) \right), \quad \forall \mathbf{u} \in \mathbb{R}^d, \quad (2.1) \end{aligned}$$

where $\rho_{ij,lk} = \max(|\sigma_{ij,lk}^{(0)}|, |\sigma_{ij,lk}^{(1)}|)$. If further

$$\sigma_{ij,ik}^{(1)} = \sigma_{ij,ik}^{(0)}, \quad 1 \leq i \leq d, \quad 1 \leq j, k \leq n, \quad (2.2)$$

then

$$\Delta_{(r)}(\mathbf{u}) \leq \frac{1}{2\pi} \sum_{\substack{1 \leq i < l \leq d \\ 1 \leq j, k \leq n}} Q_{ij, lk}^+ \exp\left(-\frac{u_i^2 + u_l^2}{2(1 + \rho_{ij, lk})}\right), \quad \forall \mathbf{u} \in \mathbb{R}^d. \quad (2.3)$$

Remark 2.2. For $r = 1$ and $r = n$ the claims in (2.1) reduce to Lemma 11 in Dębicki et al. (2014a). Note that for $1 < r < n$ our results are derived using a different technique. Furthermore, using in addition similar arguments as in Theorem 1.2 in Piterbarg (1996), one can establish for any $[\mathbf{a}, \mathbf{b}] \subset [-\infty, \infty]^d$ the following comparison inequality

$$\begin{aligned} & \left| \mathbb{P}\{\mathbf{X}_{(r)} \in [\mathbf{a}, \mathbf{b}]\} - \mathbb{P}\{\mathbf{Y}_{(r)} \in [\mathbf{a}, \mathbf{b}]\} \right| \\ & \leq \frac{1}{\pi} \left(\sum_{\substack{1 \leq i \leq d \\ 1 \leq j < k \leq n}} Q_{ij, ik} \exp\left(-\frac{u_i^2}{1 + \rho_{ij, ik}}\right) + \sum_{\substack{1 \leq i < l \leq d \\ 1 \leq j, k \leq n}} Q_{ij, lk} \exp\left(-\frac{u_i^2 + u_l^2}{2(1 + \rho_{ij, lk})}\right) \right), \end{aligned}$$

with $u_i = \min(|a_i|, |b_i|)$, $1 \leq i \leq d$.

A direct consequence of Theorem 2.1 is the following Slepian's inequality for the order statistics of Gaussian arrays, which for $r = 1$ is, however, weaker than Theorem 1.1 in Gordon (1985).

Corollary 2.3. *Suppose that (2.2) is satisfied and $\sigma_{ij, lk}^{(0)} \geq \sigma_{ij, lk}^{(1)}$ holds for $1 \leq i < l \leq d, 1 \leq j, k \leq n$. Then*

$$\mathbb{P}\left\{\bigcup_{i=1}^d \{X_{i(r)} > u_i\}\right\} \geq \mathbb{P}\left\{\bigcup_{i=1}^d \{Y_{i(r)} > u_i\}\right\}, \quad \forall \mathbf{u} \in \mathbb{R}^d. \quad (2.4)$$

Note that the bounds in Theorem 2.1 do not depend on r , which indicates that in some cases they may not be sharp enough. Below we present a sharper result which holds under the assumption that the columns of both \mathcal{X} and \mathcal{Y} are mutually independent and identically distributed, i.e.,

$$\sigma_{ij, lk}^{(\kappa)} = \sigma_{il}^{(\kappa)} \mathbb{I}\{j = k\}, \quad 1 \leq i, l \leq d, 1 \leq j, k \leq n, \kappa = 0, 1, \quad (2.5)$$

with some $\sigma_{il}^{(\kappa)} \in (-1, 1)$, $1 \leq i, l \leq d, \kappa = 0, 1$, where $\mathbb{I}\{\cdot\}$ stands for the indicator function. This result is useful for establishing mixed Gumbel limit theorems; see Section 3.

In order to simplify the presentation, we shall define

$$c_{n,r} = \frac{n!}{r!(n-r)!}, \quad 0 \leq r \leq n, \quad \rho_{il} = \max(|\sigma_{il}^{(0)}|, |\sigma_{il}^{(1)}|), \quad 1 \leq i, l \leq d$$

and

$$A_{il}^{(r)} = \int_{\sigma_{il}^{(0)}}^{\sigma_{il}^{(1)}} \frac{(1 + |h|)^{2(n-r)}}{(1 - h^2)^{(n-r+1)/2}} dh, \quad 1 \leq i, l \leq d, 1 \leq r \leq n.$$

Theorem 2.4. *Under the assumptions of Theorem 2.1, if further (2.5) is satisfied, then for any $\mathbf{u} \in (0, \infty)^d$*

$$\Delta_{(r)}(\mathbf{u}) \leq \frac{n(c_{n-1,r-1})^2}{(2\pi)^{n-r+1}} u^{-2(n-r)} \sum_{1 \leq i < l \leq d} (A_{il}^{(r)})_+ \exp\left(-\frac{(n-r+1)u^2}{1 + \rho_{il}}\right) \quad (2.6)$$

and

$$|\Delta_{(r)}(\mathbf{u})| \leq \frac{n(c_{n-1,r-1})^2}{(2\pi)^{n-r+1}} u^{-2(n-r)} \sum_{1 \leq i < l \leq d} |A_{il}^{(r)}| \exp\left(-\frac{(n-r+1)u^2}{1+\rho_{il}}\right)$$

hold with $u = \min_{1 \leq i \leq d} u_i$.

As in Theorem 2.4 we also have the following bounds, without introducing $u = \min_{1 \leq i \leq d} u_i$.

Proposition 2.5. *Under the assumptions of Theorem 2.1, if further (2.5) is satisfied, then for any $\mathbf{u} \in (0, \infty)^d$*

$$\begin{aligned} \Delta_{(r)}(\mathbf{u}) &\leq \frac{n(c_{n-1,r-1})^2}{2(\sqrt{\pi})^{n-r+2}} \sum_{1 \leq i < l \leq d} \frac{(B_{il}^{(r)})_+}{(u_i + u_l)^{n-r}} \\ &\quad \times \exp\left(-\frac{(n-r)(u_i + u_l)^2 + 2(u_i^2 + u_l^2)}{4(1+\rho_{il})}\right), \end{aligned} \quad (2.7)$$

with

$$B_{il}^{(r)} = \int_{\sigma_{il}^{(0)}}^{\sigma_{il}^{(1)}} \frac{(1+|h|)^{(n-r)/2}}{(1-h^2)^{1/2}} dh, \quad 1 \leq i < l \leq d, \quad 1 \leq r \leq n.$$

If additionally $D_{il} = \min(u_i - \rho_{il}u_l, u_l - \rho_{il}u_i) > 0$ for all $1 \leq i < l \leq d$, then

$$\begin{aligned} \Delta_{(r)}(\mathbf{u}) &\leq \frac{n(c_{n-1,r-1})^2}{(2\pi)^{n-r+1}} \sum_{1 \leq i < l \leq d} (\tilde{A}_{il}^{(r)})_+ \left(\frac{u_i + u_l}{2} D_{il}\right)^{-(n-r)} \\ &\quad \times \exp\left(-\frac{(n-r+1)(u_i^2 + u_l^2)}{2(1+\rho_{il})}\right), \end{aligned} \quad (2.8)$$

where

$$\tilde{A}_{il}^{(r)} = \int_{\sigma_{il}^{(0)}}^{\sigma_{il}^{(1)}} \frac{(1+|h|)^{2(n-r)}(1-|h|)^{(n-r)}}{(1-h^2)^{(n-r+1)/2}} dh, \quad 1 \leq i < l \leq d, \quad 1 \leq r \leq n.$$

Motivated by, e.g., Li and Shao (2002); Lu and Wang (2014); Yan (2009), we obtain next an upper bound for $\Theta_{(r)}(\mathbf{u}) := \mathbb{P}\{\mathbf{X}_{(r)} \leq \mathbf{u}\} / \mathbb{P}\{\mathbf{Y}_{(r)} \leq \mathbf{u}\}$.

Proposition 2.6. *Under the assumptions of Theorem 2.1, if further (2.2) holds and $0 \leq \sigma_{ij,lk}^{(0)} \leq \sigma_{ij,lk}^{(1)} < 1$ for $1 \leq i < l \leq d, 1 \leq j, k \leq n$, then for any $\mathbf{u} \in [0, \infty)^d$*

$$1 \leq \Theta_{(r)}(\mathbf{u}) \leq \exp\left(\frac{1}{\sqrt{2\pi}} \sum_{\substack{1 \leq i < l \leq d \\ 1 \leq j, k \leq n}} \frac{C_{ij,lk} e^{-(u_i+u_l)^2/8}}{\mathbb{E}\{(W + (u_i + u_l)/2)_+\}}\right), \quad (2.9)$$

with W an $N(0, 1)$ random variable and

$$C_{ij,lk} = \ln\left(\frac{\pi - 2 \arcsin(\sigma_{ij,lk}^{(0)})}{\pi - 2 \arcsin(\sigma_{ij,lk}^{(1)})}\right), \quad 1 \leq i < l \leq d, \quad 1 \leq j, k \leq n.$$

3. Applications and Discussions

3.1. *Limit theorems for stationary order statistics processes.* When dealing with supremum of Gaussian processes on large intervals, the so-called Gumbel limit theorems are of interest for statistical applications.

Next, let $\{X_{n-r+1:n}(t), t \geq 0\}$ be the r th upper order statistics process generated by a centered stationary Gaussian process $\{X(t), t \geq 0\}$ with a.s. continuous sample paths, unit variance and correlation function $\rho(\cdot)$ satisfying for some $\alpha \in (0, 2]$

$$\rho(t) = 1 - |t|^\alpha + o(|t|^\alpha), \quad t \rightarrow 0, \quad \text{and} \quad \rho(t) < 1, \quad \forall t \neq 0. \quad (3.1)$$

From Theorem 1 in Dębicki et al. (2015a) or Theorem 2.2 in Dębicki et al. (2014b), we have for any $T > 0$ and $u \rightarrow \infty$

$$\mathbb{P} \left\{ \sup_{t \in [0, T]} X_{n-r+1:n}(t) > u \right\} \sim T \mathcal{A}_{r,\alpha} c_{n,r} (2\pi)^{-\frac{r}{2}} u^{\frac{2}{\alpha}-r} \exp \left(-\frac{ru^2}{2} \right), \quad (3.2)$$

where $\mathcal{A}_{r,\alpha} \in (0, \infty)$ is given explicitly as a limit and \sim means asymptotic equivalence. As a continuation of Dębicki et al. (2015a) we establish below a limit theorem for the r th upper order statistics process $X_{n-r+1:n}$.

Theorem 3.1. *Let $\{X_{n-r+1:n}(t), t \geq 0\}$ be the r th upper order statistics process generated by X , a centered stationary Gaussian process with a.s. continuous sample paths. Suppose that (3.1) holds and further $\lim_{t \rightarrow \infty} \rho(t) \ln t = \gamma \in [0, \infty]$.*

a) *If $\gamma = 0$, then*

$$\lim_{T \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| \mathbb{P} \left\{ a_{r,T} \left(\sup_{t \in [0, T]} X_{n-r+1:n}(t) - b_{r,T} \right) \leq x \right\} - \exp(-e^{-x}) \right| = 0,$$

where, with $D = c_{n,r} \mathcal{A}_{r,\alpha} (r/2)^{r/2-1/\alpha} (2\pi)^{-r/2}$

$$a_{r,T} = \sqrt{2r \ln T}, \quad b_{r,T} = \sqrt{\frac{2 \ln T}{r}} + \frac{1}{\sqrt{2r \ln T}} \left(\left(\frac{1}{\alpha} - \frac{r}{2} \right) \ln \ln T + \ln D \right). \quad (3.3)$$

b) *If $\gamma = \infty$, and $\alpha \in (0, 1]$, $\rho(t)$ is convex for $t \geq 0$ with $\lim_{t \rightarrow \infty} \rho(t) = 0$ and further $\rho(t) \ln t$ is monotone for large t , then with Φ the df of an $N(0, 1)$ random variable*

$$\lim_{T \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| \mathbb{P} \left\{ \frac{1}{\sqrt{\rho(T)}} \left(\sup_{t \in [0, T]} X_{n-r+1:n}(t) - \sqrt{1 - \rho(T)} b_{r,T} \right) \leq x \right\} - \Phi(x) \right| = 0.$$

c) *If $\gamma \in (0, \infty)$, then, with W an $N(0, 1)$ random variable*

$$\begin{aligned} \lim_{T \rightarrow \infty} \sup_{x \in \mathbb{R}} \left| \mathbb{P} \left\{ a_{r,T} \left(\sup_{t \in [0, T]} X_{n-r+1:n}(t) - b_{r,T} \right) \leq x \right\} \right. \\ \left. - \mathbb{E} \left\{ \exp \left(-e^{-(x+\gamma-\sqrt{2\gamma r}W)} \right) \right\} \right| = 0. \end{aligned}$$

The proof of Theorem 3.1 is presented in Appendix.

3.2. *Lower tail probability for order statistics processes.* The seminal contributions [Li and Shao \(2004, 2005\)](#) show that the investigation of the lower tail probability of Gaussian processes is of special interest in many applied fields, including the study of real zeros of random polynomials, the study of Gaussian pursuit problem, and the study of the first-passage time for the Slepian process. In this section, we aim at generalizing some results in [Li and Shao \(2004, 2005\)](#), by considering order statistics processes instead of Gaussian processes.

Our first result is concerned with extension of the celebrated Slepian inequality for order statistics processes. Let $\{Y(t), t \geq 0\}$ and $\{Z(t), t \geq 0\}$ be two centered Gaussian processes with a.s. continuous sample paths, and $\{Y_{r:n}(t), t \geq 0\}$, $\{Z_{r:n}(t), t \geq 0\}$ be the corresponding r th lower order statistics processes. Applying the standard discrete-continuous approximation technique (cf. [Adler and Taylor, 2007](#)) to [Corollary 2.3](#) one can easily verify the following proposition.

Proposition 3.2. *If for all $s, t \geq 0$*

$$\mathbb{E}\{Y(t)^2\} = \mathbb{E}\{Z(t)^2\} \quad \text{and} \quad \mathbb{E}\{Y(s)Y(t)\} \leq \mathbb{E}\{Z(s)Z(t)\},$$

then for any $T > 0$ and $u \in \mathbb{R}$ we have

$$\mathbb{P}\left\{\sup_{t \in [0, T]} Y_{r:n}(t) > u\right\} \geq \mathbb{P}\left\{\sup_{t \in [0, T]} Z_{r:n}(t) > u\right\}.$$

Remark 3.3. In view of [Proposition 3.2](#) for any $x \in \mathbb{R}$ (cf. [Li and Shao, 2004](#))

$$p_r(x) := \lim_{T \rightarrow \infty} \frac{1}{T} \ln \mathbb{P}\left\{\sup_{0 \leq t \leq T} Y_{r:n}(t) \leq x\right\} = \sup_{T > 0} \frac{1}{T} \ln \mathbb{P}\left\{\sup_{0 \leq t \leq T} Y_{r:n}(t) \leq x\right\}$$

exists and $p_r(x)$, $x \in \mathbb{R}$ is left-continuous, provided that $\{Y(t), t \geq 0\}$ is a centered stationary Gaussian processes with $\mathbb{E}\{Y(0)Y(t)\} \geq 0$ for all $t \geq 0$.

4. Proofs

Hereafter, we write $\stackrel{d}{=}$ for equality of the distribution functions. A vector $\mathbf{z} = (z_1, \dots, z_{dn})$ will also be denoted by

$$\mathbf{z} = (\mathbf{z}_1, \dots, \mathbf{z}_d), \quad \text{with } \mathbf{z}_i = (z_{i1}, \dots, z_{in}), \quad 1 \leq i \leq d,$$

where $z_{ij} = z_{(i-1)n+j}$, $1 \leq i \leq d$, $1 \leq j \leq n$. Note that for any $p = (i-1)n + j$, $q = (l-1)n + k$, $1 \leq i, l \leq d$, $1 \leq j, k \leq n$

$$\{p < q\} = \{i < l, \text{ or } i = l \text{ and } j < k\}.$$

Denote

$$\mathbf{z}/\mathbf{z}_i = (\mathbf{z}_1, \dots, \mathbf{z}_{i-1}, \mathbf{z}_{i+1}, \dots, \mathbf{z}_d), \quad 1 \leq i \leq d.$$

Furthermore, for any $\mathbf{x} \in \mathbb{R}^n$ we denote

$$\begin{aligned} \mathbf{x}/x_i &= (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n), \\ \frac{d\mathbf{x}}{dx_i} &= dx_1 dx_2 \cdots dx_{i-1} dx_{i+1} \cdots dx_n, \quad 1 \leq i \leq n, \end{aligned}$$

and for $1 \leq i < j \leq n$

$$\frac{d\mathbf{x}}{dx_i dx_j} = dx_1 dx_2 \cdots dx_{i-1} dx_{i+1} \cdots dx_{j-1} dx_{j+1} \cdots dx_n.$$

Proof of [Theorem 2.1](#): We shall first establish [\(2.1\)](#) by considering $r = 1$, $r = 2$ and $2 < r \leq n$ separately.

Case $r = 1$. Note that $\mathcal{X} \stackrel{d}{=} -\mathcal{X}$ for the standard Gaussian array \mathcal{X} . It follows from Theorem 2.1 in [Lu and Wang \(2014\)](#) that

$$\begin{aligned} |\Delta_{(1)}(\mathbf{u})| &= \left| \mathbb{P}\left\{ \bigcup_{i=1}^d \bigcap_{j=1}^n \{-Y_{ij} < -u_i\} \right\} - \mathbb{P}\left\{ \bigcup_{i=1}^d \bigcap_{j=1}^n \{-X_{ij} < -u_i\} \right\} \right| \\ &\leq \frac{1}{2\pi} \sum_{(i-1)n+j < (l-1)n+k} Q_{ij,lk} \exp\left(-\frac{u_i^2 + u_l^2}{2(1 + \rho_{ij,lk})}\right) \end{aligned}$$

establishing (2.1) for $r = 1$.

Next, by a standard approximation procedure we may assume that both $\Sigma^{(1)}$ and $\Sigma^{(0)}$ are positive definite. Let further $\mathcal{Z} = (Z_{ij})_{d \times n}$ be a standard Gaussian array with covariance matrix

$$\Gamma^h = h\Sigma^{(1)} + (1-h)\Sigma^{(0)} = (\delta_{ij,lk}^h)_{dn \times dn},$$

where by our notation $\delta_{ij,lk}^h = \mathbb{E}\{Z_{ij}Z_{lk}\}$. Clearly, Γ^h is also positive definite for any $h \in [0, 1]$. Denote below by $g_h(\mathbf{z})$ the probability density function (pdf) of \mathcal{Z} . It is known that the Plackett's partial differential equation holds as (see e.g., [Leadbetter et al., 1983](#), p. 82, or [Lu and Wang, 2014](#))

$$\frac{\partial g_h(\mathbf{z})}{\partial \delta_{ij,lk}^h} = \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{lk}}, \quad 1 \leq i, l \leq d, 1 \leq j, k \leq n, (i, j) \neq (l, k). \quad (4.1)$$

Case $r = 2$. Hereafter, we write $\boldsymbol{\lambda} = -\mathbf{u}$ and set

$$Q(\mathcal{Z}; \Gamma^h) = \mathbb{P}\{\mathbf{Z}_{(n-1)} > \boldsymbol{\lambda}\} = \int_{\bigcap_{i=1}^d \bigcup_{j,j'=1;j \neq j'}^n \{z_{ij} > \lambda_i, z_{ij'} > \lambda_i\}} g_h(\mathbf{z}) d\mathbf{z}. \quad (4.2)$$

Since $\mathbf{X}_{(2)} \stackrel{d}{=} -\mathbf{X}_{(n-1)}$ we have

$$\Delta_{(2)}(\mathbf{u}) = Q(\mathcal{Z}; \Gamma^1) - Q(\mathcal{Z}; \Gamma^0) = \int_0^1 \frac{\partial Q(\mathcal{Z}; \Gamma^h)}{\partial h} dh. \quad (4.3)$$

Note that the quantities $Q(\mathcal{Z}; \Gamma^h)$ and $g_h(\mathbf{z})$ depend on h only through the entries $\delta_{ij,lk}^h$ of Γ^h . Hence we have by (4.1)

$$\begin{aligned} \frac{\partial Q}{\partial h}(\mathcal{Z}; \Gamma^h) &= \sum_{(i-1)n+j < (l-1)n+k} \frac{\partial Q(\mathcal{Z}; \Gamma^h)}{\partial \delta_{ij,lk}^h} \frac{\partial \delta_{ij,lk}^h}{\partial h} \\ &= \sum_{(i-1)n+j < (l-1)n+k} (\sigma_{ij,lk}^{(1)} - \sigma_{ij,lk}^{(0)}) E_{il}(j, k), \end{aligned} \quad (4.4)$$

where

$$E_{il}(j, k) := \int_{\bigcap_{s=1}^d \bigcup_{t,t'=1;t \neq t'}^n \{z_{st} > \lambda_s, z_{st'} > \lambda_s\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{lk}} d\mathbf{z}. \quad (4.5)$$

Next, in order to establish (2.1) we shall show that

$$|E_{il}(j, k)| \leq \varphi(\lambda_i, \lambda_l; \delta_{ij,lk}^h), \quad (i-1)n+j < (l-1)n+k, \quad (4.6)$$

where $\varphi(\cdot, \cdot; \cdot)$ is the pdf of (Z_{ij}, Z_{lk}) , given by

$$\varphi(x, y; \delta_{ij,lk}^h) = \frac{1}{2\pi \sqrt{1 - (\delta_{ij,lk}^h)^2}} \exp\left(-\frac{x^2 - 2\delta_{ij,lk}^h xy + y^2}{2(1 - (\delta_{ij,lk}^h)^2)}\right), \quad x, y \in \mathbb{R}.$$

We consider below two sub-cases: a) $i = l$ and b) $i < l$.

a) Proof of (4.6) for $i = l$. Let

$$\begin{aligned} A'_i &= \bigcap_{s=1; s \neq i}^d \bigcup_{t, t'=1; t \neq t'}^n \{z_{st} > \lambda_s, z_{st'} > \lambda_s\} \\ &:= \{\mathbf{z}/\mathbf{z}_i \in \mathbb{R}^{(d-1)n} : \text{for any } 1 \leq s (\neq i) \leq d, \\ &\quad \text{there exist } 1 \leq t, t' \leq n : z_{st} > \lambda_s, z_{st'} > \lambda_s\}, \end{aligned} \quad (4.7)$$

$$\begin{aligned} A_i &= \bigcup_{t, t'=1; t \neq t'}^n \{z_{it} > \lambda_i, z_{it'} > \lambda_i\} \\ &:= \{\mathbf{z}_i \in \mathbb{R}^n : \text{there exist } 1 \leq t, t' \leq n : z_{it} > \lambda_i, z_{it'} > \lambda_i\}. \end{aligned} \quad (4.8)$$

We can rewrite $E_{ii}(j, k)$ as

$$E_{ii}(j, k) = \int_{A'_i} \int_{A_i} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}, \quad 1 \leq i \leq d, \quad 1 \leq j < k \leq n. \quad (4.9)$$

Next, we decompose the integral region A_i according to

- $a_1) \{z_{ij} > \lambda_i, z_{ik} > \lambda_i\} := \{\mathbf{z}_i \in \mathbb{R}^n : z_{ij} > \lambda_i, z_{ik} > \lambda_i\};$
- $a_2) \{z_{ij} > \lambda_i, z_{ik} \leq \lambda_i\} := \{\mathbf{z}_i \in \mathbb{R}^n : z_{ij} > \lambda_i, z_{ik} \leq \lambda_i\};$
- $a_3) \{z_{ij} \leq \lambda_i, z_{ik} > \lambda_i\} := \{\mathbf{z}_i \in \mathbb{R}^n : z_{ij} \leq \lambda_i, z_{ik} > \lambda_i\};$
- $a_4) \{z_{ij} \leq \lambda_i, z_{ik} \leq \lambda_i\} := \{\mathbf{z}_i \in \mathbb{R}^n : z_{ij} \leq \lambda_i, z_{ik} \leq \lambda_i\}.$

For case $a_1)$ we have

$$\int_{A_i \cap \{z_{ij} > \lambda_i, z_{ik} > \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}_i = \int_{\mathbb{R}^{n-2}} g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i) \frac{d\mathbf{z}_i}{dz_{ij} dz_{ik}}, \quad (4.10)$$

where $g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i)$ denotes a function of $dn - 2$ variables formed from $g_h(\mathbf{z})$ by putting $z_{ij} = \lambda_i, z_{ik} = \lambda_i$. Similarly, for cases $a_2)$ and $a_3)$

$$\begin{aligned} \int_{A_i \cap \{z_{ij} > \lambda_i, z_{ik} \leq \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}_i &= \int_{A_i \cap \{z_{ij} \leq \lambda_i, z_{ik} > \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}_i \\ &= - \int_{\bigcup_{t=1; t \neq j, k}^n \{z_{it} > \lambda_i\}} g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i) \frac{d\mathbf{z}_i}{dz_{ij} dz_{ik}}, \end{aligned} \quad (4.11)$$

where

$$\begin{aligned} &\bigcup_{t=1; t \neq j, k}^n \{z_{it} > \lambda_i\} \\ &:= \{\mathbf{z}_i / (z_{ij} z_{ik}) \in \mathbb{R}^{n-2} : \text{it exists } 1 \leq t (\neq j, k) \leq n \text{ such that } z_{it} > \lambda_i\}. \end{aligned}$$

Finally, for case $a_4)$

$$\begin{aligned} &\int_{A_i \cap \{z_{ij} \leq \lambda_i, z_{ik} \leq \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}_i \\ &= \int_{\bigcup_{t, t'=1; t, t' \neq j, k; t \neq t'}^n \{z_{it} > \lambda_i, z_{it'} > \lambda_i\}} g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i) \frac{d\mathbf{z}_i}{dz_{ij} dz_{ik}}, \end{aligned}$$

where

$$\begin{aligned} &\bigcup_{t, t'=1; t, t' \neq j, k; t \neq t'}^n \{z_{it} > \lambda_i, z_{it'} > \lambda_i\} \\ &:= \left\{ \frac{\mathbf{z}_i}{z_{ij} z_{ik}} \in \mathbb{R}^{n-2} : \text{there exist } 1 \leq t, t' (\neq j, k) \leq n \text{ such that } z_{it}, z_{it'} > \lambda_i \right\}. \end{aligned}$$

This together with (4.9)–(4.11) yields

$$\begin{aligned}
& \frac{E_{ii}(j, k)}{\varphi(\lambda_i, \lambda_i; \delta_{ij, ik}^h)} = \int_{A'_i} \int_{\mathbb{R}^{n-2} - \cup_{t=1, t \neq j, k}^n \{z_{it} > \lambda_i\}} \frac{g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i)}{\varphi(\lambda_i, \lambda_i; \delta_{ij, ik}^h)} \frac{dz}{dz_{ij} dz_{ik}} \\
& - \int_{A'_i} \int_{\cup_{t=1, t \neq j, k}^n \{z_{it} > \lambda_i\} - \cup_{t, t'=1, t, t' \neq j, k; t \neq t'}^d \{z_{it} > \lambda_i, z_{it'} > \lambda_i\}} \frac{g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i)}{\varphi(\lambda_i, \lambda_i; \delta_{ij, ik}^h)} \frac{dz}{dz_{ij} dz_{ik}} \\
& = \mathbb{P} \left\{ \cap_{s \neq i} \{Z_{s(n-1)} > \lambda_s\}, \mathbf{Z}_i'' \in \{w_{i1}'' = \infty\} \mid Z_{ij} = Z_{ik} = \lambda_i \right\} \\
& - \mathbb{P} \left\{ \cap_{s \neq i} \{Z_{s(n-1)} > \lambda_s\}, \mathbf{Z}_i'' \in \{w_{i1}'' \leq n, w_{i2}'' = \infty\} \mid Z_{ij} = Z_{ik} = \lambda_i \right\}, \quad (4.12)
\end{aligned}$$

with \mathbf{Z}_i'' the $(n-2)$ -dimensional components of \mathbf{Z}_i obtained by deleting Z_{ij} and Z_{ik} , and w_{i1}'', w_{i2}'' given by

$$\begin{aligned}
w_{i1}'' &= \inf\{t : 1 \leq t (\neq j, k) \leq n, z_{it} > \lambda_i\} \\
w_{i2}'' &= \inf\{t : w_{i1}'' < t (\neq j, k) \leq n, z_{it} > \lambda_i\}. \quad (4.13)
\end{aligned}$$

Hereafter we use the convention that $\inf\{\emptyset\} = \infty$. For instance,

$$\begin{aligned}
\{w_{i1}'' = \infty\} &= \{\mathbf{z}_i / (z_{ij} z_{ik}) \in \mathbb{R}^{n-2} : z_{it} \leq \lambda_i \text{ for all } 1 \leq t (\neq j, k) \leq n\} \\
\{w_{i1}'' \leq n, w_{i2}'' = \infty\} &= \{\mathbf{z}_i / (z_{ij} z_{ik}) \in \mathbb{R}^{n-2} : \text{it exists } 1 \leq l (\neq j, k) \leq n \\
&\text{such that } z_{il} > \lambda_i, \text{ and } z_{it} \leq \lambda_i \text{ for all } 1 \leq t (\neq j, k, l) \leq n\}.
\end{aligned}$$

Consequently, it follows thus from (4.12) that (4.6) holds for $i = l$.

b) Proof of (4.6) for $i < l$. Denote $A_{il}'' = \cap_{s=1, s \neq i, l}^n \cup_{t, t'=1, t \neq t'}^n \{z_{st} > \lambda_s, z_{st'} > \lambda_s\} \subset \mathbb{R}^{(d-2)n}$ parallel to (4.7) and recall A_i in (4.8). We have

$$E_{il}(j, k) = \int_{A_{il}''} \int_{A_i} \int_{A_i} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{lk}} dz. \quad (4.14)$$

Next, we decompose the integral region A_i according to $\{\mathbf{z}_i \in \mathbb{R}^n : z_{ij} > \lambda_i\}$ and $\{\mathbf{z}_i \in \mathbb{R}^n : z_{ij} \leq \lambda_i\}$. We have

$$\begin{aligned}
& \int_{A_i \cap \{\mathbf{z}_i \in \mathbb{R}^n : z_{ij} > \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{lk}} dz_i + \int_{A_i \cap \{\mathbf{z}_i \in \mathbb{R}^n : z_{ij} \leq \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{lk}} dz_i \\
& = - \int_{\cup_{t=1, t \neq j}^n \{z_{it} > \lambda_i\} - \cup_{t, t'=1, t, t' \neq j; t \neq t'}^n \{z_{it} > \lambda_i, z_{it'} > \lambda_i\}} \frac{\partial g_h(z_{ij} = \lambda_i)}{\partial z_{lk}} \frac{dz_i}{dz_{ij}} \\
& = - \int_{\{w_{i1}' \leq n, w_{i2}' = \infty\}} \frac{\partial g_h(z_{ij} = \lambda_i)}{\partial z_{lk}} \frac{dz_i}{dz_{ij}},
\end{aligned}$$

where w_{i1}', w_{i2}' are defined by (similar notation below for w_{l1}', w_{l2}' with respect to k instead of j)

$$\begin{aligned}
w_{i1}' &= \inf\{t : 1 \leq t (\neq j) \leq n, z_{it} > \lambda_i\} \\
w_{i2}' &= \inf\{t : w_{i1}' < t (\neq j) \leq n, z_{it} > \lambda_i\}. \quad (4.15)
\end{aligned}$$

Using similar arguments for the integral with region A_l , we have by (4.14)

$$\begin{aligned} E_{il}(j, k) &= \int_{A''_{il}} \int_{\{w'_{i1} \leq n, w'_{i2} = \infty\}} \int_{\{w'_{i1} \leq n, w'_{i2} = \infty\}} g_h(z_{ij} = \lambda_i, z_{lk} = \lambda_l) \frac{dz}{dz_{ij} dz_{lk}} \\ &= \varphi(\lambda_i, \lambda_l; \delta_{ij, lk}^h) \mathbb{P}\left\{\bigcap_{s \neq i, l} \{Z_{s(n-1)} > \lambda_s\}, \mathbf{Z}'_i \in \{w'_{i1} \leq n, w'_{i2} = \infty\},\right. \\ &\quad \left. \mathbf{Z}'_l \in \{w'_{l1} \leq n, w'_{l2} = \infty\} \mid \{Z_{ij} = \lambda_i, Z_{lk} = \lambda_l\}\right\}, \end{aligned} \quad (4.16)$$

where \mathbf{Z}'_i and \mathbf{Z}'_l are the $(n-1)$ -dimensional components of \mathbf{Z}_i and \mathbf{Z}_l obtained by deleting Z_{ij} and Z_{lk} , respectively. Consequently, by (4.12) and (4.16) the validity of (4.6) follows.

Next, by combining (4.3)–(4.6), the claim in (2.1) for $r = 2$ follows by the fact that (see Li and Shao, 2002)

$$\int_0^1 \varphi(\lambda_i, \lambda_l; \delta_{ij, lk}^h) dh \leq \frac{\arcsin(\sigma_{ij, lk}^{(1)}) - \arcsin(\sigma_{ij, lk}^{(0)})}{2\pi(\sigma_{ij, lk}^{(1)} - \sigma_{ij, lk}^{(0)})} \exp\left(-\frac{\lambda_i^2 + \lambda_l^2}{2(1 + \rho_{ij, lk})}\right). \quad (4.17)$$

Case $2 < r \leq n$. Letting $\tilde{Q}(\mathcal{Z}; \Gamma^h) = \mathbb{P}\{\mathbf{Z}_{(n-r+1)} > \boldsymbol{\lambda}\}$ we have

$$\Delta_{(r)}(\mathbf{u}) = \int_0^1 dh \left(\sum_{(i-1)n+j < (l-1)n+k} (\sigma_{ij, lk}^{(1)} - \sigma_{ij, lk}^{(0)}) \tilde{E}_{il}(j, k) \right), \quad (4.18)$$

where

$$\tilde{E}_{il}(j, k) := \int_{\bigcap_{s=1}^d \cup_{t_1, \dots, t_r=1; t_l \neq t_j} \{z_{st_1} > \lambda_s, \dots, z_{st_r} > \lambda_s\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{lk}} d\mathbf{z}.$$

With the aid of (4.17), it suffices to show that

$$\left| \tilde{E}_{il}(j, k) \right| \leq \varphi(\lambda_i, \lambda_l; \delta_{ij, lk}^h), \quad (i-1)n+j < (l-1)n+k. \quad (4.19)$$

Similarly as above, two sub-cases : a) $i = l$ and b) $i < l$ need to be considered separately.

a) Proof of (4.19) for $i = l$. Similarly to $E_{ii}(j, k)$, we rewrite $\tilde{E}_{ii}(j, k)$ as

$$\tilde{E}_{ii}(j, k) = \int_{\tilde{A}'_i} \int_{\tilde{A}_i} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}, \quad (4.20)$$

with

$$\begin{aligned} \tilde{A}'_i &= \bigcap_{s=1; s \neq i}^d \cup_{t_1, \dots, t_r=1; t_l \neq t_j} \{\mathbf{z}/\mathbf{z}_i \in \mathbb{R}^{(d-1)n} : z_{st_1} > \lambda_s, \dots, z_{st_r} > \lambda_s\}, \\ \tilde{A}_i &= \cup_{t_1, \dots, t_r=1; t_l \neq t_j} \{\mathbf{z}_i \in \mathbb{R}^n : z_{it_1} > \lambda_i, \dots, z_{it_r} > \lambda_i\}. \end{aligned}$$

Next, we decompose the integral region \tilde{A}_i according to the four cases $a_1)$ – $a_4)$ as introduced for A_i (see the lines right above (4.10)).

For case $a_1)$

$$\int_{\tilde{A}_i \cap \{z_{ij} > \lambda_i, z_{ik} > \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}_i = \int_{\{w''_{i, r-2} \leq n\}} g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i) \frac{dz_i}{dz_{ij} dz_{ik}}, \quad (4.21)$$

where w''_{i1} is given by (4.13) and (notation: $w''_{i, t} = w''_{it}$)

$$w''_{it} = \inf\{s : w''_{i, t-1} < s (\neq j, k) \leq n, z_{is} > \lambda_i\}, \quad 2 \leq t \leq r, \quad 1 \leq i \leq d.$$

Next, for cases $a_2)$ and $a_3)$

$$\begin{aligned} \int_{\tilde{A}_i \cap \{z_{ij} > \lambda_i, z_{ik} \leq \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}_i &= \int_{\tilde{A}_i \cap \{z_{ij} \leq \lambda_i, z_{ik} > \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}_i \\ &= - \int_{\{w''_{i,r-1} \leq n\}} g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i) \frac{d\mathbf{z}_i}{dz_{ij} dz_{ik}}. \end{aligned} \quad (4.22)$$

Finally, for case $a_4)$

$$\int_{\tilde{A}_i \cap \{z_{ij} \leq \lambda_i, z_{ik} \leq \lambda_i\}} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{ik}} d\mathbf{z}_i = \int_{\{w''_{i,r} \leq n\}} g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i) \frac{d\mathbf{z}_i}{dz_{ij} dz_{ik}}.$$

This together with (4.20)–(4.22) yields that

$$\begin{aligned} \frac{\tilde{E}_{ii}(j, k)}{\varphi(\lambda_i, \lambda_i; \delta_{ij,ik}^h)} &= \int_{\tilde{A}'_i} \int_{\{w''_{i,r-2} \leq n, w''_{i,r-1} = \infty\}} \frac{g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i)}{\varphi(\lambda_i, \lambda_i; \delta_{ij,ik}^h)} \frac{d\mathbf{z}}{dz_{ij} dz_{ik}} \\ &\quad - \int_{\tilde{A}'_i} \int_{\{w''_{i,r-1} \leq n, w''_{i,r} = \infty\}} \frac{g_h(z_{ij} = \lambda_i, z_{ik} = \lambda_i)}{\varphi(\lambda_i, \lambda_i; \delta_{ij,ik}^h)} \frac{d\mathbf{z}}{dz_{ij} dz_{ik}} \\ &= \mathbb{P}\left\{ \cap_{s \neq i} \{Z_{s(n-r+1)} > \lambda_s\}, \mathbf{Z}''_i \in \{w''_{i,r-2} \leq n, w''_{i,r-1} = \infty\} \mid Z_{ij} = Z_{ik} = \lambda_i \right\} \\ &\quad - \mathbb{P}\left\{ \cap_{s \neq i} \{Z_{s(n-r+1)} > \lambda_s\}, \mathbf{Z}''_i \in \{w''_{i,r-1} \leq n, w''_{i,r} = \infty\} \mid Z_{ij} = Z_{ik} = \lambda_i \right\} \end{aligned} \quad (4.23)$$

establishing (4.19) for $i = l$.

b) Proof of (4.19) for $i < l$. By \tilde{A}_i as in (4.20) and

$$\tilde{A}''_{il} = \cap_{s=1; s \neq i, l}^d \cup_{t_1, \dots, t_r=1; t_l \neq t_j}^n \{\mathbf{z}/(\mathbf{z}_i \mathbf{z}_l) : z_{st_1} > \lambda_s, \dots, z_{st_r} > \lambda_s\},$$

we have

$$\tilde{E}_{il}(j, k) = \int_{\tilde{A}''_{il}} \int_{\tilde{A}_i} \int_{\tilde{A}_l} \frac{\partial^2 g_h(\mathbf{z})}{\partial z_{ij} \partial z_{lk}} d\mathbf{z}. \quad (4.24)$$

By decomposing the integral regions \tilde{A}_i and \tilde{A}_l according to $z_{ij} >, \leq \lambda_i$ and $z_{lk} >, \leq \lambda_l$ in \mathbb{R}^n , respectively, we obtain by similar arguments as for $E_{il}(j, k)$ that

$$\begin{aligned} \frac{\tilde{E}_{il}(j, k)}{\varphi(\lambda_i, \lambda_l; \delta_{ij,lk}^h)} &= \mathbb{P}\left\{ \cap_{s \neq i, l} \{Z_{s(n-r+1)} > \lambda_s\}, \mathbf{Z}'_i \in \{w'_{i,r-1} \leq n, w'_{i,r} = \infty\}, \right. \\ &\quad \left. \mathbf{Z}'_l \in \{w'_{l,r-1} \leq n, w'_{l,r} = \infty\} \mid Z_{ij} = Z_{lk} = \lambda_l \right\}, \end{aligned} \quad (4.25)$$

where w'_{i1} is introduced in (4.15) and (similar notation for w'_{lt})

$$w'_{it} = \inf\{s : w'_{i,t-1} < s (\neq j) \leq n, z_{is} > \lambda_i\}, \quad 2 \leq t \leq r, \quad 1 \leq i \leq d.$$

It follows then from (4.25) that (4.19) holds. Consequently, the desired result (2.1) follows for $2 < r \leq n$.

Finally, in view of (2.2) we see that the indices over the sum in (4.4) and (4.18) are simplified to $1 \leq i < l \leq d, 1 \leq j, k \leq n$. Then the claim in (2.3) follows immediately from (4.16), (4.17) and (4.25). This completes the proof of Theorem 2.1. \square

Proof of Theorem 2.4: It is sufficient to prove (2.6) since it implies the second result of Theorem 2.4. Note that (2.6) holds for $r = 1$ by the argument of Lemma 12 in Dębicki et al. (2014a), and in view of (2.5), (4.3), (4.4) and (4.20), we have

$$\Delta_{(r)}(\mathbf{u}) = \begin{cases} n \sum_{1 \leq i < l \leq d} (\sigma_{il}^{(1)} - \sigma_{il}^{(0)}) \int_0^1 E_{il} dh, & r = 2, \\ n \sum_{1 \leq i < l \leq d} (\sigma_{il}^{(1)} - \sigma_{il}^{(0)}) \int_0^1 \tilde{E}_{il} dh, & 2 < r \leq n, \end{cases} \quad (4.26)$$

where $E_{il} := E_{il}(1, 1)$, $\tilde{E}_{il} := \tilde{E}_{il}(1, 1)$ with $E_{il}(1, 1)$, $\tilde{E}_{il}(1, 1)$ given by (4.5) and (4.16), respectively. Therefore, we shall present next the proofs of (2.6) for a) $r = 2$ and b) $2 < r \leq n$, and assume in the following that $\sigma_{il}^{(1)} \neq \sigma_{il}^{(0)}$ for all $1 \leq i < l \leq d$.

a) Proof of (2.6) for $r = 2$. It follows by (2.5) and (4.16) that, with $\delta_{il}^h := \delta_{i1, l1}^h$ (recall $\lambda_i := -u_i$, $1 \leq i \leq d$)

$$\begin{aligned} 0 &\leq \frac{E_{il}}{\varphi(-u_i, -u_l; \delta_{il}^h)} \\ &\leq \mathbb{P}\left\{\mathbf{Z}'_i \in \{w'_{i1} \leq n, w'_{i2} = \infty\}, \mathbf{Z}'_l \in \{w'_{l1} \leq n, w'_{l2} = \infty\}\right\}. \end{aligned} \quad (4.27)$$

Note that hereafter w'_{i1}, w'_{i2} and w'_{l1}, w'_{l2} are defined as in (4.15) with respect to $j = k = 1$.

Next, let $(\tilde{Z}_i, \tilde{Z}_l)$ be a bivariate standard normal random vector with correlation $|\delta_{il}^h|$ and $u = \min_{1 \leq i \leq d} u_i > 0$. It follows by Slepian's inequality in Slepian (1962) and Lemma 2.3 in Pickands (1969) that, for $1 \leq j, k \leq n$

$$\begin{aligned} \mathbb{P}\left\{Z_{ij} < -u_i, Z_{lk} < -u_l\right\} &\leq \mathbb{P}\left\{\tilde{Z}_i < -u_i, \tilde{Z}_l < -u_l\right\} \\ &\leq \mathbb{P}\left\{-\tilde{Z}_i > u, -\tilde{Z}_l > u\right\} \leq \frac{(1 + |\delta_{il}^h|)^2}{u^2} \varphi(u, u; |\delta_{il}^h|) \end{aligned} \quad (4.28)$$

implying thus

$$\begin{aligned} &\mathbb{P}\left\{\mathbf{Z}'_i \in \{(w'_{i1}, w'_{i2}) = (2, \infty)\}, \mathbf{Z}'_l \in \{(w'_{l1}, w'_{l2}) = (2, \infty)\}\right\} \\ &= \mathbb{P}\left\{Z_{i2} > -u_i, Z_{l2} > -u_l\right\} \prod_{j=3}^n \mathbb{P}\left\{Z_{ij} \leq -u_i, Z_{lj} \leq -u_l\right\} \\ &\leq \left(\frac{(1 + |\delta_{il}^h|)^2}{u^2} \varphi(u, u; |\delta_{il}^h|)\right)^{n-2} \end{aligned}$$

and

$$\begin{aligned} &\mathbb{P}\left\{\mathbf{Z}'_i \in \{(w'_{i1}, w'_{i2}) = (3, \infty)\}, \mathbf{Z}'_l \in \{(w'_{l1}, w'_{l2}) = (2, \infty)\}\right\} \\ &= \mathbb{P}\left\{Z_{i2} < -u_i, Z_{l2} > -u_l, Z_{i3} > -u_i, Z_{l3} < -u_l\right\} \prod_{j=4}^n \mathbb{P}\left\{Z_{ij} < -u_i, Z_{lj} < -u_l\right\} \\ &\leq \left(\frac{(1 + |\delta_{il}^h|)^2}{u^2} \varphi(u, u; |\delta_{il}^h|)\right)^{n-2}. \end{aligned}$$

Similarly, we may consider all $(n-1)^2$ cases in (4.27) for $w'_{i1} = w'_{l1}$ and $w'_{i1} \neq w'_{l1}$. Therefore, using further (4.6) in Li and Shao (2002) we have

$$\begin{aligned} E_{il} &\leq (n-1)^2 \left(\frac{(1+|\delta_{il}^h|)^2}{u^2} \varphi(u, u; |\delta_{il}^h|) \right)^{n-2} \varphi(-u_i, -u_l; \delta_{il}^h) \\ &\leq \frac{(n-1)^2}{(2\pi)^{n-1}} u^{-2(n-2)} \frac{(1+|\delta_{il}^h|)^{2(n-2)}}{(1-|\delta_{il}^h|^2)^{(n-1)/2}} \exp\left(-\frac{(n-1)u^2}{1+|\delta_{il}^h|}\right). \end{aligned}$$

Consequently, by (4.26) we have

$$\begin{aligned} \Delta_{(2)}(\mathbf{u}) &\leq n \sum_{1 \leq i < l \leq d} (\sigma_{il}^{(1)} - \sigma_{il}^{(0)})_+ \int_0^1 E_{il} dh \\ &\leq \frac{n(n-1)^2}{(2\pi)^{n-1}} u^{-2(n-2)} \sum_{1 \leq i < l \leq d} (\sigma_{il}^{(1)} - \sigma_{il}^{(0)})_+ \\ &\quad \times \exp\left(-\frac{(n-1)u^2}{1+\rho_{il}}\right) \int_0^1 \frac{(1+|\delta_{il}^h|)^{2(n-2)}}{(1-|\delta_{il}^h|^2)^{(n-1)/2}} dh \\ &= \frac{n(n-1)^2}{(2\pi)^{n-1}} u^{-2(n-2)} \sum_{1 \leq i < l \leq d} (A_{il}^{(2)})_+ \exp\left(-\frac{(n-1)u^2}{1+\rho_{il}}\right). \end{aligned}$$

The last step follows since $\rho_{il} = \max(|\sigma_{il}^{(0)}|, |\sigma_{il}^{(1)}|) \geq \delta_{il}^h$ and (recall $\delta_{il}^h = h(\sigma_{il}^{(1)} - \sigma_{il}^{(0)}) + \sigma_{il}^{(0)}$)

$$\int_0^1 \frac{(1+|\delta_{il}^h|)^{2(n-2)}}{(1-|\delta_{il}^h|^2)^{(n-1)/2}} dh = \frac{1}{\sigma_{il}^{(1)} - \sigma_{il}^{(0)}} \int_{\sigma_{il}^{(0)}}^{\sigma_{il}^{(1)}} \frac{(1+|h|)^{2(n-2)}}{(1-h^2)^{(n-1)/2}} dh. \quad (4.29)$$

b) Proof of (2.6) for $2 < r \leq n$. Clearly, from (4.25) we have $\tilde{E}_{il} \geq 0$. Further, similar arguments as for E_{il} (consider the number of $w'_{it} = w'_{is}$, $s, t < r$) yield that

$$\begin{aligned} \frac{\tilde{E}_{il}}{\varphi(-u_i, -u_l; \delta_{il}^h)} &\leq \mathbb{P}\left\{ \mathbf{Z}'_i \in \{w'_{i,r-1} \leq n, w'_{ir} = \infty\}, \mathbf{Z}'_l \in \{w'_{l,r-1} \leq n, w'_{lr} = \infty\} \right\} \\ &\leq (c_{n-1,r-1})^2 \left(\frac{(1+|\delta_{il}^h|)^2}{u^2} \varphi(u, u; |\delta_{il}^h|) \right)^{n-r}. \end{aligned}$$

Consequently, the claim in (2.6) for $2 < r \leq n$ follows, establishing the proof. \square

We give next a result which extends Lemma 2.3 in Pickands (1969) needed for the proof of Proposition 2.5.

Lemma 4.1. *Let (X, Y) be a bivariate standard normal random vector with correlation $\rho \in (-1, 1)$. For any $x, y > 0$, if $\rho < \max(x/y, y/x)$, then*

$$\mathbb{P}\{X > x, Y > y\} \leq \frac{2(1+\rho)^2(1-\rho)}{(x+y)\min(x-\rho y, y-\rho x)} \varphi(x, y; \rho). \quad (4.30)$$

Proof: The proof follows with similar arguments as in [Pickands \(1969\)](#). By a change of variable $x' = x + u/x, y' = y + v/y$, we have

$$\begin{aligned}
\mathbb{P}\{X > x, Y > y\} &= \int_x^\infty \int_y^\infty \varphi(x', y'; \rho) dx' dy' \\
&= \frac{\varphi(x, y; \rho)}{xy} \int_0^\infty \int_0^\infty \exp\left(-\frac{u(1 - \rho y/x) + v(1 - \rho x/y)}{1 - \rho^2}\right) \\
&\quad \times \exp\left(-\frac{(u/x)^2 - 2\rho(u/x)(v/y) + (y/v)^2}{2(1 - \rho^2)}\right) dudv \\
&\leq \frac{\varphi(x, y; \rho)}{xy} \int_0^\infty \int_0^\infty \exp\left(-\frac{(u/x)(x - \rho y) + (v/y)(y - \rho x)}{1 - \rho^2}\right) \\
&\quad \times \exp\left(-\frac{(u/x - y/v)^2}{2(1 - \rho^2)}\right) dudv \\
&=: \frac{\varphi(x, y; \rho)}{xy} J(x, y, \rho).
\end{aligned}$$

Next, let $s = ((u/x)(x - \rho y) + (v/y)(y - \rho x))/(1 - \rho^2), t = (u/x - v/y)/\sqrt{1 - \rho^2}$. Clearly,

$$\left| \frac{\frac{\partial s(u, v)}{\partial u}}{\frac{\partial t(u, v)}{\partial u}} \quad \frac{\partial s(u, v)}{\partial v} \quad \frac{\partial t(u, v)}{\partial v} \right| = -\frac{x + y}{xy(1 + \rho)\sqrt{1 - \rho^2}}. \quad (4.31)$$

Further, since $\rho < \max(x/y, y/x)$, we have

$$\frac{s(1 - \rho^2)}{\min(x - \rho y, y - \rho x)} \geq \frac{u}{x} + \frac{v}{y} \geq |t|\sqrt{1 - \rho^2}, \quad -\infty < t < \infty.$$

Consequently, with $m_{x, y} := \min(x - \rho y, y - \rho x)/\sqrt{1 - \rho^2}$

$$\begin{aligned}
J(x, y, \rho) &\leq \frac{xy(1 + \rho)\sqrt{1 - \rho^2}}{x + y} \int_{-\infty}^\infty e^{-t^2/2} \int_{|t|m_{x, y}}^\infty e^{-s} ds dt \\
&= \frac{2xy(1 + \rho)\sqrt{1 - \rho^2}}{x + y} \int_0^\infty \exp\left(-\frac{t^2}{2} - m_{x, y}t\right) dt \\
&= \frac{2xy(1 + \rho)\sqrt{1 - \rho^2}}{x + y} \frac{(1 - \Phi(m_{x, y}))}{\varphi(m_{x, y})},
\end{aligned}$$

where $\varphi(x)$ and $\Phi(x)$ are the pdf and df of the standard normal random variable, respectively. Hence the well-known inequality

$$1 - \Phi(x) \leq \varphi(x)/x, \quad x > 0 \quad (4.32)$$

establishes the proof. \square

Proof of Proposition 2.5: We adopt the same notation as in Theorem 2.4. It follows by Slepian's inequality and (4.32) that

$$\begin{aligned} \mathbb{P}\{Z_{ij} < -u_i, Z_{lk} < -u_l\} &\leq \mathbb{P}\{\tilde{Z}_i < -u_i, \tilde{Z}_l < -u_l\} \\ &\leq \mathbb{P}\left\{\frac{\tilde{Z}_i + \tilde{Z}_l}{\sqrt{2(1 + |\delta_{il}^h|)}} > \frac{u_i + u_l}{\sqrt{2(1 + |\delta_{il}^h|)}}\right\} \\ &\leq \frac{\sqrt{2(1 + |\delta_{il}^h|)}}{u_i + u_l} \varphi\left(\frac{u_i + u_l}{\sqrt{2(1 + |\delta_{il}^h|)}}\right). \end{aligned} \quad (4.33)$$

Furthermore, we have by Lemma 4.1

$$\begin{aligned} \mathbb{P}\{Z_{ij} < -u_i, Z_{lk} < -u_l\} &\leq \mathbb{P}\{-\tilde{Z}_i > u_i, -\tilde{Z}_l > u_l\} \\ &\leq \frac{2(1 + |\delta_{il}^h|)^2(1 - |\delta_{il}^h|)}{(u_i + u_l) \min(u_i - |\delta_{il}^h|u_l, u_l - |\delta_{il}^h|u_i)} \varphi(u_i, u_l; |\delta_{il}^h|) \\ &\leq \frac{2(1 + |\delta_{il}^h|)^2(1 - |\delta_{il}^h|)}{(u_i + u_l) \min(u_i - \rho_{il}u_l, u_l - \rho_{il}u_i)} \varphi(u_i, u_l; |\delta_{il}^h|). \end{aligned} \quad (4.34)$$

Hence (2.7) and (2.8) are established by replacing (4.28) with (4.33) and (4.34), respectively, and utilising similar arguments as in the proof of Theorem 2.4. \square

Proof of Proposition 2.6: The lower bound follows directly from Corollary 2.3. Next we focus on the upper bound. We present below the proof for $r = 2$. Hereafter, we adopt the same notation as in the proof of Theorem 2.1. Further, define

$$f(h) = \exp\left(\sum_{\substack{1 \leq i < l \leq d \\ 1 \leq j, k \leq n}} \frac{1}{H((u_i + u_l)/2)} \mathcal{C}_{ij, lk}^h\right), \quad h \in [0, 1],$$

where

$$\mathcal{C}_{ij, lk}^h = \ln\left(\frac{\pi - 2 \arcsin(\sigma_{ij, lk}^{(0)})}{\pi - 2 \arcsin(\delta_{ij, lk}^h)}\right), \quad H(x) = \sqrt{2\pi}e^{x^2/2} \mathbb{E}\{(W + x)_+\},$$

with W an $N(0, 1)$ random variable. It suffices to show that $Q(\mathcal{Z}; \Gamma^h)/f(h)$ is non-increasing in h , i.e.,

$$\frac{\partial Q(\mathcal{Z}; \Gamma^h)/\partial h}{Q(\mathcal{Z}; \Gamma^h)} \leq \frac{\partial f(h)/\partial h}{f(h)}, \quad h \in [0, 1]. \quad (4.35)$$

We have

$$\frac{\partial f(h)/\partial h}{f(h)} = \sum_{\substack{1 \leq i < l \leq d \\ 1 \leq j, k \leq n}} \frac{2(\sigma_{ij, lk}^{(1)} - \sigma_{ij, lk}^{(0)})}{(\pi - 2 \arcsin(\delta_{ij, lk}^h)) \sqrt{1 - (\delta_{ij, lk}^h)^2} H((u_i + u_l)/2)} \quad (4.36)$$

and by (4.4)

$$\frac{\partial Q(\mathcal{Z}; \Gamma^h)}{\partial h} = \sum_{\substack{1 \leq i < l \leq d \\ 1 \leq j, k \leq n}} (\sigma_{ij, lk}^{(1)} - \sigma_{ij, lk}^{(0)}) E_{il}(j, k). \quad (4.37)$$

Therefore, by the assumption that $0 < \sigma_{ij,lk}^{(0)} \leq \sigma_{ij,lk}^{(1)} < 1$ for $1 \leq i < l \leq d, 1 \leq j, k \leq n$, it is sufficient to show that

$$E_{il}(j, k) \leq \frac{2Q(\mathcal{Z}; \Gamma^h)}{(\pi - 2 \arcsin(\delta_{ij,lk}^h)) \sqrt{1 - (\delta_{ij,lk}^h)^2}} \frac{1}{H((u_i + u_l)/2)}. \quad (4.38)$$

From (4.16) we have (recall $\mathbf{u} = -\boldsymbol{\lambda}$)

$$\begin{aligned} & \frac{E_{il}(j, k)}{\varphi(u_i, u_l; \delta_{ij,lk}^h)} \\ & \leq \mathbb{P} \left\{ \bigcap_{s \neq i, l} \{Z_{s(n-1)} > \lambda_s\}, (\mathbf{Z}'_i, \mathbf{Z}'_l) \in \{w'_{i1}, w'_{l1} \leq n\} \mid (Z_{ij}, Z_{lk}) = (\lambda_i, \lambda_l) \right\} \\ & = \mathbb{P} \left\{ \bigcap_{s \neq i, l} \{Z_{s(2)} < u_s\}, (\mathbf{Z}'_i, \mathbf{Z}'_l) \in \{v'_{i1}, v'_{l1} \leq n\} \mid (Z_{ij}, Z_{lk}) = (u_i, u_l) \right\}, \end{aligned} \quad (4.39)$$

where v'_{i1}, v'_{l1} are defined by

$$v'_{i1} = \inf \{t : 1 \leq t (\neq j) \leq n, z_{it} < u_i\}, \quad v'_{l1} = \inf \{t : 1 \leq t (\neq k) \leq n, z_{lt} < u_l\}.$$

Define next

$$T_{ij} = \frac{(Z_{ij} - u_i) - \delta_{ij,lk}^h(Z_{lk} - u_l)}{1 - (\delta_{ij,lk}^h)^2}, \quad T_{lk} = \frac{(Z_{lk} - u_l) - \delta_{ij,lk}^h(Z_{ij} - u_i)}{1 - (\delta_{ij,lk}^h)^2}.$$

Since (Z_{ij}, Z_{lk}) is a bivariate Gaussian random vector with $N(0, 1)$ marginals and correlation $\delta_{ij,lk}^h$, we have

$$\mathbb{E} \{T_{ij} Z_{ij}\} = \mathbb{E} \{T_{lk} Z_{lk}\} = 1, \quad \mathbb{E} \{T_{ij} Z_{lk}\} = \mathbb{E} \{T_{ij} Z_{lk}\} = 0.$$

Then it follows that the random vectors

$$\begin{aligned} \mathbf{Z}_v^* &= (Z_{vw} - \delta_{vw,ij}^h T_{ij} - \delta_{vw,lk}^h T_{lk}, 1 \leq w \leq n), \quad v \neq i, l \\ \mathbf{Z}_i^{*'} &= (Z_{it} - \delta_{it,ij}^h T_{ij} - \delta_{it,lk}^h T_{lk}, 1 \leq t (\neq j) \leq n) \\ \mathbf{Z}_l^{*'} &= (Z_{lt} - \delta_{it,ij}^h T_{ij} - \delta_{lt,lk}^h T_{lk}, 1 \leq t (\neq k) \leq n) \end{aligned}$$

are independent of (Z_{ij}, Z_{lk}) and further independent of (T_{ij}, T_{lk}) . Thus, by (4.39) and the fact that $0 \leq \delta_{ij,lk}^h < 1, 1 \leq i < l \leq d, 1 \leq j, k \leq n, h \in [0, 1]$, we have as in Lemma 2.1 in Yan (2009)

$$\begin{aligned} & E_{il}(j, k) \frac{\mathbb{P} \{T_{ij} < 0, T_{lk} < 0\}}{\varphi(u_i, u_l; \delta_{ij,lk}^h)} \\ & \leq \mathbb{P} \left\{ \bigcap_{s \neq i, l} \{Z_{s(2)}^* < u_s\}, \mathbf{Z}_i^{*'} \in \{v'_{i1} \leq n\}, \mathbf{Z}_l^{*'} \in \{v'_{l1} \leq n\}, T_{ij} < 0, T_{lk} < 0 \right\} \\ & \leq \mathbb{P} \left\{ \bigcap_{s \neq i, l} \{Z_{s(2)} < u_s\}, \mathbf{Z}'_i \in \{v'_{i1} \leq n\}, \mathbf{Z}'_l \in \{v'_{l1} \leq n\}, Z_{ij} < u_i, Z_{lk} < u_l \right\} \\ & = Q(\mathcal{Z}; \Gamma^h). \end{aligned} \quad (4.40)$$

Moreover, by Lemma 2.2 in Yan (2009)

$$\frac{\mathbb{P} \{T_{ij} < 0, T_{lk} < 0\}}{\varphi(u_i, u_l; \delta_{ij,lk}^h)} \geq \frac{\pi - 2 \arcsin(\delta_{ij,lk}^h)}{2} \sqrt{1 - (\delta_{ij,lk}^h)^2} H\left(\frac{u_i + u_l}{2}\right),$$

which together with (4.40) implies (4.38), hence the proof for $r = 2$ is complete.

For $2 < r \leq n$, we need to show that (4.38) holds for $\tilde{E}_{il}(j, k)$. This follows by similar arguments as for $r = 2$, using the inequality (4.25) instead of (4.16). \square

5. Appendix

We give the detailed proof of Theorem 3.1, which is based on the two lemmas below. For notational simplicity, we set $q = q(u) = u^{-2/\alpha}$, $u > 0$ and write $[x]$ for the integer part of $x \in \mathbb{R}$.

Lemma 5.1. *Under the assumptions of Theorem 3.1 with $\gamma = 0$, for any $a, T > 0$ and any positive integer $k \leq n$ we have*

$$\limsup_{u \rightarrow \infty} \sum_{j=[T/(aq)]}^{[\varepsilon/\mathbb{P}\{X_{k:n}(0) > u\}]} \mathbb{P} \left\{ X_{k:n}(aqj) > u \mid X_{k:n}(0) > u \right\} \rightarrow 0, \quad \varepsilon \downarrow 0. \quad (5.1)$$

Proof: By Lemma 2 in Dębicki et al. (2015a) (see the proof of (3.5) therein), for sufficiently large u

$$a_u(t) := \mathbb{P} \left\{ X_{n-r+1:n}(t) > u \mid X_{n-r+1:n}(0) > u \right\} \leq 2\mathbb{P} \left\{ X_{1:r}(t) > u \mid X_{1:r}(0) > u \right\}.$$

Since further $X(t) - \rho(t)X(0)$, $t > 0$ is independent of $X(0)$, we have for some constant $K > 0$ (below the value of K might change from line to line)

$$\begin{aligned} a_u(t) &\leq 2^{r+1} \left(\mathbb{P} \left\{ X(t) > X(0) > u \mid X(0) > u \right\} \right)^r \\ &\leq 2^{r+1} \left(\mathbb{P} \left\{ X(t) - \rho(t)X(0) > u(1 - \rho(t)) \mid X(0) > u \right\} \right)^r \\ &= 2^{r+1} \left(1 - \Phi \left(u \sqrt{\frac{1 - \rho(t)}{1 + \rho(t)}} \right) \right)^r \\ &\leq K u^{-r} \left(\frac{1 - |\rho(t)|}{1 + |\rho(t)|} \right)^{-r/2} \exp \left(-\frac{r u^2}{2} \frac{1 - |\rho(t)|}{1 + |\rho(t)|} \right), \end{aligned} \quad (5.2)$$

where the last inequality follows by (4.32).

Next, let g be a positive function such that

$$\lim_{u \rightarrow \infty} g(u) = \infty, \quad |\rho(g(u))| = u^{-2}.$$

It follows from $u^{-2} \ln g(u) = o(1)$ that $g(u) \leq \exp(\epsilon' u^2)$ for some $0 < \epsilon' < r/2(1 - |\rho(T)|)/(1 + |\rho(T)|)$ and sufficiently large u (recall that $|\rho(T)| < 1$; see Leadbetter et al. (1983), p.86). Next, we split the sum in (5.1) at $aqj = g(u)$. The first term is

$$\begin{aligned} &\sum_{j=[T/(aq)]}^{[g(u)/(aq)]} \mathbb{P} \left\{ X_{n-r+1:n}(aqj) > u \mid X_{n-r+1:n}(0) > u \right\} \\ &\leq K \frac{g(u)}{aq} u^{-r} \left(\frac{1 - |\rho(T)|}{1 + |\rho(T)|} \right)^{-r/2} \exp \left(-\frac{r u^2}{2} \frac{1 - |\rho(T)|}{1 + |\rho(T)|} \right) \\ &\leq K u^{2/\alpha - r} \exp \left(\epsilon' u^2 - \frac{r u^2}{2} \frac{1 - |\rho(T)|}{1 + |\rho(T)|} \right) \rightarrow 0, \quad u \rightarrow \infty. \end{aligned}$$

For the remaining term, it follows by Lemma 1 in [Dębicki et al. \(2015a\)](#) and (5.2)

$$\begin{aligned}
& \sum_{j=[g(u)/(aq)]}^{[\varepsilon/\mathbb{P}\{X_{n-r+1:n}(0) > u\}]} \mathbb{P}\left\{X_{n-r+1:n}(aqj) > u \mid X_{n-r+1:n}(0) > u\right\} \\
& \leq K \frac{\varepsilon}{\mathbb{P}\{X_{n-r+1:n}(0) > u\}} u^{-r} \left(\frac{1-u^{-2}}{1+u^{-2}}\right)^{-r/2} \exp\left(-\frac{ru^2}{2} \frac{1-u^{-2}}{1+u^{-2}}\right) \\
& \leq K\varepsilon \exp\left(-\frac{ru^2}{2} \left(\frac{1-u^{-2}}{1+u^{-2}} - 1\right)\right) \\
& \leq K\varepsilon, \quad u \rightarrow \infty.
\end{aligned}$$

Therefore, the claim follows by letting $\varepsilon \downarrow 0$. \square

Next, with the notation as in (3.2) we set

$$T = T(u) = \frac{1}{c_{n,r}\mathcal{A}_{r,\alpha}} (2\pi)^{\frac{r}{2}} u^{r-\frac{2}{\alpha}} \exp\left(\frac{ru^2}{2}\right), \quad u > 0. \quad (5.3)$$

Lemma 5.2. *Let $T = T(u)$ be defined as in (5.3) and $a > 0, 0 < \lambda < 1$ be given constants. Under the assumptions of Lemma 5.1 for any $0 \leq s_1 < \dots < s_p < t_1 < \dots < t_{p'}$ in $\{aqj : j \in \mathbb{Z}, 0 \leq aqj \leq T\}$ with $t_1 - s_p \geq \lambda T$, we have as $u \rightarrow \infty$*

$$\begin{aligned}
& \left| \mathbb{P}\left\{\bigcap_{i=1}^p \{X_{n-r+1:n}(s_i) \leq u\}, \bigcap_{j=1}^{p'} \{X_{n-r+1:n}(t_j) \leq u\}\right\} \right. \\
& \left. - \mathbb{P}\left\{\bigcap_{i=1}^p \{X_{n-r+1:n}(s_i) \leq u\}\right\} \mathbb{P}\left\{\bigcap_{j=1}^{p'} \{X_{n-r+1:n}(t_j) \leq u\}\right\} \right| \rightarrow 0. \quad (5.4)
\end{aligned}$$

Proof: Denote

$$X_{ij} = X_j(s_i)\mathbb{I}\{i \leq p\} + X_j(t_{i-p})\mathbb{I}\{p < i \leq p+p'\}, \quad 1 \leq i \leq p+p', \quad 1 \leq j \leq n$$

and $\{Y_{ij}, 1 \leq i \leq p, 1 \leq j \leq n\} \stackrel{d}{=} \{X_{ij}, 1 \leq i \leq p, 1 \leq j \leq n\}$, independent of $\{Y_{ij}, p+1 \leq i \leq p+p', 1 \leq j \leq n\} \stackrel{d}{=} \{X_{ij}, p+1 \leq i \leq p+p', 1 \leq j \leq n\}$. Applying Theorem 2.4 with

$$X_{i(n-r+1)} = X_{n-r+1:n}(s_i)\mathbb{I}\{i \leq p\} + X_{n-r+1:n}(t_{i-p})\mathbb{I}\{p < i \leq p+p'\}$$

and

$$Y_{i(n-r+1)} = Y_{n-r+1:n}(s_i)\mathbb{I}\{i \leq p\} + Y_{n-r+1:n}(t_{i-p})\mathbb{I}\{p < i \leq p+p'\},$$

it follows that, the left-hand side of (5.4) is bounded from above by

$$\begin{aligned}
& Ku^{-2(r-1)} \frac{T}{q} \sum_{\lambda T \leq t_j - s_i \leq T} \exp\left(-\frac{ru^2}{1+|\rho(t_j - s_i)|}\right) \int_0^{|\rho(t_j - s_i)|} \frac{(1+|h|)^{2(r-1)}}{(1-h^2)^{r/2}} dh \\
& \leq Ku^{-2(r-1)} \frac{T}{q} \sum_{\lambda T \leq aqj \leq T} |\rho(aqj)| \exp\left(-\frac{ru^2}{1+|\rho(aqj)|}\right) \quad (5.5)
\end{aligned}$$

for sufficiently large u . Here K is some constant. The rest of the proof consists of the similar arguments as that of Lemma 12.3.1 in [Leadbetter et al. \(1983\)](#). Indeed, letting $\gamma(t) = \sup\{|\rho(s)| \ln s : s \geq t\}, t > 1$, we have that $|\rho(t)| \leq \gamma(t)/\ln t$ and $\gamma(t) \leq M$ for some positive constant M and all sufficiently large t . Recalling (5.3), we have

$$u^2 = \frac{2}{r} \ln T + \left(\frac{2}{r\alpha} - 1\right) \ln \ln T + \ln \left(\left(\frac{r}{2}\right)^{1-2/(r\alpha)} \frac{(c_{n,r}\mathcal{A}_{r,\alpha})^{2/r}}{2\pi} \right) (1 + o(1)),$$

which implies that

$$\begin{aligned} \exp\left(-\frac{ru^2}{1+|\rho(aqj)|}\right) &\leq \exp\left(-ru^2\left(1-\frac{\gamma(\lambda T)}{\ln(\lambda T)}\right)\right) \\ &\leq K \exp(-ru^2) \leq KT^{-2}(\ln T)^{r-2/\alpha} \end{aligned}$$

for all T large. Consequently, the right-hand side of (5.5) is bounded from above by

$$\begin{aligned} &Ku^{-2(r-1)}\left(\frac{T}{q}\right)^2\left(\frac{1}{T/q}\sum_{\lambda T \leq aqj \leq T}|\rho(aqj)|\ln(aqj)\right)\frac{1}{\ln(\lambda T)}T^{-2}(\ln T)^{r-2/\alpha} \\ &\leq K\frac{q}{T}\sum_{\lambda T \leq aqj \leq T}|\rho(aqj)|\ln(aqj), \end{aligned}$$

which tends to 0 as $T \rightarrow \infty$ since $\rho(t)\ln t = o(1)$. Hence the proof is complete. \square

Below W denotes an $N(0, 1)$ random variable which is independent of any other random element involved.

Proof of Theorem 3.1: a) Note that (3.2) and Lemmas 5.1 and 5.2 hold for the r th upper order statistics process $\{X_{n-r+1:n}(t), t \geq 0\}$. In view of Theorem 10 in Albin (1990) we have for $T = T(u)$ defined as in (5.3)

$$\lim_{u \rightarrow \infty} \mathbb{P}\left\{\sup_{t \in [0, T(u)]} X_{n-r+1:n}(t) \leq u + \frac{x}{ru}\right\} = \exp(-e^{-x}), \quad x \in \mathbb{R}.$$

Expressing u in terms of T using (5.3) we obtain the required claim for any $x \in \mathbb{R}$, with $a_{r,T}, b_{r,T}$ given as in (3.3); the uniform convergence in x follows since all functions (with respect to x) are continuous, bounded and increasing.

b) The proof follows from the main arguments of Theorem 3.1 in Mittal and Ylvisaker (1975) by showing that, for any $\varepsilon > 0$ and $x \in \mathbb{R}$

$$\begin{aligned} \Phi(x - \varepsilon) &\leq \liminf_{T \rightarrow \infty} \mathbb{P}\left\{M_X(T) \leq c_T b_{r,T} + \sqrt{\rho(T)}x\right\} \\ &\leq \limsup_{T \rightarrow \infty} \mathbb{P}\left\{M_X(T) \leq c_T b_{r,T} + \sqrt{\rho(T)}x\right\} \leq \Phi(x + \varepsilon), \end{aligned} \quad (5.6)$$

where

$$M_X(T) := \sup_{t \in [0, T]} X_{n-r+1:n}(t), \quad c_T := \sqrt{1 - \rho(T)}.$$

We start with the proof of the first inequality. Let $\rho^*(t), t \geq 0$ be a correlation function of a stationary Gaussian process such that $\rho^*(t) = 1 - 2|t|^\alpha + o(|t|^\alpha)$ as $t \rightarrow 0$. There exists some $t_0 > 0$ such that for T large

$$\rho^*(t)c_T^2 + \rho(T) \leq \rho(t), \quad 0 \leq t \leq t_0. \quad (5.7)$$

Denote by $\{Y_k(t), t \geq 0\}, k \in \mathbb{N}$ independent centered stationary Gaussian processes with a.s. continuous sample paths and common covariance function $\rho^*(\cdot)$, and define $\{Y(t), t \geq 0\}$ by

$$Y(t) = \sum_{k=1}^{\infty} Y_k(t) \mathbb{I}\{t \in [(k-1)t_0, kt_0)\}, \quad t \geq 0. \quad (5.8)$$

It follows from (5.7) that for T sufficiently large

$$\mathbb{E}\{X(s)X(t)\} \geq \mathbb{E}\left\{(c_T Y(s) + \sqrt{\rho(T)}W)(c_T Y(t) + \sqrt{\rho(T)}W)\right\}, \quad s, t \geq 0.$$

Therefore, by Proposition 3.2

$$\begin{aligned} & \mathbb{P} \left\{ M_X(T) \leq c_T b_{r,T} + \sqrt{\rho(T)} x \right\} \\ & \geq \mathbb{P} \left\{ c_T M_Y(T) + \sqrt{\rho(T)} W \leq c_T b_{r,T} + \sqrt{\rho(T)} x \right\} \\ & \geq \Phi(x - \varepsilon) \left(\mathbb{P} \left\{ \sup_{t \in [0, t_0]} Y_{n-r+1:n}(t) \leq b_{r,T} + \varepsilon \sqrt{\rho(T)} \right\} \right)^{[T/t_0]+1}. \end{aligned}$$

Noting that $a = \inf_{0 < t \leq t_0} (1 - \rho^*(t)) |t|^\alpha > 0$, we have by Theorem 1.1 in [Dębicki et al. \(2015a\)](#) (see also (3.2))

$$\lim_{T \rightarrow \infty} \frac{\mathbb{P} \left\{ \sup_{t \in [0, t_0]} Y_{n-r+1:n}(t) > b_{r,T} + \varepsilon \sqrt{\rho(T)} \right\}}{t_0 c_{n,r} b_{r,T}^{2/\alpha} (1 - \Phi(b_{r,T} + \varepsilon \sqrt{\rho(T)}))^r} = 2^{1/\alpha} \mathcal{A}_{r,\alpha}.$$

Consequently, since $\gamma = \infty$ we have

$$\begin{aligned} & \lim_{T \rightarrow \infty} ([T/t_0] + 1) \ln \mathbb{P} \left\{ \sup_{t \in [0, t_0]} Y_{n-r+1:n}(t) \leq b_{r,T} + \varepsilon \sqrt{\rho(T)} \right\} \\ & = - \lim_{T \rightarrow \infty} \frac{T}{t_0} \mathbb{P} \left\{ \sup_{t \in [0, t_0]} Y_{n-r+1:n}(t) > b_{r,T} + \varepsilon \sqrt{\rho(T)} \right\} \\ & = - \lim_{T \rightarrow \infty} T c_{n,r} 2^{1/\alpha} \mathcal{A}_{r,\alpha} b_{r,T}^{2/\alpha} (1 - \Phi(b_{r,T} + \varepsilon \sqrt{\rho(T)}))^r \\ & = 0 \end{aligned}$$

establishing the first inequality in (5.6).

Next, we consider the last inequality in (5.6). Note that, by the convexity of $\rho(\cdot)$, there exists a separable stationary Gaussian process $\{Y(t), t \in [0, T]\}$ with correlation function given by (using the well-known Polya criteria, see e.g., [Gneiting, 2001](#))

$$\tilde{\rho}(t) = \frac{\rho(t) - \rho(T)}{1 - \rho(T)}, \quad t \in [0, T]. \quad (5.9)$$

We have the equality in distribution

$$M_X(T) \stackrel{d}{=} c_T M_Y(T) + \sqrt{\rho(T)} W$$

implying

$$\begin{aligned} & \mathbb{P} \left\{ M_X(T) \leq c_T b_{r,T} + \sqrt{\rho(T)} x \right\} \\ & = \int_{-\infty}^{\infty} \mathbb{P} \left\{ M_Y(T) \leq b_{r,T} + \frac{\sqrt{\rho(T)}}{c_T} (x - u) \right\} \varphi(u) du \\ & \leq \Phi(x + \varepsilon) + \mathbb{P} \left\{ M_Y(T) \leq b_{r,T} - \varepsilon \frac{\sqrt{\rho(T)}}{c_T} \right\}. \end{aligned} \quad (5.10)$$

Consequently, we only need to prove that

$$\lim_{T \rightarrow \infty} \mathbb{P} \left\{ M_Y(T) \leq b_{r,T} - \varepsilon \sqrt{\rho(T)} \right\} = 0.$$

To this end, using again the convexity of $\tilde{\rho}(\cdot)$, we construct a separable stationary Gaussian process $\{Z(t), t \in [0, T]\}$ with the correlation function (recall $\tilde{\rho}(\cdot)$ in (5.9))

$$\sigma(t) = \max\left(\tilde{\rho}(t), \tilde{\rho}(T \exp(-\sqrt{\ln T}))\right), \quad t \in [0, T]. \quad (5.11)$$

Again by Proposition 3.2, we have

$$\mathbb{P}\left\{M_Y(T) \leq b_{r,T} - \varepsilon\sqrt{\rho(T)}\right\} \leq \mathbb{P}\left\{M_Z(T) \leq b_{r,T} - \varepsilon\sqrt{\rho(T)}\right\}. \quad (5.12)$$

Next, we construct a grid of intervals as follows. Let $I_1, \dots, I_{[T]}$ be $[T]$ consecutive unit intervals with an interval of length δ removed from the right-hand side of each one with $\delta \in (0, 1)$ given, and

$$\mathcal{G}_T = \{k(2 \ln T)^{-3/\alpha}, k \in \mathbb{N}\} \cap \left(\cup_{i=1}^{[T]} I_i\right).$$

It follows from Theorem 10 in Albin (1990) and Theorem 1.1 in Dębicki et al. (2015a) that, $\sup_{t \in [0, T]} Z_{n-r+1:n}(t)$ and $\sup_{t \in \mathcal{G}_T} Z_{n-r+1:n}(t)$ have the same asymptotic distribution and thus we only need to show that

$$\lim_{T \rightarrow \infty} \mathbb{P}\left\{\sup_{t \in \mathcal{G}_T} Z_{n-r+1:n}(t) \leq b_{r,T} - \varepsilon\sqrt{\rho(T)}\right\} = 0.$$

Let $\{Z'_{n-r+1:n}(t), t \geq 0\}$ be generated by $\{Z'(t), t \in [0, T]\}$ which is again a separable stationary process with the correlation function (recall $\sigma(\cdot)$ in (5.11))

$$\sigma^*(t) = \frac{\sigma(t) - \sigma(T)}{1 - \sigma(T)}, \quad t \in [0, T].$$

Analogously to the derivation of (5.10) we obtain

$$\begin{aligned} & \mathbb{P}\left\{\sup_{t \in \mathcal{G}_T} Z_{n-r+1:n}(t) \leq b_{r,T} - \varepsilon\sqrt{\rho(T)}\right\} \\ &= \mathbb{P}\left\{\sqrt{1 - \sigma(T)} \max_{t \in \mathcal{G}_T} Z'_{n-r+1:n}(t) + \sqrt{\sigma(T)}W \leq b_{r,T} - \varepsilon\sqrt{\rho(T)}\right\} \\ &\leq \Phi\left(-\frac{1}{2}\varepsilon\left(\frac{\rho(T)}{\sigma(T)}\right)^{1/2}\right) \\ &+ \mathbb{P}\left\{\max_{t \in \mathcal{G}_T} Z'_{n-r+1:n}(t) \leq b_{r,T} + \frac{b_{r,T}\sigma(T)}{\sqrt{1 - \sigma(T)}(1 + \sqrt{1 - \sigma(T)})} - \frac{\varepsilon\sqrt{\rho(T)}}{2\sqrt{1 - \sigma(T)}}\right\}, \end{aligned}$$

which tends to 0 as $T \rightarrow \infty$. The proof of it is the same as that of Theorem 3.1 in Mittal and Ylvisaker (1975), by using instead Theorem 1.1 in Dębicki et al. (2015a) and our Theorem 2.4. Consequently, the last inequality in (5.6) follows by (5.10) and (5.12). We complete the proof for $\gamma = \infty$.

c) Given $\delta \in (0, 1)$, take $I_1, \dots, I_{[T]}$ as in b). For $\{Y_k(t), t \geq 0\}, k \in \mathbb{N}$ independent copies of X define

$$Y(t) := \sum_{k=1}^{\infty} Y_k(t) \mathbb{I}\{t \in [k-1, k)\}, \quad t \geq 0$$

and

$$X_*(t) := \sqrt{1 - \rho_*(T)}Y(t) + \sqrt{\rho_*(T)}W, \quad t \in \cup_{k=1}^{[T]} I_k,$$

where $\rho_*(T) = \gamma/\ln T$. The rest of the proof is similar to that as of Theorem 2.1 in Tan et al. (2012) by using our Theorem 2.4 instead of Berman's inequality. We

omit the details.

Combining all the arguments for the three cases above, we complete the proof of Theorem 3.1. \square

Acknowledgements

We are thankful to the Editor-in-chief and the referees for several constructive suggestions which greatly improved the manuscript.

References

- R. J. Adler and J. E. Taylor. *Random fields and geometry*. Springer Monographs in Mathematics. Springer, New York (2007). ISBN 978-0-387-48112-8. [MR2319516](#).
- J. M. P. Albin. On extremal theory for stationary processes. *Ann. Probab.* **18** (1), 92–128 (1990). [MR1043939](#).
- M. T. Alodat. An approximation to cluster size distribution of two Gaussian random fields conjunction with application to fMRI data. *J. Statist. Plann. Inference* **141** (7), 2331–2347 (2011). [MR2775212](#).
- M. T. Alodat, M. Al-Rawwash and M. A. Jebrini. Duration distribution of the conjunction of two independent F processes. *J. Appl. Probab.* **47** (1), 179–190 (2010). [MR2654766](#).
- A. Aue, L. Horváth and M. Hušková. Extreme value theory for stochastic integrals of Legendre polynomials. *J. Multivariate Anal.* **100** (5), 1029–1043 (2009). [MR2498730](#).
- S. M. Berman. Limit theorems for the maximum term in stationary sequences. *Ann. Math. Statist.* **35**, 502–516 (1964). [MR0161365](#).
- S. M. Berman. Sojourns and extremes of stationary processes. *Ann. Probab.* **10** (1), 1–46 (1982). [MR637375](#).
- S. M. Berman. *Sojourns and extremes of stochastic processes*. The Wadsworth & Brooks/Cole Statistics/Probability Series. Wadsworth & Brooks/Cole Advanced Books & Software, Pacific Grove, CA (1992). ISBN 0-534-13932-9. [MR1126464](#).
- V. Chernozhukov, D. Chetverikov and K. Kato. Comparison and anti-concentration bounds for maxima of Gaussian random vectors. *Probab. Theory Related Fields* **162** (1-2), 47–70 (2015). [MR3350040](#).
- K. Dębicki, E. Hashorva, L. Ji and C. Ling. Extremes of order statistics of stationary processes. *ArXiv Mathematics e-prints* (2014a). [arXiv: 1403.7354](#).
- K. Dębicki, E. Hashorva, L. Ji and C. Ling. Extremes of order statistics of stationary processes. *TEST* **24** (2), 229–248 (2015a). [MR3348209](#).
- K. Dębicki, E. Hashorva, L. Ji and K. Tabiś. On the probability of conjunctions of stationary Gaussian processes. *Statist. Probab. Lett.* **88**, 141–148 (2014b). [MR3178344](#).
- K. Dębicki, E. Hashorva, L. Ji and K. Tabiś. Extremes of vector-valued Gaussian processes: exact asymptotics. *Stochastic Process. Appl.* **125** (11), 4039–4065 (2015b). [MR3385594](#).
- K. Dębicki, E. Hashorva and N. Soja-Kukieła. Extremes of homogeneous Gaussian random fields. *J. Appl. Probab.* **52** (1), 55–67 (2015c). [MR3336846](#).
- T. Gneiting. Criteria of Pólya type for radial positive definite functions. *Proc. Amer. Math. Soc.* **129** (8), 2309–2318 (2001). [MR1823914](#).

- Y. Gordon. Some inequalities for Gaussian processes and applications. *Israel J. Math.* **50** (4), 265–289 (1985). [MR800188](#).
- A. J. Harper. Bounds on the suprema of Gaussian processes, and omega results for the sum of a random multiplicative function. *Ann. Appl. Probab.* **23** (2), 584–616 (2013). [MR3059269](#).
- A. J. Harper. Pickands' constant H_α does not equal $1/\Gamma(1/\alpha)$, for small α . *Bernoulli* **23** (1), 582–602 (2017). [MR3556785](#).
- E. Hashorva and Z. Weng. Berman's inequality under random scaling. *Stat. Interface* **7** (3), 339–349 (2014). [MR3262770](#).
- D. Jarušková. Detecting non-simultaneous changes in means of vectors. *TEST* **24** (4), 681–700 (2015). [MR3414512](#).
- M. R. Leadbetter, G. Lindgren and H. Rootzén. *Extremes and related properties of random sequences and processes*. Springer Series in Statistics. Springer-Verlag, New York-Berlin (1983). ISBN 0-387-90731-9. [MR691492](#).
- W. V. Li and Q.-M. Shao. A normal comparison inequality and its applications. *Probab. Theory Related Fields* **122** (4), 494–508 (2002). [MR1902188](#).
- W. V. Li and Q.-M. Shao. Lower tail probabilities for Gaussian processes. *Ann. Probab.* **32** (1A), 216–242 (2004). [MR2040781](#).
- W. V. Li and Q.-M. Shao. Recent developments on lower tail probabilities for Gaussian processes. *Cosmos* **1** (1), 95–106 (2005). [MR2329259](#).
- C. Ling and Z. Peng. Extremes of order statistics of self-similar processes (in Chinese). *Sci Sin Math* **46** (8), 1139–1148 (2016).
- D. Lu and X. Wang. Some new normal comparison inequalities related to Gordon's inequality. *Statist. Probab. Lett.* **88**, 133–140 (2014). [MR3178343](#).
- Y. Mittal and D. Ylvisaker. Limit distributions for the maxima of stationary Gaussian processes. *Stochastic Processes Appl.* **3**, 1–18 (1975). [MR0413243](#).
- James Pickands, III. Upcrossing probabilities for stationary Gaussian processes. *Trans. Amer. Math. Soc.* **145**, 51–73 (1969). [MR0250367](#).
- V. I. Piterbarg. *Asymptotic methods in the theory of Gaussian processes and fields*, volume 148 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI (1996). ISBN 0-8218-0423-5. [MR1361884](#).
- V.I. Piterbarg. Discrete and continuous time extremes of Gaussian processes. *Extremes* **7** (2), 161–177 (2005) (2004). [MR2154366](#).
- V.I. Piterbarg. *Twenty Lectures About Gaussian Processes*. Atlantic Financial Press, London, New York (2015).
- R. L. Plackett. A reduction formula for normal multivariate integrals. *Biometrika* **41**, 351–360 (1954). [MR0065047](#).
- D. Slepian. The one-sided barrier problem for Gaussian noise. *Bell System Tech. J.* **41**, 463–501 (1962). [MR0133183](#).
- Z. Tan, E. Hashorva and Z. Peng. Asymptotics of maxima of strongly dependent Gaussian processes. *J. Appl. Probab.* **49** (4), 1106–1118 (2012). [MR3058991](#).
- K. J. Worsley and K. J. Friston. A test for a conjunction. *Statist. Probab. Lett.* **47** (2), 135–140 (2000). [MR1747100](#).
- L. Yan. Comparison inequalities for one sided normal probabilities. *J. Theoret. Probab.* **22** (4), 827–836 (2009). [MR2558653](#).