



This is a repository copy of *The volume-outcome relationship in severely injured patients: A systematic review and meta-analysis*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/139023/>

Version: Accepted Version

---

**Article:**

Sewalt, C.A., Wiegers, E.J.A., Venema, E. et al. (4 more authors) (2018) The volume-outcome relationship in severely injured patients: A systematic review and meta-analysis. *Journal of Trauma and Acute Care Surgery*, 85 (4). pp. 810-819. ISSN 2163-0755

<https://doi.org/10.1097/TA.0000000000002043>

---

© 2018 Lippincott Williams & Wilkins, Inc. This is an author produced version of a paper subsequently published in *Journal of Trauma and Acute Care Surgery*. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# **The volume-outcome relationship in severely injured patients: a systematic review and meta-analysis**

Charlie A Sewalt, BSc<sup>\*1</sup>; Eveline JA Wieggers, BSc<sup>\*1</sup>; Esmee Venema, MD<sup>1,2</sup>; Fiona E Lecky, MD, PhD<sup>3,4</sup>; Stephanie CE Schuit, MD, PhD<sup>5</sup>; Dennis Den Hartog, MD, PhD<sup>6</sup>; Hester F Lingsma, PhD<sup>1</sup>

\*both authors contributed equally

## Affiliations

1. Department of Public Health, Erasmus University Medical Center, Rotterdam, The Netherlands.
2. Department of Neurology, Erasmus University Medical Center, Rotterdam, The Netherlands.
3. School of Health and Related Research, Sheffield University. Salford Royal NHS Foundation Trust, Salford, UK.
4. Trauma Audit and Research Network, University of Manchester, Salford, Manchester, UK.
5. Department of Emergency Medicine, Erasmus University Medical Center, Rotterdam, The Netherlands.
6. Trauma Research Unit, Department of Surgery, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

Corresponding Author:

Charlie Aletta Sewalt, BSc,

Department of Public Health, Erasmus University Medical Center,

P.O. Box 2040, 3000 CA Rotterdam, The Netherlands

Email: [c.sewalt@erasmusmc.nl](mailto:c.sewalt@erasmusmc.nl)

Telephone number: +31107038460

Author disclosures: the authors have nothing to disclose.

Email addresses:

Charlie A Sewalt: [c.sewalt@erasmusmc.nl](mailto:c.sewalt@erasmusmc.nl)

Eveline JA Wiegers: [e.wiegers@erasmusmc.nl](mailto:e.wiegers@erasmusmc.nl)

Esmee Venema: [e.venema@erasmusmc.nl](mailto:e.venema@erasmusmc.nl)

Fiona E Lecky: [fiona.lecky@manchester.ac.uk](mailto:fiona.lecky@manchester.ac.uk)

Stephanie CE Schuit: [s.schuit@erasmusmc.nl](mailto:s.schuit@erasmusmc.nl)

Dennis Den Hartog: [d.denhartog@erasmusmc.nl](mailto:d.denhartog@erasmusmc.nl)

Hester F Lingsma: [h.lingsma@erasmusmc.nl](mailto:h.lingsma@erasmusmc.nl)

## **Abstract**

**Background:** The volume-outcome relationship in severely injured patients remains under debate and this has consequences for the designation of trauma centers.

**Objectives:** The aim of this study was to evaluate the relationship between hospital or surgeon volume and health outcomes in severely injured patients.

**Methods:** Six electronic databases were searched from 1980 up to January 30<sup>th</sup> 2018 to identify studies that describe the relationship between hospital or surgeon volume and health outcomes in severely injured patients (preferably Injury Severity Score (ISS) above 15). Selection of relevant studies, data extraction and critical appraisal of the methodological quality were performed by two independent reviewers. Pooled adjusted and unadjusted estimates of the effect of volume on in-hospital mortality, only in study populations with ISS > 15, were calculated with a random-effects meta-analysis. A mixed effects linear regression model was used to assess hospital volume as continuous parameter.

**Results:** Eighteen observational cohort studies were included. The majority (13/18, 72%) reported an association between higher hospital or surgeon volume and lower mortality rate. Overall, the quality of the included studies was reasonable, with insufficient adjustment as one of the most common limitations. Eight studies were included in the meta-analysis with a total of 222,418 patients. High hospital volume (>240 admitted severely injured patients per year) was associated with a lower risk of mortality (adjusted odds ratio 0.85, 95% confidence interval (CI) 0.76-0.94). Four studies were included in the regression model, providing a beta of -0.17 per 10 patients (95% CI -0.27 to -0.07). There was no clear association between surgeon volume and mortality rates based on three available studies.

**Conclusion:** Our systematic overview of the literature reveals a modest association between high volume centers and lower mortality in severely injured patients, suggesting that designation of high volume centers might improve outcomes among severely injured patients.

**Level of evidence:** level III, Systematic review and meta-analysis

**Systematic review registration number:** PROSPERO registration ID CRD42017056729

**Keywords:** Trauma, Severely Injured, Hospital volume, Volume-outcome Relationship

## **Background**

Each year, about 4.8 million people die worldwide as a result of injuries (1). In adults younger than 45 years, injury is even the major cause of death (2). Therefore, trauma imposes a substantial burden on society: In the Netherlands, the annual total costs of injuries are €3.5 billion (3). The implementation of trauma systems and dedicated level I trauma centers has reduced mortality of severely injured patients and improved functional outcome at discharge (4).

The American College of Surgeons Committee on Trauma (ACS-COT) requires a minimum of 240 admission of severely injured patients (Injury Severity Score (ISS) >15) per year for all level 1 trauma centers. Alternatively, individual trauma surgeons should admit at least 35 severely injured patients per year (5, 6). These volume requirements are originally based on the volume-outcome relationship in other surgical specialties such as cardiothoracic surgery (7-9). Two previously published systematic reviews on the impact of volume on outcome concluded that the benefit of high volume of annually admitted trauma patients in terms of health outcomes remains unclear (10, 11) .

However, these systematic reviews did not perform a meta-analysis. It is likely that a potential positive effect of volume on outcome is more visible in a meta-analysis, because it increases power. The aim of this systematic review and meta-analysis is to evaluate the volume-outcome relationship in severely injured patients.

## **Methods**

### Literature search

The search engines Embase.com (Medline and Embase), Pubmed, Web of Science, Cochrane Central and Google Scholar were searched until January 30<sup>th</sup> 2018 (date last searched) to identify published studies that examined the association between the volume of severely injured patients and different health outcomes. There were time restrictions; only articles published after 1980 were taken into account. The search was designed by an experienced biomedical information specialist. The search algorithm included subject heading (MeSH) terms and text words for injured patients, hospital volume, trauma center and different health outcomes (see appendix). In order to identify potential additional studies, we checked the reference lists of studies included and contacted experts in the field of trauma.

### Study selection and inclusion criteria

Observational cohort studies that examined the association between hospital or surgeon volume of severely injured patients and different health outcomes were included. Studies were included when a severely injured population was defined as patients with ISS >15 or, when ISS was not used, with clinical and anatomical patient characteristics comparable to the severity of ISS >15 confirmed by an experienced trauma surgeon. Studies that focused on patients with specific anatomical injuries or studies that only used pediatric cohorts were excluded. Studies not written in English were excluded.

Two reviewers (CS and EW) independently screened titles and abstracts to identify potentially eligible articles. Full-text reports of the potentially eligible articles were retrieved and these two reviewers independently screened these full-text articles to identify eligible

studies. Any disagreement was resolved through discussion or, if necessary, a third review author (HL) was consulted to reach consensus. The PRISMA flowchart was used to provide an overview of the data screening process (12).

#### Data extraction and quality assessment

Data extraction was performed by two independent reviewers (CS and EW). Extracted information included study characteristics (publication year, study design, study period, setting, sample size), patient characteristics (inclusion and exclusion criteria), type of volume (hospital or surgeon), definition of volume (unit of measurement, continuous or categorical variable with corresponding thresholds), health outcomes, and key figures (adjusted and unadjusted estimates of the effect of volume and outcome).

Based on previous literature on quality of observational studies, a quality assessment form was made to assess quality, generalizability and risk of bias of the included studies (13). The quality assessment form was applied to each study by two independent reviewers (CS and EW). Also, the risk for publication bias was assessed using funnel plots.

#### Data analysis

A meta-analysis was performed to determine the relationship between hospital volume and outcome in severely injured patients. Hospital volume was defined as the mean annual number of admitted severely injured patients. Studies were included in meta-analyses when severely injured was defined as ISS>15 to reduce heterogeneity. The outcome of the meta-analysis was mortality, defined as either in-hospital mortality or death within 30 days following trauma.



The association between patient volume per hospital and mortality was assessed in two ways: using a threshold of 240 patients to separate high volume centers from low volume centers (5, 6) and using volume as a continuous parameter.

For the meta-analysis, only studies with cut-offs close to 240 were included. Both adjusted and unadjusted outcomes were used to calculate pooled effect estimates. For the adjusted estimates, the adjusted odds ratio (OR) was used when reported. To calculate unadjusted ORs, mortality rates per study and volume group were used. The lowest volume group was used as a reference, results were transformed (1/effect size) if necessary. Studies that did not report either mortality rates or ORs were excluded from the meta-analysis. The pooled unadjusted OR was calculated with the Mantel-Haenszel method, the pooled adjusted OR was calculated with the inverse variance method. A random effects model was used to pool the estimates and to account for expected heterogeneity since studies had different study populations, were from different regions and time frames. Heterogeneity was assessed using the Q-test quantified by the  $I^2$  statistic.

To assess hospital volume in a continuous way, data from studies reporting volumes and in-hospital mortality from at least two volume groups were used. Only studies using the definition for severely injured as  $ISS > 15$  were included. Volume was calculated as the mean number of severely injured patients per hospitals per year in a specific volume group and outcome was the reported mortality rate. A random effects linear regression model, with in-hospital mortality as outcome, a random effect for study to adjust for study differences, adjustments for mean age and mean ISS and weighted for the number of patients in each volume group was used. A random effects model was used to account for heterogeneity of studies and inter-study variance (14). By weighting the number of patients in each volume group, larger studies were had more influence in the regression model compared to smaller

studies. This resulted in a beta regression coefficient for the effect of 10 additional patients per year on mortality. The effect of hospital volume was tested for non-linearity using the likelihood ratio test with natural cubic splines.

Analyses were performed with the R software environment (version 3.2.2 or higher, the R Foundation for Statistical Computing, Vienna, Austria) and Review Manager (RevMan, version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014).

## Results

In the initial search, 9,181 records were identified (Figure 1). After removing the duplicates, 5,364 records were screened on title and abstract. The remaining 202 potentially eligible articles were selected for full-text assessment and eighteen of these studies were included in the systematic review (Table 1).

### Study characteristics

Fifteen (83%) were retrospective cohort studies (15-29), one study (6%) was a secondary analysis of two randomized controlled trials (30) and two studies (11%) were prospective cohort studies (31, 32) (Table 1). Fifteen studies (83%) were conducted in the United States (15-17, 19-27, 30, 32, 33), one study (6%) was conducted in Germany (28), one study was conducted in Japan (29), and one study was conducted in the United Kingdom (31). The National Trauma Databank (NTDB) from the United States was used in three studies (18%) (15, 19, 33). Other studies selected their sample from nation, state or hospital registries. Seventeen studies (94%) evaluated the hospital volume-outcome relationship (14-16, 18-25, 27-32) and three studies (18%) examined the surgeon volume-outcome relationship (20, 23, 27).

Mortality was defined as in-hospital mortality in sixteen studies (89%) (15-17, 19-29, 32, 33), in one study (6%) as 24-hour mortality (30) and in one study as 6-months mortality (31) (Table 1). Other outcomes reported were hospital length of stay (21, 25), discharge locations (17), intensive care length of stay (18), severe disability at discharge (18), total costs per admission (29) and complication rates (30). Thirteen studies (72%) used the ISS to define the severity of injuries. In ten studies (59%) the population examined had an ISS of at least 16 (17, 21-24, 26-28, 31, 33), three studies (18%) examined patients with an ISS of 15 or more

(15, 16, 19) and one study (6%) used a threshold of 13 (20). In three studies (17%) ISS was not used to define injury severity (25, 30, 32). As an alternative, a combination of clinical and anatomical characteristics like the Abbreviated Injury Scale (AIS), Glasgow Coma Score (GCS), Systolic Blood Pressure, Trauma Score (TS) or admission to the intensive care unit were used in these three studies (25, 30, 32). These characteristics were confirmed as comparable to ISS > 15 by an experienced trauma surgeon.

Seven studies (39%) (19, 20, 23, 25, 28, 29, 31) analyzed hospital volume as a continuous parameter and used the annual volume of severely injured patients (Table 1). Three studies (17%) (15, 24, 33) used a cut-off of 240 severely injured trauma patients per year. Three studies (17%) (21, 26, 32) quantified hospital volume based on the total volume of trauma patients categorized in two groups, all defined 'high volume' as above 1200 annual patients. For surgeon volume, two studies examined the volume-outcome relationship with volume as a continuous parameter (20, 23), while in one study a total of 35 or more severely injured patients admitted per surgeon per year was defined as 'high surgeon volume'(27).

### Quality assessment

All studies reported the total number of severely injured patients and had a limited impact of loss of follow-up (Figure 2). Most studies (72%) were considered to have a representative study population and the number of patients per volume group was frequently reported (89%). In 72% of the studies the reported mortality was clearly defined as in-hospital mortality and crude ORs were reported in 94% of the studies. Shortcomings mostly concerned the adjusted analyses; only ten (56%) studies reported adjusted ORs, and when adjustments were made the type of adjustments were not always mentioned.

## Effect of hospital volume

Eleven of the seventeen (65%) studies on hospital volume reported lower mortality rates in high volume centers compared to low volume centers (15, 17, 23-26, 28-32). Five studies (28%) did not find a positive nor a negative association between hospital volume and mortality (16, 19-21, 33). One study (6%) looked at specific subgroups of ISS (16-24 and >25) and did not find any evidence that high volume hospitals perform better than low volume hospitals in terms of mortality in extremely severe injured patients (ISS>25) (34). Another study that divided its population into blunt and penetrating injury found that the relationship between volume and mortality was stronger in penetrating injuries (14). Other outcomes were reported too infrequently.

## Meta-analysis

Eight studies evaluating the relationship between trauma center volume and in-hospital mortality in severely injured patients could be included in the meta-analysis with a total of 222,418 patients (15-17, 19, 23, 24, 33). All of these studies presented crude mortality rates, and adjusted ORs were reported in four out of eight studies (50%). The pooled effect estimate showed no association between volume and mortality when using the unadjusted ORs (OR 1.00, 95% confidence interval (CI): 0.92-1.10,  $p = 0.93$ , Figure 3A), with large heterogeneity ( $I^2 = 84%$ ,  $p < 0.001$ ). When including adjusted estimates only, high volume was associated with lower mortality (OR 0.85, 95% CI: 0.76-0.94,  $p = 0.003$ , Figure 3B), with moderate heterogeneity ( $I^2 = 44%$ ,  $p = 0.13$ ). There was no suggestion for publication bias (Figure 5). Four studies could be used for the analysis of the effect of continuous hospital volume on mortality and showed a similar association (Beta: -0.17 per 10 patients, 95% CI -0.27 to -0.07,  $p < 0.01$ , Figure 4). The effect of hospital volume was considered to be linear ( $p=0.998$ ).

### Surgeon volume

Three of the eighteen studies (17%) examined the relationship between surgeon volume and in-hospital mortality in severely injured patients (20, 23, 27). One study reported that increased per-surgeon volume in the treatment of seriously injured patients is associated with lower mortality (20). The authors suggest that a surgeon should treat at least 35 severely injured patients per surgeon per year. The other two studies found no relationship (23, 27).

## Discussion

This study aimed to evaluate the relationship between hospital or surgeon volume and health outcomes after severe trauma. The systematic review included eighteen studies, of which eight studies were included in the meta-analysis and four studies were included in the regression analysis. Our results indicate that a significant association between hospital volume and mortality exists, although the effect is modest. Overall, the quality of the included articles was reasonable, with insufficient adjustment as one of the most common limitations. No clear conclusions can be drawn with respect to surgeon volume due to an insufficient number of studies.

There might be several reasons explaining the fact that high volume hospitals perform better. The most obvious declaration is the 'practice makes perfect' hypothesis, which suggests that physicians and nurses can develop higher proficiency in the care of severely injured patients in case of high exposure (35). Although trauma patients all present in different ways, trauma care is a highly standardized in terms of initial approach in the Emergency Department (36), where lifesaving interventions are performed even before the precise injuries are known.

Personalized care is needed once the full extent of injury, and the patient's pre-existing health status is known - but this requires competence in complex operations and procedures and high fidelity team work. More exposure to this process may result in more efficiency, which is of extreme importance in the care for severely injured patients.

Another possible explanation might be the infrastructure of high volume trauma centers. The development of designated trauma services or trauma teams for the management of severely injured patients in high volume centers decreases mortality (37, 38). Also, the presence of in-house coverage by trauma surgeons at high volume centers is believed to improve outcomes, although there is no empirical evidence showing an actual decrease in mortality (39, 40).

Another possibility is the ability of each center to treat the range of injuries presenting, high volume centers could have more skills on site compared to small volume centers. There is also a possibility that high volume centers implement significant changes in trauma care like new technological improvements earlier. High volume centers have the commitment and resources to implement advances in trauma care which could lead to a decrease in mortality.

Some hospitals are more likely to receive more severely injured patients (i.e. with higher ISS) than other hospitals due to trauma center designation or hospital location. Furthermore, it is evident that the risk of mortality increases with higher ISS (41). When examining the relationship between volume of severely injured patients and mortality it is of high importance to adjust for this case-mix differences. Nevertheless, in the studies included in this systematic review, the severity of injury is not always taken into account. Due to the lack of these adjustments, it is hard to correctly interpret the results of these studies.

Most studies used of a cut-off value of 240 severely injured patients per year. This cut-off value has been arbitrarily chosen by the ACS-COT and was originally based on studies from other surgical specialties like cardiothoracic surgery (7-9). Although several studies with a continuous analysis of the volume-outcome relationship have been published, the ideal cut-off value remains unclear. As suggested in Figure 4, the optimal cut-off may be higher than 240, but more extensive continuous analysis of the volume-outcome relationship is needed. Also, it might be possible that there is an inflection point where mortality begins to increase, but extreme high volume centers are needed to examine this association and these were not included in our meta-analysis. Furthermore, using a higher cut-off value will lead to more centralization of trauma care. This may increase transport times of severely injured patients, which is associated with increased mortality rates in trauma patients (42-44). However, this might be less important in small countries like the Netherlands and other European countries.



Two previously published systematic reviews on the impact of volume on outcome did not find a clear benefit for severely injured patients in high volume centers (10, 11). However, both studies did not perform a meta-analysis and looked at a smaller amount of studies. Our meta-analysis and regression analysis revealed new insights and showed more consistent results that severely injured patients benefit from treatment in high volume centers.

A limitation of our study is that only few studies could be included for the meta-analysis. As a result, pooled findings might be largely influenced by the results of one study. There was no indication of publication bias. Furthermore, we only assessed the relationship of volume with mortality. Other outcomes like hospital length of stay and quality of life after trauma are also important indicators of quality of care and might as well be influenced by hospital or surgeon volume. Another limitation is that a majority of our included studies, almost 90%, were conducted in the United States. Variations in the structure of trauma centers might make our results less generalizable to trauma centers in Europe. The high amount of studies conducted in the United States was also the cause for including studies with the same data source in the meta-analysis. This makes it possible that patients are included in the meta-analysis more than once. In addition, a clear definition of severely injured patients was lacking. Most studies included in our systematic review used ISS to define injury severity, since it is the universal injury severity measure in trauma registries and research. But we also had to include three studies with other definitions of injury severity, like clinical and anatomical characteristics. Although an experienced trauma surgeon confirmed these studies as comparable to ISS > 15, it still increases heterogeneity of the study population. In addition, the ISS has been regarded as the golden standard to define injury severity over the last decades (45-47). However, questions about the accuracy of the ISS have been raised. First, it does not account for multiple injuries in the same body region (48, 49). This could cause underscoring of the

overall injury severity in patients with severe injury in one body region. Second, an equal AIS score in different body regions is assumed to be equal in injury severity (48, 50).

To the best of our knowledge, our search identified all studies that examined the relationship between volume of severely injured patients and mortality. Although most articles reported a positive relationship between volume and outcome, the cut-off of high volume and low volume hospitals was inconsistent across different studies. Some studies adopted the ACS-COT volume requirements of 240 or more severely injured patients per year, while other studies appeared to arbitrarily defined the volume cut-offs. This variance in cut-offs might be a good explanation for the difference in results. Although we tried to make equal volume groups, inconsistency in volume groups remained due to the limited available data. However, excluding even more studies in our meta-analysis would make our findings less generalizable.

## **Conclusion**

In conclusion, this systematic review and meta-analysis revealed an association between larger hospital volume and lower mortality in severely injured patients. Our findings suggest that designation of high volume centers can improve outcomes among severely injured patients. Future studies with more rigorous methodological case mix adjustment, additional outcome measures and standardized cut-off values for high volume in combination with continuous analyses are needed to further define the effect of hospital volume on mortality and outcome of severely injured patients.

## **Acknowledgements**

The authors would like to thank Wichor Bramer, biomedical information specialist from Erasmus MC, University Medical Center Rotterdam, for developing and assisting with the literature search strategy.

## **Author Contributions**

CS: literature search, study selection, data extraction, data-analysis, drafting the manuscript

EW: literature search, study selection, data extraction, data-analysis, drafting the manuscript

EV: study selection, data-analysis, critical revision of the manuscript

FL: critical revision of the manuscript

SS: critical revision of the manuscript

DDH: study selection, critical revision of the manuscript

HL: study selection, data-analysis, critical revision of the manuscript

## References

1. Haagsma JA, Graetz N, Bolliger I, Naghavi M, Higashi H, Mullany EC, Abera SF, Abraham JP, Adofo K, Alsharif U, et al. The global burden of injury: incidence, mortality, disability-adjusted life years and time trends from the Global Burden of Disease study 2013. *Inj Prev*. 2016;22(1):3-18.
2. Polinder S, Haagsma JA, Toet H, van Beeck EF. Epidemiological burden of minor, major and fatal trauma in a national injury pyramid. *Br J Surg*. 2012;99 Suppl 1:114-21.
3. Polinder S, Haagsma J, Panneman M, Scholten A, Brugmans M, Van Beeck E. The economic burden of injury: Health care and productivity costs of injuries in the Netherlands. *Accid Anal Prev*. 2016;93:92-100.
4. Celso B, Tepas J, Langland-Orban B, Pracht E, Papa L, Lottenberg L, Flint L. A systematic review and meta-analysis comparing outcome of severely injured patients treated in trauma centers following the establishment of trauma systems. *J Trauma*. 2006;60(2):371-8; discussion 8.
5. Eastman AB. Resources for optimal care of the injured patient--1993. *Bull Am Coll Surg*. 1994;79(5):21-7.
6. Trauma ACoSCo. Resources for optimal care of the injured patient: Amer College of Surgeons; 2014.
7. Adams DF, Fraser DB, Abrams HL. The complications of coronary arteriography. *Circulation*. 1973;48(3):609-18.
8. Hannan EL, Racz M, Ryan TJ, McCallister BD, Johnson LW, Arani DT, Guerci AD, Sosa J, Topol EJ. Coronary angioplasty volume-outcome relationships for hospitals and cardiologists. *JAMA*. 1997;277(11):892-8.
9. Luft HS, Bunker JP, Enthoven AC. Should operations be regionalized? The empirical relation between surgical volume and mortality. *N Engl J Med*. 1979;301(25):1364-9.
10. Kim YJ. Relationship of trauma centre characteristics and patient outcomes: a systematic review. *J Clin Nurs*. 2014;23(3-4):301-14.
11. Caputo LM, Salottolo KM, Slone DS, Mains CW, Bar-Or D. The relationship between patient volume and mortality in American trauma centres: A systematic review of the evidence. *Injury*. 2014;45(3):478-86.
12. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339:b2535.
13. Sanderson S, Tatt ID, Higgins JP. Tools for assessing quality and susceptibility to bias in observational studies in epidemiology: a systematic review and annotated bibliography. *Int J Epidemiol*. 2007;36(3):666-76.
14. DerSimonian R, Kacker R. Random-effects model for meta-analysis of clinical trials: an update. *Contemp Clin Trials*. 2007;28(2):105-14.
15. Bennett KM, Vaslef S, Pappas TN, Scarborough JE. The volume-outcomes relationship for United States Level I trauma centers. *J Surg Res*. 2011;167(1):19-23.
16. Cooper A, Hannan EL, Bessey PQ, Farrell LS, Cayten CG, Mottley L. An examination of the volume-mortality relationship for New York State trauma centers. *J Trauma*. 2000;48(1):16-23; discussion -4.

17. Cudnik MT, Newgard CD, Sayre MR, Steinberg SM. Level I versus Level II trauma centers: an outcomes-based assessment. *J Trauma*. 2009;66(5):1321-6.
18. Demetriades D, Martin M, Salim A, Rhee P, Brown C, Chan L. The effect of trauma center designation and trauma volume on outcome in specific severe injuries. *Ann Surg*. 2005;242(4):512-7; discussion 7-9.
19. Glance LG, Osler TM, Dick A, Mukamel D. The relation between trauma center outcome and volume in the National Trauma Databank. *J Trauma*. 2004;56(3):682-90.
20. Konvolinka CW, Copes WS, Sacco WJ. Institution and per-surgeon volume versus survival outcome in Pennsylvania's trauma centers. *Am J Surg*. 1995;170(4):333-40.
21. London JA, Battistella FD. Is there a relationship between trauma center volume and mortality? *J Trauma*. 2003;54(1):16-24; discussion -5.
22. Lucas CE, Buechter KJ, Coscia RL, Hurst JM, Lane V, Meredith JW, Middleton JD, Mitchell FL, Jr., Rinker CF, Tuggle D, et al. The effect of trauma program registry on reported mortality rates. *J Trauma*. 2001;51(6):1122-6; discussion 6-7.
23. Margulies DR, Cryer HG, McArthur DL, Lee SS, Bongard FS, Fleming AW. Patient volume per surgeon does not predict survival in adult level I trauma centers. *J Trauma*. 2001;50(4):597-601; discussion -3.
24. Marx WH, Simon R, O'Neill P, Shapiro MJ, Cooper AC, Farrell LS, McCormack JE, Bessey PQ, Hannan E. The relationship between annual hospital volume of trauma patients and in-hospital mortality in New York State. *J Trauma*. 2011;71(2):339-45; discussion 45-6.
25. Nathens AB, Jurkovich GJ, Maier RV, Grossman DC, MacKenzie EJ, Moore M, Rivara FP. Relationship between trauma center volume and outcomes. *JAMA*. 2001;285(9):1164-71.
26. Olufajo OA, Metcalfe D, Rios-Diaz A, Lilley E, Havens JM, Kelly E, Weissman JS, Haider AH, Salim A, Cooper Z. Does Hospital Experience Rather than Volume Improve Outcomes in Geriatric Trauma Patients? *J Am Coll Surg*. 2016;223(1):32-40 e1.
27. Sava J, Kennedy S, Jordan M, Wang D. Does volume matter? The effect of trauma surgeons' caseload on mortality. *J Trauma*. 2003;54(5):829-33; discussion 33-4.
28. Zacher MT, Kanz KG, Hanschen M, Haberle S, van Griensven M, Lefering R, Buhren V, Biberthaler P, Huber-Wagner S, TraumaRegister DGU. Association between volume of severely injured patients and mortality in German trauma hospitals. *Br J Surg*. 2015;102(10):1213-9.
29. Endo A, Shiraishi A, Fushimi K, Murata K, Otomo Y. Increased Severe Trauma Patient Volume is Associated With Survival Benefit and Reduced Total Health Care Costs: A Retrospective Observational Study Using a Japanese Nationwide Administrative Database. *Ann Surg*. 2017.
30. Minei JP, Fabian TC, Guffey DM, Newgard CD, Bulger EM, Brasel KJ, Sperry JL, MacDonald RD. Increased trauma center volume is associated with improved survival after severe injury: results of a Resuscitation Outcomes Consortium study. *Ann Surg*. 2014;260(3):456-64; discussion 64-5.
31. Freeman J, Nicholl J, Turner J. Does size matter? The relationship between volume and outcome in the care of major trauma. *J Health Serv Res Policy*. 2006;11(2):101-5.
32. Pasquale MD, Peitzman AB, Bednarski J, Wasser TE. Outcome analysis of Pennsylvania trauma centers: factors predictive of nonsurvival in seriously injured patients. *J Trauma*. 2001;50(3):465-72; discussion 73-4.

33. Demetriades D, Martin M, Salim A, Rhee P, Brown C, Chan L, Britt LD, Flint Jr LM, Maier RV, Meredith JW. The effect of trauma center designation and trauma volume on outcome in specific severe injuries. *Ann Surg.* 2005;242(4):512-9.
34. Cooper A, Hannan EL, Bessey PQ, Farrell LS, Cayten CG, Mottley L. An examination of the volume-mortality relationship for New York State trauma centers. *J Trauma Inj Infect Crit Care.* 2000;48(1):16-24.
35. Luft HS, Hunt SS, Maerki SC. The volume-outcome relationship: practice-makes-perfect or selective-referral patterns? *Health Serv Res.* 1987;22(2):157-82.
36. Araujo R, Della Corte F, Dick W, Driscoll P, Girbes R, Lorenzo AE, Lipp M, Nardi G, Oakley P, Sabbe M, et al. European comprehensive training course on prehospital advanced trauma life support in adults. *Eur J Emerg Med.* 2002;9(3):280-2.
37. Baker CC, Degutis LC, Desantis J, Baue AE. Impact of a Trauma Service on Trauma Care in a University Hospital. *American Journal of Surgery.* 1985;149(4):453-8.
38. Lansink KW, Leenen LP. Do designated trauma systems improve outcome? *Curr Opin Crit Care.* 2007;13(6):686-90.
39. Cox JA, Bernard AC, Bottiggi AJ, Chang PK, Talley CL, Tucker B, Davenport DL, Kearney PA. Influence of In-House Attending Presence on Trauma Outcomes and Hospital Efficiency. *J Am Coll Surgeons.* 2014;218(4):734-8.
40. Helling TS, Nelson PW, Shook JW, Lainhart K, Kintigh D. The presence of in-house attending trauma surgeons does not improve management or outcome of critically injured patients. *Journal of Trauma-Injury Infection and Critical Care.* 2003;55(1):20-5.
41. Baker SP, O'Neill B, Haddon W, Jr., Long WB. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma.* 1974;14(3):187-96.
42. Grossman DC, Kim A, Macdonald SC, Klein P, Copass MK, Maier RV. Urban-rural differences in prehospital care of major trauma. *J Trauma.* 1997;42(4):723-9.
43. Kristiansen T, Rehn M, Gravseth HM, Lossius HM, Kristensen P. Paediatric trauma mortality in Norway: a population-based study of injury characteristics and urban-rural differences. *Injury.* 2012;43(11):1865-72.
44. Wisborg T, Hoylo T, Siem G. Death after injury in rural Norway: high rate of mortality and prehospital death. *Acta Anaesthesiol Scand.* 2003;47(2):153-6.
45. Lavoie A, Moore L, LeSage N, Liberman M, Sampalis JS. The Injury Severity Score or the New Injury Severity Score for predicting intensive care unit admission and hospital length of stay? *Injury.* 2005;36(4):477-83.
46. Palmer C. Major trauma and the injury severity score--where should we set the bar? *Annu Proc Assoc Adv Automot Med.* 2007;51:13-29.
47. Rutledge R, Hoyt DB, Eastman AB, Sise MJ, Velky T, Canty T, Wachtel T, Osler TM. Comparison of the Injury Severity Score and ICD-9 diagnosis codes as predictors of outcome in injury: analysis of 44,032 patients. *J Trauma.* 1997;42(3):477-87; discussion 87-9.
48. Koksall O, Ozdemir F, Bulut M, Aydin S, Almacioglu ML, Ozguc H. Comparison of trauma scoring systems for predicting mortality in firearm injuries. *Ulus Travma Acil Cerrahi Derg.* 2009;15(6):559-64.

49. Tamim H, Al Hazzouri AZ, Mahfoud Z, Atoui M, El-Chemaly S. The injury severity score or the new injury severity score for predicting mortality, intensive care unit admission and length of hospital stay: experience from a university hospital in a developing country. *Injury*. 2008;39(1):115-20.
50. Chawda MN, Hildebrand F, Pape HC, Giannoudis PV. Predicting outcome after multiple trauma: which scoring system? *Injury*. 2004;35(4):347-58.





Author, year	Unit	Study design	Country	Period	Data Source	Patients	Volume definition	Outcomes	Key findings (unadjusted and adjusted estimates)
Bennett, 2011	Hospital	RCS	USA	2001-2006	NTDB	115,538	Categorical, 3 groups  <240  240-480  >480	In-hospital mortality	There is a complex volume-outcome relationship for level I trauma centers in the United States.  +
Cooper, 2000	Hospital	RCS	USA	1994-1995	NYTR	26,793	Categorical, 3 groups  <151  151-250  >250	In-hospital mortality	No association between hospital volume and health outcomes for trauma centers in New York State could be made.  -
Cudnik, 2009	Hospital	RCS	USA	2003-2006	OHTR	18,103	Level I vs Level II  Level I: average of 110 patients per year  Level II: average of 36 patients per year	In-hospital mortality, discharge locations	Severely injured patients have improved survival when taken to a Level I trauma center compared to those taken to level II trauma centers, which have a lower annual volume of trauma patients.  +

Demetriades, 2005	Hospital	RCS	USA	1996-2003	NTDB	12,254	Categorical, 2 groups  <240  ≥240  Level I vs Level II	In-hospital mortality, intensive care unit stay, severe disability at discharge	Health outcomes are not influenced by the volume of major trauma admissions.  -
Endo, 2017	Hospital	RCS	Japan	2010 - 2015	DPC	116,329	Continuous + categorical, 5 groups  1-50  51-100  101-150  151-200  >200	Survival to hospital discharge, total health care cost per admission	Higher hospital volume was significantly associated with a survival benefit and lower total costs per admission in severe trauma patients.  +
Freeman, 2006	Hospital	PCS	United Kingdom	1990-1993	RHD	2,190	Continuous	Mortality at 6 months	Severely injured patients had better health outcomes in higher volume departments.  +

Glance, 2002	Hospital	RCS	USA	1999	NTDB	7,371	Continuous and categorical, 4 groups  <140  140-261  262-462  >462	In-hospital mortality	Higher trauma center volumes are not associated with improved patients outcomes.  -
Konvolinka , 1995	Hospital + Surgeon	RCS	USA	1988 - 1989	PTOS	13,002	Continuous	In-hospital mortality	Higher hospital volume does not contribute to better survival rates.  -  To achieve reasonable survival rates, surgeons need to treat at least approximately 35 seriously injured patients.  +
London, 2003	Hospital	RCS	USA	1998 - 1999	RHD	98,245	Categorical and dichotomous (total trauma)  <1200  ≥1200	In-hospital mortality, hospital length of stay	Health outcomes in low-volume centers were comparable to higher- volume centers.  -

Lucas, 2001	Hospital	RCS	USA	Feb 1997- June 2000	TCSR	25,020	Continuous	In-hospital mortality	Mortality decreased when centers admitted more severely injured patients.  +
Marx, 2001	Hospital	RCS	USA	2003- 2006	NYTR	52,838	Categorical  <180 vs ≥180  <240 vs ≥240	In-hospital mortality	An higher volume of severely injured patients contributes to lower in- hospital mortality.  +
Margulies, 2000	Hospital + surgeon	RCS	USA	1998- 1999	DHS- EMS TR	1,754	Continuous	In-hospital mortality	The volume of trauma institutions is associated with better survival in severely injured patients.  +  Surgeon volume, however does not contribute to better survival.  -
Minei, 2014	Hospital	ARCT	USA & Canada	2006 - 2009	ROC	2222	Continuous and categorical (total trauma), 4 groups  ≤1000  1001-1999	24 hour mortality, 28 day mortality, complicatio ns	Mortality decreased with increased total trauma center volume.  +

							2000-2999 ≥3000		
Nathens, 2001	Hospital	RCS	USA	Nov 1997- Jul 1998	UHC	1019	Continuous	In-hospital mortality and hospital length of stay (LOS)	Patients outcomes strongly improve with higher trauma center volume. +
Olufajo, 2015	Hospital	RCS	USA	2007-2011	CSID	61,915	Categorical (total trauma), 2 groups <1200 ≥1200 Geriatric trauma volume per 100 increase	In-hospital mortality, Failure to rescue rate (FTR)	Higher geriatric trauma volume is associated with lower hospital mortality among geriatric patients. +
Pasquale, 2001	Hospital	PCS	USA	1992-1996	PTR	13,942	Categorical (total trauma), 2 groups <1200 ≥1200	In-hospital mortality	There exists an association between volume and in-hospital mortality. +

Sava, 2003	Surgeon	RCS	USA	1990- 2001	RHD	20,695	Categorical, 2 groups ≤35 >35	In-hospital mortality	Surgeon volume appeared to not influence outcome in severely injured patients.  -
Zacher, 2015	Hospital	RCS	Germany	2009- 2013	DGU	39,289	Continuous + categorical, 5 groups  1-19  20-39  40-59  60-79  80-99  ≥100	In-hospital mortality	Hospital volume of severely injured patients was identified as a predictor of survival.  +

**Table 1** Overview of included articles

Abbreviations:

RCS = Retrospective Cohort Study

PCS = Prospective Cohort Study

USA = United States

ARCT = Analysis of Randomized Controlled Trials

NTDB = National Trauma Data Bank

NYTR = New York's Trauma Registry

OHTR = State of Ohio Trauma Registry

RHD = Regional Hospital Discharge

PTOS = Pennsylvania Trauma Outcome Study

TCSR = Trauma Center Survey Reports

ROC = multicenter trials network Resuscitation Outcomes Consortium?

DHS-EMS TR = Department of Health Services-Emergency Medical Services trauma registry

PTR = Pennsylvania Trauma Registry

UHC = University Healthsystem Consortium


CSID = California State Inpatient Database

DGU = Deutsche Gesellschaft für Unfallchirurgie

DPC = Japanese Diagnosis Procedure Combination



## PRISMA 2009 Flow Diagram




From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097



Figure 2: quality assessment

Study	Population-based	Nationwide	In- and exclusion criteria clearly described	Representative study population	Total number of severely injured patients reported	Number of patients per volume group reported	Cut-off volume groups clearly reported	OR/HR hospital-volume reported	OR/HR reported + 95% CI	Limited impact of loss of follow-up	Clearly stated In-hospital mortality	Crude ORs reported/ % mortality reported	Adjusted ORs reported	All confounders mentioned for adjusted OR	Adjusted for patients demographic characteristics	Adjusted for severity	Adjusted for trauma center level	Adjusted for mechanism of injury	None declared conflict of interests	Funding sources identified
Bennett et al.	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	+	-	-
Cooper et al.	+	-	+	+	+	+	+	-	-	+	+	+	-	-	-	-	-	-	-	-
Cudnik et al.	+	-	+	+	+	+	-	+	+	+	+	+	+	+	-	-	+	-	-	-
Demetriades et al.	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	+	-	-
Endo et al.	+	+	+	-	+	+	+	+	+	+	+	+	+	+	-	+	+	-	-	-
Freeman et al.	+	-	+	+	+	+	+	-	-	+	-	+	-	-	-	-	-	-	-	-
Glance et al.	+	+	+	+	+	+	+	-	-	+	+	+	-	-	-	-	-	-	-	-
Konvolinka et al.	+	-	-	-	+	-	-	-	-	+	+	-	-	-	-	-	-	-	-	-
London et al.	+	-	+	+	+	+	+	-	-	+	+	-	+	+	+	+	+	-	-	-
Lucas et al.	+	-	-	+	+	+	-	-	-	+	+	+	-	-	-	-	-	-	-	-
Marx et al.	+	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-	-	-	-
Margulies et al.	+	-	+	+	+	+	+	-	-	+	-	+	-	-	-	-	-	-	-	-
Minei et al.	-	-	+	-	+	+	+	+	+	+	-	+	+	+	+	+	-	+	+	-
Nathens et al.	+	+	+	-	+	+	+	-	-	+	+	+	-	-	-	-	-	-	-	-
Olufajo et al.	+	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	-
Pasquale et al.	+	-	+	-	+	+	+	+	+	+	-	+	+	+	+	+	-	-	-	-
Sava et al.	+	-	-	+	+	+	+	-	-	+	-	+	-	-	-	-	-	-	-	-
Zacher et al.	+	+	+	+	+	+	+	-	+	+	+	+	+	-	+	+	-	-	-	-

Figure 3: In-hospital mortality unadjusted (A) and adjusted (B)






Figure 5

