



This is a repository copy of *Next generation sequencing (NGS) to improve the diagnosis and management of patients with disorders of sex development (DSD)*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/142190/>

Version: Accepted Version

---

**Article:**

Hughes, L.A., McKay Bounford, K., Webb, E. et al. (9 more authors) (2019) Next generation sequencing (NGS) to improve the diagnosis and management of patients with disorders of sex development (DSD). *Endocrine Connections*. ISSN 2049-3614

<https://doi.org/10.1530/EC-18-0376>

---

© 2019 The authors. This is an author produced version of a paper subsequently published in *Endocrine Connections*. Uploaded in accordance with the publisher's self-archiving policy.

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

1 **Next Generation Sequencing (NGS) to improve the diagnosis and management**  
2 **of patients with Disorders of Sex Development (DSD).**

3 Hughes, L.A.<sup>1</sup>, McKay-Bounford, K.<sup>1</sup>, Webb, E.A.<sup>2</sup>, Dasani P.<sup>1</sup>, Clokie, S.<sup>1</sup>, Chandran, H, McCarthy,  
4 L.<sup>2</sup>, Mohamed,Z.<sup>2</sup>, Kirk, J.M.W.<sup>2</sup>, Krone, N.P.<sup>2</sup>, Allen, S.<sup>1</sup>, Cole, T.R.P.<sup>1</sup>.

5 <sup>1</sup>West Midlands Regional Genetics Service, Birmingham Women's and Children's NHS Foundation Trust, Birmingham, B15  
6 2TG, United Kingdom

7 <sup>2</sup>Department of Endocrinology & Diabetes, Birmingham Women's and Children's Hospital, Steelhouse Lane, Birmingham, B4  
8 6NH, United Kingdom

9 Corresponding author and to whom requests for reprints should be addressed:

10 Dr Trevor Cole, Clinical Genetics Unit, West Midlands Regional Genetics Service, Birmingham  
11 Women's and Children's NHS Foundation Trust and Birmingham Health Partners, Mindelsohn Way,  
12 Birmingham, B15 2TG

13 Phone: 0121 335 8024 Fax: 0121 627 2618 Email: trevor.cole1@nhs.net

14

15 Keywords: Next generation sequencing (NGS), disorders of sex development (DSD), gene, variant

16

17 The authors declare that there is no conflict of interest that could be perceived as prejudicing the  
18 impartiality of the research reported

19

20 This research did not receive any specific grant from any funding agency in the public, commercial or  
21 not-for-profit sector.

22

23 **Abstract**

24 Disorders of sex development (DSDs) are a diverse group of conditions where the chromosomal,  
25 gonadal or anatomical sex can be atypical. The highly heterogeneous nature of this group of  
26 conditions often makes determining a genetic diagnosis challenging. Prior to Next Generation  
27 Sequencing (NGS) technologies, genetic diagnostic tests were only available for a few of the many  
28 DSD associated genes, which consequently had to be tested sequentially. Genetic testing is key in  
29 establishing the diagnosis, allowing for personalised management of these patients. Pinpointing the  
30 molecular cause of a patient's DSD can significantly impact patient management by informing future

31 development needs, altering management strategies and identifying correct inheritance pattern when  
32 counselling family members.

33 We have developed a 30 gene NGS panel, designed to be used as a frontline test for all suspected  
34 cases of DSD (both 46,XX and 46,XY cases). We have confirmed a diagnosis in 25 of the 80 patients  
35 tested to date. Confirmed diagnoses were linked to mutations in *AMH*, *AMHR2*, *AR*, *HSD17B3*,  
36 *HSD3B2*, *MAMLD1*, *NR5A1*, *SRD5A2* & *WT1* which have resulted in changes to patient  
37 management. The minimum diagnostic yield for patients with 46,XY DSD is 25/73. In 34/80 patients  
38 only benign or likely benign variants were identified, and in 21/80 patients only variants of uncertain  
39 significance, (VOUS) were identified, resulting in a diagnosis not being confirmed in these individuals.  
40 Our data supports previous studies, that an NGS panel approach is a clinically useful and cost  
41 effective frontline test for patients with DSDs.

42

### 43 **Introduction**

44 Disorders of sex development (DSD) encompass a wide range of conditions with diverse clinical  
45 features, pathophysiology and clinical management<sup>1-3</sup>. The recently revised stratified DSD diagnostic  
46 pathway consists of clinical examination, biochemical investigations and karyotype determination<sup>4,5</sup>.  
47 Once a presumptive diagnosis has been made, targeted sequencing of candidate genes may then be  
48 performed at a later stage<sup>5</sup>. Whilst reaching the correct diagnosis impacts significantly on  
49 management decisions, determining the aetiology of genital ambiguity in patients with DSD on the  
50 basis of clinical and biochemical assessment remains challenging<sup>6</sup>. In DSD where no clear  
51 abnormality in the steroidogenesis pathway is present, the yield from genetic testing had historically  
52 remained low, and with single-gene sequencing was both costly and time consuming<sup>4</sup>.

53 Improvements in gene sequencing technology in conjunction with rapidly falling costs have led to the  
54 use of targeted next-generation sequencing (NGS) assays. These enable multiple known disease  
55 causative genes to be sequenced in parallel alongside initial clinical assessment and biochemical  
56 investigations, potentially avoiding the need for additional expensive biochemical and radiological  
57 investigations<sup>7</sup>. Reaching a timely diagnosis is extremely important as it ends diagnostic uncertainty,  
58 avoids further unnecessary investigations, enables appropriate disease-specific counseling (including  
59 assessment of future fertility potential and malignancy risk) and implementation of personalised

60 medical plans in accordance with current disease specific consensus guidelines<sup>8</sup>. Additionally, in the  
61 long-term accurate early diagnosis will support the development of better designed outcome studies.  
62 The use of targeted next generation sequencing panels for molecular diagnosis of DSD patients has  
63 already been reported successfully in several previous publications<sup>9–12</sup>. These publications show  
64 diagnostic yield and clinical utility in predominantly 46,XY DSD cohorts using panels of 64-219 genes.  
65 We present our data showing sequence analysis of a cohort of 80 DSD patients using a 30 gene  
66 panel.

## 67 **Materials & Methods**

### 68 **Patients**

69 Eighty patients with a DSD were referred to the West Midlands Regional Genetics Laboratory  
70 (WMRGL) at the Birmingham Women's and Children's NHS Foundation Trust for diagnostic DSD  
71 testing between March 2014 and March 2017, comprising of 73 patients with 46,XY DSD and 7 with  
72 46,XX DSD. Referrals were from Clinical Genetics, Urology or Endocrinology specialists. Karyotyping  
73 and/or microarray results were typically available to confirm the patient's karyotype. Single gene  
74 testing may have also been performed in advance of the NGS screen but a pathogenic mutation had  
75 not been identified. Where DNA was available, cascade testing was performed on parental or sibling  
76 samples to confirm segregation or to confirm a diagnosis in the proband's similarly affected siblings.  
77 Data from cascade testing is not included here; all figures therefore represent only probands referred  
78 for diagnostic testing who underwent analysis via the NGS panel. Consent was obtained for clinical  
79 testing from all patients in this study. Patients undertaking routine clinical testing in this report are not  
80 identifiable. This report has been registered with the audit committee at the Birmingham Women's and  
81 Children' Hospital NHS Foundation trust (CARMS-30120).

### 82 **Gene selection**

83 Thirty genes with a reported clinical association with a DSD were selected following discussion  
84 between the WMRGL and clinical specialists in Genetics and Endocrinology (table 1). Genes include  
85 those thought to be involved in 46,XY DSD and 46,XX DSD, and are tested as a single panel pipeline  
86 covering both of these groups of patients. The *CYP21A2* gene associated with 95% of cases of  
87 Congenital Adrenal Hyperplasia (CAH) is not included in this panel. This is because this patient group  
88 typically have a clinical diagnosis prior to genetic testing, and also the *CYP21A2* pseudogene makes  
89 accurate mapping of short reads to the functional gene very difficult.

## 90 **Sample preparation**

91 Genomic DNA was typically extracted from peripheral blood samples using Qiasymphony technology  
92 (Qiagen) following the manufacturer's instructions. The concentration of all genomic DNA samples  
93 were assessed using a Qubit (Life technologies) prior to sequencing.

## 94 **Next Generation Sequencing (NGS)**

95 Library preparation was initially performed by a customised TruSeq Custom Amplicon (TSCA, Illumina  
96 Inc, San Diego, California) 30 gene panel run on the MiSeq (Illumina Inc, San Diego). Exons were  
97 targeted with 25 base pairs of padding on either side, resulting in 431 amplicons of 425bp.  
98 Enrichment was performed on 250ng of genomic DNA, and sequencing using 250 base paired-end  
99 reads. TruSight One (TSO) technology was then used to capture exonic regions of 4,813 genes,  
100 sequencing 24 samples on a HiSeq 2500 (Illumina Inc, San Diego, California). The same 30 DSD  
101 genes were analysed as a virtual panel using TSO. A depth of coverage of 20x was considered  
102 sufficient for either approach, and a technical report was generated indicating the proportion of each  
103 gene covered to this level. All patients that were initially run by TSCA but where a diagnosis was not  
104 confirmed were subsequently retested using TSO. NGS was performed on probands only and  
105 analysis for copy number variation in these genes was not performed.

## 106 **Bioinformatic analysis**

107 Bioinformatic analysis was performed using an in-house pipeline where sequence reads were  
108 mapped to the human genome hg19 reference. Several programmes are incorporated in the pipeline;  
109 Trimmomatic (quality trimming of reads), BWA mem (alignment to hg19), Samblaster (duplicate  
110 marking), Abra (realignment), Platypus (variant calling), Annovar (variant annotation) PLINK (IBS  
111 calculation) and Picard (calculating hybridisation and mapping metrics). Custom python code and  
112 bedtools were used to calculate coverage and the Python module pandas to produce patient specific  
113 Excel files.

## 114 **Variant interpretation and reporting**

115 Variants were classified following the Association of Clinical Genetic Science (ACGS,  
116 [www.acgs.uk.com](http://www.acgs.uk.com)) best practice guidelines, based on the American College of Medical Genetics and  
117 genomics recommendations<sup>13</sup>. This included utilisation of in-house frequency data, population  
118 frequency data (dbSNP, 1000 genomes & EXAC), in-silico tools including Polyphen, Align GVGD and  
119 splice tools (searched through the alamut interface), the Human Gene Mutation Database (HGMD  
120 Professional, Biobase Corporation), and evidence from peer-reviewed literature. The five classes are  
121 described in table 2. For suspected compound heterozygous mutations, parental samples were

122 requested to confirm that the mutations were on opposite alleles (in trans). Regions of interest were  
123 all exonic regions plus 30bp upstream and 10bp downstream of each exon. Intronic variants outside  
124 of these regions were considered as deep intronic variants (DIVs) and no further investigation was  
125 undertaken. In some cases, Sanger sequencing was performed to complete gene coverage to a  
126 depth of 20x. For example, where a single heterozygous mutation in a likely candidate gene  
127 associated with a recessive condition had been identified.

128

## 129 **Results**

130 The DSD NGS panel provided a molecular diagnosis in 25 out of the 80 patients tested (table 3). A  
131 diagnosis was deemed as confirmed where variants of class 4 or class 5 were identified which were  
132 consistent with the inheritance pattern for that gene, and where disruption of the gene was in keeping  
133 with the patient's phenotype. On 4 occasions a class 3 variant was found in combination with a class  
134 5 mutation (patients 2, 11, 15 & 20). Although technically class 3 variants are of uncertain  
135 significance, their presence in combination with a class 5 variant in these patients, when considered  
136 with the clinical information provided led us to believe that these findings were causally related to the  
137 clinical features. The overall diagnostic yield for this panel is therefore currently 25/80 for all samples  
138 and 25/73 for 46,XY DSD. This figure represents the likely minimum detection rate of this panel as  
139 some samples were received with an initial request for Sanger sequencing of a specific gene, and  
140 typically only included for panel testing if negative on Sanger sequencing.

141

142 Class 3 variants (listed in table 4) were typically missense mutations which had not been previously  
143 reported, and therefore no clinical information was available. These were included in the clinical report  
144 with a statement that a diagnosis had not been confirmed due to the uncertainty around the  
145 pathogenicity of such variants. Where only class 1 and/or 2 variants were identified, patient reports  
146 stated that no evidence of a pathogenic mutation had been identified. A summary of the findings can  
147 be seen in figure 1. Both previously reported and novel pathogenic mutations and variants were  
148 identified in *AMH*, *AMHR2*, *AR*, *DHCR7*, *HSD17B3*, *HSD3B2*, *LHCGR*, *MAMLD1*, *NR5A1*, *SRD5A2*  
149 *and WT1* (table 5). Diagnosis due to mutations in the *AR* gene (7 patients) were the most commonly  
150 observed (table 3) followed by diagnosis due to HSD17B3 (5 patients) and SRD5A2 (4 patients).

151

## 152 **Discussion**

153 DSDs, estimated to be present in 1.7% of live births<sup>14</sup> are a diagnostic challenge due to variable  
154 expressivity and pleiotrophy, clinical overlap of the different DSD, and their significant aetiological  
155 heterogeneity. Historically a genetic diagnosis was made in as few as 13% of cases<sup>15</sup>. We present  
156 data from 80 patients who underwent routine diagnostic testing for DSD using a 30 gene NGS panel.  
157 This diagnostic DSD panel was utilised irrespective of clinical and biochemical features, unless a  
158 specific single Sanger sequence request was made based on phenotypic assessment. The diagnostic  
159 yield of this DSD panel was shown to be 25/80 for all DSD's, higher for 46,XY DSD (25/73), and  
160 would have been higher in this cohort if all cases with a suspected diagnosis, (all subsequently  
161 confirmed on Sanger Sequencing) , had not been filtered out prior to implementation of the panel  
162 test. Pathogenic (or likely pathogenic) mutations in the *AMH*, *AMHR2*, *AR*, *HSD17B3*, *HSD3B2*,  
163 *MAMLD1*, *NR5A1*, *SRD5A2* and *WT1* genes were identified. Our detection rate and findings are  
164 similar to previous studies, summarised in Table 6. Dong *et al* demonstrated an increased detection  
165 rate of 9/13 in 46,XY DSD patients using a panel of 219 genes<sup>10</sup>, however the study included small  
166 patient numbers and so may not be representative.

167

168 Separating out analysis of 46,XY DSD from those with 46,XX DSD results in an improved 46,XY yield  
169 to 25/73 but highlights the 0/7 diagnostic yield of individuals with a 46,XX DSD. Difficulty in confirming  
170 a molecular diagnosis in those with an 46,XX DSD has also been seen in other studies<sup>11,12</sup>. Sample  
171 numbers for those with 46,XX DSD are very small in this study and therefore are unlikely to be  
172 representative of the true diagnostic capability of the panel for these patients. In addition, other  
173 causes of 46,XX DSD such as translocation of SRY to the X chromosome, duplications of *SOX9* or  
174 CAH due to *CYP21A2* deficiency are not detectable by this method. Patients would typically have had  
175 karyotype and/or microarray prior to testing on the panel and would have been tested separately for  
176 *CYP21A2* deficiency if CAH was suspected. Increased sample numbers and incorporation of more  
177 46,XX DSD associated genes as they are identified may allow a more accurate estimate of the  
178 panel's usefulness for those with 46,XX DSD. It will also be important to include new 46,XX DSD  
179 genes that are likely to be identified in current international exome/genome sequencing projects such  
180 as the 100,000 genome project in England.<sup>16</sup>

181

182 Novel mutations and variants in several genes were identified where functional studies were not  
183 available. Variants were considered likely to be causative if they were observed in trans (on opposite  
184 chromosome alleles) with a known pathogenic mutation, in a disease gene showing autosomal

185 recessive inheritance (4 patients in our cohort). In some cases, segregation studies confirming the bi-  
186 allelic nature of the findings also supported a likely pathogenic role. This information has expanded  
187 our knowledge of likely diagnostic DSD variants for future investigation of DSD patients. Novel VOUS  
188 in the absence of a confirmed diagnosis were also seen. Whilst their significance currently remains  
189 uncertain, wider data sharing through publication of studies such as this is crucial to further our  
190 understanding of such variants. The large number of VOUS in this cohort is predominantly due to  
191 limited clinical information related to some DSD related genes, for example only VOUS were detected  
192 in the *CBX2* gene. Clinicians will have to manage any patient confusion or anxiety within the current  
193 uncertainty until more data is available. When designing future NGS DSD panels it will remain  
194 important to recognise that increasing the number of genes, especially those with limited data, will  
195 generate greater numbers of VOUS with increased cost and complexity of analysis. This should be  
196 balanced against the potential for gene discovery.

197

198 Segregation studies have been helpful in determination of pathogenicity in several families, as  
199 illustrated by patient 23, who had an Androgen Receptor (*AR*) variant initially reported as a VOUS.  
200 The patient's older brother, previously shown to have a 47,XXY karyotype and a "milder" phenotype  
201 was subsequently shown to have the same *AR* variant. Given his additional X chromosome, without  
202 skewed X inactivation a milder phenotype would be expected. This increased our confidence in calling  
203 the variant a "likely pathogenic" mutation and highlights the importance of reviewing interpretation of  
204 variants when new information becomes available.

205

206 NGS technology has also allowed the identification of mosaic mutations in the *AR* gene, which may  
207 have gone undetected by other methods. Identification of an accurate number of reads enhances our  
208 knowledge of the level of mosaicism present. It should be noted, however, that the results represent  
209 the mutation load in peripheral blood (70% patient 8 and 30% patient 24) and not necessarily other  
210 relevant tissues. Of the mutations identified, both had been previously reported in non-mosaic form in  
211 the literature<sup>17,18</sup>. Mosaic mutations have however been reported in the *AR* gene<sup>19,20</sup> and taken  
212 together with the clinical features, allowed us to conclude that these mutations in mosaic form were  
213 likely to be contributing to the phenotype in these patients. Importantly for patient management, *AR*  
214 mutations in mosaic form are believed to pose a risk of virilisation in patients due to the presence of  
215 wild type androgen receptor<sup>19</sup>. Distinguishing mosaic from non-mosaic forms therefore can have  
216 significant consequences for patient management and genetic recurrence risk.



217

218 Identifying the correct genetic diagnosis modifies the patient management and impacts on the  
219 accuracy of information and choices available to family members. The former is clearly illustrated in  
220 case 21 where mutations in two different DSD related genes were present in the same family;  
221 *HSD17B3* and *AR*. The family were requesting gonadectomy pre-adolescence in a 46,XY DSD  
222 female, on the basis that her cousins who were 46,XY DSD females, due to a homozygous *HSD17B3*  
223 mutations, had virilised and been managed with gonadectomy in another centre. Identification of a  
224 previously unidentified mutation in *AR* in this family meant 3 girls with 46,XY changed their  
225 subsequent management. In 4 cases the diagnosis in affected siblings has been confirmed (listed in  
226 table 3 patients 7, 17, 21 & 23) including cases enabling early prenatal (patient 20) or neonatal  
227 (patient 17) diagnosis, and thus implementation of appropriate management from birth. This highlights  
228 the importance of identifying the molecular diagnosis not only for the proband but also for the wider  
229 family.

230 The panel also identified pathogenic mutations which were thought to be co-incidental and not to be  
231 related to the initial clinical presentation in the proband. In patient 18 with Congenital Adrenal  
232 Hyperplasia (CAH) due to *HSD3B2* deficiency, the patient was also shown to be a carrier for the  
233 common splice mutation c.964-1G>C in the *DHCR7* gene which is linked to Smith Lemli Opitz  
234 syndrome (SLOS). Whilst such incidental findings can be challenging for patient counselling, the  
235 information provided may also be of great significance. This will be especially true when previously  
236 undetected autosomal recessive mutations are uncovered in highly consanguineous families, which  
237 are common within many DSD cohorts.

238

239 Where a clinician has a strong suspicion of the involvement of a specific gene, Sanger sequencing  
240 may be more cost effective, especially where the number of amplicons is relatively small. For  
241 example pathogenic mutations in the *AMH*, *AMHR2*, *AR* and *SRD5A2* gene were identified in this  
242 way. When the original clinical diagnosis is incorrect however, the potential cost savings rapidly  
243 disappear if sequential Sanger sequencing is required, and therefore any benefit is highly dependent  
244 on the clinical expertise and the specificity of the additional non-genetic investigations. There may  
245 also be atypical presentations not yet recognised for mutations in some genes, and therefore a wider  
246 panel approach has the potential to address this.

247

248 Despite the diagnostic rate of 25/73 for 46,XY DSD, the failure to achieve a diagnosis in 55 patients  
249 confirms the need for further development. Of note, the panel cannot currently detect copy number  
250 changes (CNVs), and so further development should include detection of CNVs that are below the  
251 resolution of the current chromosomal microarray assay. The 30 genes in our panel were selected  
252 due to published evidence of their involvement in DSD, but new gene discoveries in studies such as  
253 the 100,000 genome project<sup>16</sup> should enhance the diagnostic utility, especially for 46,XX DSD. Eggers  
254 et al demonstrated a 118/278 diagnostic rate for 46,XY DSD patients by including 64 DSD genes<sup>11</sup>,  
255 and Dong et al reported a 6/13 rate by including 219 genes<sup>10</sup>, thus diagnostic capability may be  
256 improved by increasing the gene number. These benefits may be marginal however as only one likely  
257 pathogenic mutation in the Dong series was in the additional 179 genes not included in this 30 gene  
258 panel. Comparison of these studies could indicate that a greater impact on the detection rate may be  
259 due to patient selection. It will be important, therefore, that further candidate gene inclusion is critically  
260 evaluated as the addition of genes without clear clinical utility will likely result in increasing cost and  
261 numbers of VOUS without necessarily increasing diagnostic capability. The future of this investigative  
262 pathway may well be transformed by implementation of whole exome or whole genome sequencing,  
263 but any benefits of diagnostic detection will have to be weighed against increased cost and clinical  
264 complexities resulting from VOUS and co-incidental findings. It therefore remains important to  
265 optimise such NGS panels for DSD so that a valid comparison can be made in future.

266

267 This data demonstrates clear advantages of an NGS panel approach for highly heterogenous  
268 conditions such as DSD. Despite the limitations of the panel including incomplete coverage and  
269 inability to detect copy number changes, the results presented here demonstrate that an NGS based  
270 panel approach is a useful frontline tool for diagnosing DSDs. In addition to a diagnostic yield of at  
271 least 25/80 we have shown examples of cases where the information provided from the panel has  
272 identified diagnoses in complex families with the potential for multiple aetiologies, cases where panel  
273 findings have significantly impacted management and treatment decisions, and examples of novel  
274 variants being identified, thus expanding our current knowledge. As more and more patients are  
275 tested, the information provided by such panels will continue to grow and improve our understanding  
276 of these complex conditions and hopefully improve the diagnostic capability of such tests. Despite its  
277 limitations, the clinical benefit of this approach is clearly demonstrated for DSD patients allowing for  
278 timely accurate diagnoses, more informed management strategies and improved counselling for  
279 patients and their families.

280

## 281 Declaration of Interest

282 The authors declare that there is no conflict of interest that could be perceived as prejudicing the  
283 impartiality of the research reported

284

## 285 Funding

286 This work did not receive any specific grant from any funding agency in the public, commercial, or not-  
287 for-profit sector.

288

289

## 290 References

- 291 1. Biason-Lauber, A. Control of sex development. *Best Pr. Res Clin Endocrinol Metab* **24**, 163–  
292 186 (2010).
- 293 2. Higashi, Y., Yoshioka, H., Yamane, M., Gotoh, O. & Fujii-Kuriyama, Y. Complete nucleotide  
294 sequence of two steroid 21-hydroxylase genes tandemly arranged in human chromosome: a  
295 pseudogene and a genuine gene. *Proc Natl Acad Sci U S A* **83**, 2841–2845 (1986).
- 296 3. Arboleda, V. A., Sandberg, D. E. & Vilain, E. DSDs: genetics, underlying pathologies and  
297 psychosexual differentiation. *Nat. Rev. Endocrinol.* **10**, 603–15 (2014).
- 298 4. Ahmed, S. F., Achermann, J. C., Arlt, W., Balen, A. H., Conway, G., Edwards, Z. L., Elford, S.,  
299 Hughes, I. A., Izatt, L., Krone, N. *et al.* UK guidance on the initial evaluation of an infant or an  
300 adolescent with a suspected disorder of sex development. *Clin. Endocrinol. (Oxf)*. **75**, 12–26  
301 (2011).
- 302 5. Lee, P. A., Nordenström, A., Houk, C. P., Ahmed, S. F., Auchus, R., Baratz, A., Baratz Dalke,  
303 K., Liao, L.-M., Lin-Su, K., Looijenga, L. H. J. *et al.* Global Disorders of Sex Development  
304 Update since 2006: Perceptions, Approach and Care. *Horm. Res. Paediatr.* **85**, 158–80  
305 (2016).
- 306 6. Lee, Y. S., Kirk, J. M., Stanhope, R. G., Johnston, D. I., Harland, S., Auchus, R. J., Andersson,  
307 S. & Hughes, I. A.. Phenotypic variability in 17beta-hydroxysteroid dehydrogenase-3 deficiency  
308 and diagnostic pitfalls. *Clin Endocrinol* **67**, 20–28 (2007).

- 309 7. Ahmed, S. F., Bashamboo, A., Lucas-Herald, A. & McElreavey, K. Understanding the genetic  
310 aetiology in patients with XY DSD. *Br. Med. Bull.* **106**, 67–89 (2013).
- 311 8. Lee, P. A., Houk, C. P., Ahmed, S. F., Hughes, I. A. & Endocrinology, I. C. C. on I. organized  
312 by the L. W. P. E. S. and the E. S. for P. Consensus statement on management of intersex  
313 disorders. International Consensus Conference on Intersex. *Pediatrics* **118**, e488-500 (2006).
- 314 9. Baxter, R. M., Arboleda, V. A., Lee, H., Barseghyan, H., Adam, M. P., Fechner, P. Y.,  
315 Bargman, R., Keegan, C., Travers, S., Schelley, S. *et al.* Exome sequencing for the diagnosis  
316 of 46,XY disorders of sex development. *J. Clin. Endocrinol. Metab.* **100**, E333-44 (2015).
- 317 10. Dong, Y., Yi, Y., Yao, H., Yang, Z., Hu, H., Liu, J., Gao, C., Zhang, M., Zhou, L., Asan, Yi, X. &  
318 Liang, Z. Targeted next-generation sequencing identification of mutations in patients with  
319 disorders of sex development. *BMC Med. Genet.* **17**, 23 (2016).
- 320 11. Eggers, S., Sadedin, S., van den Bergen, J. A., Robevska, G., Ohnesorg, T., Hewitt, J.,  
321 Lambeth, L., Bouty, A., Knarston, I. M., Tan, T. Y. *et al.* Disorders of sex development: insights  
322 from targeted gene sequencing of a large international patient cohort. *Genome Biol.* **17**, 243  
323 (2016).
- 324 12. Kim, J. H., Kang, E., Heo, S. H., Kim, G.-H., Jang, J.-H., Cho, E.-H., Lee, B. H., Yoo, H.-W. &  
325 Choi, J.-H. Diagnostic yield of targeted gene panel sequencing to identify the genetic etiology  
326 of disorders of sex development. *Mol. Cell. Endocrinol.* **444**, 19–25 (2017).
- 327 13. Richards, S., Aziz, N., Bale, S., Bick, D., Das, S., Gastier-Foster, J., Grody, W. W., Hegde, M.,  
328 Lyon, E., Spector, E. *et al.* Standards and guidelines for the interpretation of sequence  
329 variants: a joint consensus recommendation of the American College of Medical Genetics and  
330 Genomics and the Association for Molecular Pathology. *Genet. Med.* **17**, 405–423 (2015).
- 331 14. Blackless, M., Charuvastra, A., Derryck, A., Fausto-Sterling, A., Lauzanne, K. & Lee, E. How  
332 sexually dimorphic are we? Review and synthesis. *Am. J. Hum. Biol.* **12**, 151–166 (2000).
- 333 15. Arboleda, V. A., Lee, H., Sánchez, F. J., Délot, E. C., Sandberg, D. E., Grody, W. W., Nelson,  
334 S. F. & Vilain, E. Targeted massively parallel sequencing provides comprehensive genetic  
335 diagnosis for patients with disorders of sex development. *Clin. Genet.* **83**, 35–43 (2013).
- 336 16. Siva, N. UK gears up to decode 100,000 genomes from NHS patients. *Lancet* **385**, 103–104

- 337 (2015).
- 338 17. Galani, A., Sofocleous, C., Karahaliou, F., Papathanasiou, A., Kitsiou-Tzeli, S. & Kalpini-  
339 Mavrou, A. Sex-reversed phenotype in association with two novel mutations c.2494delA and  
340 c.T3004C in the ligand-binding domain of the androgen receptor gene. *Fertil. Steril.* **90**,  
341 2008.e1-2008.e4 (2008).
- 342 18. Marcelli, M., Tilley, W. D., Wilson, C. M., Wilson, J. D., Griffin, J. E. & McPhaul, M. J. A single  
343 nucleotide substitution introduces a premature termination codon into the androgen receptor  
344 gene of a patient with receptor-negative androgen resistance. *J. Clin. Invest.* **85**, 1522–8  
345 (1990).
- 346 19. Holterhus, P. M., Wiebel, J., Sinnecker, G. H., Brüggewirth, H. T., Sippell, W. G., Brinkmann,  
347 A. O., Kruse, K. & Hiort, O. Clinical and molecular spectrum of somatic mosaicism in androgen  
348 insensitivity syndrome. *Pediatr. Res.* **46**, 684–90 (1999).
- 349 20. Köhler, B., Lumbroso, S., Leger, J., Audran, F., Grau, E. S., Kurtz, F., Pinto, G., Salerno, M.,  
350 Semitcheva, T., Czernichow, P. *et al.* Androgen Insensitivity Syndrome: Somatic Mosaicism of  
351 the Androgen Receptor in Seven Families and Consequences for Sex Assignment and  
352 Genetic Counseling. *J. Clin. Endocrinol. Metab.* **90**, 106–111 (2005).

353


354

355

356 **Figure 1: Summary of findings of DSD panel.**

357 Results of the panel are separated by karyotype (XX or XY) and by result. ‘Diagnosis  
358 confirmed’ indicates patients where a pathogenic variant was detected compatible with the  
359 patient’s phenotype. ‘VOUS (Variant Of Uncertain Significance) only’ indicates solely Class 3  
360 variants were detected and therefore a diagnosis could not be confirmed. ‘No mutation’  
361 indicates only Class 1 (polymorphisms) or Class 2 variants (unlikely to be pathogenic) were  
362 detected.

363



184x126mm (96 x 96 DPI)

46,XY DSD	Gene Name	Location	
Disorders of testicular development	ARX	Xp22.13	
	ATRX	Xq13.3	
	CBX2	17q25	
	DHH	12q13.1	
	DMRT1	9p24.3	
	MAMLD1	Xq28	
	NR0B1	Xp21.3	
	NR5A1	9q33	
	SOX9	17q24-q25	
	SRY	Yp11.3	
	TSPYL1	6q22-23	
	WNT4	1p35	
	WT1	11p13	
	Disorders of Hormone synthesis or action	AMH	19p13.3-p13.2
		AMHR2	12q13
AR		Xq11-q12	
CYB5A		18q23	
CYP11A1		15q23-24	
CYP17A1		10q24.3	
DHCR7		11q12-q13	
HSD3B2		1p13.1	
HSD17B3		9q22	
LHCGR		2p21	
POR		7q11.2	
SRD5A2		2p23	
StAR		8p11.2	

**Table 1a**

**Table 1: Genes included in the DSD panel**

A summary of all genes and their chromosomal location which are included in the panel for both 46,XY DSD (table 1a) and 46,XX DSD (table 1b). CAG repeat in *AR* not analysed to avoid incidental diagnosis of Spinal bulbar muscular atrophy.

46,XX DSD	Gene Name	Location
Disorders of ovarian development	RSPO1	1p34.3
	SOX9	17q24
	SRY	Yp11.3
	WNT4	1p35
Androgen Excess	CYP11B1	8q21-q22
	CYP19A1	15q21
	HSD3B2	1p13
	NR3C1	5q31
	POR	7q11.2

**Table 1b**

Variant class	Variant description	Confirmation by Sanger	Reported
1	Polymorphism	No	No
2	Unlikely to be pathogenic	No	Yes
3	Variant of uncertain significance (VOUS)	Yes	Yes
4	Likely to be pathogenic	Yes	Yes
5	Clearly pathogenic	Yes	Yes

**Table 2: Classification and follow up of variants.** All variants determined to be class 3-5 were confirmed using Sanger sequencing and all were included in the clinical reports. Variants considered to be unlikely to be pathogenic (class 2) were not confirmed by Sanger sequencing and were recorded for information only in the clinical report. Class 1 variants were not reported.



Patient	Sex	Reason for referral	Gene	Allele 1	Allele 2	Results reported and follow up
1	XY M	?PAIS	AR	c.2402C>T p.(Thr801Ile)		Confirmed diagnosis of PAIS.
2	XY M	?PMDS	AMH	c.283C>T p.(Arg95*)	c.905G>A p.(Arg302Gln)	Consistent with diagnosis of PMDS.
3	XY F	?46,XY DSD	NR5A1	c.1171A>T p.(Lys391*)	Normal	Consistent with a diagnosis of a 46,XY DSD. <b>Novel mutation.</b>
4	XY <sup>P</sup> F	?46,XY DSD	HSD17B3	c.614T>A p.(Val205Glu)	c.645A>T p.(Glu215Asp)	Supports diagnosis of 46,XY DSD due to HSD17B3 deficiency.
5	XY <sup>P</sup> F	?46,XY DSD	HSD17B3	c.194C>T p.(Ser65Leu)	c.729_735del7 p.(Ile244Argfs*11)	Supports diagnosis of 46,XY DSD due to HSD17B3 deficiency.
6	XY <sup>P</sup> M	?46,XY DSD	SRD5A2	c.698+1G>T	c.698+1G>T	Consistent with diagnosis of 46,XY due to SRD5A2 deficiency.
7	XY <sup>P</sup> M	X-linked hypospadias	MAMLD1	c.1366C>T p.(Arg456*)		Consistent with MAMLD1 associated hypospadias. Confirmed in 2 affected brothers and mother (carrier). Carrier of PMDS (c.35T>G p.(Val12Gly) in AMH).
8	XY <sup>P</sup> M	?46,XY DSD	AR	c.2391G>A p.(Trp797*)		Mosaic (70% of reads). Likely causally related to clinical features.
9	XY F	?XY DSD	NR5A1	c.69 C>A p.(Tyr23*)	Normal	Consistent with diagnosis of 46,XY DSD. <b>Novel mutation.</b>
10	XY F	?46,XY DSD	HSD17B3	c.695C>T p.(Ser232Leu)	c.695C>T p.(Ser232Leu)	Confirms diagnosis 46,XY DSD due to HSD17B3 deficiency.
11	XY <sup>P</sup> M	?PMDS	AMHR2	c.813_817delGCTCT, p.(Leu272Trpfs*24)	c.931G>A, p.(Gly311Ser)	Consistent with features of PMDS. <b>Novel mutation and novel variant.</b>
12	XY F	?46,XY DSD	SRD5A2	c.737G>A, p.(Arg246Gln)	c.737G>A, p.(Arg246Gln)	Consistent with diagnosis of SRD5A2 deficiency.
13	XY <sup>P</sup> M	Penoscrotal hypospadias	SRD5A2	c.586G>A, p.(Gly196Ser)	c.586G>A, p.(Gly196Ser)	Consistent with diagnosis of SRD5A2 deficiency assuming XY.
14	XY M	Gynaecomastia, Hypospadias, micropenis	AR	c.2057T>C p.(Val686Ala)		Consistent with clinical features. Confirmed inherited from mother.
15	XY M	Ambiguous genitalia	HSD17B3	c.277+4A>T	c.133C>T p.(Arg45Trp)	Consistent with clinical features. c.13C>T p.(Arg45Trp) is <b>novel variant.</b>
16	XY F	?AIS	AR	c.2343G>A p.(Met 781Ile)		Consistent with diagnosis of AIS.
17	XY M	?PMDS	AMHR2	c.289C>T p.(Arg97*)	c.289C>T p.(Arg97*)	Confirms diagnosis of PMDS. Both parents carriers. Also had another child affected child who was homozygous for the same mutation (detected in neonatal period).
18	XY M	?XY DSD	HSD3B2	c.518T>G p.(Leu173Arg)	c.518T>G p.(Leu173Arg)	Confirms diagnosis of CAH due to HSD3B2 deficiency. Both parents are carriers. Patient also a carrier of the c.964-1G>C splice mutation in DHCR7.
19	XY <sup>P</sup> F	?46, XY DSD	HSD17B3	c.277+4A>T	c.645A>T p.(Glu215Asp)	Confirmed diagnosis of 46, XY DSD due to 17-Beta Hydroxysteroid dehydrogenase deficiency. Each parent carries 1 mutation.
20	XY M	Undervirilised male	SRD5A2	c.307C>T p.(Arg103*)	c.107A>G, p.(His36Arg)	Consistent with clinical features. Parental samples confirmed compound heterozygous. Follow up biochemical testing confirmed SRD5A2 deficiency. <b>Novel variant.</b>
21	XY F	?XY DSD	AR	c.2407dupC p.(Gln803Profs*27)		Confirms diagnosis of AIS. Two affected siblings also have mutation. <b>Novel mutation.</b> Also heterozygous for HSD17B3 familial mutation. c.803G>A p.(Cys268Tyr).
22	XY M	Ambiguous genitalia	WT1	c.1087A>T p.(Arg363*)	Normal	May be contributing to features. Confirmed de-novo. Tumour screening initiated.
23	XY M	Severe hypospadias	AR	c.2384T>A p.(Phe795Tyr)		Initially reported as VOUS. Once identified in affected (milder) brother who's karyotype was 47,XXY more confident that linked to features. <b>Novel variant.</b>
24	XY M	Severe hypospadias and penile transposition	AR	c.2645T>C p.(Leu882Pro)		Mosaic (30% of reads). Likely causally related to phenotype.
25	XY M	?PMDS	AMH	c.649C>T p.(Gln217*)	c.649C>T p.(Gln217*)	Consistent with diagnosis of PMDS. <b>Novel mutation.</b>

**Table 3: Patient details with a confirmed molecular diagnosis**

Details of mutations and variants found in patient reported with a confirmed molecular diagnosis. Details of the karyotypic and phenotypic sex are in the second column with M and F representing phenotypic sex. P indicates presumed karyotype (reports not seen) from SRY sequence reads. Pathogenic mutations linked to the diagnoses listed in “gene column”. Allele 1 and 2 describe the mutations in the different alleles, a black box indicates an absent second X allele in XY individuals. Green = Normal (wildtype), Yellow = Class 3 variant (Variant Of Uncertain Significance (VOUS)), Orange = Class 4 variant (Likely pathogenic), Red = Class 5 variant (Clearly pathogenic). AIS = Androgen Insensitivity Syndrome, CAH = Congenital Adrenal Hyperplasia, PAIS = Partial Androgen Insensitivity Syndrome, PMDS = Persistent Mullerian Duct Syndrome.

Patient	Sex	Reason for referral	Gene	Allele 1	Allele 2	Results reported and follow up
26	XY ?	N/A	POR	c.948-30G>A	Normal	A molecular diagnosis has not been confirmed
27	XY <sup>P</sup> M	1° gonadal failure, short stature	WT1	c.11C>G, p.(Pro4Arg)	Normal	A molecular diagnosis has not been confirmed
28	XY M	Mullerian resistant disorder	HSD17B3	c.133C>T p.(Arg45Trp)	Normal	A molecular diagnosis has not been confirmed
29	XY M	?46,XY DSD	RSPO1	c.658C>T p.(Arg220Trp)	Normal	A molecular diagnosis has not been confirmed
30	XY M	Hypogonadism	HSD3B2	c.809T>C p.(Ile270Thr)	Normal	A molecular diagnosis has not been confirmed
31	XY F	Facial dysmorphism, gastric motility issues, undescended testes, adrenal insufficiency	CBX2	c.1411C>G p.(Pro471Ala)	Normal	A molecular diagnosis has not been confirmed. #Variant found in alternative transcript.
			CBX2#	c.616C>T p.(Gln206*)	Normal	
32	XX M	Hypospadias	NR5A1	c.275G>A p.(Arg92Gln)	Normal	A molecular diagnosis has not been confirmed
33	XY F	? Gonadal dysgenesis	CYP11A1	c.1250T>G p.(Val417Gly)	Normal	A molecular diagnosis has not been confirmed
34	XY F	? CYP11A1 imbalance	CYP11A1	c.989C>T p.(Thr330Met)	Normal	A molecular diagnosis has not been confirmed
			MAMLDI	c.2009C>T p.(Thr670Ile)		
35	XY F	Primary ovarian failure	WT1	c.1493A>G p.(Glu498Gly)	Normal	A molecular diagnosis has not been confirmed
36	XY M	Severe penoscrotal hypospadias.	CBX2	c.1416C>G p.(Asp472Glu)	Normal	A molecular diagnosis has not been confirmed
			HSD3B2	c.500C>T p.(Ala167Val)	c.500C>T p.(Ala167Val)	
37	XY F	Tall stature, uterus present, no obvious ovaries	CBX2	c.1411C>G p.Pro471Ala	Normal	A molecular diagnosis has not been confirmed
			AMH	c.53C>T p.(Ala18Val)	Normal	
			AMH	c.1556C>T p.(Ala519Val)	Normal	
38	XX F	Premature ovarian failure	CYP11B1	c.1451T>A p.(Val484Asp)	Normal	A molecular diagnosis has not been confirmed
39	XY F	Clitoromegaly, no vaginal opening	AR	c.1174C>T p.(Pro392Ser)		Pathogenicity of variant uncertain due to conflicting evidence
40	XY M	Penoscrotal hypospadias, micropenis & undescended testes	CBX2	c.785G>A p.(Arg262Gln)	Normal	A molecular diagnosis has not been confirmed
41	XY M	Hypospadias and penoscrotal transposition	CYP11A1	c.940G>A p.(Glu314Lys)	Normal	A molecular diagnosis has not been confirmed
			HSD17B3	c.133C>T p.(Arg45Trp)	c.133C>T p.(Arg45Trp)	
42	XY M	?46,XY DSD	LHCGR	c.828delC p.(Ser277Alafs*32)	Normal	A molecular diagnosis has not been confirmed. Variant likely to be pathogenic but absence of second mutation leads to uncertain significance.
			CBX2	c.785G>A p.(Arg262Gln)	Normal	
43	XY M	Ambiguous genitalia	NR5A1	c.146G>A p.(Cys49Tyr)	Normal	A molecular diagnosis has not been confirmed. De novo variant.
44	XY <sup>P</sup> F	?46,XY DSD	NR5A1	c.1019C>T p.(Ala340Val)	Normal	A molecular diagnosis has not been confirmed. Maternally inherited.
45	XY M	Ambiguous genitalia	HSD17B3	c.202-22G>A	c.202-22G>A	A molecular diagnosis has not been confirmed. Recommend biochemical testing.
46	XX F	Ambiguous genitalia, complete labial fusion	LHCGR	c.458+3A>G	Normal	A molecular diagnosis has not been confirmed
			NR5A1	c.486C>T p.(=)	Normal	

Table 4a: Patients with Variants of Uncertain Significance (VOUS) where a diagnosis was not confirmed

Patient	Sex	Reason for referral	Gene	Allele 1	Allele 2	Results reported and follow up
3	XY F	?46,XY DSD	ATRX	c.2595C>G p.(His865Gln)		ATRX & AMH variants found in addition to NR5A1 class 4 mutation (table 3).
			AMH	c.-2C>T	Normal	
4	XY F	? 46,XY DSD	MAMLD1	c.2744A>C p.(Asp915Ala)		MAMLD1 variant found in addition to HSD17B3 mutations (table 3)
7	XY M	X-linkedhypospadias	AMH	c.35T>G p.(Val12Gly)	Normal	AMH & CBX2 variants found in addition to MAMLD1 class 4 mutation (table 3)
			AMH	c.-74C>G	Normal	
			CBX2	c.565G>A p.(Ala189Thr)	Normal	
19	XY F	?46,XY DSD	MAMLD1	c.728G>A p.(Cys243Tyr)	Normal	MAMLD1 variant found in addition to HSD17B3 mutations (table 3)
22	XY M	Ambiguous genitalia	ATRX	c.546A>G p.(=)	Normal	ATRX variant found in addition to WT1 mutation (table 3)

**Table 4b: Patients with variants of Uncertain Significance where a diagnosis has been confirmed.**

#### **Table 4: Details of Variants of Uncertain Significance (VOUS) identified.**

Details of all VOUS found in this study. Table 4a indicates patients where only VOUS were found i.e. no pathogenic or likely pathogenic variants were identified. As such a diagnosis could not be confirmed in these patients. Table 4b indicates patients where VOUS were found in addition to the pathogenic/likely pathogenic mutations which were believed to be causative of the patients phenotype. Details of the karyotypic and phenotypic sex are in the second column with M and F representing phenotypic sex. P indicates presumed karyotype (reports not seen) from SRY sequence reads. Allele 1 and 2 describe the variants in the different alleles, a black box indicates an absent second X allele in XY individuals. Green = Normal (wildtype), Yellow = Class 3 variant (Variant Of Uncertain Significance (VOUS)), Red = Class 5 variant (Clearly pathogenic).

**Table 5: Summary of the frequency of mutations and variants of uncertain significance found for each gene.**

The total number of pathogenic mutations (class 4 and 5 variants) and Variants of Uncertain Significance (VOUS, class 3) variants identified in the patients tested. Each mutated allele is given a score of 1 therefore a patient homozygous for a pathogenic mutation (score=2) would be equally represented in the table as a patient who is compound heterozygous for 2 pathogenic mutations.

Gene	Pathogenic (Class 4 or 5)	VOUS Class (3)	Gene	Pathogenic (Class 4 or 5)	VOUS
AMH	3	3	HSD3B2	2	3
AMHR2	3	1	LHCGR	1	1
AR	7	1	MAMLD1	1	1
ARX	0	0	NR0B1	0	0
ATRX	0	0	NR3C1	0	0
CBX2	0	6	NR5A1	2	4
CYB5A	0	0	POR	0	1
CYP11A1	0	3	RSPO1	0	1
CYP11B1	0	1	SOX9	0	0
CYP17A1	0	0	SRD5A2	7	1
CYP19A1	0	0	SRY	0	0
DHCR7	1	0	STAR	0	0
DHH	0	0	TSPYL1	0	0
DMRT1	0	0	WNT4	0	0
HSD17B3	9	6	WT1	1	2

Paper	Number of genes in panel	Number of patients tested and karyotypic sex	Diagnostic Yield	Diagnostic Yield	Diagnostic Yield
			Overall	46,XY	46,XX
<b>Baxter, 2015 (ref 9)</b>	64	40 XY	-	14/40	-
<b>Dong, 2016 (ref 10)</b>	219	13 XY, 8 XX	8/21*	6/13	2/8
<b>Eggers, 2016 (ref 11)</b>	64	278 XY, 48 XX	126/326	118/278	8/48
<b>Kim, 2017 (ref 12)</b>	67	37 XY, 7 XX	13/44	13/37	0/7
<b>Hughes, 2018</b>	30	73 XY, 7 XX	25/80	25/73	0/7

**Table 6: Summary of previous studies using NGS analysis for DSDs**

\*Dong et al report their detection rate of 46,XY DSD as 9/13 however 3 of these patients had Variants of Uncertain Significance (VOUS) only and therefore using the same parameters as in this study a diagnosis would not be confirmed. Numbers modified above to allow more accurate comparison with this study.