



This is a repository copy of *Low-value clinical practices in injury care: a scoping review and expert consultation survey*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/145013/>

Version: Accepted Version

---

**Article:**

Moore, L., Lauzier, F., Tardif, P.-A. et al. (18 more authors) (2019) Low-value clinical practices in injury care: a scoping review and expert consultation survey. *Journal of Trauma and Acute Care Surgery*, 86 (6). pp. 983-993. ISSN 2163-0755

<https://doi.org/10.1097/TA.0000000000002246>

---

© 2019 Wolters Kluwer Health, Inc. This is an author-produced version of a paper subsequently published in *Journal of Trauma and Acute Care Surgery*. Uploaded in accordance with the publisher's self-archiving policy. Moore, L. et al, Low-value clinical practices in injury care: a scoping review and expert consultation survey, *Journal of Trauma and Acute Care Surgery*, 2019, 86 (6), 983-993.  
<https://doi.org/10.1097/TA.0000000000002246>

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## LOW-VALUE CLINICAL PRACTICES IN INJURY CARE: A SCOPING REVIEW AND EXPERT CONSULTATION SURVEY

Lynne Moore PhD<sup>1,2</sup>; François Lauzier MD MSc<sup>2,3</sup>; Pier-Alexandre Tardif MA MSc<sup>2</sup>; Khadidja Malloum Boukar MSc<sup>1,2</sup>; Imen Farhat MSc<sup>1,2</sup>; Patrick Archambault MD MSc<sup>4</sup>; Éric Mercier MD MSc<sup>2</sup>; François Lamontagne MD MSc<sup>5</sup>; Michael Chassé MD PhD<sup>6</sup>; Henry T Stelfox MD PhD<sup>7</sup>; Simon Berthelot MD MSc<sup>2</sup>; Belinda Gabbe PhD<sup>8</sup>; Fiona Lecky MD MSc<sup>10</sup>; Natalie Yanchar MD MSc<sup>11</sup>; Howard Champion MD FCRS<sup>12</sup>; John Kortbeek MD<sup>13</sup>; Peter Cameron MD<sup>14</sup>; Paule Lessard Bonaventure MD MSc<sup>2,15</sup>; Jérôme Paquet MD<sup>15</sup>; Catherine Truchon PhD<sup>16</sup>; Alexis F Turgeon MD MSc<sup>2,3</sup>; on behalf of  
the Canadian Traumatic brain injury Research Consortium

<sup>1</sup>Department of Social and Preventative Medicine, Université Laval, Québec (Qc), Canada;

<sup>2</sup>Population Health and Optimal Health Practices Research Unit, Trauma – Emergency – Critical Care Medicine, Centre de Recherche du CHU de Québec – Université Laval (Hôpital de l'Enfant-Jésus),  
Université Laval, Québec (Qc), Canada

<sup>3</sup>Department of Anesthesiology and Critical Care Medicine, Division of Critical Care Medicine,  
Université Laval, Québec City, Québec, Canada

<sup>4</sup>Population Health and Optimal Health Practices Research Unit, Transfert des connaissances et évaluation des technologies et modes d'intervention en santé, Centre de Recherche du CHU de Québec –  
Université Laval (Hôpital St François d'Assise), Université Laval, Québec (Qc), Canada

<sup>5</sup>Department of Medicine, Université de Sherbrooke, 3001, 12th Avenue North, Sherbrooke, Québec,  
Canada

<sup>6</sup>Department of Medicine, Université de Montréal, Montréal, Québec, Canada

<sup>7</sup>Departments of Critical Care Medicine, Medicine and Community Health Sciences, O'Brien Institute  
for Public Health, University of Calgary

<sup>8</sup>School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia

<sup>9</sup>Farr Institute, Swansea University Medical School, Swansea University, UK

<sup>10</sup>Emergency Medicine, University of Sheffield, United Kingdom; Trauma Audit and Research Network,  
United Kingdom

<sup>11</sup>Department of Surgery, Dalhousie University, Halifax, Nova Scotia

<sup>12</sup>Department of Surgery, Uniformed Services University of the Health Sciences, Bethesda, Maryland,  
USA

<sup>13</sup>Department of Surgery, University of Calgary, Calgary, Alberta, Canada

<sup>14</sup>The Alfred Hospital, Monash University, Melbourne, Australia

<sup>15</sup>Department of Surgery, Division of Neurosurgery, Université Laval, Québec (Qc), Canada

<sup>16</sup>Institut national d'excellence en santé et en services sociaux, Québec (Qc), Canada

**Corresponding author and address for reprints:**

Lynne Moore

CHU de Québec Research Center (Enfant-Jésus Hospital)

Axe Santé des Populations et Pratiques Optimales en Santé (Population Health and Optimal Health  
Practices Research Unit), Traumatologie – Urgence - Soins intensifs (Trauma – Emergency – Critical  
Care Medicine)

1401, 18e rue, local H-012a, Québec (Québec), G1J 1Z4

Tel. 418-649-0252 #3366

Fax: 418-649-5733

Email: lynne.moore @fmed.ulaval.ca

**Conflict of interest and source of funding:** This research is funded by the Canadian Institutes of Health Research (Foundation grant, #353374) and the Fonds de Recherche du Québec – Santé (career award, LM, FLau, FLam, MC). Patrick Archambault is supported by a Clinical-Embedded Scientist Award from the CIHR. Dr Turgeon is the Canada Research Chair in Critical Care Neurology and Trauma. For the remaining authors, no conflicts were declared.

**Meetings:**

The results of this study were presented at the 77<sup>th</sup> Annual Meeting of AAST and Clinical Congress of Acute Care Surgery and 4th World Trauma Congress, September 26<sup>th</sup>-29<sup>th</sup>, 2018 in San Diego, California

## **ABSTRACT**

**Background:** Tests and treatments that are not supported by evidence and could expose patients to unnecessary harm, referred to here as low-value clinical practices, consume up to 30% of healthcare resources. Choosing Wisely and other organisations have published lists of clinical practices to be avoided. However, few apply to injury and most are based uniquely on expert consensus. We aimed to identify low-value clinical practices in acute injury care.

**Methods:** We conducted a scoping review targeting articles, reviews and guidelines that identified low-value clinical practices specific to injury populations. Thirty-six experts rated clinical practices on a 5-point Likert scale from clearly low-value to clearly beneficial. Clinical practices reported as low-value by at least one level I, II or III study and considered clearly or potentially low-value by at least 75% of experts were retained as candidates for low-value injury care.

**Results:** Of 50,695 citations, 815 studies were included and led to the identification of 150 clinical practices. Of these 63 were considered candidates for low-value injury care; 33 in the emergency room, 9 in trauma surgery, 15 in the intensive care unit and 5 in orthopaedics. We also identified 87 ‘grey zone’ practices, which did not meet our criteria for low-value care.

**Conclusions:** We identified 63 low-value clinical practices in acute injury care that are supported by empirical evidence and expert opinion. Conditional on future research, they represent potential targets for guidelines, overuse metrics and de-implementation interventions. We also identified 87 ‘grey zone’ practices, which may be interesting targets for value-based decision-making. Our study represents an important step towards the de-implementation of low-value clinical practices in injury care.

**Level of evidence:** III

**Keywords:** Low-value care, trauma systems, scoping review, expert survey

## BACKGROUND

Injuries led to 192,000 deaths, 3 million hospitalizations and 27 million emergency department visits in the USA in 2013 and generated medical and work loss costs of \$671 billion USD.<sup>(1)</sup> In Canada, injury deaths increased by 23% from 13,000 in 2004 to 16,000 in 2010 while costs increased by 35% and are projected to reach \$75 billion CAN by 2035.<sup>(2)</sup> Given the huge burden of injury and evidence of unwarranted variation in injury outcomes across healthcare providers,<sup>(3-5)</sup> efforts to optimize care has the potential to yield major dividends.

Rapid innovation in imaging and therapeutic techniques has led to an exponential rise in the use of tests and treatments that are not supported by evidence and could expose patients to unnecessary harm,<sup>(6, 7)</sup> referred to here as low-value clinical practices.<sup>(8-15)</sup> Low-value clinical practices have been estimated to consume up to 30% of healthcare resources<sup>(10, 12, 14, 16)</sup> but little is known about this issue in the context of injury care. Low-value clinical practices have multiple negative consequences. From a healthcare system perspective, they strain healthcare budgets and decrease the availability of resources. From a patient and caregiver perspective, they expose patients to physical and psychological harm, delay effective treatment, and increase direct and indirect expenses.<sup>(8-10, 12, 14)</sup> Finally, from a societal perspective, low-value clinical practices threaten the sustainability of affordable, accessible healthcare. Interventions targeting the de-implementation of low-value clinical practices therefore have the potential to reduce waste and improve patient outcomes.<sup>(15, 17)</sup>

Physicians report overusing resources for fear of legal actions but also because of lack of guidelines on low-value clinical practices.<sup>(12-14, 18)</sup> Choosing Wisely has developed lists of commonly used tests or procedures whose necessity should be questioned including top five lists for emergency medicine,

radiology, pediatric orthopaedics, neurology, and surgery.<sup>(11)</sup> However, few apply to injury care and most are based solely on expert consensus. Previous systematic reviews aiming to identify low-value clinical practices have not been specific to injury but have underlined the importance of targeting diagnostic groups to improve feasibility and subsequent knowledge transfer.<sup>(15, 19-22)</sup> We aimed to identify low-value clinical practices in acute, intrahospital injury care.

## **METHODS**

Our study was conducted in 6 stages following published guidelines for scoping reviews and comprised a literature review followed by a web-based survey consultation with clinical experts.<sup>(23, 24)</sup> The protocol has been published previously.<sup>(25)</sup> Ethics approval was obtained from the institutional research ethics committee.

### **1. Identify research questions and develop definitions**

First, using an iterative approach, the interdisciplinary and intersectorial project steering committee comprising clinicians, allied health professionals and policy and decision-makers identified the following research question for our review: Which clinical practices are considered low-value in acute injury care? Second, the committee used highly-cited literature on healthcare overuse<sup>(7, 13, 14, 17)</sup> to establish the following working definition of low-value clinical practices: A test or treatment (i.e. admission, monitoring, diagnostic interventions, therapeutic interventions, consultation) that is used in practice but is ineffective or its harm/cost outweighs its benefits. Third, the committee consulted UCLA/RAND recommendations to establish the following criteria for identifying candidates for low-value injury care: clinical practices identified as low-value in at least one level I, II or III study AND

considered to be clearly/potentially low-value by at least 75% of experts and not considered clearly beneficial by any expert.

## **2. Identify relevant studies**

### Eligibility criteria

We included original research, literature reviews, recommendations and guidelines that identified at least one low-value clinical practice specific to injury populations according to the definition given above.<sup>(11)</sup> We included studies on clinical practices specific to intrahospital acute care (in the emergency department or following hospital admission). We excluded: i) studies with no clear indication for the low-value practice (e.g. based on physician gestalt), ii) studies based exclusively on populations with combat injuries, osteoporotic fractures, burns, bites, or foreign bodies, iii) case reports, animal and cadaver studies, iv) studies on pre-hospital or post-acute clinical practices.

### Information sources

We systematically searched MEDLINE, EMBASE, Cochrane CENTRAL, BIOSIS/Web of Science, ClinicalTrials and ISRCTN; Thesis repositories (Thesis portal Canada, EtHOS, DART-Europe E-Theses Portal, the National Library of Australia's Trove and ProQuest Dissertations & Theses Global); Websites of healthcare quality organizations (Agency for Healthcare Research and Quality, Australasian Association for Quality in Healthcare, Canadian Institutes for Health Information, Choosing Wisely, Lown Institute, National Association for Healthcare Quality, National Institute of Health and Care Excellence, National Quality Forum, and World Health Organization) and injury organisations (American Association for the Surgery of Trauma, American Association of Orthopaedic Surgeons, American College of Surgeons, American Trauma Society, Australasian Trauma Society, Brain Trauma



Foundation, British Trauma Society, Eastern Association for the Surgery of Trauma, International Association for Trauma Surgery and Intensive Care, International Trauma Anesthesia and Critical Care Society, Orthopaedic Trauma Association, The Society of Trauma Nurses, Trauma Association of Canada, Trauma Audit Research Network, Trauma.org, and Western Trauma Association.); and patient advocacy organizations including Safer Healthcare Now!

#### Search strategy

We developed a systematic search strategy with an information specialist.<sup>(26)</sup> The strategy was developed for MEDLINE and EMBASE using keywords covering combinations of search terms under the themes injury and low-value clinical practices (Supplemental Digital Content 3, Table 1, <http://links.lww.com/TA/B326>). This search strategy was then adapted for the other databases.

### **3. Select studies**

#### Data management

Citations were managed using EndNote software (version X7.0.1, New York City: Thomson Reuters, 2011). Duplicates were identified and eliminated using electronic and manual screening. Multiple publications based on the same dataset were identified by crosschecking authors, dates and settings. In the case of replication, we identified only one publication for analyses using criteria based on study dates (most recent) and sample size (largest).

#### Selection process

Pairs of reviewers with methodological and content expertise (two of four reviewers LM, KMB, PAT, IF) independently evaluated all citations for eligibility. Consecutive samples of 500 citations were

independently assessed by each reviewer until high agreement was achieved on study inclusion (3 samples for kappa>0.8). Any further disagreement on study eligibility was resolved by consensus and a fifth reviewer adjudicated when necessary (FL).

#### **4. Chart material**

A standard electronic data abstraction form and a detailed instruction manual were developed and piloted independently by all reviewers on a representative sample of five publications. Pairs of reviewers (LM, KMB, PAT, IF) independently extracted information on the study design, setting (country, year, language, funding), study objective, study population, low-value clinical practices, and primary outcomes when appropriate. Any discrepancies between reviewers was resolved by consensus and a fifth reviewer adjudicated when necessary (FL).

#### **5. Collate, summarize, and report on results**

Clinical practices were classified according to the type of practice and the clinical speciality.<sup>(19)</sup> Classifications were conducted independently by two reviewers (KMB, PAT) and then checked independently by a third reviewer (LM). Any disagreements were adjudicated by a fourth reviewer (FL). As is common in scoping reviews, the methodological quality of included studies was not evaluated.<sup>(27)</sup> We summarized the level of evidence for each practice by calculating the number of studies by type using an adaptation of Oxford Center for Evidence-based Medicine classifications:<sup>(28)</sup> randomized controlled trials (RCTs) or systematic review of RCTs (I), prospective cohort studies or systematic review of RCTs and prospective cohort studies (II), retrospective cohort, case-control, cross sectional and case series studies or systematic review of any of the former (III), expert consensus and other (IV).

## 6. Consultation

We recruited four groups of experts for the consultation phase using a snowball technique based on the following criteria: representation of clinical expertise involved in acute intrahospital injury care, actively involved in injury research (knowledge of the evidence base for clinical practices) and geographical diversity.<sup>(29)</sup> Recruitment was independent of scoping review results and authorship status to minimize the influence of intellectual or academic biases. Groups were formed according to clinical specialty: emergency physicians, critical care physicians/neurosurgeons, trauma surgeons and orthopaedic/spine surgeons. Each group reviewed clinical practices within their area of expertise. For the main objective, we used two phases of consultation. First, we consulted a subgroup of 8 experts (two from each specialty) to regroup overlapping clinical practices, harmonize terminology and develop and test our survey. Second, we administered a web-based survey<sup>(30)</sup> asking experts to rate each clinical practice on a 5-point Likert scale from clearly low-value to clearly beneficial (see Supplemental Digital Content 1, Figure 2, <http://links.lww.com/TA/B324>). These categories mirror the ‘clearly ineffective, grey zone, and clearly effective’ classifications described in the Lancet Right Care series.<sup>(14, 31)</sup>

After the consultation phase, we applied the a priori criteria described above to identify candidate low-value clinical practices for injury care, i.e. practices reported as low-value in at least one level I, II or III study AND considered to be clearly/potentially low-value by at least 75% of experts and not considered clearly beneficial by any expert.

## RESULTS

Of 77,733 citations, 1,593 studies were retained for full text review and 815 were included (Supplemental Digital Content 2, Figure 1, <http://links.lww.com/TA/B325>). Data extraction led to the

identification of 965 clinical practices (Table 1). Over one half were prospective or retrospective cohort studies, 22% were reviews (one third of these systematic), 5% were based on expert opinion and less than 5% were RCTs. The majority of studies aimed to evaluate the effectiveness of the clinical practice (55%) whereas one quarter aimed to develop guidelines or derive/validate a clinical decision rule. Seventeen percent aimed to evaluate the prevalence of overuse or the efficacy of a de-implementation intervention. Less than 1% aimed to derive or validate quality indicators. More than one third of low-value practices pertained to the treatment of head injury and most were specific to adult (37%) or pediatric (12%) populations. One half of clinical practices targeted diagnostic interventions, 40% targeted therapeutic interventions and 5% targeted ICU or hospital admission.

We approached 39 experts of whom 36 (92%) agreed to participate and completed the survey including 8/9 emergency physicians, 9/9 critical care physicians, 1/1 neurosurgeon, 10/12 trauma surgeons and 8/8 orthopaedic/spine surgeons from Canada, US, Australia and the UK. After the first consultation phase, we identified 150 clinical practices (Tables 2-5 and Supplemental Digital Content 4, Table 2, <http://links.lww.com/TA/B327>). In the web-based survey, 66 clinical practices were considered clearly or potentially low-value by at least 75% of respondents. Thereafter, we identified 63 clinical practices that met our criteria as candidates for low-value injury care, i.e. they were reported as low-value in at least one level I, II or III study, considered clearly or potentially low-value by at least 75% of respondents and not considered clearly beneficial by any of the experts (Tables 2-5). Among these clinical practices, 13 were supported by do-not-do recommendations in internationally recognized clinical practice guidelines (i.e. indications were the same or very similar). Nine practices included as do-not-do recommendations in clinical guidelines were not selected by our criteria (Supplemental Digital Content 4, Table 2, <http://links.lww.com/TA/B327>).

We identified 33 candidates for low-value injury care in the emergency room of which five were related to hospital admission for abdominal trauma or mild TBI and 20 were related to imaging including CT or X-ray for mild TBI, ankle, knee, chest and cervical spine injuries (Table 2). We also identified 15 ED practices in the grey zone including repeat head CT in adult mild complicated TBI and hospital admission in pediatric isolated skull fracture (Supplemental Digital Content 4, Table 2, <http://links.lww.com/TA/B327>). Nine low-value practices were selected for general trauma surgery, 6 of which were related to operative management of liver, renal, splenic, and neck injuries (Table 3). In addition, we identified 15 practices in the grey zone including follow-up imaging for nonoperative blunt renal injury and surgical management of high-grade pancreatic or renal injuries (Supplemental Digital Content 4, Table 2, <http://links.lww.com/TA/B327>). We identified 15 low-value practices in the intensive care unit of which 8 targeted TBI (Table 4). Four were related to medications (corticosteroids, antibiotics and antiseizure prophylaxis) and four were related to fluids and blood products (albumin, colloids, platelet and red blood cell transfusion). Twenty-six (63%) of ICU clinical practices were in the grey zone (Supplemental Digital Content 4, Table 2, <http://links.lww.com/TA/B327>) including neurosurgical consultation in acute mild complicated TBI, decompressive craniotomy and hourly neurological assessments >24h for stable TBI. Five low-value practices were identified in orthopaedics targeting follow-up consultation, spine service consultation, repeat X-ray, orthosis for thoracolumbar burst fractures and pre-operative blood tests (Table 5). Thirty-one (86%) orthopaedic practices in acute injury care were classed in the grey zone of which 6 targeted follow-up consultation, 9 imaging and 5 immobilization (Supplemental Digital Content 4, Table 2, <http://links.lww.com/TA/B327>).

## DISCUSSION

We identified 63 clinical practices that met criteria for low-value intrahospital injury care. These potential low-value practices are supported by empirical evidence and expert opinion. Conditional on the results of future research, they represent potential targets for guidelines, overuse metrics and de-implementation interventions. We also identified 87 clinical practices in the grey zone, which are not consistently supported by empirical studies and expert opinion. While these practices require more evidence before being labelled low-value, they may be interesting targets for value-based decision-making.

The literature on low-value clinical practices in injury care is scarce. Internationally recognized medical associations publish guidelines on injury care.<sup>(32-35)</sup> However, few pertain to clinical practices that should be avoided. Healthcare quality organisations including Choosing Wisely and the National Institute for Health and Care Excellence publish recommendations specific to low-value practices but few target injury care.<sup>(36, 37)</sup> In addition, these recommendations are often based only on expert consensus.<sup>(20)</sup> Three previous literature reviews on low-value care across a range of diagnostic groups identified 9 low-value practices specific to injury care.<sup>(14, 19, 20, 38)</sup> We were able to identify many more practices because targeting a specific diagnostic group allows for a much more sensitive review strategy.<sup>(31)</sup> With over 50,000 citations to screen and more than 1400 documents to extract in our study, a similar search strategy with no restrictions on diagnosis would have been unfeasible.

Twenty-six percent of low-value practices identified in our review were related to imaging. This is consistent with a previous review of low-value care measures<sup>(20)</sup> and may be because the value of imaging is relatively easy to evaluate retrospectively. Unnecessary imaging generates important costs<sup>(14,</sup>

<sup>39)</sup> and may expose patients to high doses of radiation with non-negligible long term risks of cancer.<sup>(40-42)</sup> We retained 12 low-value practices on imaging which are already supported by guidelines and/or widely used clinical decision rules and 8 additional clinical practices which are potential targets for low-value imaging. We identified 21 low-value practices related to operative (versus non operative) management of which two are included in EAST guidelines.<sup>(32)</sup> A recent review found 71 low-value practices in general surgery representing an estimated annual cost of 153 million euros per year in the UK.<sup>(43)</sup> However, none of these practices pertained to injury. Seventeen practices identified in our review pertained to medications of which five were supported by do-not-do recommendations in clinical guidelines.<sup>(32, 34, 36, 37)</sup> There is a large body of literature on overprescribing in primary care.<sup>(14, 44-46)</sup> However, an important knowledge gap on in-hospital medication exists, probably in part due to the fact that hospital prescriptions are not recorded in administrative databases. Other low-value practices identified in our review were hospital and ICU admission (n=11) and follow-up consultation (n=7). Literature on overuse in these areas is sparse, possibly because they are very context-specific. Nine practices included in internationally-recognised guidelines as practices to avoid were not retained in our study, all because less than 75% of experts identified them as clearly or potentially low-value. This discordance could be due to our strict selection criteria based on literature evidence and agreement of more than 75% of experts. Guidelines are often based on few, low-quality studies or expert consensus, but rarely both.<sup>(47)</sup> It may also be explained by differing influences of local context, industry pressure or single highly-mediatized studies.<sup>(13, 15, 21, 48, 49)</sup> It does suggest that moving forward, guidelines/metrics on low-value injury care should be based both on evidence from high-quality experimental or observational studies AND expert opinion and should account for the possible influence of local context. Also, the consensus process should strive to minimize intellectual, academic and financial biases.

## **Strengths and limitations**

This study represents a rigorous, exhaustive review of the literature on low-value clinical practices in injury care. Results from our scoping review are supported by a consultation study with 36 experts representing the clinical specialties involved in trauma care on three continents. The participation rate of over 90% demonstrates the high level of knowledge-user interest in this topic. In addition, experts are all involved in clinical research in acute injury care so are likely to have good knowledge of the evidence-base on clinical practices for injury admissions.

This study does have limitations that should be considered in the interpretation of results. First, for feasibility reasons, our search strategy was based on key words related to low-value care and was therefore dependent on authors' judgement of the value of clinical practices. This may have led us to miss some low-value practices. For example, authors of the Randomised Evaluation of Surgery with Craniectomy of Uncontrollable Elevation of Intracranial Pressure (RESCUEicp) trial that observed lower mortality but worse functional outcomes in the intervention group did not clearly identify decompressive craniectomy as a low-value practice.<sup>(50)</sup> However, by thoroughly screening article references, grey literature including injury organisations and healthcare quality websites, and consulting experts for further references, we are confident that we captured a large proportion of potentially low-value clinical practices that have been reported in the literature. Second, for feasibility reasons, we restricted the review to studies published since 2006. We may therefore have missed some important RCTs published earlier, for example the National Acute Spinal Cord Injury Studies I on high-dose steroids for spinal cord injury<sup>(51)</sup> and the Harborview trial on antiseizure prophylaxis in traumatic brain injury.<sup>(52)</sup> However, both these practices were captured through review of guidelines. Fourth, due to the scoping design of our review, we did not evaluate methodological quality. Strength of evidence was only



based on study design. Fifth, the last phase of the review was based on a single web survey therefore represents the results of a consultation rather than expert consensus. In addition, we used a convenience sample and only one neurosurgeon was surveyed. Finally, to identify targets for de-implementation we will need data on frequency (how frequently is the clinical practice actually used?), inter-provider variations (is there evidence of practice variation?) and economic impact (would de-adoption lead to important savings?).<sup>(53, 54)</sup> These aspects will be incorporated into the following subsequent phases of the Canadian Program for Monitoring Overuse in Injury Care; a systematic review to GRADE evidence for low-value clinical practices identified in this review,<sup>(55)</sup> a RAND-UCLA expert consensus study to develop a set of quality indicators targeting low-value practices, a multicenter retrospective cohort study to derive and validate metrics for the quality indicators and a cluster randomized controlled trial to evaluate the effectiveness of quality indicators in an audit-feedback intervention. The research program will also allow us to take into account the specificities of low-frequency, high-risk injuries.

## **CONCLUSIONS**

This study fills a major knowledge gap on medical procedure overuse in acute injury care. Results will inform research priorities and the development of metrics to measure overuse. This knowledge will provide a solid basis for the development of interventions targeting de-implementation, such as clinical decision rules and shared decision-making tools. This has the potential to decrease costs, increase resource availability, reduce mortality and morbidity due to unnecessary tests and treatments and reduce patient stress and physicians' workload.

**Conflict of interest and source of funding:** This research is funded by the Canadian Institutes of Health Research (Foundation grant, #353374) and the Fonds de Recherche du Québec – Santé (career award, LM, FLau, FLam, MC). Patrick Archambault is supported by a Clinical-Embedded Scientist Award from the CIHR. Dr Turgeon is the Canada Research Chair in Critical Care Neurology and Trauma. For the remaining authors, no conflicts were declared.

### **Authorship statement**

Lynne Moore led the conception and design of the study, acquisition of data, analysis and interpretation of data, and drafted the article.

François Lauzier made substantial contributions to the conception and design, the acquisition of the data, the analysis and the interpretation of data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Pier-Alexandre Tardif made substantial contributions to the acquisition of the data, the analysis and the interpretation of data. He participated in drafting the article and gave final approval of the version to be published

Khadidja Malloum Boukar made substantial contributions to the acquisition of the data. She revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Imen Farhat made substantial contributions to the acquisition of the data. She revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Patrick Archambault made substantial contributions to the conception and design, the acquisition of the data, the analysis and the interpretation of data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Éric Mercier made substantial contributions to the conception and design and the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

François Lamontagne made substantial contributions to the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Michael Chassé made substantial contributions to the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Henry T Stelfox made substantial contributions to the conception and design and the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Simon Berthelot made substantial contributions to the conception and design and the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Belinda Gabbe made substantial contributions to conception and design. She revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Fiona Lecky made substantial contributions to the acquisition of the data. She revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Natalie Yanchar made substantial contributions to the the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Howard Champion made substantial contributions to the the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

John Kortbeek made substantial contributions to the the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Peter Cameron made substantial contributions to the the acquisition of the data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Paule Lessard Bonaventure made substantial contributions to the acquisition of the data and the analysis and the interpretation of data. Shee revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Jérôme Paquet made substantial contributions to the acquisition of the data and the analysis and the interpretation of data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Catherine Truchon made substantial contributions to the acquisition of the data. She revised the manuscript critically for important intellectual content and gave final approval of the version to be published

Alexis F Turgeon made substantial contributions to the conception and design, the acquisition of the data, the analysis and the interpretation of data. He revised the manuscript critically for important intellectual content and gave final approval of the version to be published

ACCEPTED

## REFERENCES

1. Centers for Disease Control and Prevention. Injury Prevention & Control 2018. Available at: <https://www.cdc.gov/injury/index.html>. Accessed Mai 14, 2018.
2. Egi M, Bellomo R, Stachowski E, French CJ, Hart G, Stow P, Li WQ, Bates S. Intensive insulin therapy in postoperative intensive care unit patients - A decision analysis. *Am J Respir Crit Care Med*. 2006;173(4):407-13.
3. Birkmeyer JD, Reames BN, McCulloch P, Carr AJ, Campbell WB, Wennberg JE. Understanding of regional variation in the use of surgery. *Lancet*. 2013;382(9898):1121-9.
4. Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, Pinder EL. The implications of regional variations in Medicare spending. Part 2: health outcomes and satisfaction with care. *Ann Intern Med*. 2003;138(4):288-98.
5. Moore L, Evans D, Hameed SM, Yanchar NL, Stelfox HT, Simons RK, Kortbeek J, Bourgeois G, Clément J, Lauzier F, et al. Mortality in Canadian trauma systems: a multicenter cohort study. *Ann Surg*. 2016;Epub ahead of print.
6. Barker P, Buchanan-Barker P. More harm than good. *Nursing standard (Royal College of Nursing (Great Britain) : 1987)*. 2006;20(30):28-9.
7. Institute of Medicine Committee on Quality of Health Care in A. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington (DC): National Academies Press (US); 2001.
8. Berwick DM, Hackbarth AD. Eliminating waste in US health care. *JAMA*. 2012;307(14):1513-6.
9. Boat TF, Chao SM, O'Neill PH. From waste to value in health care. *JAMA*. 2008;299(5):568-71.
10. Reilly BM, Evans AT. Much ado about (doing) nothing. *Ann Intern Med*. 2009;150(4):270-1.
11. Morgan DJ, Dhruva SS, Wright SM, Korenstein D. 2016 Update on Medical Overuse: A Systematic Review. *JAMA Intern Med*. 2016;176(11):1687-92.

12. Hauser CJ, Adams CA, Jr., Eachempati SR, Council of the Surgical Infection S. Surgical Infection Society guideline: prophylactic antibiotic use in open fractures: an evidence-based guideline. *Surg Infect (Larchmt)*. 2006;7(4):379-405.
13. Berwick DM. Avoiding overuse-the next quality frontier. *Lancet*. 2017.
14. Brownlee S, Chalkidou K, Doust J, Elshaug AG, Glasziou P, Heath I, Nagpal S, Saini V, Srivastava D, Chalmers K, et al. Evidence for overuse of medical services around the world. *Lancet*. 2017.
15. Saini V, Brownlee S, Elshaug AG, Glasziou P, Heath I. Addressing overuse and underuse around the world. *Lancet*. 2017.
16. Bennett MH, Trytko BE, Jonker B. A systematic review of the use of hyperbaric oxygen therapy in the treatment of acute traumatic brain injury. *Diving Hyperb Med*. 2006;36(1):39-46.
17. Niven DJ, Mrklas KJ, Holodinsky JK, Straus SE, Hemmelgarn BR, Jeffs LP, Stelfox HT. Towards understanding the de-adoption of low-value clinical practices: a scoping review. *BMC Med*. 2015;13:255.
18. Emanuel EJ, Fuchs VR. The perfect storm of overutilization. *JAMA*. 2008;299(23):2789-91.
19. Chan KS, Chang E, Nassery N, Chang HY, Segal JB. The state of overuse measurement: a critical review. *Medical care research and review : MCRR*. 2013;70(5):473-96.
20. de Vries EF, Struijs JN, Heijink R, Hendrikx RJ, Baan CA. Are low-value care measures up to the task? A systematic review of the literature. *BMC Health Serv Res*. 2016;16(1):405.
21. Elshaug AG, Rosenthal MB, Lavis JN, Brownlee S, Schmidt H, Nagpal S, Littlejohns P, Srivastava D, Tunis S, Saini V. Levers for addressing medical underuse and overuse: achieving high-value health care. *Lancet*. 2017.

22. Segal JB, Bridges JF, Chang HY, Chang E, Nassery N, Weiner J, Chan KS. Identifying possible indicators of systematic overuse of health care procedures with claims data. *Med Care*. 2014;52(2):157-63.
23. Arksey H, O'Malley L. Scoping studies: Towards a Methodological Framework. *Int J Soc Res Methodol*. 2005;8:19-32.
24. Tricco AC, Lillie E, Zarin W, O'Brien K, Colquhoun H, Kastner M, Levac D, Ng C, Sharpe JP, Wilson K, et al. A scoping review on the conduct and reporting of scoping reviews. *BMC Med Res Methodol*. 2016;16:15.
25. Moore L, Boukar KM, Tardif PA, Stelfox HT, Champion H, Cameron P, Gabbe B, Yanchar N, Kortbeek J, Lauzier F, et al. Low-value clinical practices in injury care: a scoping review protocol. *BMJ open*. 2017;7(7):e016024.
26. Dretzke J, Burls A, Bayliss S, Sandercock J. The clinical effectiveness of pre-hospital intravenous fluid replacement in trauma patients without head injury: A systematic review. *Trauma*. 2006;8(3):131-41.
27. Guttman A, Razzaq A, Lindsay P, Zagorski B, Anderson GM. Development of measures of the quality of emergency department care for children using a structured panel process. *Pediatrics*. 2006;118(1):114-23.
28. OCEBM Levels of Evidence Working Group. The Oxford 2011 Levels of Evidence: Oxford Centre for Evidence-Based Medicine; 2011. Available at: <http://www.cebm.net/index.aspx?o=5653>. Accessed January 25, 2019.
29. Morgan D. Snowball Sampling. In: Given L, editor. *The SAGE Encyclopedia of Qualitative Research Methods*: SAGE Publications Inc., Thousand Oaks; 2008. p. 816-7.



30. Limesurvey GmbH. LimeSurvey: An Open Source survey tool LimeSurvey GmbH, Hamburg, Germany. Available at: <http://www.limesurvey.org>. Accessed January 25, 2019.
31. Saini V, Brownlee S, Elshaug AG, Glasziou P, Heath I. Addressing overuse and underuse around the world. *Lancet*. 2017;390(10090):105-7.
32. Eastern Association for the Surgery of Trauma. EAST Practice Management Guidelines 2018. Available at: <https://www.east.org/education/practice-management-guidelines>. Accessed Mai 3, 2018.
33. Carney N, Ghajar J, Jagoda A, Bedrick S, Davis-O'Reilly C, du Coudray H, Hack D, Helfand N, Huddleston A, Nettleton T, et al. Concussion guidelines step 1: systematic review of prevalent indicators. *Neurosurgery*. 2014;75 Suppl 1:S3-15.
34. Carney N, Totten AM, O'Reilly C, Ullman JS, Hawryluk GW, Bell MJ, Bratton SL, Chesnut R, Harris OA, Kissoon N, et al. Guidelines for the Management of Severe Traumatic Brain Injury, Fourth Edition. *Neurosurgery*. 2017;80(1):6-15.
35. American College of Surgeons. Resources for Optimal Care of the Injured Patient 2014/Resources Repository 2018. Available at: <https://www.facs.org/quality-programs/trauma/vrc/resources>. Accessed May 4, 2018.
36. Choosing Wisely. Clinician Lists: ABIM Foundation. Available at: <http://www.choosingwisely.org/clinician-lists/>. Accessed May 4, 2018.
37. National Institute for Health and Care Excellence (NICE). Trauma quality standard 2018. Available at: <https://www.nice.org.uk/guidance/qs166>. Accessed May 4, 2018.
38. Elshaug AG. Over 150 potentially low-value health care practices: an Australian study. *Med J Aust*. 2013;198(2):85.

39. Schoen C, Osborn R, Doty MM, Bishop M, Peugh J, Murukutla N. Toward higher-performance health systems: adults' health care experiences in seven countries, 2007. *Health Aff (Millwood)*. 2007;26(6):w717-34.
40. Brenner DJ, Hricak H. Radiation exposure from medical imaging: time to regulate? *JAMA*. 2010;304(2):208-9.
41. Hendee WR, O'Connor MK. Radiation risks of medical imaging: separating fact from fantasy. *Radiology*. 2012;264(2):312-21.
42. Royal HD. Effects of low level radiation-what's new? *Semin Nucl Med*. 2008;38(5):392-402.
43. Malik HT, Marti J, Darzi A, Mossialos E. Savings from reducing low-value general surgical interventions. *Br J Surg*. 2018;105(1):13-25.
44. Morgan SG, Hunt J, Rioux J, Proulx J, Weymann D, Tannenbaum C. Frequency and cost of potentially inappropriate prescribing for older adults: a cross-sectional study. *CMAJ open*. 2016;4(2):E346-51.
45. Makary MA, Overton HN, Wang P. Overprescribing is major contributor to opioid crisis. *BMJ (Clinical research ed)*. 2017;359:j4792.
46. Bentley TG, Effros RM, Palar K, Keeler EB. Waste in the U.S. Health care system: a conceptual framework. *Milbank Q*. 2008;86(4):629-59.
47. Chen Y, Yang K, Marusic A, Qaseem A, Meerpohl JJ, Flottorp S, Akl EA, Schunemann HJ, Chan ES, Falck-Ytter Y, et al. A Reporting Tool for Practice Guidelines in Health Care: The RIGHT Statement. *Ann Intern Med*. 2017;166(2):128-32.
48. Kleinert S, Horton R. From universal health coverage to right care for health. *Lancet*. 2017;390(10090):101-2.

49. Saini V, Garcia-Armesto S, Klemperer D, Paris V, Elshaug AG, Brownlee S, Ioannidis JPA, Fisher ES. Drivers of poor medical care. *Lancet*. 2017;390(10090):178-90.
50. Hutchinson PJ, Koliass AG, Timofeev IS, Corteen EA, Czosnyka M, Timothy J, Anderson I, Bulters DO, Belli A, Eynon CA, et al. Trial of Decompressive Craniectomy for Traumatic Intracranial Hypertension. *N Engl J Med*. 2016;375(12):1119-30.
51. Bracken MB, Shepard MJ, Hellenbrand KG, Collins WF, Leo LS, Freeman DF, Wagner FC, Flamm ES, Eisenberg HM, Goodman JH, et al. Methylprednisolone and neurological function 1 year after spinal cord injury. Results of the National Acute Spinal Cord Injury Study. *J Neurosurg*. 1985;63(5):704-13.
52. Temkin NR, Dikmen SS, Wilensky AJ, Keihm J, Chabal S, Winn HR. A randomized, double-blind study of phenytoin for the prevention of post-traumatic seizures. *N Engl J Med*. 1990;323(8):497-502.
53. Pandya A. Adding Cost-effectiveness to Define Low-Value Care. *JAMA*. 2018.
54. Ryan AM, Tompkins CP. Efficiency and Value in Healthcare: Linking Cost and Quality Measures. National Quality Forum. 2014.
55. Atkins D, Best D, Briss PA, Eccles M, Falck-Ytter Y, Flottorp S, Guyatt GH, Harbour RT, Haugh MC, Henry D, et al. Grading quality of evidence and strength of recommendations. *BMJ (Clinical research ed)*. 2004;328(7454):1490.

**Table 1.** Overview of included studies (n=815)

Country		N (%)
USA		397 (48.7)
UK		86 (10.6)
Canada		61 (7.5)
Australia		39 (4.8)
Netherlands		23 (2.8)
Turkey		19 (2.3)
Other		190 (23.3)
Year of publication		
2006-2007		105 (12.9)
2008-2009		119 (14.6)
2010-2011		121 (14.9)
2012-2013		148 (18.2)
2014-2015		161 (19.8)
2016-Mar2018		152 (18.6)
Study design		
Experimental	randomized controlled trial	38 (4.7)
	quasi-randomized controlled trial	7 (0.9)
Observational	retrospective cohort	266 (32.6)
	prospective cohort	156 (19.1)
	case series	104 (12.8)
	cross-sectional	8 (0.9)

Review	narrative review	110 (13.5)
	systematic review with meta-analysis	33 (4.1)
	systematic review without meta-analysis	35 (4.3)
Expert opinion		44 (5.4)
Other		14 (1.7)
<b>Main study objective</b>		
	Effectiveness of clinical practice	448 (55.0)
	Development/validation of a clinical decision rule	119 (14.6)
	Guidelines/recommendations	75 (9.2)
	Prevalence of overuse	74 (9.1)
	Efficacy of a deimplementation intervention	68 (8.3)
	Safety	14 (1.7)
	Development/validation of indicators	5 (0.6)
	Other	12 (1.5)
<b>Injury type*</b>		
	Head	326 (33.8)
	Thoracoabdominal	258 (26.7)
	Orthopaedic	155 (16.1)
	Spine	120 (12.4)
	All injury types	94 (9.7)
	Other	12 (1.2)
<b>Age group*</b>		
	Adult	356 (36.9)


Pediatric		113 (11.7)
Geriatric		8 (0.8)
All		281 (29.1)
Not reported		207 (21.5)
Type of clinical practice*		
Diagnostic		496 (51.4)
Therapeutic	surgical	157 (16.3)
	medical	86 (8.9)
	drugs	104 (10.8)
	device	40 (4.2)
Admission		44 (4.6)
Consultation		21 (2.2)
Monitoring		9 (0.9)
Transfer		8 (0.8)

\*Based on the number of low value clinical practices (n=965)


**Table 2.** Low value clinical practices in the **emergency department** according to level of evidence (review phase) and expert opinion (consultation phase)

Practices in the emergency department	Level of evidence† I-RCT to IV-expert consensus Number of studies	Expert opinion‡ 1-clearly low value to 5-beneficial Number of experts																						
Admission in adult blunt abdominal trauma with normal physical exam and negative FAST or CT[1-3]◇	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>10</td></tr> <tr><td>II</td><td>5</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of studies	I	10	II	5	III	0	IV	0	<table border="1"> <tr><th>Rating</th><th>Number of experts</th></tr> <tr><td>1</td><td>4</td></tr> <tr><td>2</td><td>3</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> </table>	Rating	Number of experts	1	4	2	3	3	0	4	1	5	0
Level	Number of studies																							
I	10																							
II	5																							
III	0																							
IV	0																							
Rating	Number of experts																							
1	4																							
2	3																							
3	0																							
4	1																							
5	0																							
Admission in pediatric blunt abdominal trauma with normal physical exam and asymptomatic and negative FAST or CT[4-6]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>1</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Level	Number of studies	I	0	II	1	III	0	IV	2	<table border="1"> <tr><th>Rating</th><th>Number of experts</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>5</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> </table>	Rating	Number of experts	1	1	2	5	3	0	4	1	5	0
Level	Number of studies																							
I	0																							
II	1																							
III	0																							
IV	2																							
Rating	Number of experts																							
1	1																							
2	5																							
3	0																							
4	1																							
5	0																							
Admission in stable anterior abdominal stab wound, negative on CT and negative local wound exploration[7-9]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>1</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Level	Number of studies	I	0	II	0	III	1	IV	2	<table border="1"> <tr><th>Rating</th><th>Number of experts</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> </table>	Rating	Number of experts	1	2	2	2	3	1	4	0	5	0
Level	Number of studies																							
I	0																							
II	0																							
III	1																							
IV	2																							
Rating	Number of experts																							
1	2																							
2	2																							
3	1																							
4	0																							
5	0																							
Admission in adult mild TBI, negative on a validated clinical decision rule (e.g. CATCH, NEXUS II) or normal CT and normal clinical exam, not on observation therapy[2 10-18]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>2</td></tr> <tr><td>III</td><td>5</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Level	Number of studies	I	0	II	2	III	5	IV	2	<table border="1"> <tr><th>Rating</th><th>Number of experts</th></tr> <tr><td>1</td><td>6</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> </table>	Rating	Number of experts	1	6	2	1	3	1	4	0	5	0
Level	Number of studies																							
I	0																							
II	2																							
III	5																							
IV	2																							
Rating	Number of experts																							
1	6																							
2	1																							
3	1																							
4	0																							
5	0																							
Admission in pediatric mild TBI, negative on validated clinical decision rule (e.g. CATCH, PECARN, CHALICE) or normal CT and normal clinical exam[19-21]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>1</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Level	Number of studies	I	0	II	0	III	1	IV	2	<table border="1"> <tr><th>Rating</th><th>Number of experts</th></tr> <tr><td>1</td><td>5</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> </table>	Rating	Number of experts	1	5	2	2	3	1	4	0	5	0
Level	Number of studies																							
I	0																							
II	0																							
III	1																							
IV	2																							
Rating	Number of experts																							
1	5																							
2	2																							
3	1																							
4	0																							
5	0																							
Admission in suspected scaphoid fracture with negative CT or MRI[22-24]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>2</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>1</td></tr> <tr><td>IV</td><td>1</td></tr> </table>	Level	Number of studies	I	2	II	0	III	1	IV	1	<table border="1"> <tr><th>Rating</th><th>Number of experts</th></tr> <tr><td>1</td><td>3</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> </table>	Rating	Number of experts	1	3	2	2	3	1	4	0	5	0
Level	Number of studies																							
I	2																							
II	0																							
III	1																							
IV	1																							
Rating	Number of experts																							
1	3																							
2	2																							
3	1																							
4	0																							
5	0																							


in adult mild TBI, negative on a validated clinical decision rule (e.g. HIP, NEXUS II)<sup>CW, EAST, NQF, CIHI</sup> [13 15 25-88]



in pediatric mild TBI, negative on a validated clinical decision rule (e.g. PECARN, CATCH, CHALICE)<sup>CW, CIHI</sup> [19-21 25 38 89-124]



and CT in pediatric mild TBI, positive initial CT and no clinical signs[125-134]



and CT in adult trauma, negative on a validated clinical decision rule (e.g. Canadian C-Spine Rule, NEXUS)<sup>CW, NQF, NICE</sup> [47 58 135-149]





and CT in pediatric trauma, able to co-operate and communicate and negative on a validated clinical decision rule (e.g. NEXUS)[109 124 150-154]





and CT of the neck in suspected blunt cerebrovascular injury, negative on a validated clinical decision rule (e.g. DENVER)[160-162]



and CT in adult blunt thoracic trauma, negative on a validated clinical decision rule (e.g. NEXUS-Chest)[163-172]



and CT in pediatric blunt abdominal trauma, negative on a validated clinical decision rule (e.g. PECARN, BATiC) and negative FAST[6 109 123 124]



and CT in pediatric multiple trauma, no pain, normal exam of pelvis/hip, no neurologic deficit, no hematuria or abdominal pain/tenderness, GCS>13 and hemodynamically stable[183]





CT in minor or single-system trauma <sup>CW, NICE</sup> [25 172 184-187]		
CT in pediatric trauma for injuries that the facility does not have ability to treat[6 188-191]		
CT repeat CT in transferred trauma patient with imaging performed at center, no disease progression or additional details needed[88 192-		
CT in pediatric minor head injury, negative on a validated clinical decision rule (e.g. C3PO)[124 197-199]		
CT in blunt trauma, hemodynamically stable with normal physical exam[200-205]		
CT in adult wrist injury with normal physical exam[206]		
CT in pediatric wrist injury, >2 years of age and normal physical exam[208]		
CT in blunt trauma, stable with negative physical exam for pelvic injury[205 209-213]		
CT in adult trauma, negative on a validated clinical decision rule (e.g. the Ottawa Rule, Pittsburgh)[214-217]		

<p>ay in adult trauma, negative on a validated clinical decision rule (e.g. Ottawa Ankle Rule)[218-239]</p>		
<p>ay in pediatric trauma, &gt;2 years of age and negative on a validated clinical decision rule (e.g. Ottawa Ankle Rule)[240-248]</p>		
<p>ood tests in trauma, &lt;60 years old, no regular medications, isolated trauma or low-energy injury and no significant medical history[249]</p>		
<p>zymes in sternal fractures[250]</p>		
<p>ostomy in pediatric blunt trauma with small hemothorax or occult fracture[251]</p>		
<p>ic acid &gt;3h in trauma<sup>NICE</sup> [172 252 253]</p>		
<p>ant factor VIIa (rFVIIa) in isolated TBI with intracerebral hemorrhage[254 255]</p>		
<p>ny in penetrating trauma with CPR &gt;15 minutes and no signs of life (no pupal response, respiratory effort, or motor activity)[256-259]</p>		
<p>ny in blunt trauma with CPR &gt; 10 minutes, no signs of life or normalizing the presenting rhythm and no pericardial tamponade[257-259]</p>		

\*Review phase: at least one Level I, II or III study (review phase) AND Consultation phase:  $\geq 75\%$  of experts who responded to the question classified the practice as clearly or potentially low value and no experts classified it as clearly beneficial















†**Level of evidence of clinical practices based on study design**, I, RCT or SR of RCT; II, prospective studies, quasi-randomized studies, SR of level II studies; III, case-control, case series, cross-sectional, retrospective, SR of level III studies; IV, expert consensus, narrative review, other

‡**Level of agreement of consulted experts on the value of clinical practices**, 1, clearly low-value; 2, possibly low-value; 3, controversial; 4, possibly beneficial; 5, clearly beneficial; 6, undecided

◇See eReferences for table's references

BATIC, Blunt Abdominal Trauma in Children; CATCH, Canadian Assessment of Tomography for Childhood Head injury; CCHR, Canadian CT Head Rule; CHALICE, Children's Head Injury Algorithm for the prediction of Important Clinical Events; CHIP, CT in Head Injury Patients; CIHI, Canadian Institute for Health Information; CPR, cardiopulmonary resuscitation; CT, computed tomography; CW, Choosing Wisely; EAST, Eastern Association for the Surgery of Trauma; FAST, Focused Assessment with Sonography in Trauma; GCS, Glasgow Coma Scale; MRI, magnetic resonance imaging; NEXUS, National Emergency X-Ray Utilization; NICE, National Institute for Health and Care Excellence; NQF, National Quality Forum; PECARN, Pediatric Emergency Care Applied Research Network; RCT, randomized controlled trial; SR, systematic review; TBI, traumatic brain injury

**Table 3.** Low value clinical practices in **general trauma surgery** according to level of evidence (review phase) and expert opinion (consultation phase)

Practices in surgery*	Level of evidence† I-RCT to IV-expert consensus Number of studies	Expert opinion‡ 1-clearly low value to 5 beneficial Number of experts
bedrest for pediatric blunt splenic or liver injury; >1 night for grade I-II and >2 nights for grade III[1 2]◇		
Observation for grade I-III renal injuries[3]		
Control laparotomy for resuscitated trauma patients who are hemodynamically restored and not massively transfused[4]		
Observation of grade IV-V liver injury in patients who are hemodynamically stable with no indication for surgical treatment of associated injuries† [5-9]		
Observation of pediatric liver injury[10 11]		
Observation of penetrating neck injury with soft signs on clinical examination and negative on multidetector CT angiography[12-16]		
Observation of penetrating renal injury in patients who are hemodynamically stable, have no contrast blush indicating arterial injury, have a viable kidney and have no gross extravasation[17 18]		

<p>management of blunt isolated splenic injury in patients who are hemodynamically stable<sup>EAST</sup>[19-24]</p>	<table border="1"> <thead> <tr> <th>Level</th> <th>Number of Studies</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>1</td> </tr> <tr> <td>II</td> <td>1</td> </tr> <tr> <td>III</td> <td>4</td> </tr> <tr> <td>IV</td> <td>1</td> </tr> </tbody> </table>	Level	Number of Studies	I	1	II	1	III	4	IV	1	<table border="1"> <thead> <tr> <th>Level</th> <th>Number of Experts</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>7</td> </tr> <tr> <td>2</td> <td>1</td> </tr> <tr> <td>3</td> <td>0</td> </tr> <tr> <td>4</td> <td>1</td> </tr> <tr> <td>5</td> <td>0</td> </tr> </tbody> </table>	Level	Number of Experts	1	7	2	1	3	0	4	1	5	0
Level	Number of Studies																							
I	1																							
II	1																							
III	4																							
IV	1																							
Level	Number of Experts																							
1	7																							
2	1																							
3	0																							
4	1																							
5	0																							
<p>management of pediatric splenic injury in children who are monitored hemodynamically stable[25-28]</p>	<table border="1"> <thead> <tr> <th>Level</th> <th>Number of Studies</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>0</td> </tr> <tr> <td>II</td> <td>0</td> </tr> <tr> <td>III</td> <td>3</td> </tr> <tr> <td>IV</td> <td>1</td> </tr> </tbody> </table>	Level	Number of Studies	I	0	II	0	III	3	IV	1	<table border="1"> <thead> <tr> <th>Level</th> <th>Number of Experts</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>7</td> </tr> <tr> <td>2</td> <td>1</td> </tr> <tr> <td>3</td> <td>0</td> </tr> <tr> <td>4</td> <td>0</td> </tr> <tr> <td>5</td> <td>0</td> </tr> </tbody> </table>	Level	Number of Experts	1	7	2	1	3	0	4	0	5	0
Level	Number of Studies																							
I	0																							
II	0																							
III	3																							
IV	1																							
Level	Number of Experts																							
1	7																							
2	1																							
3	0																							
4	0																							
5	0																							

\*Review phase: at least one Level I, II or III study (review phase) AND Consultation phase:  $\geq 75\%$  of experts who responded to the question classified the practice as clearly or potentially low value and no experts classified it as clearly beneficial

†**Level of evidence of clinical practices based on study design**, I, RCT or SR of RCT; II, prospective studies, quasi-randomized studies, SR of level II studies; III, case-control, case series, cross-sectional, retrospective, SR of level III studies; IV, expert consensus, narrative review, other



















‡**Level of agreement of consulted experts on the value of clinical practices**, 1, clearly low-value; 2, possibly low-value; 3, controversial; 4, possibly beneficial; 5, clearly beneficial; 6, undecided

◇See eReferences for table's references

CT, computed tomography; EAST, Eastern Association for the Surgery of Trauma; RCT, randomized controlled trial; SR, systematic review

**Table 4.** Low value clinical practices in the **intensive care unit** according to level of evidence (review phase) and expert opinion (consultation phase)

Clinical practices in the intensive care unit*	Level of evidence† I-RCT to IV-expert consensus Number of studies	Expert opinion‡ 1-clearly low value to 5 beneficial Number of experts
Anticoagulation in adults with acute mild complicated TBI who are not on mechanical ventilation[1-5]◇		
Medical consultation in adults with acute mild TBI and a negative CT[6]		
Inferior vena cava filter for prevention of PE in acute spinal cord injury with mechanical ventilation and no contraindications for low-molecular weight heparin[8 9]		
Use of pneumatic devices for thromboprophylaxis in nonambulatory patients admitted to the trauma service with no contraindications for low-molecular weight heparin[10]		
Oxygenation after chest tube removal in patients with thoracic trauma who are mechanically ventilated and have appropriate mental status to undergo a physical examination without new symptoms[11]		
Anticoagulation for thromboprophylaxis in basal skull fractures without evidence of CSF leakage[12-14]		

corticosteroids in spinal cord injury[15-20]		
corticosteroids in adults with TBI <sup>BTF, CW, NICE</sup> [21-32]		
e prophylaxis >1 week in adults with severe TBI <sup>BTF</sup> [32-36]		
n severe TBI[37-39]		
colloids (dextran, gelatin, hydroxyethyl starch) in trauma [40-46]		
fusion in adults with TBI on antiplatelet therapy[47-51]		
fusion in adult trauma patients above the transfusion threshold (hemoglobin >7 gram/deciliter) with no ongoing or suspected uncontrolled intracranial hemorrhage, no TBI and no coronary heart disease[52-66]		
c hypothermia in adults with TBI and ICP responding to other stage of care <sup>ACS, BTF</sup> [32 67-74]		
ic hyperventilation in adults with severe TBI <sup>BTF</sup> [22 28 32 67 75 76]		

\*Review phase: at least one Level I, II or III study (review phase) AND Consultation phase:  $\geq 75\%$  of experts who responded to the question classified the practice as clearly or potentially low value and no experts classified it as clearly beneficial

†**Level of evidence of clinical practices based on study design**, I, RCT or SR of RCT; II, prospective studies, quasi-randomized studies, SR of level II studies; III, case-control, case series, cross-sectional, retrospective, SR of level III studies; IV, expert consensus, narrative review, other

‡**Level of agreement of consulted experts on the value of clinical practices**, 1, clearly low-value; 2, possibly low-value; 3, controversial; 4, possibly beneficial; 5, clearly beneficial; 6, undecided

◇See eReferences for table's references

ACS, American College of Surgeons; BTF, Brain Trauma Foundation; CSF, cerebral spinal fluid; CT, computed tomography; CW, Choosing Wisely; ICP, intracranial pressure; NICE, National Institute for Health and Care Excellence; RCT, randomized controlled trial; SR, systematic review; TBI, traumatic brain injury; RBC: red blood cell



**Table 5.** Low value clinical practices in orthopaedics according to level of evidence (review phase) and expert opinion (consultation phase)

Practices in orthopaedics	Level of evidence† I-RCT to IV-expert consensus Number of studies	Expert opinion‡ 1-clearly low value to 5 beneficial Number of experts
consultation for pediatric closed isolated uncomplicated zone 2 fracture[1]◇		
ce consultation for isolated thoracolumbar transverse process		
Ray for isolated closed Mason-Johnson type-I radial head/neck with no clinical complaints[3]		
or A0-A3 thoracolumbar burst fracture with kyphotic deformity <35° associated posterior ligamentous complex injury and no symptoms[4-7]		
ve blood tests for American Society of Anaesthesiologists (ASA) classification of orthopedic injury requiring minor surgery[8]		

\*Review phase: at least one Level I, II or III study (review phase) AND Consultation phase:  $\geq 75\%$  of experts who responded to the question classified the practice as clearly or potentially low value and no experts classified it as clearly beneficial

†**Level of evidence of clinical practices based on study design**, I, RCT or SR of RCT; II, prospective studies, quasi-randomized studies, SR of level II studies; III, case-control, case series, cross-sectional, retrospective, SR of level III studies; IV, expert consensus, narrative review, other

‡**Level of agreement of consulted experts on the value of clinical practices**, 1, clearly low-value; 2, possibly low-value; 3, controversial; 4, possibly beneficial; 5, clearly beneficial; 6, undecided

◇See eReferences for table's references

RCT, randomized controlled trial; SR, systematic review

ACCEPTED

eTable 1. Ovid search strategies

eFigure 1. Preferred Reporting Items for Systematic Reviews and Meta-analysis flow diagram

eFigure 2. Extract from the on-line survey

eTable 2a. Grey zone clinical practices in the **emergency department** according to level of evidence (review phase) and expert opinion (consultation phase)

eTable 2b. Grey zone clinical practices in **general trauma surgery** according to level of evidence (review phase) and expert opinion (consultation phase)

eTable 2c. Grey zone clinical practices in the **intensive care unit** according to level of evidence (review phase) and expert opinion (consultation phase)

eTable 2d. Grey zone clinical practices in **orthopaedics** according to level of evidence (review phase) and expert opinion (consultation phase)

eReferences.

**eTable 1. Ovid search strategies**

MEDLINE SEARCH STRATEGY
<p><b>1. Trauma</b></p> <p>exp "Craniocerebral Trauma"/ OR "Craniocerebral Trauma".ti,ab. OR "head injur\$.ti,ab. OR "traumatic brain injur\$.ti,ab. OR Fracture.ti,ab. OR Injur\$.ti,ab. OR exp "Motor Vehicles"/ OR "motor vehicle collision".ti,ab. OR "motor vehicle crash".ti,ab. OR "Traffic accidents".ti,ab. OR Spinal Cord Injuries/ OR Spinal Cord Injur\$.ti,ab. OR Spinal cord trauma?.ti,ab. OR Trauma?.ti,ab. OR Wound\$.ti,ab. OR exp "Wounds and Injuries"/</p>
<p><b>2. Criteria to evaluate overuse</b></p> <p>De-adopt\$.ti,ab. OR Decommission\$.ti,ab. OR de-commission\$.ti,ab. OR Deimplent\$.ti,ab. OR De-list\$.ti,ab. OR Disinvest\$.ti,ab. OR dis-invest\$.ti,ab. OR Do-not-do.ti,ab. OR Harm\$.ti,ab. OR "patient harm"/ OR Inappropriate\$.ti,ab. OR Ineffective\$.ti,ab. OR "low quality".ti,ab. OR "low-value".ti,ab. OR Misuse.ti,ab. OR "Health Services Misuse"/ OR (overuse\$.ti,ab. not "overuse injury".ti,ab.) OR "medical overuse"/ OR "poor quality".ti,ab. OR "practice reversal".ti,ab. OR "medical reversal".ti,ab. OR Unnecessary.ti,ab. OR "Unnecessary Procedures"/ OR Unneeded.ti,ab. OR Wasteful.ti,ab.</p>
<p><b>3. Human animals only</b></p> <p>Animals/ NOT humans/</p>
<p><b>4. Years</b></p> <p>("2006" or "2007" or "2008" or "2009" or "2010" or "2011" or "2012" or "2013" or "2014" or "2015" or "2016" or "2017" or "2018").yr.</p>
<p><b>Finalization</b></p> <p><b>5. (1 AND 2 AND 4) NOT 3</b></p> <p><b>6. Limit 5 to English language</b></p>
EMBASE SEARCH STRATEGY
<p><b>1. Trauma</b></p>

exp "Craniocerebral Trauma"/ OR "Craniocerebral Trauma".ti,ab. OR "head injur\$.ti,ab. OR "traumatic brain injur\$.ti,ab. OR Fracture.ti,ab. OR Injur\$.ti,ab. OR exp "Motor Vehicles"/ OR "motor vehicle collision".ti,ab. OR "motor vehicle crash".ti,ab. OR "Traffic accidents".ti,ab. OR Spinal Cord Injuries/ OR Spinal Cord Injur\$.ti,ab. OR Spinal cord trauma?.ti,ab. OR Trauma?.ti,ab. OR Wound\$.ti,ab. OR exp "Wounds and Injuries"/

**2. Criteria to evaluate overuse**

De-adopt\$.ti,ab. OR Decommission\$.ti,ab. OR de-commission\$.ti,ab. OR Deimplent\$.ti,ab. OR De-list\$.ti,ab. OR Disinvest\$.ti,ab. OR dis-invest\$.ti,ab. OR Do-not-do.ti,ab. OR Harm\$.ti,ab. OR "patient harm"/ OR Inappropriate\$.ti,ab. OR Ineffective\$.ti,ab. OR "low quality".ti,ab. OR "low-value".ti,ab. OR Misuse.ti,ab. OR "Health Services Misuse"/ OR (overuse\$.ti,ab. not "overuse injury".ti,ab.) OR "medical overuse"/ OR "poor quality".ti,ab. OR "practice reversal".ti,ab. OR "medical reversal".ti,ab. OR Unnecessary.ti,ab. OR "Unnecessary Procedures"/ OR Unneeded.ti,ab. OR Wasteful.ti,ab.

**3. Human animals only**

Animals/ NOT humans/

**4. Years**


("2006" or "2007" or "2008" or "2009" or "2010" or "2011" or "2012" or "2013" or "2014" or "2015" or "2016" or "2017" or "2018").yr.

**Finalization**

**5. (1 AND 2 AND 4) NOT 3**

**6. limit 5 to English language**

**eFigure 1. Preferred Reporting Items for Systematic Reviews and Meta-analysis flow diagram**



**eFigure 2. Extract from the on-line survey**















See Online Supplements 2

ACCEPTED

**eTable 2a. Grey zone clinical practices in the emergency department according to level of evidence (review phase) and expert opinion (consultation phase)**

Clinical practices in the emergency department	Level of evidence† I-RCT to IV-expert consensus Number of studies	Expert opinion‡ 1-clearly low value to 5-clearly beneficial Number of experts																								
Hospital admission in isolated sternal fractures with normal cardiac enzymes (troponin) and normal ECG[1]◇	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>1</td></tr> </table>	Level	Number of studies	I	0	II	0	III	0	IV	1	<table border="1"> <tr><th>Expert opinion</th><th>Number of experts</th></tr> <tr><td>1</td><td>5</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Expert opinion	Number of experts	1	5	2	2	3	0	4	0	5	0	6	0
Level	Number of studies																									
I	0																									
II	0																									
III	0																									
IV	1																									
Expert opinion	Number of experts																									
1	5																									
2	2																									
3	0																									
4	0																									
5	0																									
6	0																									
Hospital admission in pediatric isolated skull fracture with GCS=15, normal neurological exam and low-energy injury mechanism[2-7]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>1</td></tr> <tr><td>III</td><td>3</td></tr> <tr><td>IV</td><td>1</td></tr> </table>	Level	Number of studies	I	0	II	1	III	3	IV	1	<table border="1"> <tr><th>Expert opinion</th><th>Number of experts</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>3</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Expert opinion	Number of experts	1	1	2	2	3	3	4	0	5	0	6	0
Level	Number of studies																									
I	0																									
II	1																									
III	3																									
IV	1																									
Expert opinion	Number of experts																									
1	1																									
2	2																									
3	3																									
4	0																									
5	0																									
6	0																									
Cervical collar retention in obtunded or intubated trauma patient with no injuries detected on cervical spine CT[8-10]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>3</td></tr> <tr><td>III</td><td>3</td></tr> <tr><td>IV</td><td>3</td></tr> </table>	Level	Number of studies	I	0	II	3	III	3	IV	3	<table border="1"> <tr><th>Expert opinion</th><th>Number of experts</th></tr> <tr><td>1</td><td>0</td></tr> <tr><td>2</td><td>3</td></tr> <tr><td>3</td><td>3</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Expert opinion	Number of experts	1	0	2	3	3	3	4	0	5	0	6	0
Level	Number of studies																									
I	0																									
II	3																									
III	3																									
IV	3																									
Expert opinion	Number of experts																									
1	0																									
2	3																									
3	3																									
4	0																									
5	0																									
6	0																									
Thoracolumbar spine X-Ray in patients with no complaints of thoracolumbar spinal pain, normal mental status and normal neurological and physical examination[11]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>1</td></tr> </table>	Level	Number of studies	I	0	II	0	III	0	IV	1	<table border="1"> <tr><th>Expert opinion</th><th>Number of experts</th></tr> <tr><td>1</td><td>5</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Expert opinion	Number of experts	1	5	2	2	3	0	4	0	5	0	6	0
Level	Number of studies																									
I	0																									
II	0																									
III	0																									
IV	1																									
Expert opinion	Number of experts																									
1	5																									
2	2																									
3	0																									
4	0																									
5	0																									
6	0																									



Repeat head CT in adult mild TBI with negative initial CT and on anticoagulant and/or antiplatelet therapy[12-24]	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>4</td></tr> <tr><td>III</td><td>8</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of Studies	I	0	II	4	III	8	IV	0	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>3</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Level	Number of Studies	1	2	2	1	3	3	4	2	5	0	6	0
Level	Number of Studies																									
I	0																									
II	4																									
III	8																									
IV	0																									
Level	Number of Studies																									
1	2																									
2	1																									
3	3																									
4	2																									
5	0																									
6	0																									
Repeat head CT in adult mild complicated TBI[12 25-30]	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>2</td></tr> <tr><td>III</td><td>5</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of Studies	I	0	II	2	III	5	IV	0	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>1</td><td>0</td></tr> <tr><td>2</td><td>4</td></tr> <tr><td>3</td><td>3</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Level	Number of Studies	1	0	2	4	3	3	4	1	5	0	6	0
Level	Number of Studies																									
I	0																									
II	2																									
III	5																									
IV	0																									
Level	Number of Studies																									
1	0																									
2	4																									
3	3																									
4	1																									
5	0																									
6	0																									
Chest CT in pediatric blunt thoracic trauma with normal mediastinal silhouette on X-Ray <sup>NICE</sup> [31 32]	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>5</td></tr> <tr><td>IV</td><td>5</td></tr> </table>	Level	Number of Studies	I	0	II	0	III	5	IV	5	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>5</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Level	Number of Studies	1	1	2	1	3	5	4	1	5	0	6	0
Level	Number of Studies																									
I	0																									
II	0																									
III	5																									
IV	5																									
Level	Number of Studies																									
1	1																									
2	1																									
3	5																									
4	1																									
5	0																									
6	0																									
Abdominal CT in adult blunt abdominal trauma with normal physical exam and negative FAST[33-43]	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>6</td></tr> <tr><td>III</td><td>3</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Level	Number of Studies	I	0	II	6	III	3	IV	2	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>3</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Level	Number of Studies	1	1	2	2	3	3	4	2	5	0	6	0
Level	Number of Studies																									
I	0																									
II	6																									
III	3																									
IV	2																									
Level	Number of Studies																									
1	1																									
2	2																									
3	3																									
4	2																									
5	0																									
6	0																									
Routine panels in pediatric blunt abdominal trauma[44]	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of Studies	I	0	II	0	III	2	IV	0	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>3</td></tr> <tr><td>3</td><td>2</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Level	Number of Studies	1	2	2	3	3	2	4	1	5	0	6	0
Level	Number of Studies																									
I	0																									
II	0																									
III	2																									
IV	0																									
Level	Number of Studies																									
1	2																									
2	3																									
3	2																									
4	1																									
5	0																									
6	0																									
Head MRI in adult TBI who received timely helical CT with a new generation scanner <sup>NQF, NICE</sup> [45-49]	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>1</td></tr> <tr><td>III</td><td>1</td></tr> <tr><td>IV</td><td>3</td></tr> </table>	Level	Number of Studies	I	0	II	1	III	1	IV	3	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>1</td><td>3</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Level	Number of Studies	1	3	2	2	3	1	4	1	5	0	6	1
Level	Number of Studies																									
I	0																									
II	1																									
III	1																									
IV	3																									
Level	Number of Studies																									
1	3																									
2	2																									
3	1																									
4	1																									
5	0																									
6	1																									
Aerodigestive tract endoscopy in penetrating neck injury with negative neck exploration[50]	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of Studies	I	0	II	0	III	2	IV	0	 <table border="1"> <tr><th>Level</th><th>Number of Studies</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>3</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Level	Number of Studies	1	1	2	2	3	1	4	3	5	0	6	1
Level	Number of Studies																									
I	0																									
II	0																									
III	2																									
IV	0																									
Level	Number of Studies																									
1	1																									
2	2																									
3	1																									
4	3																									
5	0																									
6	1																									

Esophagography in esophageal injury with pneumomediastinum but a negative CT[51]		
Massive transfusion in trauma, negative on a validated score (e.g. TASH, revised MTS, ABC)[52-53]		
Thoracotomy in pediatric blunt trauma with cardiac arrest[54]		
Cardiopulmonary resuscitation in trauma, resuscitation >15 mins and no immediate reversible cause[55]		

†Level of evidence of clinical practices based on study design, I, RCT or SR of RCT; II, prospective studies, quasi-randomized studies, SR of level II studies; III, case-control, case series, cross-sectional, retrospective, SR of level III studies; IV, expert consensus, narrative review, other

‡Level of agreement of consulted experts on the value of clinical practices, 1, clearly low-value; 2, possibly low-value; 3, controversial; 4, possibly beneficial; 5, clearly beneficial; 6, undecided















◇See eReferences for table's references

ABC, Assessment of Blood Consumption; CT, computed tomography; ECG, electrocardiogram; FAST, Focused Assessment with Sonography in Trauma; GCS, Glasgow Coma Scale; MRI, magnetic resonance imaging; MTS, Massive Transfusion Score; NICE, National Institute for Health and Care Excellence; NQF, National Quality Forum; RCT, randomized controlled trial; SR, systematic review; TASH, Trauma Associated Severe Hemorrhage; TBI, traumatic brain injury

**eTable 2b. Grey zone clinical practices in general trauma surgery according to level of evidence (review phase) and expert opinion (consultation phase)**

Clinical practices in surgery	Level of evidence†	Expert opinion‡
-------------------------------	--------------------	-----------------

	<b>I-RCT to IV-expert consensus Number of studies</b>	<b>1-clearly low value to 5-clearly beneficial Number of experts</b>
Hospital admission for stable patients with an abdominal anterior stab wound, negative FAST and negative wound exploration <sup>EAST</sup> [1-3]◇		
Hospitalisation > 24 hours for penetrating abdominal trauma with non-operative management, reliable abdominal examination, and minimal or no abdominal tenderness <sup>EAST</sup> [1]		
Follow-up imaging for blunt grade IV renovascular renal injury with non-operative management and no clinical deterioration[4]		
Follow-up imaging for blunt grade I-III renal injury with non-operative management and no clinical deterioration[4 5]		
Stent graft for minimal aortic injury with regression on follow-up CTA[6]		
Decompression, diversion, exclusion for full thickness duodenal laceration managed with damage control surgery[7]		

Foley catheter for temporary hemostasis in gaping cardiac injury[8]	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>0</td> </tr> <tr> <td>II</td> <td>0</td> </tr> <tr> <td>III</td> <td>0</td> </tr> <tr> <td>IV</td> <td>1</td> </tr> </tbody> </table>	Category	Count	I	0	II	0	III	0	IV	1	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>2</td> <td>1</td> </tr> <tr> <td>3</td> <td>2</td> </tr> <tr> <td>4</td> <td>5</td> </tr> <tr> <td>5</td> <td>0</td> </tr> <tr> <td>6</td> <td>0</td> </tr> </tbody> </table>	Category	Count	1	1	2	1	3	2	4	5	5	0	6	0
Category	Count																									
I	0																									
II	0																									
III	0																									
IV	1																									
Category	Count																									
1	1																									
2	1																									
3	2																									
4	5																									
5	0																									
6	0																									
Prophylactic nasogastric decompression following emergency laparotomy for abdominal injury[9]	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>1</td> </tr> <tr> <td>II</td> <td>0</td> </tr> <tr> <td>III</td> <td>0</td> </tr> <tr> <td>IV</td> <td>0</td> </tr> </tbody> </table>	Category	Count	I	1	II	0	III	0	IV	0	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>0</td> </tr> <tr> <td>2</td> <td>6</td> </tr> <tr> <td>3</td> <td>1</td> </tr> <tr> <td>4</td> <td>2</td> </tr> <tr> <td>5</td> <td>0</td> </tr> <tr> <td>6</td> <td>0</td> </tr> </tbody> </table>	Category	Count	1	0	2	6	3	1	4	2	5	0	6	0
Category	Count																									
I	1																									
II	0																									
III	0																									
IV	0																									
Category	Count																									
1	0																									
2	6																									
3	1																									
4	2																									
5	0																									
6	0																									
Complex surgery for duodenal injury from low-velocity gunshot wound with <50% circumference[10]	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>0</td> </tr> <tr> <td>II</td> <td>0</td> </tr> <tr> <td>III</td> <td>1</td> </tr> <tr> <td>IV</td> <td>1</td> </tr> </tbody> </table>	Category	Count	I	0	II	0	III	1	IV	1	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>1</td> </tr> <tr> <td>2</td> <td>4</td> </tr> <tr> <td>3</td> <td>1</td> </tr> <tr> <td>4</td> <td>0</td> </tr> <tr> <td>5</td> <td>1</td> </tr> <tr> <td>6</td> <td>1</td> </tr> </tbody> </table>	Category	Count	1	1	2	4	3	1	4	0	5	1	6	1
Category	Count																									
I	0																									
II	0																									
III	1																									
IV	1																									
Category	Count																									
1	1																									
2	4																									
3	1																									
4	0																									
5	1																									
6	1																									
Damage control laparotomy for pediatric trauma[11 12]	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>0</td> </tr> <tr> <td>II</td> <td>0</td> </tr> <tr> <td>III</td> <td>2</td> </tr> <tr> <td>IV</td> <td>1</td> </tr> </tbody> </table>	Category	Count	I	0	II	0	III	2	IV	1	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>0</td> </tr> <tr> <td>2</td> <td>1</td> </tr> <tr> <td>3</td> <td>0</td> </tr> <tr> <td>4</td> <td>2</td> </tr> <tr> <td>5</td> <td>1</td> </tr> <tr> <td>6</td> <td>2</td> </tr> </tbody> </table>	Category	Count	1	0	2	1	3	0	4	2	5	1	6	2
Category	Count																									
I	0																									
II	0																									
III	2																									
IV	1																									
Category	Count																									
1	0																									
2	1																									
3	0																									
4	2																									
5	1																									
6	2																									
Surgical management of penetrating zone II neck injury without hard signs <sup>EAST</sup> [13-16]	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>0</td> </tr> <tr> <td>II</td> <td>1</td> </tr> <tr> <td>III</td> <td>2</td> </tr> <tr> <td>IV</td> <td>1</td> </tr> </tbody> </table>	Category	Count	I	0	II	1	III	2	IV	1	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>3</td> </tr> <tr> <td>2</td> <td>1</td> </tr> <tr> <td>3</td> <td>0</td> </tr> <tr> <td>4</td> <td>2</td> </tr> <tr> <td>5</td> <td>0</td> </tr> <tr> <td>6</td> <td>1</td> </tr> </tbody> </table>	Category	Count	1	3	2	1	3	0	4	2	5	0	6	1
Category	Count																									
I	0																									
II	1																									
III	2																									
IV	1																									
Category	Count																									
1	3																									
2	1																									
3	0																									
4	2																									
5	0																									
6	1																									
Surgical management of grade III-IV pancreatic injury in patients who are hemodynamically stable and have no hollow organ injuries[17 18]	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>1</td> </tr> <tr> <td>II</td> <td>0</td> </tr> <tr> <td>III</td> <td>1</td> </tr> <tr> <td>IV</td> <td>0</td> </tr> </tbody> </table>	Category	Count	I	1	II	0	III	1	IV	0	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>0</td> </tr> <tr> <td>2</td> <td>0</td> </tr> <tr> <td>3</td> <td>3</td> </tr> <tr> <td>4</td> <td>2</td> </tr> <tr> <td>5</td> <td>2</td> </tr> <tr> <td>6</td> <td>0</td> </tr> </tbody> </table>	Category	Count	1	0	2	0	3	3	4	2	5	2	6	0
Category	Count																									
I	1																									
II	0																									
III	1																									
IV	0																									
Category	Count																									
1	0																									
2	0																									
3	3																									
4	2																									
5	2																									
6	0																									
Surgical management of blunt grade IV-V renal injury in patients who are hemodynamically stable[2 18-23]	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>1</td> </tr> <tr> <td>II</td> <td>0</td> </tr> <tr> <td>III</td> <td>3</td> </tr> <tr> <td>IV</td> <td>3</td> </tr> </tbody> </table>	Category	Count	I	1	II	0	III	3	IV	3	 <table border="1"> <thead> <tr> <th>Category</th> <th>Count</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>4</td> </tr> <tr> <td>2</td> <td>3</td> </tr> <tr> <td>3</td> <td>0</td> </tr> <tr> <td>4</td> <td>0</td> </tr> <tr> <td>5</td> <td>1</td> </tr> <tr> <td>6</td> <td>0</td> </tr> </tbody> </table>	Category	Count	1	4	2	3	3	0	4	0	5	1	6	0
Category	Count																									
I	1																									
II	0																									
III	3																									
IV	3																									
Category	Count																									
1	4																									
2	3																									
3	0																									
4	0																									
5	1																									
6	0																									

Surgical management of blunt isolated splenic or liver injury in patients with no peritonitis who are hemodynamically stable or unstable but responsive[20 22 24-27]		
Surgical management of penetrating transmediastinal injury in patients who are hemodynamically stable and are either negative on CT or positive on CT but negative on esophagoscopy/esophagography, bronchoscopy or angiography[28]		

†Level of evidence of clinical practices based on study design, I, RCT or SR of RCT; II, prospective studies, quasi-randomized studies, SR of level II studies; III, case-control, case series, cross-sectional, retrospective, SR of level III studies; IV, expert consensus, narrative review, other















‡Level of agreement of consulted experts on the value of clinical practices, 1, clearly low-value; 2, possibly low-value; 3, controversial; 4, possibly beneficial; 5, clearly beneficial; 6, undecided















◇See eReferences for table’s references















CT, computed tomography; CTA, CT angiography; EAST, Eastern Association for the Surgery of Trauma; FAST, Focused Assessment with Sonography in Trauma; RCT, randomized controlled trial; SR, systematic review

**eTable 2c. Grey zone clinical practices in the intensive care unit according to level of evidence (review phase) and expert opinion (consultation phase)**

Clinical practices in the intensive care unit	Level of evidence† I-RCT to IV-expert consensus Number of studies	Expert opinion‡ 1-clearly low value to 5-clearly beneficial Number of experts
---	--	--

Neurosurgical consultation in adults with acute mild complicated TBI[1]◇	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	0	II	0	III	2	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>4</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Category	Count	1	2	2	4	3	1	4	1	5	0	6	1
Category	Count																									
I	0																									
II	0																									
III	2																									
IV	0																									
Category	Count																									
1	2																									
2	4																									
3	1																									
4	1																									
5	0																									
6	1																									
Decompressive craniectomy in severe TBI with diffuse injury and refractory ICP[2-5]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>2</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>3</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Category	Count	I	2	II	0	III	3	IV	2	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>2</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>1</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Category	Count	1	2	2	1	3	2	4	1	5	1	6	0
Category	Count																									
I	2																									
II	0																									
III	3																									
IV	2																									
Category	Count																									
1	2																									
2	1																									
3	2																									
4	1																									
5	1																									
6	0																									
Decompressive craniectomy in severe TBI as a standard of care <sup>ACS, BTF</sup> [2-6]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>2</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>3</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Category	Count	I	2	II	0	III	3	IV	2	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>4</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Category	Count	1	4	2	1	3	1	4	2	5	0	6	0
Category	Count																									
I	2																									
II	0																									
III	3																									
IV	2																									
Category	Count																									
1	4																									
2	1																									
3	1																									
4	2																									
5	0																									
6	0																									
Inferior vena cava filter for prevention of PE in isolated acute TBI with intracerebral hemorrhage and no DVT[7]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Category	Count	I	0	II	0	III	0	IV	2	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>5</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Category	Count	1	5	2	2	3	0	4	1	5	0	6	0
Category	Count																									
I	0																									
II	0																									
III	0																									
IV	2																									
Category	Count																									
1	5																									
2	2																									
3	0																									
4	1																									
5	0																									
6	0																									
ICP monitoring in adults with severe TBI, normal CT and not more than one of the following criteria: aged>40, unilateral or bilateral posturing, systolic blood pressure <90 mmHg <sup>ACS</sup> [8-10]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>3</td></tr> </table>	Category	Count	I	0	II	0	III	2	IV	3	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>0</td></tr> <tr><td>2</td><td>4</td></tr> <tr><td>3</td><td>2</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>2</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Category	Count	1	0	2	4	3	2	4	1	5	2	6	0
Category	Count																									
I	0																									
II	0																									
III	2																									
IV	3																									
Category	Count																									
1	0																									
2	4																									
3	2																									
4	1																									
5	2																									
6	0																									
Neurological assessments hourly >24h in adults admitted to the ICU with mild or moderate TBI who are stable[11]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	0	II	0	III	2	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>4</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>1</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Category	Count	1	1	2	4	3	0	4	0	5	1	6	1
Category	Count																									
I	0																									
II	0																									
III	2																									
IV	0																									
Category	Count																									
1	1																									
2	4																									
3	0																									
4	0																									
5	1																									
6	1																									
Neurological assessments hourly >24h in adults admitted to the ICU with severe TBI who are stable[11]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	0	II	0	III	2	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>0</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>2</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Category	Count	1	0	2	2	3	1	4	1	5	2	6	0
Category	Count																									
I	0																									
II	0																									
III	2																									
IV	0																									
Category	Count																									
1	0																									
2	2																									
3	1																									
4	1																									
5	2																									
6	0																									

<p>Antibiotic combination therapy to cover gram negative bacilli as standard of care in trauma patients with ventilator-associated pneumonia[12]</p>		
<p>Antibiotic combination therapy to cover gram negative bacilli and MRSA as standard of care in trauma patients with ventilator-associated pneumonia[12]</p>		
<p>Postoperative antibiotic prophylaxis in penetrating abdominal trauma with no hollow viscus injury[13]</p>		
<p>Antibiotic prophylaxis in basal skull fractures with evidence of CSF leakage[14-16]</p>		
<p>Antibiotic prophylaxis &gt;24h post-operation in penetrating abdominal trauma with or without hollow viscus injury<sup>EAST</sup>[17]</p>		
<p>Antibiotic prophylaxis for external ventricular drain placement in adults with TBI[18]</p>		
<p>Barbiturates in adults with severe TBI<sup>BTF</sup>[5 18-21]</p>		

Dopamine antagonists (methylphenidate, amantadine, and bromocriptine) in adults with severe TBI[22]		
Antiseizure prophylaxis <1 week in adults with severe TBI and no seizure activity[18 23 24]		
Neuromuscular blocking agents in TBI with no refractory intracranial hypertension[25]		
Octreotide as routine post-operative prophylaxis to prevent fistula in pancreatic injuries[26]		
Hypertonic saline solution in severe TBI[7]		
Early hypertonic saline solution in TBI when intracranial pressure is not monitored[27]		
Plasma transfusion with international normalized ratio <1.3 in TBI[28]		



Therapeutic hypothermia in spinal cord injury[29]		
Hyperbaric oxygen therapy in TBI[19 30-32]		
Parenteral nutrition in trauma patients with no contraindications for enteral nutrition[25]		
Immunisation following angiographic embolization in splenic injury[33]		
Bed rest immobilization in blunt renal, hepatic or splenic injury[34]		

†Level of evidence of clinical practices based on study design, I, RCT or SR of RCT; II, prospective studies, quasi-randomized studies, SR of level II studies; III, case-control, case series, cross-sectional, retrospective, SR of level III studies; IV, expert consensus, narrative review, other

‡Level of agreement of consulted experts on the value of clinical practices, 1, clearly low-value; 2, possibly low-value; 3, controversial; 4, possibly beneficial; 5, clearly beneficial; 6, undecided















◇See eReferences for table's references













ACS, American College of Surgeons; BTF, Brain Trauma Foundation; CSF, cerebral spinal fluid; CT, computed tomography; DVT, deep vein thrombosis; EAST, Eastern Association for the Surgery of Trauma; ICP, intracranial pressure; MRSA, Methicillin-Resistant Staphylococcus Aureus; PE, pulmonary embolism; RBC, red blood cells; RCT, randomized controlled trial; SR, systematic review; TBI, traumatic brain injury

**eTable 2d. Grey zone clinical practices in orthopaedics according to level of evidence (review phase) and expert opinion (consultation phase)**

Clinical practices in orthopedics	Level of evidence† I-RCT to IV-expert consensus Number of studies	Expert opinion‡ 1-clearly low value to 5-clearly beneficial Number of experts																								
Follow-up consultation for adults with adequately aligned fifth metacarpal fracture[1-3]◇	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>3</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of studies	I	0	II	3	III	0	IV	0	<table border="1"> <tr><th>Value</th><th>Number of experts</th></tr> <tr><td>1</td><td>4</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Value	Number of experts	1	4	2	1	3	0	4	2	5	0	6	1
Level	Number of studies																									
I	0																									
II	3																									
III	0																									
IV	0																									
Value	Number of experts																									
1	4																									
2	1																									
3	0																									
4	2																									
5	0																									
6	1																									
Follow-up consultation for adult with fifth metatarsal fracture[4]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of studies	I	0	II	0	III	2	IV	0	<table border="1"> <tr><th>Value</th><th>Number of experts</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Value	Number of experts	1	2	2	1	3	1	4	1	5	0	6	1
Level	Number of studies																									
I	0																									
II	0																									
III	2																									
IV	0																									
Value	Number of experts																									
1	2																									
2	1																									
3	1																									
4	1																									
5	0																									
6	1																									
Follow-up consultation for adult with non-displaced or minimally displaced distal radius fracture[3]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>2</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of studies	I	0	II	2	III	0	IV	0	<table border="1"> <tr><th>Value</th><th>Number of experts</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>0</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Value	Number of experts	1	2	2	0	3	1	4	2	5	0	6	1
Level	Number of studies																									
I	0																									
II	2																									
III	0																									
IV	0																									
Value	Number of experts																									
1	2																									
2	0																									
3	1																									
4	2																									
5	0																									
6	1																									
Follow-up consultation for adult with Mason I radial head and neck fracture[5]	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of studies	I	0	II	0	III	2	IV	0	<table border="1"> <tr><th>Value</th><th>Number of experts</th></tr> <tr><td>1</td><td>4</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Value	Number of experts	1	4	2	1	3	1	4	1	5	0	6	1
Level	Number of studies																									
I	0																									
II	0																									
III	2																									
IV	0																									
Value	Number of experts																									
1	4																									
2	1																									
3	1																									
4	1																									
5	0																									
6	1																									
Hand surgery consultation for adult hand injury without injury to the nerves, tendons or joints, skin loss or complex fractures or injuries requiring skin grafting or	<table border="1"> <tr><th>Level</th><th>Number of studies</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Number of studies	I	0	II	0	III	2	IV	0	<table border="1"> <tr><th>Value</th><th>Number of experts</th></tr> <tr><td>1</td><td>3</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Value	Number of experts	1	3	2	2	3	0	4	1	5	0	6	1
Level	Number of studies																									
I	0																									
II	0																									
III	2																									
IV	0																									
Value	Number of experts																									
1	3																									
2	2																									
3	0																									
4	1																									
5	0																									
6	1																									

reconstruction[6 7]		
Follow-up consultation for pediatric distal radial metaphysis buckle fracture[8]		
Follow-up consultation for uncomplicated pediatric toddler fractures[9]		
Repeat X-Ray for fractures with fixation repair and no clinical complaints[10]		
Repeat X-Ray for torus or buckle distal radial fracture[11]		
X-Ray on cast removal for adult $\geq 50$ years old with a closed distal radius fracture, $<2$ cm from the distal end of the radius, living independently before the fracture[12]		
Post-operative X-Ray for pediatric forearm fracture treated with manipulation under anesthesia with fluoroscopic guidance[13]		

<p>Post-operative X-Ray for pediatric pin-fixed displaced supracondylar humeral fracture[14]</p>	 <p>10 5 0</p> <p>I II III IV</p>	 <p>8 4 0</p> <p>1 2 3 4 5 6</p>
<p>Post-operative X-Ray of fractures treated by operative fixation with a load-sharing construct in good quality bone[15]</p>	 <p>10 5 0</p> <p>I II III IV</p>	 <p>8 4 0</p> <p>1 2 3 4 5 6</p>
<p>Post splinting X-Ray of non-displaced and minimally displaced fractures with no manipulation before or during immobilization[16 17]</p>	 <p>10 5 0</p> <p>I II III IV</p>	 <p>8 4 0</p> <p>1 2 3 4 5 6</p>
<p>Magnetic resonance imaging for suspected scaphoid fracture[18]</p>	 <p>10 5 0</p> <p>I II III IV</p>	 <p>8 4 0</p> <p>1 2 3 4 5 6</p>
<p>Routine in-hospital post-operative X-Ray for surgically treated thoracolumbar injuries with no clinical deterioration[19]</p>	 <p>10 5 0</p> <p>I II III IV</p>	 <p>8 4 0</p> <p>1 2 3 4 5 6</p>
<p>Cast immobilization for adult fifth metacarpal neck fracture[1 20]</p>	 <p>10 5 0</p> <p>I II III IV</p>	 <p>8 4 0</p> <p>1 2 3 4 5 6</p>
<p>Immobilization for suspected scaphoid fractures with negative computed tomography or magnetic resonance imaging[21 22]</p>	 <p>10 5 0</p> <p>I II III IV</p>	 <p>8 4 0</p> <p>1 2 3 4 5 6</p>

Reduction and cast immobilization in fifth metacarpal neck fracture with initial angulation of less than 70 degrees[20]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>10</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	10	II	0	III	0	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>4</td></tr> <tr><td>3</td><td>2</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>2</td></tr> </table>	Category	Count	1	2	2	4	3	2	4	2	5	0	6	2
Category	Count																									
I	10																									
II	0																									
III	0																									
IV	0																									
Category	Count																									
1	2																									
2	4																									
3	2																									
4	2																									
5	0																									
6	2																									
Percutaneous pin fixation for adults with unstable, extra-articular distal radial fracture[23]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>10</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	10	II	0	III	0	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>0</td></tr> <tr><td>2</td><td>0</td></tr> <tr><td>3</td><td>2</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>4</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Category	Count	1	0	2	0	3	2	4	2	5	4	6	1
Category	Count																									
I	10																									
II	0																									
III	0																									
IV	0																									
Category	Count																									
1	0																									
2	0																									
3	2																									
4	2																									
5	4																									
6	1																									
Syndesmotic screw removal for adult surgical ankle fracture without persistent hardware complaints (asymptotic)[24-26]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>4</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	0	II	0	III	4	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>3</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>2</td></tr> </table>	Category	Count	1	3	2	2	3	1	4	0	5	0	6	2
Category	Count																									
I	0																									
II	0																									
III	4																									
IV	0																									
Category	Count																									
1	3																									
2	2																									
3	1																									
4	0																									
5	0																									
6	2																									
Radial head prosthesis in adult Mason IV radial head fracture-dislocation[27]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>1</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	0	II	0	III	1	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>0</td></tr> <tr><td>2</td><td>0</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>5</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Category	Count	1	0	2	0	3	0	4	0	5	5	6	1
Category	Count																									
I	0																									
II	0																									
III	1																									
IV	0																									
Category	Count																									
1	0																									
2	0																									
3	0																									
4	0																									
5	5																									
6	1																									
Long arm cast for pediatric (>4 years old) displaced distal third radius and ulna fractures[28]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>10</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	10	II	0	III	0	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>0</td></tr> <tr><td>2</td><td>5</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>1</td></tr> <tr><td>6</td><td>2</td></tr> </table>	Category	Count	1	0	2	5	3	0	4	0	5	1	6	2
Category	Count																									
I	10																									
II	0																									
III	0																									
IV	0																									
Category	Count																									
1	0																									
2	5																									
3	0																									
4	0																									
5	1																									
6	2																									
Rigid cast for pediatric isolated distal fibular fracture[29 30]	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>I</td><td>10</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Category	Count	I	10	II	0	III	0	IV	0	 <table border="1"> <tr><th>Category</th><th>Count</th></tr> <tr><td>1</td><td>2</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>2</td></tr> </table>	Category	Count	1	2	2	2	3	1	4	1	5	0	6	2
Category	Count																									
I	10																									
II	0																									
III	0																									
IV	0																									
Category	Count																									
1	2																									
2	2																									
3	1																									
4	1																									
5	0																									
6	2																									

Halo vest for geriatric type II odontoid fracture[31]	<table border="1"> <tr><th>Level</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>10</td></tr> </table>	Level	Count	I	0	II	0	III	0	IV	10	<table border="1"> <tr><th>Study</th><th>Count</th></tr> <tr><td>1</td><td>8</td></tr> <tr><td>2</td><td>0</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Study	Count	1	8	2	0	3	1	4	0	5	0	6	0
Level	Count																									
I	0																									
II	0																									
III	0																									
IV	10																									
Study	Count																									
1	8																									
2	0																									
3	1																									
4	0																									
5	0																									
6	0																									
Open Reduction and Internal Fixation (ORIF) in Mason II radial head fractures[32 33]	<table border="1"> <tr><th>Level</th><th>Count</th></tr> <tr><td>I</td><td>2</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Level	Count	I	2	II	0	III	0	IV	2	<table border="1"> <tr><th>Study</th><th>Count</th></tr> <tr><td>1</td><td>0</td></tr> <tr><td>2</td><td>1</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>1</td></tr> <tr><td>6</td><td>2</td></tr> </table>	Study	Count	1	0	2	1	3	0	4	2	5	1	6	2
Level	Count																									
I	2																									
II	0																									
III	0																									
IV	2																									
Study	Count																									
1	0																									
2	1																									
3	0																									
4	2																									
5	1																									
6	2																									
Hemiarthroplasty in patients 65 years of age and over with a proximal, four-part humeral fracture[34]	<table border="1"> <tr><th>Level</th><th>Count</th></tr> <tr><td>I</td><td>2</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Level	Count	I	2	II	0	III	0	IV	2	<table border="1"> <tr><th>Study</th><th>Count</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>0</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>2</td></tr> </table>	Study	Count	1	1	2	2	3	1	4	0	5	0	6	2
Level	Count																									
I	2																									
II	0																									
III	0																									
IV	2																									
Study	Count																									
1	1																									
2	2																									
3	1																									
4	0																									
5	0																									
6	2																									
Supplementary cancellous bone graft in femoral, tibial or humeral fractures during renailling surgery when adequate reaming and a larger nail are used[35]	<table border="1"> <tr><th>Level</th><th>Count</th></tr> <tr><td>I</td><td>0</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Count	I	0	II	0	III	2	IV	0	<table border="1"> <tr><th>Study</th><th>Count</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>2</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Study	Count	1	1	2	2	3	0	4	1	5	0	6	1
Level	Count																									
I	0																									
II	0																									
III	2																									
IV	0																									
Study	Count																									
1	1																									
2	2																									
3	0																									
4	1																									
5	0																									
6	1																									
Surgical management in thoracolumbar burst fractures with no more than minor neurologic deficit[36 37]	<table border="1"> <tr><th>Level</th><th>Count</th></tr> <tr><td>I</td><td>2</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Count	I	2	II	0	III	2	IV	0	<table border="1"> <tr><th>Study</th><th>Count</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>0</td></tr> <tr><td>3</td><td>1</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Study	Count	1	1	2	0	3	1	4	2	5	0	6	1
Level	Count																									
I	2																									
II	0																									
III	2																									
IV	0																									
Study	Count																									
1	1																									
2	0																									
3	1																									
4	2																									
5	0																									
6	1																									
Spinal fusion for thoracolumbar and lumbar burst fractures requiring surgery[38-41]	<table border="1"> <tr><th>Level</th><th>Count</th></tr> <tr><td>I</td><td>2</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>2</td></tr> <tr><td>IV</td><td>0</td></tr> </table>	Level	Count	I	2	II	0	III	2	IV	0	<table border="1"> <tr><th>Study</th><th>Count</th></tr> <tr><td>1</td><td>1</td></tr> <tr><td>2</td><td>0</td></tr> <tr><td>3</td><td>2</td></tr> <tr><td>4</td><td>1</td></tr> <tr><td>5</td><td>1</td></tr> <tr><td>6</td><td>1</td></tr> </table>	Study	Count	1	1	2	0	3	2	4	1	5	1	6	1
Level	Count																									
I	2																									
II	0																									
III	2																									
IV	0																									
Study	Count																									
1	1																									
2	0																									
3	2																									
4	1																									
5	1																									
6	1																									
Daily pin site care for fractures with an external fixation device[42 43]	<table border="1"> <tr><th>Level</th><th>Count</th></tr> <tr><td>I</td><td>2</td></tr> <tr><td>II</td><td>0</td></tr> <tr><td>III</td><td>0</td></tr> <tr><td>IV</td><td>2</td></tr> </table>	Level	Count	I	2	II	0	III	0	IV	2	<table border="1"> <tr><th>Study</th><th>Count</th></tr> <tr><td>1</td><td>4</td></tr> <tr><td>2</td><td>0</td></tr> <tr><td>3</td><td>0</td></tr> <tr><td>4</td><td>2</td></tr> <tr><td>5</td><td>0</td></tr> <tr><td>6</td><td>0</td></tr> </table>	Study	Count	1	4	2	0	3	0	4	2	5	0	6	0
Level	Count																									
I	2																									
II	0																									
III	0																									
IV	2																									
Study	Count																									
1	4																									
2	0																									
3	0																									
4	2																									
5	0																									
6	0																									

†Level of evidence of clinical practices based on study design, I, RCT or SR of RCT; II, prospective studies, quasi-randomized studies, SR of level II studies; III, case-control, case series, cross-sectional, retrospective, SR of level III studies; IV, expert consensus, narrative review, other

‡**Level of agreement of consulted experts on the value of clinical practices**, 1, clearly low-value; 2, possibly low-value; 3, controversial; 4, possibly beneficial; 5, clearly beneficial; 6, undecided

◇See eReferences for table's references

RCT, randomized controlled trial; SR, systematic review

ACCEPTED

## eReferences

## Index

References for Table 2.....	18
References for Table 3.....	31
References for Table 4.....	33
References for Table 5.....	34
References for eTable 2a.....	35
References for eTable 2b.....	36
References for eTable 2c.....	38
References for eTable 2d.....	40



## References for Table 2

1. Diercks DB, Mehrotra A, Nazarian DJ, et al. Clinical policy: critical issues in the evaluation of adult patients presenting to the emergency department with acute blunt abdominal trauma. *Annals of emergency medicine* 2011;**57**(4):387-404 doi: 10.1016/j.annemergmed.2011.01.013[published Online First: Epub Date]].
2. Holmes JF, McGahan JP, Wisner DH. Rate of intra-abdominal injury after a normal abdominal computed tomographic scan in adults with blunt trauma. *Am J Emerg Med* 2012;**30**(4):574-79
3. Chardoli M, Rezvani S, Mansouri P, et al. Is it safe to discharge blunt abdominal trauma patients with normal initial findings? *Acta Chir Belg* 2017;**117**(4):211-15
4. Awasthi S, Mao A, Wooton-Gorges SL, et al. Is Hospital Admission and Observation Required after a Normal Abdominal Computed Tomography Scan in Children with Blunt Abdominal Trauma? *Academic Emergency Medicine* 2008;**15**(10):895-99
5. Schonfeld D, Lee LK. Blunt abdominal trauma in children. *Current opinion in pediatrics* 2012;**24**(3):314-8 doi: 10.1097/MOP.0b013e328352de97[published Online First: Epub Date]].
6. Schacherer N, Miller J, Petronis K. Pediatric blunt abdominal trauma in the emergency department: evidence-based management techniques. *Pediatr Emerg Med Pract* 2014;**11**(10):1-23; quiz 23-4
7. Como JJ, Bokhari F, Chiu WC, et al. Practice Management Guidelines for Selective Nonoperative Management of Penetrating Abdominal Trauma. *J Trauma-Injury Infect Crit Care* 2010;**68**(3):721-33

8. Biffl WL, Leppaniemi A. Management guidelines for penetrating abdominal trauma. *World journal of surgery* 2015;**39**(6):1373-80 doi: 10.1007/s00268-014-2793-7[published Online First: Epub Date]].
9. Kevric J, O'Reilly GM, Gocentas RA, et al. Management of haemodynamically stable patients with penetrating abdominal stab injuries: review of practice at an Australian major trauma centre. *Eur J Trauma Emerg Surg* 2016;**42**(6):671-75
10. af Geijerstam JL, Oredsson S, Britton M. Medical outcome after immediate computed tomography or admission for observation in patients with mild head injury: randomised controlled trial. *BMJ (Clinical research ed)* 2006;**333**(7566):465 doi: 10.1136/bmj.38918.669317.4F[published Online First: Epub Date]].
11. Norlund A, Marke LA, af Geijerstam JL, et al. Immediate computed tomography or admission for observation after mild head injury: cost comparison in randomised controlled trial. *BMJ (Clinical research ed)* 2006;**333**(7566):469 doi: 10.1136/bmj.38918.659120.4F[published Online First: Epub Date]].
12. Crandon IW, Harding-Goldson HE, Benaris M, et al. Unnecessary admissions of head-injured patients at the University Hospital of the West Indies. *The West Indian medical journal* 2007;**56**(3):226-9
13. Heskestad B, Baardsen R, Helseth E, et al. Guideline Compliance in Management of Minimal, Mild, and Moderate Head Injury: High Frequency of Noncompliance Among Individual Physicians Despite Strong Guideline Support From Clinical Leaders. *J Trauma-Injury Infect Crit Care* 2008;**65**(6):1309-13
14. Schaller B, Evangelopoulos DS, Muller C, et al. Do we really need 24-h observation for patients with minimal brain injury and small intracranial bleeding? The Bernese Trauma Unit Protocol. *Emergency Medicine Journal* 2010;**27**(7):537-39
15. Richardson J, May D, Gore M, et al. Admission for Neurological Observation following Minor Head Injury is Unnecessary. *Br J Surg* 2012;**99**:39-39

16. Rendell S, Sultan L. Towards evidence-based emergency medicine: Best BETs from the Manchester Royal Infirmary. BET 3: Observation is unnecessary following a normal CT brain in warfarinised head injuries: an update. *Emergency medicine journal* : EMJ 2014;**31**(4):339-42 doi: 10.1136/emered-2014-203646.3[published Online First: Epub Date]].
17. Joseph B, Pandit V, Haider AA, et al. Improving Hospital Quality and Costs in Nonoperative Traumatic Brain Injury The Role of Acute Care Surgeons. *JAMA Surg* 2015;**150**(9):866-72
18. Howard BM, Rindler RS, Holland CM, et al. Management and Outcomes of Isolated Tentorial and Parafalcine "Smear" Subdural Hematomas at a Level-1 Trauma Center: Necessity of High Acuity Care. *Journal of neurotrauma* 2017;**34**(1):128-36 doi: 10.1089/neu.2015.4270[published Online First: Epub Date]].
19. Tavaréz MM, Atabaki SM, Teach SJ. Acute evaluation of pediatric patients with minor traumatic brain injury. *Current opinion in pediatrics* 2012;**24**(3):307-13
20. Simma B, Lutschg J, Callahan JM. Mild head injury in pediatrics: algorithms for management in the ED and in young athletes. *The American journal of emergency medicine* 2013;**31**(7):1133-8 doi: 10.1016/j.ajem.2013.04.007[published Online First: Epub Date]].
21. Beaudin M, Saint-Vil D, Ouimet A, et al. Clinical algorithm and resource use in the management of children with minor head trauma. *Journal of pediatric surgery* 2007;**42**(5):849-52 doi: 10.1016/j.jpedsurg.2006.12.038[published Online First: Epub Date]].
22. Pincus S, Weber M, Meakin A, et al. Introducing a Clinical Practice Guideline Using Early CT in the Diagnosis of Scaphoid and Other Fractures. *The western journal of emergency medicine* 2009;**10**(4):227-32
23. Patel NK, Davies N, Mirza Z, et al. Cost and clinical effectiveness of MRI in occult scaphoid fractures: a randomised controlled trial. *Emergency medicine journal* : EMJ 2013;**30**(3):202-7 doi: 10.1136/emered-2011-200676[published Online First: Epub Date]].

24. Buijze GA, Goslings JC, Rhemrev SJ, et al. Cast immobilization with and without immobilization of the thumb for nondisplaced and minimally displaced scaphoid waist fractures: a multicenter, randomized, controlled trial. *The Journal of hand surgery* 2014; 39(4). <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/617/CN-00984617/frame.html>.
25. Choosing Wisely. Clinician Lists. Secondary Clinician Lists. <http://www.choosingwisely.org/clinician-lists/>.
26. Clement CM, Stiell IG, Schull MJ, et al. Clinical features of head injury patients presenting with a Glasgow Coma Scale score of 15 and who require neurosurgical intervention. *Annals of emergency medicine* 2006;**48**(3):245-51 doi: 10.1016/j.annemergmed.2006.04.008[published Online First: Epub Date]].
27. Elgamal EA, El-Watidy SM, Jamjoom ZA, et al. Evaluation of the Canadian CT head rule for minor head trauma in a tertiary referral institution. *European Journal of Trauma* 2006;**32**(6):527-32 doi: <http://dx.doi.org/10.1007/s00068-006-5156-8>[published Online First: Epub Date]].
28. Smits M, Dippel DW, Steyerberg EW, et al. Predicting intracranial traumatic findings on computed tomography in patients with minor head injury: the CHIP prediction rule. *Annals of internal medicine* 2007;**146**(6):397-405
29. Stiell IG, Bennett C. Implementation of clinical decision rules in the emergency department. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2007;**14**(11):955-9 doi: 10.1197/j.aem.2007.06.039[published Online First: Epub Date]].
30. Turedi S, Hasanbasoglu A, Gunduz A, et al. Clinical decision instruments for CT scan in minor head trauma. *The Journal of emergency medicine* 2008;**34**(3):253-9 doi: 10.1016/j.jemermed.2007.05.055[published Online First: Epub Date]].
31. Stein SC, Fabbri A, Servadei F, et al. A Critical Comparison of Clinical Decision Instruments for Computed Tomographic Scanning in Mild Closed Traumatic Brain Injury in Adolescents and Adults. *Annals of emergency medicine* 2009;**53**(2):180-88
32. Unden J, Romner B. A new objective method for CT triage after minor head injury - serum S100B. *Scand J Clin Lab Invest* 2009;**69**(1):13-17

33. Wilson JRF, Green A. Acute Traumatic Brain Injury: A Review of Recent Advances in Imaging and Management. *Eur J Trauma Emerg Surg* 2009;**35**(2):176-85
34. Dehdari A, Sutherland J. Impact of implementing Canadian CT head rules on number of CT head studies in emergency department. *Journal of Medical Imaging and Radiation Oncology* 2010;**54**:A97 doi: <http://dx.doi.org/10.1111/%28ISSN%291754-9485>[published Online First: Epub Date]].
35. Stiell IG, Clement CM, Grimshaw JM, et al. A prospective cluster-randomized trial to implement the Canadian CT Head Rule in emergency departments. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* 2010;**182**(14):1527-32 doi: 10.1503/cmaj.091974[published Online First: Epub Date]].
36. Harnan SE, Pickering A, Pandor A, et al. Clinical decision rules for adults with minor head injury: a systematic review. *The Journal of trauma* 2011;**71**(1):245-51 doi: 10.1097/TA.0b013e31820d090f[published Online First: Epub Date]].
37. Muller B, Evangelopoulos DS, Bias K, et al. Can S-100B serum protein help to save cranial CT resources in a peripheral trauma centre? A study and consensus paper. *Emergency Medicine Journal* 2011;**28**(11):938-40
38. Pandor A, Goodacre S, Harnan S, et al. Diagnostic management strategies for adults and children with minor head injury: a systematic review and an economic evaluation. *Health Technol Assess* 2011;**15**(27):1-+
39. Peck KA, Sise CB, Shackford SR, et al. Delayed Intracranial Hemorrhage After Blunt Trauma: Are Patients on Preinjury Anticoagulants and Prescription Antiplatelet Agents at Risk? *J Trauma-Injury Infect Crit Care* 2011;**71**(6):1600-04
40. Swensen SJ, Kaplan GS, Meyer GS, et al. Controlling healthcare costs by removing waste: what American doctors can do now. *BMJ Qual Saf* 2011;**20**(6):534-37
41. Calcagnile O, Uden L, Uden J. Clinical validation of S100B use in management of mild head injury. *BMC Emerg Med* 2012;**12**:13 doi: 10.1186/1471-227x-12-13[published Online First: Epub Date]].

42. Melnick ER, Szlezak CM, Bentley SK, et al. Overuse of CT for mild traumatic brain injury. *Academic Emergency Medicine* 2012;**19**:S42 doi: <http://dx.doi.org/10.1111/j.1553-2712.2012.01332.x>[published Online First: Epub Date]].
43. Morton MJ, Korley FK. Head computed tomography use in the emergency department for mild traumatic brain injury: integrating evidence into practice for the resident physician. *Annals of emergency medicine* 2012;**60**(3):361-7 doi: 10.1016/j.annemergmed.2011.12.026[published Online First: Epub Date]].
44. Rohacek M, Albrecht M, Kleim B, et al. Reasons for ordering computed tomography scans of the head in patients with minor brain injury. *Injury-Int J Care Inj* 2012;**43**(9):1415-18
45. Albers CE, von Allmen M, Evangelopoulos DS, et al. What is the incidence of intracranial bleeding in patients with mild traumatic brain injury? A retrospective study in 3088 Canadian CT head rule patients. *BioMed research international* 2013;**2013**:453978 doi: 10.1155/2013/453978[published Online First: Epub Date]].
46. Bouida W, Marghli S, Souissi S, et al. Prediction value of the Canadian CT head rule and the New Orleans criteria for positive head CT scan and acute neurosurgical procedures in minor head trauma: a multicenter external validation study. *Annals of emergency medicine* 2013;**61**(5):521-7 doi: 10.1016/j.annemergmed.2012.07.016[published Online First: Epub Date]].
47. Chan KS, Chang E, Nassery N, et al. The state of overuse measurement: a critical review. *Medical care research and review : MCRR* 2013;**70**(5):473-96 doi: 10.1177/1077558713492202[published Online First: Epub Date]].
48. Curran JA, Brehaut J, Patey AM, et al. Understanding the Canadian adult CT head rule trial: use of the theoretical domains framework for process evaluation. *Implement Sci* 2013;**8**:25 doi: <http://dx.doi.org/10.1186/1748-5908-8-25>[published Online First: Epub Date]].
49. Haydon NB. Head injury: audit of a clinical guideline to justify head CT. *J Med Imaging Radiat Oncol* 2013;**57**(2):161-8 doi: <http://dx.doi.org/10.1111/1754-9485.12007>[published Online First: Epub Date]].

50. Huang X, Zhou JC, Pan KH, et al. Awareness and use of the Canadian computed tomography head rule for mild head injury patients among Chinese emergency physicians. *Pak J Med Sci* 2013;**29**(4):951-56
51. Melnick ER, Szlezak CM, Dziura JD, et al. Risk estimates for the canadian CT head rule in patients with minor head injury. *Academic Emergency Medicine* 2013;**1**:S239 doi: <http://dx.doi.org/10.1111/acem.12115>[published Online First: Epub Date]].
52. Gupta A, Ip IK, Raja AS, et al. Effect of clinical decision support on documented guideline adherence for head CT in emergency department patients with mild traumatic brain injury. *Journal of the American Medical Informatics Association : JAMIA* 2014;**21**(e2):e347-51 doi: 10.1136/amiajnl-2013-002536[published Online First: Epub Date]].
53. Jame SZB, Majdzadeh R, Sari AA, et al. Indications and Overuse of Computed Tomography in Minor Head Trauma. *Iran Red Crescent Med J* 2014;**16**(5)
54. Jones LA, Morley EJ, Grant WD, et al. Adherence to head computed tomography guidelines for mild traumatic brain injury. *The western journal of emergency medicine* 2014;**15**(4):459-64 doi: <http://dx.doi.org/10.5811/westjem.2014.1.19898>[published Online First: Epub Date]].
55. Kavalci C, Aksel G, Salt O, et al. Comparison of the Canadian CT head rule and the new orleans criteria in patients with minor head injury. *World journal of emergency surgery : WJES* 2014;**9**:31 doi: 10.1186/1749-7922-9-31[published Online First: Epub Date]].
56. Marin JR, Shofer FS, Chang IC, et al. Emergency department clinician adherence to clinical decision policy for head computed tomography in adult traumatic brain injury. *Academic Emergency Medicine* 2014;**1**:S190 doi: <http://dx.doi.org/10.1111/acem.12365>[published Online First: Epub Date]].
57. Parma C, Carney D, Grim R, et al. Unnecessary Head Computed Tomography Scans: A Level 1 Trauma Teaching Experience. *Am Surg* 2014;**80**(7):664-68

58. Schuur JD, Carney DP, Lyn ET, et al. A top-five list for emergency medicine: a pilot project to improve the value of emergency care. *JAMA internal medicine* 2014;**174**(4):509-15 doi: 10.1001/jamainternmed.2013.12688[published Online First: Epub Date]].
59. Altman J, Neustadtl A, Rao S, et al. Lack of utility of head ct in concussive injury in non-geriatric ED patients. *Academic Emergency Medicine* 2015;**1**:S255 doi: <http://dx.doi.org/10.1111/acem.12644>[published Online First: Epub Date]].
60. Doherty S. CT scanning for head injury in the emergency department of a rural hospital. *EMA - Emergency Medicine Australasia* 2015;**27**:36-37 doi: <http://dx.doi.org/10.1111/17426723.12416>[published Online First: Epub Date]].
61. Eiting E, Park E, Rifembark N, et al. Validation of computerized tomography head rules in correctional patients after minor head trauma. *Academic Emergency Medicine* 2015;**1**:S412 doi: <http://dx.doi.org/10.1111/acem.12644>[published Online First: Epub Date]].
62. Ip IK, Raja AS, Gupta A, et al. Impact of clinical decision support on head computed tomography use in patients with mild traumatic brain injury in the ED. *The American journal of emergency medicine* 2015;**33**(3):320-5 doi: 10.1016/j.ajem.2014.11.005[published Online First: Epub Date]].
63. Melnick ER, Keegan J, Taylor RA. Redefining Overuse to Include Costs: A Decision Analysis for Computed Tomography in Minor Head Injury. *Joint Commission journal on quality and patient safety* 2015;**41**(7):313-22
64. Melnick ER, Shafer K, Rodulfo N, et al. Understanding Overuse of Computed Tomography for Minor Head Injury in the Emergency Department: A Triangulated Qualitative Study. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2015;**22**(12):1474-83 doi: 10.1111/acem.12824[published Online First: Epub Date]].
65. Shafer K, Kochert E, Stahlman B, et al. Utility of the canadian head CT rule in reducing CT use in mvc patients presenting to the ED. *Academic emergency medicine* 2015; 22(5 suppl. 1). <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/171/CN-01080171/frame.html>.



66. Uden L, Calcagnile O, Uden J, et al. Validation of the Scandinavian guidelines for initial management of minimal, mild and moderate traumatic brain injury in adults. *BMC Med* 2015;**13**
67. Granata RT, Castillo EM, Vilke GM. Safety of deferred computed tomographic imaging of intoxicated patients presenting with possible traumatic brain injury. *Am J Emerg Med* 2016;**30**:30 doi: <http://dx.doi.org/10.1016/j.ajem.2016.09.069>[published Online First: Epub Date]].
68. Kemp A, Nickerson E, Trefan L, et al. Selecting children for head CT following head injury. *Archives of disease in childhood* 2016;**101**(10):929-34
69. Lin MP, Nguyen T, Richardson LD, et al. Emergency physician attitudes and knowledge of ACEP's choosing wisely recommendations. *Academic Emergency Medicine* 2016;**23**:S40 doi: <http://dx.doi.org/10.1111/acem.12974>[published Online First: Epub Date]].
70. Melnick ER. Big vs. Small Data and the Generalizability of the Rate of CT Overuse in Minor Head Injury. *Academic Emergency Medicine* 2016;**12**:12 doi: <http://dx.doi.org/10.1111/acem.13084>[published Online First: Epub Date]].
71. Zakhari R, Sterrett SE. Attitudes toward evidence-based clinical decision support tools to reduce exposure to ionizing radiation: The Canadian CT Head Rule. *J Am Assoc Nurs Pract* 2016;**28**(12):659-67
72. C. W, D. W, S. F, et al. The impact of alcohol intoxication on the utility and effectiveness of the Canadian CT head rule. *Journal of Medical Imaging and Radiation Oncology* 2017;**Conference**:68th Annual Scientific Meeting of the Royal Australian and New Zealand College of Radiologists, RANZCR 2017. Australia. 61 (Supplement 1) (pp 192-193) doi: <http://dx.doi.org/10.1111/1754-9485.12657>[published Online First: Epub Date]].
73. Canadian Institute for Health Information. Unnecessary Care in Canada: Technical Report. Ottawa, ON: CIHI 2017
74. Granata RT, Castillo EM, Vilke GM. Safety of deferred CT imaging of intoxicated patients presenting with possible traumatic brain injury. *Am J Emerg Med* 2017;**35**(1):51-54

75. Klang E, Beytelman A, Greenberg D, et al. Overuse of Head CT Examinations for the Investigation of Minor Head Trauma: Analysis of Contributing Factors. *Journal of the American College of Radiology : JACR* 2017;**14**(2):171-76 doi: 10.1016/j.jacr.2016.08.032[published Online First: Epub Date]].
76. Lin S, Ashkenazi M, Karawan M, et al. Management of Ankylotic Root Resorption Following Dental Trauma: A Short Review and Proposal of a Treatment Protocol. *Oral Health Prev Dent* 2017;**15**(5):467-74 doi: 10.3290/j.ohpd.a38736[published Online First: Epub Date]].
77. Luoto T, Iverson G, Pauniah SL, et al. The Scandinavian guidelines for initial management of minimal, mild and moderate traumatic brain injury in adults: A prospective validation study. *Brain injury* 2017;**31** (6-7):876-77 doi: <http://dx.doi.org/10.1080/02699052.2017.1312145>[published Online First: Epub Date]].
78. Melnick ER. Big Versus Small Data and the Generalizability of the Rate of Computed Tomography Overuse in Minor Head Injury. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2017;**24**(3):391-92 doi: 10.1111/acem.13084[published Online First: Epub Date]].
79. Melnick ER, Hess EP, Guo G, et al. Patient-Centered Decision Support: Formative Usability Evaluation of Integrated Clinical Decision Support With a Patient Decision Aid for Minor Head Injury in the Emergency Department. *J Med Internet Res* 2017;**19**(5):e174 doi: <https://dx.doi.org/10.2196/jmir.7846>[published Online First: Epub Date]].
80. National Institute for Health and Care Excellence (NICE). Head injury: assessment and early management (CG176). Secondary Head injury: assessment and early management (CG176) 2017. <https://www.nice.org.uk/guidance/qs166>.
81. R. L, A. J, I. F, et al. Mindless scanning: Does every trauma patient need a head CT? *Journal of the American College of Surgeons* 2017;**Conference**:2017 Clinical Congress of the 3rd Owen H Wangesteen Scientific Forum. United States. 225 (4 Supplement 1) (pp S57-S58)

82. Raja AS, Venkatesh AK, Mick N, et al. "Choosing Wisely" Imaging Recommendations: Initial Implementation in New England Emergency Departments. *The western journal of emergency medicine* 2017;**18**(3):454-58 doi: 10.5811/westjem.2017.1.32677[published Online First: Epub Date]].
83. Sharp AL, Nagaraj G, Rippberger EJ, et al. Computed Tomography Use for Adults With Head Injury: Describing Likely Avoidable Emergency Department Imaging Based on the Canadian CT Head Rule. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2017;**24**(1):22-30 doi: 10.1111/acem.13061[published Online First: Epub Date]].
84. Singh N, Hess E, Guo G, et al. Tablet-Based Patient-Centered Decision Support for Minor Head Injury in the Emergency Department: Pilot Study. *JMIR mHealth uHealth* 2017;**5**(9):e144 doi: 10.2196/mhealth.8732[published Online First: Epub Date]].
85. Smith TM, Milzman D. Lack of utility of head CT in concussive head injury amongst non-geriatric patients. *Acta radiologica Conference: 62nd nordic congress of radiology and 23rd nordic congress of radiography Iceland* 2017; 58(1 Supplement 1). <http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/052/CN-01397052/frame.html>.
86. Webster NJ, Moore N, Stewart F. Reducing Unnecessary Head Computed Tomography in Mild Traumatic Brain Injury. *Adv Emerg Nurs J* 2017;**39**(4):300-08 doi: 10.1097/TME.000000000000169[published Online First: Epub Date]].
87. Cellina M, Panzeri M, Floridi C, et al. Overuse of computed tomography for minor head injury in young patients: an analysis of promoting factors. *Radiol Med* 2018;**07**:07 doi: <https://dx.doi.org/10.1007/s11547-018-0871-x>[published Online First: Epub Date]].
88. Tung M, Sharma R, Hinson JS, et al. Factors associated with imaging overuse in the emergency department: A systematic review. *Am J Emerg Med* 2018;**36**(2):301-09 doi: <https://dx.doi.org/10.1016/j.ajem.2017.10.049>[published Online First: Epub Date]].
89. Dunning J, Daly JP, Lomas JP, et al. Derivation of the children's head injury algorithm for the prediction of important clinical events decision rule for head injury in children. *Archives of disease in childhood* 2006;**91**(11):885-91 doi: 10.1136/adc.2005.083980[published Online First: Epub Date]].

90. Oman JA, Cooper RJ, Holmes JF, et al. Performance of a decision rule to predict need for computed tomography among children with blunt head trauma. *Pediatrics* 2006;**117**(2):e238-46 doi: 10.1542/peds.2005-1063[published Online First: Epub Date]].
91. Atabaki SM, Stiell IG, Bazarian JJ, et al. A clinical decision rule for cranial computed tomography in minor pediatric head trauma. *Arch Pediatr Adolesc Med* 2008;**162**(5):439-45
92. Kuppermann N, Holmes JF, Dayan PS, et al. Identification of children at very low risk of clinically-important brain injuries after head trauma: a prospective cohort study. *Lancet (London, England)* 2009;**374**(9696):1160-70 doi: 10.1016/s0140-6736(09)61558-0[published Online First: Epub Date]].
93. Osmond MH, Klassen TP, Wells GA, et al. CATCH: a clinical decision rule for the use of computed tomography in children with minor head injury. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* 2010;**182**(4):341-8 doi: 10.1503/cmaj.091421[published Online First: Epub Date]].
94. Goldberg J, McClaine RJ, Cook B, et al. Use of a mild traumatic brain injury guideline to reduce inpatient hospital imaging and charges. *Journal of pediatric surgery* 2011;**46**(9):1777-83 doi: 10.1016/j.jpedsurg.2011.02.052[published Online First: Epub Date]].
95. Schachar JL, Zampolin RL, Miller TS, et al. External validation of the New Orleans Criteria (NOC), the Canadian CT Head Rule (CCHR) and the National Emergency X-Radiography Utilization Study II (NEXUS II) for CT scanning in pediatric patients with minor head injury in a non-trauma center. *Pediatr Radiol* 2011;**41**(8):971-9 doi: 10.1007/s00247-011-2032-4[published Online First: Epub Date]].
96. Bressan S, Romanato S, Mion T, et al. Implementation of adapted PECARN decision rule for children with minor head injury in the pediatric emergency department. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2012;**19**(7):801-7 doi: 10.1111/j.1553-2712.2012.01384.x[published Online First: Epub Date]].
97. Gerdung C, Dowling S, Lang E. Review of the CATCH study: a clinical decision rule for the use of computed tomography in children with minor head injury. *Cjem* 2012;**14**(4):243-7

98. Klement W, Wilk S, Michalowski W, et al. Predicting the need for CT imaging in children with minor head injury using an ensemble of Naive Bayes classifiers. *Artificial intelligence in medicine* 2012;**54**(3):163-70 doi: 10.1016/j.artmed.2011.11.005[published Online First: Epub Date]].
99. Natale JE, Joseph JG, Rogers AJ, et al. Cranial computed tomography use among children with minor blunt head trauma: association with race/ethnicity.[Erratum appears in *Arch Pediatr Adolesc Med*. 2014 Jun;168(6):586]. *Arch Pediatr Adolesc Med* 2012;**166**(8):732-7 doi: <http://dx.doi.org/10.1001/archpediatrics.2012.307>[published Online First: Epub Date]].
100. Osmond MH, Correll R, Klassen T, et al. Multicentre prospective validation of the Canadian assessment of tomography for childhood head injury (CATCH) rule. *Canadian Journal of Emergency Medicine* 2012;**14**:S6-S7
101. Hennelly KE, Mannix R, Nigrovic LE, et al. Pediatric traumatic brain injury and radiation risks: a clinical decision analysis. *The Journal of pediatrics* 2013;**162**(2):392-7 doi: 10.1016/j.jpeds.2012.07.018[published Online First: Epub Date]].
102. Linscott LL, Kessler MM, Kitchin DR, et al. CT for Pediatric, Acute, Minor Head Trauma: Clinician Conformity to Published Guidelines. *Am J Neuroradiol* 2013;**34**(6):1252-56
103. Parri N, Crosby BJ, Glass C, et al. Ability of Emergency Ultrasonography to Detect Pediatric Skull Fractures: A Prospective, Observational Study. *J Emerg Med* 2013;**44**(1):135-41
104. Maskiell C, Stapleton T, Kreltzheim M. An audit of CT head imaging in paediatric patients presenting to the Department of Emergency Department (DEM) at Nambour General Hospital (NGH) & Caloundra Hospital. *Journal of Medical Imaging and Radiation Oncology* 2014;**58**:259 doi: <http://dx.doi.org/10.1111/1754-9485.12223>[published Online First: Epub Date]].
105. Mihindu E, Bhullar I, Tepas J, et al. Computed Tomography of the Head in Children with Mild Traumatic Brain Injury. *Am Surg* 2014;**80**(9):841-43

106. Schonfeld D, Bressan S, Da Dalt L, et al. Pediatric Emergency Care Applied Research Network head injury clinical prediction rules are reliable in practice. *Archives of disease in childhood* 2014;**99**(5):427-31 doi: 10.1136/archdischild-2013-305004[published Online First: Epub Date].
107. Stanley RM, Hoyle JD, Dayan PS, et al. Emergency Department Practice Variation in Computed Tomography Use for Children with Minor Blunt Head Trauma. *J Pediatr* 2014;**165**(6):1201-+
108. Engineer R, Ferguson S, Podolsky S, et al. Reduced head computed tomography utilization for pediatric mild head injury after implementation of decision support tools and shared decisionmaking. *Annals of emergency medicine* 2015;**1**):S4
109. Pierce D, Mangona KL, Bisset G, et al. Computed Tomography in the Evaluation of Pediatric Trauma. *Clinical Pediatric Emergency Medicine* 2015;**16**(4):220-29 doi: <http://dx.doi.org/10.1016/j.cpem.2015.11.001>[published Online First: Epub Date].
110. Atabaki SM, Hoyle JD, Jr., Schunk JE, et al. Comparison of Prediction Rules and Clinician Suspicion for Identifying Children With Clinically Important Brain Injuries After Blunt Head Trauma. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2016;**23**(5):566-75 doi: 10.1111/acem.12923[published Online First: Epub Date].
111. Bharadwaj S, Rucker J. Minor head injury: limiting patient exposure to ionizing radiation, risk stratification, and concussion management. *Current opinion in pediatrics* 2016;**28**(1):121-31
112. Eisenmenger LB, Anzai Y. Computed Tomography in Pediatric Traumatic Brain Injury: Who Needs It and How Is It Scored? *Journal of Pediatric Neuroradiology* 2016;**5**(1):13-19 doi: <http://dx.doi.org/10.1055/s-0036-1584243>[published Online First: Epub Date].
113. Harwayne-Gidansky I, Olvet D, Chandran L, et al. Concordance of clinical criteria between resident and attending using the PECARN tool. *Critical Care Medicine* 2016;**44** (12 Supplement 1):182 doi: <http://dx.doi.org/10.1097/01.ccm.0000509099.72473.b6>[published Online First: Epub Date].

114. Hurley T, Curran P. G335(P) The use of CT brain in children with head injuries in a general hospital compared to PECARN guidelines. Archives of disease in childhood 2016;**101**(Suppl 1):A195
115. Bozan O, Aksel G, Kahraman HA, et al. Comparison of PECARN and CATCH clinical decision rules in children with minor blunt head trauma. European journal of trauma and emergency surgery : official publication of the European Trauma Society 2017 doi: 10.1007/s00068-017-0865-8[published Online First: Epub Date]].
116. Gokharman FD, Aydin S, Fatihoglu E, et al. Pediatric Emergency Care Applied Research Network head injury prediction rules: on the basis of cost and effectiveness. Turk J Med Sci 2017;**47**(6):1770-77 doi: 10.3906/sag-1703-206[published Online First: Epub Date]].
117. Ide K, Uematsu S, Tetsuhara K, et al. External Validation of the PECARN Head Trauma Prediction Rules in Japan. Academic emergency medicine : official journal of the Society for Academic Emergency Medicine 2017;**24**(3):308-14 doi: 10.1111/acem.13129[published Online First: Epub Date]].
118. Jennings RM, Burtner JJ, Pellicer JF, et al. Reducing Head CT Use for Children With Head Injuries in a Community Emergency Department. Pediatrics 2017;**139**(4) doi: 10.1542/peds.2016-1349[published Online First: Epub Date]].
119. K. M, N. K, A. W, et al. Adherence to the pediatric emergency care applied research network head CT rule: 2013 to 2015. Annals of emergency medicine 2017;**Conference**:American College of Emergency Physicians, ACEP 2017 Research Forum. United States. 70 (4 Supplement 1) (pp S88)
120. Lal D. Vomiting as an indication for imaging in paediatric traumatic brain injury; questionable significance! Developmental Medicine & Child Neurology 2017;**59** (Suppl. 1):pp. 44-45
121. Nakhjavan-Shahraki B, Yousefifard M, Oraili A, et al. Prediction of Clinically Important Traumatic Brain Injury in Pediatric Minor Head Trauma; proposing Pediatric Traumatic Brain Injury (PTBI) Prognostic Rule. Int J Pediatr-Masshad 2017;**5**(1):4127-35

122. Zulfiqar M, Kim S, Lai JP, et al. The role of computed tomography in following up pediatric skull fractures. *Am J Surg* 2017;**214**(3):483-88  
doi: 10.1016/j.amjsurg.2016.07.020[published Online First: Epub Date]].
123. Halaweish I, Riebe-Rodgers J, Randall A, et al. Compliance with evidence-based guidelines for computed tomography of children with head and abdominal trauma. *Journal of pediatric surgery* 2018;**53**(4):748-51 doi: 10.1016/j.jpedsurg.2017.07.008[published Online First: Epub Date]].
124. Hiscock H, Neely RJ, Warren H, et al. Reducing Unnecessary Imaging and Pathology Tests: A Systematic Review. *Pediatrics* 2018;**141**(2)
125. Durham SR, Liu KC, Selden NR. Utility of serial computed tomography imaging in pediatric patients with head trauma. *Journal of neurosurgery* 2006;**105**(5 Suppl):365-9 doi: 10.3171/ped.2006.105.5.365[published Online First: Epub Date]].
126. da Silva PS, Reis ME, Aguiar VE. Value of repeat cranial computed tomography in pediatric patients sustaining moderate to severe traumatic brain injury. *The Journal of trauma* 2008;**65**(6):1293-7 doi: 10.1097/TA.0b013e318156866c[published Online First: Epub Date]].
127. Dawson E, Montgomery C, Koogler T, et al. Is repeat head CT necessary in mild head injury and normal neurologic exam? *Critical Care Medicine* 2010;**38**:A55 doi: <http://dx.doi.org/10.1097/01.ccm.0000390903.16849.8c>[published Online First: Epub Date]].
128. Bae KS, Park SY, Kim YJ, et al. Value of repeat brain CT in head injured children visiting ED without traumatic brain injury in initial CT. *Journal of neurotrauma* 2011;**28**:A-1-A-134 (June):A-68 (P181)
129. Dawson EC, Montgomery CP, Frim D, et al. Is Repeat Head Computed Tomography Necessary in Children Admitted with Mild Head Injury and Normal Neurological Exam? *Pediatr Neurosurg* 2012;**48**(4):221-24
130. Aziz H, Rhee P, Pandit V, et al. Mild and moderate pediatric traumatic brain injury: replace routine repeat head computed tomography with neurologic examination. *The journal of trauma and acute care surgery* 2013;**75**(4):550-4 doi: 10.1097/TA.0b013e3182a53a77[published Online First: Epub Date]].



131. Howe J, Fitzpatrick CM, Lakam DR, et al. Routine repeat brain computed tomography in all children with mild traumatic brain injury may result in unnecessary radiation exposure. *J Trauma Acute Care Surg* 2014;**76**(2):292-96
132. Hentzen AS, Helmer SD, Nold RJ, et al. Necessity of repeat head computed tomography after isolated skull fracture in the pediatric population. *Am J Surg* 2015;**210**(2):322-5 doi: 10.1016/j.amjsurg.2014.11.011[published Online First: Epub Date]].
133. Zulfiqar M, Kim S, Lai J-P, et al. The role of computed tomography in following up pediatric skull fractures. *Am J Surg* 2016;**16**:16 doi: <http://dx.doi.org/10.1016/j.amjsurg.2016.07.020>[published Online First: Epub Date]].
134. Ore CLD, Rennert R, Schupper AJ, et al. 320 The Identification of a Subgroup of Children with Traumatic Subarachnoid Hemorrhage at Low Risk of Neuroworsening. *Neurosurgery* 2017;**64**(CN\_suppl\_1):269-69 doi: 10.1093/neuros/nyx417.320[published Online First: Epub Date]].
135. Stiell IG, Clement CM, Grimshaw J, et al. Implementation of the Canadian C-Spine Rule: prospective 12 centre cluster randomised trial. *BMJ* (Clinical research ed) 2009;**339**:b4146 doi: 10.1136/bmj.b4146[published Online First: Epub Date]].
136. Kokabi N, Raper DMS, Xing M, et al. Efficacy of goergen's diagnostic algorithm in detecting cervical spine injuries in elderly victims of blunt trauma. *Emerg* 2010;**17** (6):545-46 doi: <http://dx.doi.org/10.1007/s10140-010-0904-9>[published Online First: Epub Date]].
137. Griffith B, Bolton C, Goyal N, et al. Screening cervical spine CT in a level I trauma center: overutilization? *AJR American journal of roentgenology* 2011;**197**(2):463-7 doi: 10.2214/ajr.10.5731[published Online First: Epub Date]].
138. Kokabi N, Raper DM, Xing M, et al. Application of imaging guidelines in patients with suspected cervical spine trauma: retrospective analysis and literature review. *Emerg Radiol* 2011;**18**(1):31-8 doi: 10.1007/s10140-010-0901-z[published Online First: Epub Date]].
139. Michaleff ZA, Maher CG, Verhagen AP, et al. Accuracy of the Canadian C-spine rule and NEXUS to screen for clinically important cervical spine injury in patients following blunt trauma: a systematic review. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* 2012;**184**(16):E867-76 doi: 10.1503/cmaj.120675[published Online First: Epub Date]].

140. Nicholson A, Krishnan A, Holder C. Overutilization of cervical spine CT after minor trauma. *American Journal of Roentgenology* Conference 2012;**198**(5 SUPPL. 1)
141. Sheikh K, Belfi LM, Sharma R, et al. Evaluation of acute cervical spine imaging based on ACR Appropriateness Criteria(R). *Emerg Radiol* 2012;**19**(1):11-7 doi: 10.1007/s10140-011-0994-z[published Online First: Epub Date]].
142. Griffith B, Kelly M, Vallee P, et al. Screening Cervical Spine CT in the Emergency Department, Phase 2: A Prospective Assessment of Use. *Am J Neuroradiol* 2013;**34**(4):899-903
143. Griffith B, Vallee P, Krupp S, et al. Screening cervical spine CT in the emergency department, phase 3: increasing effectiveness of imaging. *Journal of the American College of Radiology : JACR* 2014;**11**(2):139-44 doi: 10.1016/j.jacr.2013.05.026[published Online First: Epub Date]].
144. Morrison J, Jeanmonod R. Imaging in the NEXUS-negative patient: when we break the rule. *The American journal of emergency medicine* 2014;**32**(1):67-70 doi: 10.1016/j.ajem.2013.08.062[published Online First: Epub Date]].
145. Hikino K, Yamamoto LG. The benefit of neck computed tomography compared with its harm (risk of cancer). *J Trauma Acute Care Surg* 2015;**78**(1):126-31
146. Velan GM, Goergen SK, Grimm J, et al. Impact of Interactive e-Learning Modules on Appropriateness of Imaging Referrals: A Multicenter, Randomized, Crossover Study. *J Am Coll Radiol* 2015;**12**(11):1207-14
147. Benayoun MD, Allen JW, Lovasik BP, et al. Utility of computed tomographic imaging of the cervical spine in trauma evaluation of ground-level fall. *J Trauma Acute Care Surg* 2016;**81**(2):339-44
148. Uriell ML, Allen JW, Lovasik BP, et al. Yield of computed tomography of the cervical spine in cases of simple assault. *Injury* 2017;**48**(1):133-36 doi: 10.1016/j.injury.2016.10.031[published Online First: Epub Date]].

149. S. D, C. L, S.W. K, et al. Effectiveness of Implementing Evidence-based Interventions to Reduce C-spine Image Ordering in the Emergency Department: A Systematic Review. *Academic Emergency Medicine* 2018 doi: <http://dx.doi.org/10.1111/acem.13364>[published Online First: Epub Date]].
150. Daffner RH, Hackney DB. ACR Appropriateness Criteria on suspected spine trauma. *Journal of the American College of Radiology : JACR* 2007;**4**(11):762-75 doi: 10.1016/j.jacr.2007.08.006[published Online First: Epub Date]].
151. Khanna G, El-Khoury GY. Imaging of cervical spine injuries of childhood. *Skeletal radiology* 2007;**36**(6):477-94 doi: 10.1007/s00256-006-0223-0[published Online First: Epub Date]].
152. Egloff AM, Kadom N, Vezina G, et al. Pediatric cervical spine trauma imaging: a practical approach. *Pediatr Radiol* 2009;**39**(5):447-56 doi: 10.1007/s00247-008-1043-2[published Online First: Epub Date]].
153. Anderson RB, Hunt KJ, McCormick JJ. Management of Common Sports-related Injuries About the Foot and Ankle. *J Am Acad Orthop Surg* 2010;**18**(9):546-56
154. Kreykes NS, Letton RW, Jr. Current issues in the diagnosis of pediatric cervical spine injury. *Seminars in pediatric surgery* 2010;**19**(4):257-64 doi: 10.1053/j.sempedsurg.2010.06.002[published Online First: Epub Date]].
155. Booth TN. Cervical spine evaluation in pediatric trauma. *AJR American journal of roentgenology* 2012;**198**(5):W417-25 doi: 10.2214/ajr.11.8150[published Online First: Epub Date]].
156. Sun R, Skeete D, Wetjen K, et al. A pediatric cervical spine clearance protocol to reduce radiation exposure in children. *The Journal of surgical research* 2013;**183**(1):341-6 doi: 10.1016/j.jss.2012.12.048[published Online First: Epub Date]].
157. Hale DF, Fitzpatrick CM, Doski JJ, et al. Absence of clinical findings reliably excludes unstable cervical spine injuries in children 5 years or younger. *J Trauma Acute Care Surg* 2015;**78**(5):943-48

158. McMahon PM, Alwood SM, Zeretzke-Bien C, et al. Protocol to Clear Cervical Spine Injuries in Pediatric Trauma Patients. *Radiol Manage* 2015;**37**(5):42-8
159. Pepin LC, Bressler S, Brenner C, et al. Application of the national emergency x radiography utilization study criteria and the canadian c-spine rule among children aged 8 to 17 years in the emergency department: A retrospective review. *Annals of emergency medicine* 2015;**1**):S144
160. Burlew CC, Biffi WL, Moore EE, et al. Blunt cerebrovascular injuries: redefining screening criteria in the era of noninvasive diagnosis. *The journal of trauma and acute care surgery* 2012;**72**(2):330-5; discussion 36-7, quiz 539 doi: 10.1097/TA.0b013e31823de8a0[published Online First: Epub Date]].
161. Ravindra VM, Bollo RJ, Sivakumar W, et al. Predicting Blunt Cerebrovascular Injury in Pediatric Trauma: Validation of the "Utah Score". *Journal of neurotrauma* 2017;**34**(2):391-99 doi: 10.1089/neu.2016.4415[published Online First: Epub Date]].
162. Ibraheem K, Khan M, Rhee P, et al. "No zone" approach in penetrating neck trauma reduces unnecessary computed tomography angiography and negative explorations. *Journal of Surgical Research* 2018;**221**:113-20 doi: <https://dx.doi.org/10.1016/j.jss.2017.08.033>[published Online First: Epub Date]].
163. Traub M, Stevenson M, McEvoy S, et al. The use of chest computed tomography versus chest X-ray in patients with major blunt trauma. *Injury* 2007;**38**(1):43-7 doi: 10.1016/j.injury.2006.07.006[published Online First: Epub Date]].
164. Barrios C, Malinoski D, Dolich M, et al. Utility of thoracic computed tomography after blunt trauma: when is chest radiograph enough? *The American surgeon* 2009;**75**(10):966-9
165. Barrios C, Jr., Pham J, Malinoski D, et al. Ability of a chest X-ray and an abdominal computed tomography scan to identify traumatic thoracic injury. *Am J Surg* 2010;**200**(6):741-4; discussion 44-5 doi: 10.1016/j.amjsurg.2010.08.004[published Online First: Epub Date]].

166. Brink M, Deunk J, Dekker HM, et al. Criteria for the selective use of chest computed tomography in blunt trauma patients. *European radiology* 2010;**20**(4):818-28 doi: 10.1007/s00330-009-1608-y[published Online First: Epub Date]].
167. Mollberg NM, Wise SR, De Hoyos AL, et al. Chest computed tomography for penetrating thoracic trauma after normal screening chest roentgenogram. *The Annals of thoracic surgery* 2012;**93**(6):1830-5 doi: 10.1016/j.athoracsur.2012.02.095[published Online First: Epub Date]].
168. Payrastre J, Upadhye S, Worster A, et al. The SCRAP Rule: The derivation and internal validation of a clinical decision rule for computed tomography of the chest in blunt thoracic trauma. *CJEM, Can* 2012;**14**(6):344-53
169. Manka M, Moscati R, Caiola C, et al. Can sentinel clinical and CXR findings predict the likelihood of an abnormal chest CT requiring intervention following blunt trauma? *Academic Emergency Medicine* 2013;**1**:S59 doi: <http://dx.doi.org/10.1111/acem.12115>[published Online First: Epub Date]].
170. Rodriguez RM, Anglin D, Langdorf MI, et al. Validation of a Decision Instrument for Selective Chest Imaging in Blunt Trauma. *JAMA Surg* 2013;**148**(10):940-46
171. Rodriguez RM, Langdorf MI, Nishijima D, et al. Derivation and Validation of Two Decision Instruments for Selective Chest CT in Blunt Trauma: A Multicenter Prospective Observational Study (NEXUS Chest CT). *PLoS Med* 2015;**12**(10)
172. Kanani AN, Hartshorn S. NICE clinical guideline NG39: Major trauma: assessment and initial management. *Archives of disease in childhood Education and practice edition* 2017;**102**(1):20-23 doi: 10.1136/archdischild-2016-310869[published Online First: Epub Date]].
173. Nural MS, Balcik C, Elmali M, et al. Importance of free intra-peritoneal fluid at ultrasound in children with blunt abdominal trauma. *Pediatric Radiology* 2009;**39**:S568-S69 doi: <http://dx.doi.org/10.1007/s00247-009-1241-6>[published Online First: Epub Date]].
174. Chatoorgoon K, Brown RL, Garcia VF, et al. Role of Computed Tomography and Clinical Findings in Pediatric Blunt Intestinal Injury A Multicenter Study. *Pediatric emergency care* 2012;**28**(12):1338-42

175. Streck CJ, Jr., Jewett BM, Wahlquist AH, et al. Evaluation for intra-abdominal injury in children after blunt torso trauma: can we reduce unnecessary abdominal computed tomography by utilizing a clinical prediction model? *The journal of trauma and acute care surgery* 2012;**73**(2):371-6; discussion 76 doi: 10.1097/TA.0b013e31825840ab[published Online First: Epub Date]].
176. Bregstein JS, Lubell TR, Ruscica AM, et al. Nuking the radiation: minimizing radiation exposure in the evaluation of pediatric blunt trauma. *Current opinion in pediatrics* 2014;**26**(3):272-78
177. de Jong WJ, Stoepker L, Nellensteijn DR, et al. External validation of the Blunt Abdominal Trauma in Children (BATiC) score: ruling out significant abdominal injury in children. *The journal of trauma and acute care surgery* 2014;**76**(5):1282-7 doi: 10.1097/ta.000000000000175[published Online First: Epub Date]].
178. Amerstorfer EE, Haberlik A, Riccabona M. Imaging assessment of renal injuries in children and adolescents: CT or ultrasound? *Journal of pediatric surgery* 2015;**50**(3):448-55 doi: 10.1016/j.jpedsurg.2014.07.006[published Online First: Epub Date]].
179. Fallon SC, Delemos D, Akinkuotu A, et al. The use of an institutional pediatric abdominal trauma protocol improves resource use. *J Trauma Acute Care Surg* 2016;**80**(1):57-63
180. Brayboy T, Hogg MM, Dudin P, et al. Multi-center retrospective validation of the pediatric emergency care applied research network blunt abdominal trauma prediction rule. *Academic emergency medicine Conference: 2017 annual meeting of the society for academic emergency medicine, SAEM 2017 United states* 2017; 24. <http://cochranelibrary-wiley.com/o/cochrane/clcentral/articles/321/CN-01376321/frame.html>.
181. Streck CJ, Vogel AM, Zhang J, et al. Identifying Children at Very Low Risk for Blunt Intra-Abdominal Injury in Whom CT of the Abdomen Can Be Avoided Safely. *J Am Coll Surg* 2017;**224**(4):449-58.e3 doi: 10.1016/j.jamcollsurg.2016.12.041[published Online First: Epub Date]].

182. Zagory JA, Dossa A, Golden J, et al. Re-evaluation of liver transaminase cutoff for CT after pediatric blunt abdominal trauma. *Pediatric surgery international* 2017;**33**(3):311-16 doi: 10.1007/s00383-016-4026-7[published Online First: Epub Date].
183. Haasz M, Simone LA, Wales PW, et al. Which pediatric blunt trauma patients do not require pelvic imaging? *The journal of trauma and acute care surgery* 2015;**79**(5):828-32 doi: 10.1097/ta.0000000000000848[published Online First: Epub Date].
184. Huber-Wagner S, Lefering R, Qvick LM, et al. Effect of whole-body CT during trauma resuscitation on survival: a retrospective, multicentre study. *Lancet (London, England)* 2009;**373**(9673):1455-61 doi: 10.1016/s0140-6736(09)60232-4[published Online First: Epub Date].
185. Burnside N, McManus K. Blunt thoracic trauma. *Surgery (United Kingdom)* 2014;**32**(5):254-60 doi: <http://dx.doi.org/10.1016/j.mpsur.2014.02.007>[published Online First: Epub Date].
186. Davies RM, Scrimshire AB, Sweetman L, et al. A decision tool for whole-body CT in major trauma that safely reduces unnecessary scanning and associated radiation risks: An initial exploratory analysis. *Injury-Int J Care Inj* 2016;**47**(1):43-49
187. N. H, C. B. The increasing practice of whole body CT scanning in blunt trauma patients. *Annals of emergency medicine* 2017;**Conference**:American College of Emergency Physicians, ACEP 2017 Research Forum. United States. 70 (4 Supplement 1) (pp S118)
188. Chwals WJ, Robinson AV, Sivit CJ, et al. Computed tomography before transfer to a level I pediatric trauma center risks duplication with associated increased radiation exposure. *Journal of pediatric surgery* 2008;**43**(12):2268-72 doi: <http://dx.doi.org/10.1016/j.jpedsurg.2008.08.061>[published Online First: Epub Date].
189. Liepert AE, Cochran A. CT utilization in transferred trauma patients. *The Journal of surgical research* 2011;**170**(2):309-13 doi: 10.1016/j.jss.2011.06.018[published Online First: Epub Date].

190. Hartin CW, Jr., Jordan JM, Gemme S, et al. Computed tomography scanning in pediatric trauma: opportunities for performance improvement and radiation safety. *The Journal of surgical research* 2013;**180**(2):226-31 doi: 10.1016/j.jss.2012.04.020[published Online First: Epub Date]].
191. Benedict LA, Paulus JK, Rideout L, et al. Are CT scans obtained at referring institutions justified prior to transfer to a pediatric trauma center? *Journal of pediatric surgery* 2014;**49**(1):184-7; discussion 87-8 doi: <http://dx.doi.org/10.1016/j.jpedsurg.2013.09.056>[published Online First: Epub Date]].
192. Cook SH, Fielding JR, Phillips JD. Repeat abdominal computed tomography scans after pediatric blunt abdominal trauma: missed injuries, extra costs, and unnecessary radiation exposure. *Journal of pediatric surgery* 2010;**45**(10):2019-24
193. Emick DM, Carey TS, Charles AG, et al. Repeat imaging in trauma transfers: a retrospective analysis of computed tomography scans repeated upon arrival to a Level I trauma center. *The journal of trauma and acute care surgery* 2012;**72**(5):1255-62 doi: 10.1097/TA.0b013e3182452b6f[published Online First: Epub Date]].
194. Moore HB, Loomis SB, Destigter KK, et al. Airway, breathing, computed tomographic scanning: duplicate computed tomographic imaging after transfer to trauma center. *The journal of trauma and acute care surgery* 2013;**74**(3):813-7 doi: <http://dx.doi.org/10.1097/TA.0b013e3182789399>[published Online First: Epub Date]].
195. Holmes J, Siglock B, Corwin M, et al. Rate and reasons for repeat ct scanning in transferred trauma patients. *Academic Emergency Medicine* 2015;**1**:S296 doi: <http://dx.doi.org/10.1111/acem.12644>[published Online First: Epub Date]].
196. Banerjee A, Zosa BM, Allen D, et al. Implementation of an image sharing system significantly reduced repeat computed tomographic imaging in a regional trauma system. *The journal of trauma and acute care surgery* 2016;**80**(1):51-4; discussion 54-6 doi: <http://dx.doi.org/10.1097/TA.0000000000000866>[published Online First: Epub Date]].



197. Guttman A, Razzaq A, Lindsay P, et al. Development of measures of the quality of emergency department care for children using a structured panel process. *Pediatrics* 2006;**118**(1):114-23
198. Schab M, Jonczyk-Potoczna K, Jazdzewska A, et al. Diagnostic imaging in the admission room in the minor head injury in children aged 0-5 years  
Diagnostyka obrazowa w izbie przyje{ogonek}c w lekkich urazach gLowy u dzieci w wieku 0-5 lat. *Family Medicine and Primary Care Review* 2011;**13**(4):750-53
199. Gravel J, Gouin S, Chalut D, et al. Derivation and validation of a clinical decision rule to identify young children with skull fracture following isolated head trauma. *CMAJ Canadian Medical Association Journal* 2015;**187**(16):1202-8 doi: <http://dx.doi.org/10.1503/cmaj.150540>[published Online First: Epub Date].
200. Lopes JA, Frankel HL, Bokhari SJ, et al. The trauma bay chest radiograph in stable blunt-trauma patients: do we really need it? *The American surgeon* 2006;**72**(1):31-4
201. Wisbach GG, Sise MJ, Sack DI, et al. What is the role of chest X-ray in the initial assessment of stable trauma patients? *The Journal of trauma* 2007;**62**(1):74-8; discussion 78-9 doi: 10.1097/01.ta.0000251422.53368.a3[published Online First: Epub Date].
202. Rodriguez RM, Hendey GW, Mower W, et al. Derivation of a decision instrument for selective chest radiography in blunt trauma. *The Journal of trauma* 2011;**71**(3):549-53 doi: 10.1097/TA.0b013e3181f2ac9d[published Online First: Epub Date].
203. Paydar S, Johari HG, Ghaffarpassand F, et al. The role of routine chest radiography in initial evaluation of stable blunt trauma patients. *Am J Emerg Med* 2012;**30**(1):1-4
204. Ziegler K, Feeney JM, Desai C, et al. Retrospective review of the use and costs of routine chest X-rays in a trauma setting (Provisional abstract). *Journal of Trauma Management and Outcomes* 2013; 7(1). <http://onlinelibrary.wiley.com/o/cochrane/cleed/articles/NHSEED-22013022089/frame.html>.

205. Paydar S, Ahmadi A, Dalfardi B, et al. Clinical and economic effects of selective radiological evaluation of high-energy trauma patients: a prospective experience of a level 1 busy trauma centre. *Emergency Medicine Journal* 2015;**32**(7):535-38
206. Turkmen S, Cansu A, Karaca Y, et al. Determination of radiography requirement in wrist trauma. *Am J Emerg Med* 2015;**33**(9):1269-72
207. Slaar A, Walenkamp MM, Bentohami A, et al. A clinical decision rule for the use of plain radiography in children after acute wrist injury: development and external validation of the Amsterdam Pediatric Wrist Rules. *Pediatr Radiol* 2016;**46**(1):50-60 doi: 10.1007/s00247-015-3436-3[published Online First: Epub Date]].
208. Mulders MAM, Walenkamp MMJ, Dubois BFH, et al. External validation of clinical decision rules for children with wrist trauma. *Pediatric Radiology* 2017;**47**(5):590-98
209. Kessel B, Sevi R, Jeroukhimov I, et al. Is routine portable pelvic X-ray in stable multiple trauma patients always justified in a high technology era? *Injury* 2007;**38**(5):559-63 doi: 10.1016/j.injury.2006.12.020[published Online First: Epub Date]].
210. Duane TM, Dechert T, Wolfe LG, et al. Alcohol's role on the reliability of clinical examination to rule out pelvic fractures. *The American surgeon* 2009;**75**(3):257-9
211. den Boer TA, Geurts M, van Hulsteijn LT, et al. The value of clinical examination in diagnosing pelvic fractures in blunt trauma patients: a brief review. *European journal of trauma and emergency surgery : official publication of the European Trauma Society* 2011;**37**(4):373-7 doi: 10.1007/s00068-011-0076-7[published Online First: Epub Date]].
212. Langer JM, Tsai EB, Luhar A, et al. Reducing Unnecessary Portable Pelvic Radiographs in Trauma Patients: A Resident-Driven Quality Improvement Initiative. *J Am Coll Radiol* 2015;**12**(9):954-59
213. Bolt C, O'Keefe F, Finnegan P, et al. Straight leg elevation to rule out pelvic injury. *Injury-Int J Care Inj* 2018;**49**(2):279-83
214. Jalili M, Gharebaghi H. Validation of the Ottawa Knee Rule in Iran: a prospective study. *Emergency medicine journal : EMJ* 2010;**27**(11):849-51 doi: 10.1136/emj.2009.080267[published Online First: Epub Date]].

215. Yao K, Haque T. The Ottawa knee rules - a useful clinical decision tool. *Aust Fam Physician* 2012;**41**(4):223-4
216. Aagesen AL, Melek M. Choosing the Right Diagnostic Imaging Modality in Musculoskeletal Diagnosis. *Primary Care* 2013;**40**(4):849-+
217. Konan S, Zang TT, Tamimi N, et al. Can the Ottawa and Pittsburgh rules reduce requests for radiography in patients referred to acute knee clinics? *Ann R Coll Surg Engl* 2013;**95**(3):188-91
218. Yazdani S, Jahandideh H, Ghofrani H. Validation of the Ottawa Ankle Rules in Iran: a prospective survey. *BMC emerg* 2006;**6**:3
219. Marinelli M, Di Giulio A, Mancini M. Validation of the Ottawa ankle rules in a second-level trauma center in Italy. *Journal of Orthopaedics and Traumatology : Official Journal of the Italian Society of Orthopaedics and Traumatology* 2007;**8**(1):16-20 doi: 10.1007/s10195-007-0156-y[published Online First: Epub Date]].
220. Bessen T, Clark R, Shakib S, et al. A multifaceted strategy for implementation of the Ottawa ankle rules in two emergency departments. *BMJ (Clinical research ed)* 2009;**339**:b3056 doi: 10.1136/bmj.b3056[published Online First: Epub Date]].
221. Bairstow PJ, Persaud J, Mendelson R, et al. Reducing inappropriate diagnostic practice through education and decision support. *Int J Qual Health Care* 2010;**22**(3):194-200
222. Gravel J, Roy M, Carriere B. 44-55-66-PM, a mnemonic that improves retention of the Ottawa Ankle and Foot Rules: a randomized controlled trial. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2010;**17**(8):859-64 doi: 10.1111/j.1553-2712.2010.00731.x[published Online First: Epub Date]].
223. Knudsen R, Vijdea R, Damborg F. Validation of the Ottawa ankle rules in a Danish emergency department. *Danish medical bulletin* 2010;**57**(5):A4142
224. Kose O, Gokhan S, Ozhasenekler A, et al. Comparison of Ottawa Ankle Rules and Bernese Ankle Rules in Acute Ankle and Midfoot Injuries. *Turkish Journal of Emergency Medicine* 2010;**10**(3):pp. 101-05

225. Canagasabay MD, Callaghan MJ, Carley S. The sonographic Ottawa Foot and Ankle Rules study (the SOFAR study). *Emergency medicine journal* : EMJ 2011;**28**(10):838-40 doi: 10.1136/emj.2009.088286[published Online First: Epub Date]].
226. Marinelli M, Cecconi S, Busilacchi A, et al. Validation of the Ottawa ankle rules in a second level trauma center. *Journal of Orthopaedics and Traumatology* 2011;**12**:S36 doi: <http://dx.doi.org/10.1007/s10195-011-0149-8>[published Online First: Epub Date]].
227. Sorensen EL, Keeling A, Snyder A, et al. Decreasing ED length of stay with use of the Ottawa Ankle Rules among nurses. *Journal of emergency nursing: JEN* : official publication of the Emergency Department Nurses Association 2012;**38**(4):350-2 doi: 10.1016/j.jen.2011.02.014[published Online First: Epub Date]].
228. Tollefson B, Fromang G, Koury M, et al. Validation of the sonographic ottawa foot and ankle rules study in a large urban trauma center. *Annals of emergency medicine* 2012;**1**:S17 doi: <http://dx.doi.org/10.1016/j.annemergmed.2012.06.069>[published Online First: Epub Date]].
229. Baskaran D, Rahman S, Malik Q. The avoidance of radiation exposure by following RCR guidelines and ottawa rules in performing ankle radiographs. *International Journal of Surgery* 2013;**11** (8):664 doi: <http://dx.doi.org/10.1016/j.ijvs.2013.06.416>[published Online First: Epub Date]].
230. Hedelin H, Goksor LA, Karlsson J, et al. Ultrasound-assisted triage of ankle trauma can decrease the need for radiographic imaging. *The American journal of emergency medicine* 2013;**31**(12):1686-9 doi: 10.1016/j.ajem.2013.09.005[published Online First: Epub Date]].
231. Lau LH, Kerr D, Law I, et al. Nurse practitioners treating ankle and foot injuries using the Ottawa Ankle Rules: a comparative study in the emergency department. *Australasian emergency nursing journal* : AENJ 2013;**16**(3):110-5 doi: 10.1016/j.aenj.2013.05.007[published Online First: Epub Date]].
232. Wang X, Chang SM, Yu GR, et al. Clinical value of the Ottawa ankle rules for diagnosis of fractures in acute ankle injuries. *PloS one* 2013;**8**(4):e63228 doi: 10.1371/journal.pone.0063228[published Online First: Epub Date]].

233. Derksen RJ, Knijnenberg LM, Fransen G, et al. Diagnostic performance of the Bernese versus Ottawa ankle rules: Results of a randomised controlled trial. *Injury-Int J Care Inj* 2015;**46**(8):1645-49
234. Cheng C, Varma D, Smit D. How consistently are the Ottawa ankle rules applied in a major metropolitan health network's emergency departments in Australia? A retrospective review. *Journal of Medical Imaging and Radiation Oncology* 2016;**60**:152 doi: <http://dx.doi.org/10.1111/17549485.12520>[published Online First: Epub Date]].
235. Ho JKM, Chau JPC, Cheung NMC. Effectiveness of emergency nurses' use of the Ottawa Ankle Rules to initiate radiographic tests on improving healthcare outcomes for patients with ankle injuries: A systematic review. *Int J Nurs Stud* 2016;**63**:37-47
236. Tollefson B, Nichols J, Fromang S, et al. Validation of the Sonographic Ottawa Foot and Ankle Rules (SOFAR) Study in a Large Urban Trauma Center. *J Miss State Med Assoc* 2016;**57**(2):35-8
237. Clendenin BR, Conlon HA, Burns C. Overuse of Diagnostic Imaging for Work-Related Injuries. *Workplace Health Saf* 2017;**65**(2):54-56
238. MacLellan J, Smith T, Baserman J, et al. Accuracy of the Ottawa Ankle Rules applied by non-physician providers in a pediatric emergency department. *Cjem* 2017:1-7 doi: 10.1017/cem.2017.399[published Online First: Epub Date]].
239. S. F, S. F, C. W, et al. Retrospective audit of plain film imaging in acute ankle trauma: Are we choosing wisely? *Journal of Medical Imaging and Radiation Oncology* 2017;**Conference**:68th Annual Scientific Meeting of the Royal Australian and New Zealand College of Radiologists, RANZCR 2017. Australia. 61 (Supplement 1) (pp 149-150) doi: <http://dx.doi.org/10.1111/1754-9485.12657>[published Online First: Epub Date]].
240. Dowling S, Spooner CH, Liang Y, et al. Accuracy of Ottawa Ankle Rules to exclude fractures of the ankle and midfoot in children: a meta-analysis. *Academic emergency medicine : official journal of the Society for Academic Emergency Medicine* 2009;**16**(4):277-87 doi: 10.1111/j.1553-2712.2008.00333.x[published Online First: Epub Date]].

241. Gravel J, Hedrei P, Grimard G, et al. Prospective validation and head-to-head comparison of 3 ankle rules in a pediatric population. *Annals of emergency medicine* 2009;**54**(4):534-40.e1 doi: 10.1016/j.annemergmed.2009.06.507[published Online First: Epub Date]].
242. Boutis K, Grootendorst P, Willan A, et al. Effect of the Low Risk Ankle Rule on the frequency of radiography in children with ankle injuries. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne* 2013;**185**(15):E731-8 doi: 10.1503/cmaj.122050[published Online First: Epub Date]].
243. Lane N, Yorke H, Stewart C. The low risk ankle rule can safely be applied to a UK paediatric population. *Archives of disease in childhood* 2014;**99**:A163-A64 doi: <http://dx.doi.org/10.1136/archdischild-2014-307384.441>[published Online First: Epub Date]].
244. Boutis K, von Keyserlingk C, Willan A, et al. Cost Consequence Analysis of Implementing the Low Risk Ankle Rule in Emergency Departments. *Annals of emergency medicine* 2015;**66**(5):455-63.e4 doi: 10.1016/j.annemergmed.2015.05.027[published Online First: Epub Date]].
245. Loftus KV, Gittelman MA, FitzGerald M, et al. An intervention to increase knowledge and utilization of the low risk ankle rule among pediatric emergency room providers. *Academic Emergency Medicine* 2016;**23**:S197 doi: <http://dx.doi.org/10.1111/acem.12974>[published Online First: Epub Date]].
246. MacLellan J, Smith T, Baserman J, et al. Accuracy of the Ottawa Ankle Rules when applied by allied health providers in a pediatric emergency department. *Canadian Journal of Emergency Medicine* 2016;**18**:S107 doi: <http://dx.doi.org/10.1017/cem.2016.262>[published Online First: Epub Date]].
247. Ellenbogen AL, Rice AL, Vyas P. Retrospective comparison of the Low Risk Ankle Rules and the Ottawa Ankle Rules in a pediatric population. *The American journal of emergency medicine* 2017;**35**(9):1262-65 doi: 10.1016/j.ajem.2017.03.058[published Online First: Epub Date]].

248. Ramasubbu RA. Is Radiography Necessary for Pediatric Emergency Department Clinicians to Safely Manage Ankle Injuries? *Pediatric emergency care* 2017 doi: 10.1097/pec.0000000000001197[published Online First: Epub Date]].
249. Faulkner A, Reidy M, Scicluna G, et al. Blood tests: One too many? Evaluating blood requesting guidance developed for acute patients admitted to trauma and orthopaedic units. *Injury* 2015 doi: <http://dx.doi.org/10.1016/j.injury.2015.11.041>[published Online First: Epub Date]].
250. Wedde TB, Quinlan JF, Khan A, et al. Fractures of the sternum: the influence of non-invasive cardiac monitoring on management. *Archives of orthopaedic and trauma surgery* 2007;**127**(2):121-3 doi: 10.1007/s00402-006-0230-6[published Online First: Epub Date]].
251. Choi PM, Farmakis S, Desmarais TJ, et al. Management and outcomes of traumatic hemothorax in children. *Journal of emergencies, trauma, and shock* 2015;**8**(2):83-7 doi: 10.4103/0974-2700.155500[published Online First: Epub Date]].
252. Ollidashi F, Kerci M, Zhurda T, et al. The importance of early treatment with tranexamic acid in bleeding trauma patients: An exploratory analysis of the CRASH-2 randomised controlled trial. *Lancet (London, England)* 2011; 377(9771).  
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/583/CN-00888583/frame.html>.
253. Roberts I, Shakur H, Coats T, et al. The CRASH-2 trial: a randomised controlled trial and economic evaluation of the effects of tranexamic acid on death, vascular occlusive events and transfusion requirement in bleeding trauma patients. *Health technology assessment (Winchester, England)* 2013;**17**(10):1-79 doi: 10.3310/hta17100[published Online First: Epub Date]].
254. DeLoughery EP, Lenfesty B, DeLoughery TG. A retrospective case control study of recombinant factor VIIa in patients with intracranial haemorrhage caused by trauma. *British journal of haematology* 2011;**152**(5):667-9 doi: 10.1111/j.1365-2141.2010.08437.x[published Online First: Epub Date]].
255. Spahn DR, Bouillon B, Cerny V, et al. Management of bleeding and coagulopathy following major trauma: an updated European guideline. *Critical care (London, England)* 2013;**17**(2):R76 doi: 10.1186/cc12685[published Online First: Epub Date]].

256. Cothren CC, Moore EE. Emergency department thoracotomy for the critically injured patient: Objectives, indications, and outcomes. *World journal of emergency surgery* : WJES 2006;**1**:4 doi: 10.1186/1749-7922-1-4[published Online First: Epub Date]].
257. Mollberg NM, Glenn C, John J, et al. Appropriate use of emergency department thoracotomy: implications for the thoracic surgeon. *The Annals of thoracic surgery* 2011;**92**(2):455-61 doi: 10.1016/j.athoracsur.2011.04.042[published Online First: Epub Date]].
258. Moore EE, Knudson MM, Burlew CC, et al. Defining the limits of resuscitative emergency department thoracotomy: a contemporary Western Trauma Association perspective. *The Journal of trauma* 2011;**70**(2):334-9 doi: 10.1097/TA.0b013e3182077c35[published Online First: Epub Date]].
259. Seamon MJ, Haut ER, Van Arendonk K, et al. An evidence-based approach to patient selection for emergency department thoracotomy: A practice management guideline from the Eastern Association for the Surgery of Trauma. *The journal of trauma and acute care surgery* 2015;**79**(1):159-73 doi: <http://dx.doi.org/10.1097/TA.0000000000000648>[published Online First: Epub Date]].



## References for Table 3

1. St Peter SD, Sharp SW, Snyder CL, et al. Prospective validation of an abbreviated bedrest protocol in the management of blunt spleen and liver injury in children. *Journal of pediatric surgery* 2011;**46**(1):173-7 doi: 10.1016/j.jpedsurg.2010.09.079[published Online First: Epub Date].
2. St Peter SD, Aguayo P, Juang D, et al. Follow up of prospective validation of an abbreviated bedrest protocol in the management of blunt spleen and liver injury in children. *Journal of pediatric surgery* 2013;**48**(12):2437-41 doi: 10.1016/j.jpedsurg.2013.08.018[published Online First: Epub Date].
3. Gor R, Styskel B, Li T, et al. Practice patterns of angiography and angioembolization performance for isolated low grade renal trauma: Results from a large, statewide trauma registry. *Journal of Urology* 2016;**1**:e787
4. Harvin JA, Wray CJ, Steward J, et al. Control the damage: morbidity and mortality after emergent trauma laparotomy. *Am J Surg* 2016;**212**(1):34-39
5. Coimbra R, Hoyt DB, Engelhart S, et al. Nonoperative management reduces the overall mortality of grades 3 and 4 blunt liver injuries. *Int Surg* 2006;**91**(5):251-57
6. van der Wilden GM, Velmahos GC, Emhoff T, et al. Successful nonoperative management of the most severe blunt liver injuries: a multicenter study of the research consortium of new England centers for trauma. *Archives of surgery (Chicago, Ill : 1960)* 2012;**147**(5):423-8 doi: 10.1001/archsurg.2012.147[published Online First: Epub Date].
7. Zago TM, Tavares Pereira BM, Araujo Calderan TR, et al. Nonoperative management for patients with grade IV blunt hepatic trauma. *World journal of emergency surgery : WJES* 2012;**7 Suppl 1**:S8 doi: 10.1186/1749-7922-7-S1-S8[published Online First: Epub Date].

8. Azzam AZ, Gazal AH, Kassem MI, et al. The role of non-operative management (NOM) in blunt hepatic trauma. Alexandria Journal of Medicine 2013;**49**(3):223-27 doi: <http://dx.doi.org/10.1016/j.ajme.2012.10.003>[published Online First: Epub Date]].
9. Golden J, Mitchell I, Kuzniewski S, et al. Reducing scheduled phlebotomy in stable pediatric patients with blunt liver or spleen injury. Journal of pediatric surgery 2014;**49**(5):759-62 doi: 10.1016/j.jpedsurg.2014.02.062[published Online First: Epub Date]].
10. Landau A, van As AB, Numanoglu A, et al. Liver injuries in children: the role of selective non-operative management. Injury 2006;**37**(1):66-71 doi: 10.1016/j.injury.2005.07.013[published Online First: Epub Date]].
11. Inchingolo R, Ljutikov A, Deganello A, et al. Outcomes and indications for intervention in non-operative management of paediatric liver trauma: a 5 year retrospective study. Clinical radiology 2014;**69**(2):157-62
12. Vick LR, Islam S. Adding insult to injury: neck exploration for penetrating pediatric neck trauma. The American surgeon 2008;**74**(11):1104-6
13. Van Waes OJ, Cheriex KC, Navsaria PH, et al. Management of penetrating neck injuries. The British journal of surgery 2012;**99** Suppl 1:149-54 doi: 10.1002/bjs.7733[published Online First: Epub Date]].
14. Siau RTK, Moore A, Ahmed T, et al. Management of penetrating neck injuries at a London trauma centre. Eur Arch Oto-Rhino-Laryn 2013;**270**(7):2123-28
15. Prichayudh S, Choadrachata-Anun J, Pak-Art R, et al. Selective management of penetrating neck injuries using "no zone" approach. Injury 2015;**46**(9):1720-25 doi: <http://dx.doi.org/10.1016/j.injury.2015.06.019>[published Online First: Epub Date]].
16. Ibraheem K, Khan M, Rhee P, et al. "No zone" approach in penetrating neck trauma reduces unnecessary computed tomography angiography and negative explorations. The Journal of surgical research 2018;**221**:113-20 doi: 10.1016/j.jss.2017.08.033[published Online First: Epub Date]].
17. Bjurlin MA, Jeng EI, Goble SM, et al. Comparison of nonoperative management with renorrhaphy and nephrectomy in penetrating renal injuries. The Journal of trauma 2011;**71**(3):554-8 doi: 10.1097/TA.0b013e318203321a[published Online First: Epub Date]].

18. Uranues S, Popa DE, Diaconescu B, et al. Laparoscopy in penetrating abdominal trauma. *World journal of surgery* 2015;**39**(6):1381-8 doi: 10.1007/s00268-014-2904-5[published Online First: Epub Date]].
19. Eppich WJ, Zonfrillo MR. Emergency department evaluation and management of blunt abdominal trauma in children. *Current opinion in pediatrics* 2007;**19**(3):265-9 doi: 10.1097/MOP.0b013e328149af9e[published Online First: Epub Date]].
20. Weinberg JA, Magnotti LJ, Croce MA, et al. The utility of serial computed tomography imaging of blunt splenic injury: still worth a second look? *The Journal of trauma* 2007;**62**(5):1143-7; discussion 47-8 doi: 10.1097/TA.0b013e318047b7c2[published Online First: Epub Date]].
21. Bruce PJ, Helmer SD, Harrison PB, et al. Nonsurgical management of blunt splenic injury: is it cost effective? *Am J Surg* 2011;**202**(6):810-5; discussion 15-6 doi: 10.1016/j.amjsurg.2011.06.041[published Online First: Epub Date]].
22. Cirocchi R, Boselli C, Corsi A, et al. Is non-operative management safe and effective for all splenic blunt trauma? A systematic review. *Crit Care* 2013;**17**(5)
23. Tugnoli G, Bianchi E, Biscardi A, et al. Nonoperative management of blunt splenic injury in adults: there is (still) a long way to go. The results of the Bologna-Maggiore Hospital trauma center experience and development of a clinical algorithm. *Surg Today* 2015;**45**(10):1210-17
24. Khan AM, Manzoor K, Malik Z, et al. Thoracic splenosis: know it--avoid unnecessary investigations, interventions, and thoracotomy. *General thoracic and cardiovascular surgery* 2011;**59**(4):245-53 doi: 10.1007/s11748-010-0706-8[published Online First: Epub Date]].
25. Thompson BE, Munera F, Cohn SM, et al. Novel computed tomography scan scoring system predicts the need for intervention after splenic injury. *The Journal of trauma* 2006;**60**(5):1083-6 doi: 10.1097/01.ta.0000218251.67141.ef[published Online First: Epub Date]].
26. Mooney DP, Rothstein DH, Forbes PW. Variation in the management of pediatric splenic injuries in the United States. *The Journal of trauma* 2006;**61**(2):330-3; discussion 33 doi: 10.1097/01.ta.0000226167.44892.1d[published Online First: Epub Date]].

27. Davies DA, Pearl RH, Ein SH, et al. Management of blunt splenic injury in children: evolution of the nonoperative approach. *Journal of pediatric surgery* 2009;**44**(5):1005-8 doi: 10.1016/j.jpedsurg.2009.01.024[published Online First: Epub Date]].
28. Hsiao M, Sathya C, de Mestral C, et al. Population-based analysis of blunt splenic injury management in children: operative rate is an informative quality of care indicator. *Injury* 2014;**45**(5):859-63 doi: 10.1016/j.injury.2013.12.006[published Online First: Epub Date]].

ACCEPTED

## References for Table 4

1. Levy AS, Orlando A, Hawkes AP, et al. Should the management of isolated traumatic subarachnoid hemorrhage differ from concussion in the setting of mild traumatic brain injury? *The Journal of trauma* 2011;**71**(5):1199-204 doi: 10.1097/TA.0b013e31822067fc[published Online First: Epub Date]].
2. Washington CW, Grubb RL, Jr. Are routine repeat imaging and intensive care unit admission necessary in mild traumatic brain injury? *Journal of neurosurgery* 2012;**116**(3):549-57 doi: 10.3171/2011.11.jns111092[published Online First: Epub Date]].
3. C.L. DO, R. R, A.J. S, et al. The identification of a subgroup of children with traumatic subarachnoid hemorrhage at low risk of neuroworsening. *Clinical Neurosurgery* 2017;**Conference**:2017 Annual Meeting of the Congress of Neurological Surgeons. United States. 64 (Supplement 1) (pp 269) doi: <http://dx.doi.org/10.1093/neuros/nyx417>[published Online First: Epub Date]].
4. Dhillon NK, Ko A, Smith EJT, et al. Potentially Avoidable Surgical Intensive Care Unit Admissions and Disposition Delays. *JAMA Surg* 2017;**152**(11):1015-22 doi: 10.1001/jamasurg.2017.2165[published Online First: Epub Date]].
5. J. Z, P. D. Can patients who present with isolated traumatic subarachnoid hemorrhage with normal mental status be discharged from the emergency department? *Annals of emergency medicine* 2017;**Conference**:American College of Emergency Physicians, ACEP 2017 Research Forum. United States. 70 (4 Supplement 1) (pp S111-S112)
6. Huynh T, Jacobs DG, Dix S, et al. Utility of neurosurgical consultation for mild traumatic brain injury. *The American surgeon* 2006;**72**(12):1162-5; discussion66-7

7. Joseph B, Aziz H, Pandit V, et al. Prospective validation of the brain injury guidelines: managing traumatic brain injury without neurosurgical consultation. *The journal of trauma and acute care surgery* 2014;**77**(6):984-8 doi: 10.1097/ta.0000000000000428[published Online First: Epub Date]].
8. Hirsh J, Guyatt G, Albers GW, et al. Executive summary: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest* 2008;**133**(6 Suppl):71s-109s doi: 10.1378/chest.08-0693[published Online First: Epub Date]].
9. Gorman PH, Qadri SF, Rao-Patel A. Prophylactic inferior vena cava (IVC) filter placement may increase the relative risk of deep venous thrombosis after acute spinal cord injury. *The Journal of trauma* 2009;**66**(3):707-12 doi: 10.1097/TA.0b013e318188beba[published Online First: Epub Date]].
10. Macatangay C, Todd SR, Tyroch AH. Thromboembolic prophylaxis with intermittent pneumatic compression devices in trauma patients: a false sense of security? *Journal of trauma nursing : the official journal of the Society of Trauma Nurses* 2008;**15**(1):12-15
11. Goodman MD, Huber NL, Johannigman JA, et al. Omission of routine chest x-ray after chest tube removal is safe in selected trauma patients. *Am J Surg* 2010;**199**(2):199-203
12. Ratilal B, Costa J, Sampaio C. Antibiotic prophylaxis for preventing meningitis in patients with basilar skull fractures. *The Cochrane database of systematic reviews* 2006(1):Cd004884 doi: 10.1002/14651858.CD004884.pub2[published Online First: Epub Date]].
13. Ratilal BO, Costa J, Sampaio C, et al. Antibiotic prophylaxis for preventing meningitis in patients with basilar skull fractures. *The Cochrane database of systematic reviews* 2011(8):Cd004884 doi: 10.1002/14651858.CD004884.pub3[published Online First: Epub Date]].
14. Ratilal BO, Costa J, Pappamikail L, et al. Antibiotic prophylaxis for preventing meningitis in patients with basilar skull fractures. *The Cochrane database of systematic reviews* 2015(4):Cd004884 doi: 10.1002/14651858.CD004884.pub4[published Online First: Epub Date]].

15. Sayer FT, Kronvall E, Nilsson OG. Methylprednisolone treatment in acute spinal cord injury: the myth challenged through a structured analysis of published literature. *The spine journal : official journal of the North American Spine Society* 2006;**6**(3):335-43 doi: 10.1016/j.spinee.2005.11.001[published Online First: Epub Date]].

ACCEPTED

# References for Table 5

1. Matthews AH, Boyd M, Bott AR, et al. Improving emergency department management of paediatric clavicle fractures: a complete audit cycle. *British journal of hospital medicine (London, England : 2005)* 2014;**75**(5):287-9 doi: 10.12968/hmed.2014.75.5.287[published Online First: Epub Date]].
2. Bradley LH, Paullus WC, Howe J, et al. Isolated transverse process fractures: spine service management not needed. *Journal of Trauma-Injury Infection & Critical Care* 2008;**65**(4):832-6; discussion 36 doi: <http://dx.doi.org/10.1097/TA.0b013e318184d30e>[published Online First: Epub Date]].
3. Shulman BS, Lee JH, Liporace FA, et al. Minimally Displaced Radial Head/Neck Fractures (Mason Type-I, OTA Types 21A2.2 and 21B2.1): Are We "Over Treating" Our Patients? *J Orthop Trauma* 2015;**29**(2):E31-E35
4. Bailey CS, Dvorak MF, Thomas KC, et al. Comparison of thoracolumbosacral orthosis and no orthosis for the treatment of thoracolumbar burst fractures: interim analysis of a multicenter randomized clinical equivalence trial. *Journal of neurosurgery Spine* 2009;**11**(3):295-303 doi: 10.3171/2009.3.spine08312[published Online First: Epub Date]].
5. Bailey C, Dvorak M, Nadeau M, et al. No Orthosis is Equivalent to TLSO for the Treatment of Thoracolumbar Burst Fractures without Neurologic Injury: Results from a Multicenter RCT. *The Spine Journal* 2011;**11**(10):S1-S2 doi: 10.1016/j.spinee.2011.08.017[published Online First: Epub Date]].
6. Bailey CS, Urquhart JC, Dvorak MF, et al. Orthosis versus no orthosis for the treatment of thoracolumbar burst fractures without neurologic injury: a multicenter prospective randomized equivalence trial. *The spine journal : official journal of the North American Spine Society* 2014;**14**(11):2557-64 doi: 10.1016/j.spinee.2013.10.017[published Online First: Epub Date]].



7. Fournier DA, Lollis S, Hawkins HLS, et al. Multidisciplinary development of bracing protocol for stable thoracolumbar fractures. Archives of physical medicine and rehabilitation 2015; 96(10). <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/814/CN-01126814/frame.html>.
8. Alazzawi S, De Rover WBS, Leary T, et al. Patients undergoing blood tests before minor/moderate trauma surgery: a retrospective review. JRSM Short Rep 2012;**3**(6):39 doi: <http://dx.doi.org/10.1258/shorts.2011.011130>[published Online First: Epub Date].

ACCEPTED

## References for eTable 2a

1. Khoriaty AA, Rajakulasingam R, Shah R. Sternal fractures and their management. *Journal of Emergencies, Trauma and Shock* 2013;**6**(2):113-16  
doi: <http://dx.doi.org/10.4103/0974-2700.110763>[published Online First: Epub Date]].
2. Rollins MD, Barnhart DC, Greenberg RA, et al. Neurologically intact children with an isolated skull fracture may be safely discharged after brief observation. *Journal of pediatric surgery* 2011;**46**(7):1342-46
3. Metzger RR, Smith J, Wells M, et al. Impact of newly adopted guidelines for management of children with isolated skull fracture. *Journal of pediatric surgery* 2014;**49**(12):1856-60
4. Powell EC, Atabaki SM, Wootton-Gorges S, et al. Isolated Linear Skull Fractures in Children With Blunt Head Trauma. *Pediatrics* 2015;**135**(4):E851-E57
5. Blackwood BP, Bean JF, Sadecki-Lund C, et al. Observation for isolated traumatic skull fractures in the pediatric population: unnecessary and costly. *Journal of pediatric surgery* 2016;**51**(4):654-58
6. Reuveni-Salzman A, Rosenthal G, Poznanski O, et al. Evaluation of the necessity of hospitalization in children with an isolated linear skull fracture (ISF). *Childs Nerv Syst* 2016;**32**(9):1669-74
7. Coon ER, Young PC, Quinonez RA, et al. Update on Pediatric Overuse. *Pediatrics* 2017;**139**(2)
8. Miller P, Coffey F, Reid AM, et al. Can emergency nurses use the Canadian cervical spine rule to reduce unnecessary patient immobilisation? *Accident and Emergency Nursing* 2006;**14**(3):133-40 doi: <http://dx.doi.org/10.1016/j.aen.2006.03.003>[published Online First: Epub Date]].

9. Benger J, Blackham J. Why do we put cervical collars on conscious trauma patients? *Scandinavian journal of trauma, resuscitation and emergency medicine* 2009;**17**:44 doi: 10.1186/1757-7241-17-44[published Online First: Epub Date]].
10. Panczykowski DM, Tomycz ND, Okonkwo DO. Comparative effectiveness of using computed tomography alone to exclude cervical spine injuries in obtunded or intubated patients: meta-analysis of 14,327 patients with blunt trauma A review. *Journal of neurosurgery* 2011;**115**(3):541-49
11. Sixta S, Moore FO, Ditillo MF, et al. Screening for thoracolumbar spinal injuries in blunt trauma: an Eastern Association for the Surgery of Trauma practice management guideline. *The journal of trauma and acute care surgery* 2012;**73**(5 Suppl 4):S326-32 doi: 10.1097/TA.0b013e31827559b8[published Online First: Epub Date]].
12. Sifri ZC, Homnick AT, Vaynman A, et al. A prospective evaluation of the value of repeat cranial computed tomography in patients with minimal head injury and an intracranial bleed. *J Trauma-Injury Infect Crit Care* 2006;**61**(4):862-67
13. Brown CV, Zada G, Salim A, et al. Indications for routine repeat head computed tomography (CT) stratified by severity of traumatic brain injury. *The Journal of trauma* 2007;**62**(6):1339-44; discussion 44-5 doi: 10.1097/TA.0b013e318054e25a[published Online First: Epub Date]].
14. Smith JS, Chang EF, Rosenthal G, et al. The role of early follow-up computed tomography imaging in the management of traumatic brain injury patients with intracranial hemorrhage. *The Journal of trauma* 2007;**63**(1):75-82 doi: 10.1097/01.ta.0000245991.42871.87[published Online First: Epub Date]].
15. Kaen A, Jimenez-Roldan L, Arrese I, et al. The value of sequential computed tomography scanning in anticoagulated patients suffering from minor head injury. *The Journal of trauma* 2010;**68**(4):895-8 doi: 10.1097/TA.0b013e3181b28a76[published Online First: Epub Date]].

16. Connon FF, Namdarian B, Ee JLC, et al. Do routinely repeated computed tomography scans in traumatic brain injury influence management?

A prospective observational study in a level 1 trauma center. *Annals of surgery* 2011;**254**(6):1028-31 doi:

<http://dx.doi.org/10.1097/SLA.0b013e318219727f>[published Online First: Epub Date].

ACCEPTED

## References for eTable 2b

1. Como JJ, Bokhari F, Chiu WC, et al. Practice management guidelines for selective nonoperative management of penetrating abdominal trauma. *The Journal of trauma* 2010;**68**(3):721-33 doi: 10.1097/TA.0b013e3181cf7d07[published Online First: Epub Date]].
2. Kevric J, O'Reilly GM, Gocentas RA, et al. Management of haemodynamically stable patients with penetrating abdominal stab injuries: review of practice at an Australian major trauma centre. *European journal of trauma and emergency surgery : official publication of the European Trauma Society* 2016;**42**(6):671-75 doi: 10.1007/s00068-015-0605-x[published Online First: Epub Date]].
3. Biffl WL, Leppaniemi A. Management guidelines for penetrating abdominal trauma. *World journal of surgery* 2015;**39**(6):1373-80 doi: 10.1007/s00268-014-2793-7[published Online First: Epub Date]].
4. Malcolm JB, Derweesh IH, Mehrazin R, et al. Nonoperative management of blunt renal trauma: is routine early follow-up imaging necessary? *BMC urology* 2008;**8**:11 doi: 10.1186/1471-2490-8-11[published Online First: Epub Date]].
5. Breen KJ, Sweeney P, Nicholson PJ, et al. Adult blunt renal trauma: routine follow-up imaging is excessive. *Urology* 2014;**84**(1):62-7 doi: 10.1016/j.urology.2014.03.013[published Online First: Epub Date]].
6. Raupach J, Ferko A, Lojik M, et al. Endovascular treatment of acute and chronic thoracic aortic injury. *Cardiovasc Intervent Radiol* 2007;**30**(6):1117-23 doi: 10.1007/s00270-007-9053-2[published Online First: Epub Date]].
7. Mayberry J, Fabricant L, Anton A, et al. Management of full-thickness duodenal laceration in the damage control era: evolution to primary repair without diversion or decompression. *The American surgeon* 2011;**77**(6):681-5
8. Roberts DJ, Bobrovitz N, Zygun DA, et al. Indications for use of thoracic, abdominal, pelvic, and vascular damage control interventions in trauma patients: A content analysis and expert appropriateness rating study. *J Trauma Acute Care Surg* 2015;**79**(4):568-79

9. Sapkota R, Bhandari RS. Prophylactic Nasogastric Decompression after Emergency Laparotomy. *J Nepal Med Assoc* 2013;**52**(7):437-42
10. Talving P, Nicol AJ, Navsaria PH. Civilian duodenal gunshot wounds: surgical management made simpler. *World journal of surgery* 2006;**30**(4):488-94 doi: 10.1007/s00268-005-0245-0[published Online First: Epub Date]].
11. Landau A, van As AB, Numanoglu A, et al. Liver injuries in children: the role of selective non-operative management. *Injury* 2006;**37**(1):66-71 doi: 10.1016/j.injury.2005.07.013[published Online First: Epub Date]].
12. Seamon MJ, Smoger D, Torres DM, et al. A prospective validation of a current practice: the detection of extremity vascular injury with CT angiography. *The Journal of trauma* 2009;**67**(2):238-43; discussion 43-4 doi: 10.1097/TA.0b013e3181a51bf9[published Online First: Epub Date]].
13. Insull P, Adams D, Segar A, et al. Is exploration mandatory in penetrating zone II neck injuries? *ANZ journal of surgery* 2007;**77**(4):261-4 doi: 10.1111/j.1445-2197.2007.04030.x[published Online First: Epub Date]].
14. Tisherman SA, Bokhari F, Collier B, et al. Clinical practice guideline: penetrating zone II neck trauma. *The Journal of trauma* 2008;**64**(5):1392-405 doi: 10.1097/TA.0b013e3181692116[published Online First: Epub Date]].
15. Inaba K, Branco BC, Menaker J, et al. Evaluation of multidetector computed tomography for penetrating neck injury: a prospective multicenter study. *The journal of trauma and acute care surgery* 2012;**72**(3):576-83; discussion 83-4; quiz 803-4 doi: 10.1097/TA.0b013e31824badf7[published Online First: Epub Date]].
16. Hosseini SV, Sabet B, Rezaianzadeh A, et al. Role of Physical Examination in Decision Making for Selective Exploration in Patients with Penetrating Zone II Neck Injury. *Bull* 2013;**1**(2):90-2
17. Abbo O, Lemandat A, Reina N, et al. Conservative management of blunt pancreatic trauma in children: a single center experience. *European journal of pediatric surgery : official journal of Austrian Association of Pediatric Surgery [et al] = Zeitschrift fur Kinderchirurgie* 2013;**23**(6):470-3 doi: 10.1055/s-0033-1333642[published Online First: Epub Date]].

18. Cirocchi R, Trastulli S, Pressi E, et al. Non-operative management versus operative management in high-grade blunt hepatic injury. *Cochrane Database of Systematic Reviews* 2015; (8). <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD010989.pub2/abstract>.
19. Broghammer JA, Fisher MB, Santucci RA. Conservative management of renal trauma: a review. *Urology* 2007;**70**(4):623-9 doi: 10.1016/j.urology.2007.06.1085[published Online First: Epub Date]].
20. Eppich WJ, Zonfrillo MR. Emergency department evaluation and management of blunt abdominal trauma in children. *Current opinion in pediatrics* 2007;**19**(3):265-9 doi: 10.1097/MOP.0b013e328149af9e[published Online First: Epub Date]].
21. Fraser JD, Aguayo P, Ostlie DJ, et al. Review of the evidence on the management of blunt renal trauma in pediatric patients. *Pediatric surgery international* 2009;**25**(2):125-32 doi: 10.1007/s00383-008-2316-4[published Online First: Epub Date]].
22. Tugnoli G, Bianchi E, Biscardi A, et al. Nonoperative management of blunt splenic injury in adults: there is (still) a long way to go. The results of the Bologna-Maggiore Hospital trauma center experience and development of a clinical algorithm. *Surg Today* 2015;**45**(10):1210-17
23. Sujenthiran A, Elshout PJ, Veskimae E, et al. Is Nonoperative Management the Best First-line Option for High-grade Renal trauma? A Systematic Review. *European urology focus* 2017 doi: 10.1016/j.euf.2017.04.011[published Online First: Epub Date]].
24. McVay MR, Kokoska ER, Jackson RJ, et al. Throwing out the "grade" book: management of isolated spleen and liver injury based on hemodynamic status. *Journal of pediatric surgery* 2008;**43**(6):1072-6 doi: 10.1016/j.jpedsurg.2008.02.031[published Online First: Epub Date]].
25. De Vries C, Van Rensburg JJ. Liver and splenic traumatic haemorrhage: When and how to intervene. *CardioVascular and Interventional Radiology* 2010;**33**:89-90 doi: <http://dx.doi.org/10.1007/s00270-010-9954-3>[published Online First: Epub Date]].
26. Stassen NA, Bhullar I, Cheng JD, et al. Nonoperative management of blunt hepatic injury: an Eastern Association for the Surgery of Trauma practice management guideline. *The journal of trauma and acute care surgery* 2012;**73**(5 Suppl 4):S288-93 doi: 10.1097/TA.0b013e318270160d[published Online First: Epub Date]].

27. Hommes M, Navsaria PH, Schipper IB, et al. Management of blunt liver trauma in 134 severely injured patients. *Injury-Int J Care Inj* 2015;**46**(5):837-42
28. Gunn ML, Clark RT, Sadro CT, et al. Current concepts in imaging evaluation of penetrating transmediastinal injury. *Radiographics* : a review publication of the Radiological Society of North America, Inc 2014;**34**(7):1824-41 doi: 10.1148/rg.347130022[published Online First: Epub Date].

ACCEPTED



## References for eTable 2c

1. Lewis PR, Dunne CE, Wallace JD, et al. Routine neurosurgical consultation is not necessary in mild blunt traumatic brain injury. *The journal of trauma and acute care surgery* 2017;**82**(4):776-80 doi: 10.1097/ta.0000000000001388[published Online First: Epub Date]].
2. Sahuquillo J. Decompressive craniectomy for the treatment of refractory high intracranial pressure in traumatic brain injury. *Cochrane Database of Systematic Reviews* 2006; (1). <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD003983.pub2/abstract>.
3. Jamous M, Barbarawi M, Samrah S, et al. Emergency decompressive craniectomy for trauma patients with Glasgow Coma Scale of 3 and bilateral fixed dilated pupils. *Eur J Trauma Emerg Surg* 2010;**36**(5):465-69
4. Cooper DJ, Rosenfeld JV, Wolfe R. DECRA investigators' response to "The future of decompressive craniectomy for diffuse traumatic brain injury" by Honeybul et al. *Journal of neurotrauma* 2012;**29**(16):2595-6 doi: 10.1089/neu.2011.2279[published Online First: Epub Date]].
5. Carney N, Totten AM, O'Reilly C, et al. Guidelines for the Management of Severe Traumatic Brain Injury, Fourth Edition. *Neurosurgery* 2017;**80**(1):6-15 doi: 10.1227/neu.0000000000001432[published Online First: Epub Date]].
6. Kalayci M, Aktunc E, Gul S, et al. Decompressive craniectomy for acute subdural haematoma: An overview of current prognostic factors and a discussion about some novel prognostic parametres. *J Pak Med Assoc* 2013;**63**(1):38-49
7. Spahn DR, Bouillon B, Cerny V, et al. Management of bleeding and coagulopathy following major trauma: an updated European guideline. *Critical care (London, England)* 2013;**17**(2):R76 doi: 10.1186/cc12685[published Online First: Epub Date]].
8. Rangel-Castillo L, Gopinath S, Robertson CS. Management of intracranial hypertension. *Neurol Clin* 2008;**26**(2):521-+
9. Lerma JD, Pappu S. Analysis of computed tomography scan data for estimation of intracranial pressure in severe traumatic brain injury. *Journal of neurotrauma* 2013;**30** (15):A125 doi: <http://dx.doi.org/10.1089/neu.2013.9938>[published Online First: Epub Date]].

10. Stocchetti N, Picetti E, Berardino M, et al. Clinical applications of intracranial pressure monitoring in traumatic brain injury : report of the Milan consensus conference. *Acta neurochirurgica* 2014;**156**(8):1615-22 doi: 10.1007/s00701-014-2127-4[published Online First: Epub Date]].
11. Stone JJ, Childs S, Smith LE, et al. Hourly neurologic assessments for traumatic brain injury in the ICU. *Neurological research* 2014;**36**(2):164-9 doi: 10.1179/1743132813y.0000000285[published Online First: Epub Date]].
12. Magnotti LJ, Schroepfel TJ, Clement LP, et al. Efficacy of Monotherapy in the Treatment of Pseudomonas Ventilator-Associated Pneumonia in Patients With Trauma. *J Trauma-Injury Infect Crit Care* 2009;**66**(4):1052-58
13. Goldberg SR, Anand RJ, Como JJ, et al. Prophylactic antibiotic use in penetrating abdominal trauma: an Eastern Association for the Surgery of Trauma practice management guideline. *The journal of trauma and acute care surgery* 2012;**73**(5 Suppl 4):S321-5 doi: 10.1097/TA.0b013e3182701902[published Online First: Epub Date]].
14. Ratilal B, Costa J, Sampaio C. Antibiotic prophylaxis for preventing meningitis in patients with basilar skull fractures. *The Cochrane database of systematic reviews* 2006(1):Cd004884 doi: 10.1002/14651858.CD004884.pub2[published Online First: Epub Date]].
15. Ratilal BO, Costa J, Sampaio C, et al. Antibiotic prophylaxis for preventing meningitis in patients with basilar skull fractures. *The Cochrane database of systematic reviews* 2011(8):Cd004884 doi: 10.1002/14651858.CD004884.pub3[published Online First: Epub Date]].
16. Ratilal BO, Costa J, Pappamikail L, et al. Antibiotic prophylaxis for preventing meningitis in patients with basilar skull fractures. *The Cochrane database of systematic reviews* 2015(4):Cd004884 doi: 10.1002/14651858.CD004884.pub4[published Online First: Epub Date]].
17. Causey MW, Rivadeneira DE, Steele SR. Historical and current trends in colon trauma. *Clin* 2012;**25**(4):189-99 doi: <http://dx.doi.org/10.1055/s-0032-1329389>[published Online First: Epub Date]].
18. Guidelines for the management of severe traumatic brain injury. *Journal of neurotrauma* 2007;**24 Suppl 1**:S1-106 doi: 10.1089/neu.2007.9999[published Online First: Epub Date]].

19. Adamides AA, Winter CD, Lewis PM, et al. Current controversies in the management of patients with severe traumatic brain injury. ANZ journal of surgery 2006;**76**(3):163-74
20. Bratton SL, Chestnut RM, Ghajar J, et al. Guidelines for the management of severe traumatic brain injury. IX. Cerebral perfusion thresholds. Journal of neurotrauma 2007;**24 Suppl 1**:S59-64 doi: 10.1089/neu.2007.9987[published Online First: Epub Date]].
21. Roberts I, Sydenham E. Barbiturates for acute traumatic brain injury. The Cochrane database of systematic reviews 2012;**12**:Cd000033 doi: 10.1002/14651858.CD000033.pub2[published Online First: Epub Date]].
22. Frenette AJ, Kanji S, Rees L, et al. Efficacy and safety of dopamine agonists in traumatic brain injury: a systematic review of randomized controlled trials. Journal of neurotrauma 2012;**29**(1):1-18 doi: 10.1089/neu.2011.1812[published Online First: Epub Date]].
23. Bhattacharyya K, Mandal N, Paul UK, et al. Post-traumatic seizure: a multicentric epidemiological study. J Indian Med Assoc 2014;**112**(2):93-5
24. Verduzco-Gutierrez M, Reddy CC, O'Dell MW. Is There a Need for Early Seizure Prophylaxis After Traumatic Brain Injury? PM and R 2016;**8**(2):169-75 doi: <http://dx.doi.org/10.1016/j.pmrj.2016.01.005>[published Online First: Epub Date]].
25. Haddad SH, Arabi YM. Critical care management of severe traumatic brain injury in adults. Scandinavian journal of trauma, resuscitation and emergency medicine 2012;**20**:12 doi: 10.1186/1757-7241-20-12[published Online First: Epub Date]].
26. Ho VP, Patel NJ, Bokhari F, et al. Management of adult pancreatic injuries: A practice management guideline from the Eastern Association for the Surgery of Trauma. The journal of trauma and acute care surgery 2017;**82**(1):185-99 doi: 10.1097/ta.0000000000001300[published Online First: Epub Date]].
27. Edwards MR, Mythen MG. Fluid therapy in critical illness. Extreme Physiology & Medicine 2014;**3**:16-16 doi: 10.1186/2046-7648-3-16[published Online First: Epub Date]].

28. West KL, Adamson C, Hoffman M. Prophylactic correction of the international normalized ratio in neurosurgery: a brief review of a brief literature. *Journal of neurosurgery* 2011;**114**(1):9-18
29. Yilmaz T, Kaptanoglu E. Current and future medical therapeutic strategies for the functional repair of spinal cord injury. *World j* 2015;**6**(1):42-55 doi: <http://dx.doi.org/10.5312/wjo.v6.i1.42>[published Online First: Epub Date]].
30. Bennett MH, Trytko B, Jonker B. Hyperbaric oxygen therapy for the adjunctive treatment of traumatic brain injury. *The Cochrane database of systematic reviews* 2012;**12**:Cd004609 doi: 10.1002/14651858.CD004609.pub3[published Online First: Epub Date]].
31. Cadth. Optimal oxygen saturation range for adults suffering from traumatic brain injury: a review of patient benefit, harms, and guidelines (Structured abstract). *Health Technology Assessment Database* 2014; (4). <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD004609.pub3>[published Online First: Epub Date]].
32. Rincon F, Kang J, Vibbert M, et al. Significance of arterial hyperoxia and relationship with case fatality in traumatic brain injury: a multicentre cohort study. *Journal of neurology, neurosurgery, and psychiatry* 2014;**85**(7):799-805 doi: 10.1136/jnnp-2013-305505[published Online First: Epub Date]].
33. Skattum J, Naess PA, Gaarder C. Non-operative management and immune function after splenic injury. *The British journal of surgery* 2012;**99** **Suppl 1**:59-65 doi: 10.1002/bjs.7764[published Online First: Epub Date]].
34. London JA, Parry L, Galante J, et al. Safety of Early Mobilization of Patients With Blunt Solid Organ Injuries. *Arch Surg* 2008;**143**(10):972-76

## References for eTable 2d

1. Bansal R, Craigen MA. Fifth metacarpal neck fractures: is follow-up required? The Journal of hand surgery, European volume 2007;**32**(1):69-73 doi: 10.1016/j.jhsb.2006.09.021[published Online First: Epub Date]].
2. Gamble D, Jenkins PJ, Edge MJ, et al. Satisfaction and functional outcome with "self-care" for the management of fifth metacarpal fractures. Hand 2015;**10**(4):607-12 doi: <http://dx.doi.org/10.1007/s11552-015-9749-8>[published Online First: Epub Date]].
3. Finger A, Teunis T, Hageman MG, et al. Do patients prefer optional follow-up for simple upper extremity fractures: A pilot study. Injury 2016;**47**(10):2276-82 doi: <http://dx.doi.org/10.1016/j.injury.2016.06.029>[published Online First: Epub Date]].
4. Ferguson KB, McGlynn J, Jenkins P, et al. Fifth metatarsal fractures - Is routine follow-up necessary? Injury 2015;**46**(8):1664-8 doi: 10.1016/j.injury.2015.05.041[published Online First: Epub Date]].
5. Jayaram PR, Bhattacharyya R, Jenkins PJ, et al. A new "virtual" patient pathway for the management of radial head and neck fractures. Journal of shoulder and elbow surgery 2014;**23**(3):297-301 doi: 10.1016/j.jse.2013.11.006[published Online First: Epub Date]].
6. Murphy SM, Whately K, Eadie PA, et al. Unnecessary inter-hospital referral of minor hand injuries: a continuing problem. Irish J Med Sci 2010;**179**(1):123-25
7. Hartzell TL, Kuo P, Eberlin KR, et al. The overutilization of resources in patients with acute upper extremity trauma and infection. The Journal of hand surgery 2013;**38**(4):766-73 doi: 10.1016/j.jhsa.2012.12.016[published Online First: Epub Date]].
8. Randsborg PH. Fractures in children: Aspects on health service, epidemiology and risk factors. Acta Orthopaedica 2013;**84**(SUPPL.350):1-24 doi: <http://dx.doi.org/10.3109/17453674.2013.789731>[published Online First: Epub Date]].

9. Adamich JS, Camp MW. Do toddler's fractures of the tibia require evaluation and management by an orthopaedic surgeon routinely? *European journal of emergency medicine : official journal of the European Society for Emergency Medicine* 2017 doi: 10.1097/mej.0000000000000478[published Online First: Epub Date]].
10. Tufescu T. The cost of screening radiographs after stable fracture fixation. *Canadian journal of surgery Journal canadien de chirurgie* 2017;**60**(1):53-56
11. Fitzgerald E, Boran S. Management of torus fractures of the distal radius: A review of the literature. *Irish J Med Sci* 2016;**1**:S136 doi: <http://dx.doi.org/10.1007/s11845-015-1396-0>[published Online First: Epub Date]].
12. Jaremko JL, Lambert RG, Rowe BH, et al. Do radiographic indices of distal radius fracture reduction predict outcomes in older adults receiving conservative treatment? *Clinical radiology* 2007;**62**(1):65-72 doi: 10.1016/j.crad.2006.08.013[published Online First: Epub Date]].
13. Rooke G, Phillips FT. Does early radiography alter remanipulation rates in paediatric forearm fractures? *ANZ journal of surgery* 2015;**85**(1-2):38-43 doi: <http://dx.doi.org/10.1111/ans.12755>[published Online First: Epub Date]].
14. Tuomilehto N, Kivisaari R, Sommarhem A, et al. Outcome after pin fixation of supracondylar humerus fractures in children: postoperative radiographic examinations are unnecessary. *Acta Orthopaedica* 2016:1-7 doi: <http://dx.doi.org/10.1080/17453674.2016.1250058>[published Online First: Epub Date]].
15. Tufescu T. Working toward reducing postoperative fracture radiographs: a survey of Canadian surgeons. *Can J Surg* 2016;**59**(1):26-28
16. Chaudhry S, DeSole EM, Egol KA. Post-splinting radiographs of minimally displaced fractures: good medicine or medicolegal protection? *J Bone Joint Surg Am* 2012;**94**(17):e128
17. Schuld JC, Volker ML, Anderson SA, et al. Post-splinting x-rays of non-displaced hand, wrist, ankle, and foot fractures are unnecessary. *Academic Emergency Medicine* 2016;**23**:S148 doi: <http://dx.doi.org/10.1111/acem.12974>[published Online First: Epub Date]].

18. Unay K, Gokcen B, Ozkan K, et al. Examination tests predictive of bone injury in patients with clinically suspected occult scaphoid fracture. *Injury* 2009;**40**(12):1265-8 doi: 10.1016/j.injury.2009.01.140[published Online First: Epub Date]].
19. Pyun JS, Weir TB, Banagan KE, et al. The Utility of In-Hospital Postoperative Radiographs following Surgical Treatment of Traumatic Thoracolumbar Injuries. *The Spine Journal* 2017;**16**(10):S187-S88
20. Dunn JC, Kusnezov N, Orr JD, et al. The Boxer's Fracture: Splint Immobilization Is Not Necessary. *Orthopedics* 2016;**39**(3):188-92 doi: 10.3928/01477447-20160315-05[published Online First: Epub Date]].
21. Pincus S, Weber M, Meakin A, et al. Introducing a Clinical Practice Guideline Using Early CT in the Diagnosis of Scaphoid and Other Fractures. *The western journal of emergency medicine* 2009;**10**(4):227-32
22. Nirav KP, Davies N, Mirza Z, et al. Cost and clinical effectiveness of MRI in occult scaphoid fractures: A randomised controlled trial. *Emergency medicine journal* 2013; 30(3). <http://onlinelibrary.wiley.com/doi/10.1111/1524-1221.12111>
23. McFadyen I, Field J, McCann P, et al. Should unstable extra-articular distal radial fractures be treated with fixed-angle volar-locked plates or percutaneous Kirschner wires? A prospective randomised controlled trial. *Injury* 2011;**42**(2):162-6 doi: 10.1016/j.injury.2010.07.236[published Online First: Epub Date]].
24. Schepers T, Van Lieshout EM, de Vries MR, et al. Complications of syndesmotic screw removal. *Foot & ankle international* 2011;**32**(11):1040-4 doi: 10.3113/fai.2011.1040[published Online First: Epub Date]].
25. Tucker A, Street J, Kealey D, et al. Functional outcomes following syndesmotic fixation: A comparison of screws retained in situ versus routine removal - Is it really necessary? *Injury* 2013;**44**(12):1880-4 doi: 10.1016/j.injury.2013.08.011[published Online First: Epub Date]].
26. Dingemans SA, Rammelt S, White TO, et al. Should syndesmotic screws be removed after surgical fixation of unstable ankle fractures? a systematic review. *The bone & joint journal* 2016;**98-b**(11):1497-504 doi: 10.1302/0301-620x.98b11.bjj-2016-0202.r1[published Online First: Epub Date]].

27. Nestorson J, Josefsson PO, Adolfsson L. A radial head prosthesis appears to be unnecessary in Mason-IV fracture dislocation. *Acta Orthop* 2017;**88**(3):315-19 doi: 10.1080/17453674.2017.1293440[published Online First: Epub Date]].
28. Webb GR, Galpin RD, Armstrong DG. Comparison of short and long arm plaster casts for displaced fractures in the distal third of the forearm in children. *J Bone Joint Surg Am* 2006;**88**(1):9-17 doi: 10.2106/jbjs.e.00131[published Online First: Epub Date]].
29. Boutis K, Willan AR, Babyn P, et al. A randomized, controlled trial of a removable brace versus casting in children with low-risk ankle fractures. *Pediatrics* 2007;**119**(6):e1256-63 doi: 10.1542/peds.2006-2958[published Online First: Epub Date]].
30. Yeung DE, Jia X, Miller CA, et al. Interventions for treating ankle fractures in children. *Cochrane Database of Systematic Reviews* 2016;**4**:CD010836 doi: <http://dx.doi.org/10.1002/14651858.CD010836.pub2>[published Online First: Epub Date]].
31. Harrop JS. Type II Odontoid Fractures: What to Do? *World Neurosurg* 2013;**80**(3-4):313-14
32. Gao Y, Zhang W, Duan X, et al. Surgical interventions for treating radial head fractures in adults. *Cochrane Database of Systematic Reviews* 2013; (5). <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD008987.pub2/abstract>.
33. Ruchelsman DE, Christoforou D, Jupiter JB. Fractures of the radial head and neck. *J Bone Joint Surg Am* 2013;**95**(5):469-78 doi: 10.2106/jbjs.j.01989[published Online First: Epub Date]].
34. Boons HW, Goosen JH, van Grinsven S, et al. Hemiarthroplasty for humeral four-part fractures for patients 65 years and older: a randomized controlled trial. *Clinical orthopaedics and related research* 2012;**470**(12):3483-91 doi: 10.1007/s11999-012-2531-0[published Online First: Epub Date]].
35. Bhat AK, Rao SK, Bhaskaranand K. Mechanical failure in intramedullary interlocking nails. *Journal of orthopaedic surgery (Hong Kong)* 2006;**14**(2):138-41 doi: 10.1177/230949900601400206[published Online First: Epub Date]].
36. Yi L, Jingping B, Gele J, et al. Operative versus non-operative treatment for thoracolumbar burst fractures without neurological deficit. *The Cochrane database of systematic reviews* 2006(4):Cd005079 doi: 10.1002/14651858.CD005079.pub2[published Online First: Epub Date]].




37. Moller A, Hasserijs R, Redlund-Johnell I, et al. Nonoperatively treated burst fractures of the thoracic and lumbar spine in adults: a 23- to 41-year follow-up. *The spine journal : official journal of the North American Spine Society* 2007;**7**(6):701-7 doi: 10.1016/j.spinee.2006.09.009[published Online First: Epub Date]].
38. Wang ST, Ma HL, Liu CL, et al. Is fusion necessary for surgically treated burst fractures of the thoracolumbar and lumbar spine?: a prospective, randomized study. *Spine* 2006;**31**(23):2646-52; discussion 53 doi: 10.1097/01.brs.0000244555.28310.40[published Online First: Epub Date]].
39. Sargin S, Uçar BY, Necmioglu S, et al. Clinical and radiological results of posterior instrumentation without fusion for thoracolumbar fractures. *African Journal of Pharmacy and Pharmacology* 2011;**5**(7):pp. 819-22 doi: DOI: 10.5897/AJPP11.135[published Online First: Epub Date]].
40. Hwang JU, Hur JW, Lee JW, et al. Comparison of Posterior Fixation Alone and Supplementation with Posterolateral Fusion in Thoracolumbar Burst Fractures. *J Korean Neurosurg Soc* 2012;**52**(4):346-52
41. Jindal N, Sankhala SS, Bachhal V. The role of fusion in the management of burst fractures of the thoracolumbar spine treated by short segment pedicle screw fixation A prospective randomised trial. *J Bone Joint Surg-Br Vol* 2012;**94B**(8):1101-06
42. Egol KA, Paksima N, Puopolo S, et al. Treatment of external fixation pins about the wrist: a prospective, randomized trial. *J Bone Joint Surg Am* 2006;**88**(2):349-54 doi: 10.2106/jbjs.e.00011[published Online First: Epub Date]].
43. Camathias C, Valderrabano V, Oberli H. Routine pin tract care in external fixation is unnecessary: A randomised, prospective, blinded controlled study. *Injury-Int J Care Inj* 2012;**43**(11):1969-73

# Low-value clinical practices in the intensive care unit

Thanks for your participation in the **consultation phase** of our **scoping review**, the first component of the **Canadian Program on Low Value Practices in Injury Care**.

**FIGURE 1. THE CANADIAN PROGRAM ON LOW-VALUE PRACTICES IN INJURY CARE**



The objective of this survey is to **identify around 10 clinical practices** that will go on to the systematic review phase. To do so, we would like you to **rate each intervention on its potential to be labeled as a low-value clinical practice** according to the following **definition**:

**An intervention that is used in practice but is ineffective or its harm/cost outweighs its benefits**

There are 53 questions in this survey

## **ICU ADMISSION AND NEUROSURGICAL CONSULTATION (page 1/7)**

Low-value clinical practice (definition): **An intervention that is used in practice but is ineffective or its harm/cost outweighs its benefits.**

**We would like you to rate each intervention on its potential to be labeled as a low-value clinical practice according to the following definition:**

**An intervention that is used in practice but is ineffective or its harm/cost outweighs its benefits**

### **1. ICU ADMISSION**

**Indication: Adult acute mild TBI (GCS 13-15) with minimal findings on CT and not on nonreversible anticoagulation**

Please choose the appropriate response for each item:

Clearly low-value      Possibly low-value      Controversial      Possibly beneficial      Clearly beneficial      Undecided

### **2. ICU ADMISSION**

**Indication: Pediatric acute mild TBI (GCS 13-15) with minimal findings on CT but no midline shift or depressed skull fracture**

Please choose the appropriate response for each item:

Clearly low-value      Possibly low-value      Controversial      Possibly beneficial      Clearly beneficial      Undecided

### **3. NEUROSURGICAL CONSULTATION**

**Indication: Acute mild TBI with negative CT**

Please choose the appropriate response for each item:

Clearly low-value      Possibly low-value      Controversial      Possibly beneficial      Clearly beneficial      Undecided

### **4. NEUROSURGICAL CONSULTATION**

**Indication: Acute mild TBI with minimal findings on CT**

Please choose the appropriate response for each item:

Clearly low-value      Possibly low-value      Controversial      Possibly beneficial      Clearly beneficial      Undecided

**Comments:**

Please write your answer(s) here:

Question 1	
Question 2	
Question 3	
Question 4	

ACCEPTED