

This is a repository copy of *An odd oxygen framework for wintertime ammonium nitrate aerosol pollution in urban areas: NO<sub>x</sub> and VOC control as mitigation strategies*.

White Rose Research Online URL for this paper:  
<https://eprints.whiterose.ac.uk/145858/>

Version: Published Version

---

**Article:**

Womack, C. C., McDuffie, E. E., Edwards, Peter [orcid.org/0000-0002-1076-6793](https://orcid.org/0000-0002-1076-6793) et al. (27 more authors) (2019) An odd oxygen framework for wintertime ammonium nitrate aerosol pollution in urban areas: NO<sub>x</sub> and VOC control as mitigation strategies. *Geophysical Research Letters*. ISSN 0094-8276

<https://doi.org/10.1029/2019GL082028>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Geophysical Research Letters

## RESEARCH LETTER

10.1029/2019GL082028

### Key Points:

- Wintertime ammonium nitrate aerosol pollution is closely tied to photochemical ozone production through a common parameter,  $O_{x,total}$
- Box modeling reveals ammonium nitrate formation in the Salt Lake Valley is nitrate-limited but  $NO_x$ -saturated
- Mitigation strategies that focus on  $NO_x$  control in some wintertime-polluted layers may initially increase ammonium nitrate

### Supporting Information:

- Supporting Information S1

### Correspondence to:

S. S. Brown,  
steven.s.brown@noaa.gov

### Citation:






















Womack, C. C., McDuffie, E. E., Edwards, P. M., Bares, R., de Gouw, J. A., Docherty, K. S., et al. (2019). An odd oxygen framework for wintertime ammonium nitrate aerosol pollution in urban areas:  $NO_x$  and VOC control as mitigation strategies. *Geophysical Research Letters*, 46. <https://doi.org/10.1029/2019GL082028>

Received 11 JAN 2019

Accepted 3 APR 2019

Accepted article online 8 APR 2019

## An Odd Oxygen Framework for Wintertime Ammonium Nitrate Aerosol Pollution in Urban Areas: $NO_x$ and VOC Control as Mitigation Strategies

C. C. Womack<sup>1,2</sup> , E. E. McDuffie<sup>1,2,3,4</sup> , P. M. Edwards<sup>5</sup> , R. Bares<sup>6</sup> , J. A. de Gouw<sup>2,3</sup> , K. S. Docherty<sup>7</sup>, W. P. Dubé<sup>1,2</sup>, D. L. Fibiger<sup>1,2,8</sup> , A. Franchin<sup>1,2</sup> , J. B. Gilman<sup>1</sup> , L. Goldberger<sup>9,10</sup>, B. H. Lee<sup>9</sup> , J. C. Lin<sup>6</sup> , R. Long<sup>11</sup>, A. M. Middlebrook<sup>1</sup>, D. B. Millet<sup>12</sup> , A. Moravek<sup>13,14</sup> , J. G. Murphy<sup>13</sup> , P. K. Quinn<sup>15</sup>, T. P. Riedel<sup>11</sup> , J. M. Roberts<sup>1</sup> , J. A. Thornton<sup>9</sup> , L. C. Valin<sup>11</sup> , P. R. Veres<sup>1</sup> , A. R. Whitehill<sup>11</sup>, R. J. Wild<sup>1,2,16</sup> , C. Warneke<sup>1,2</sup> , B. Yuan<sup>1,2,17</sup>, M. Baasandorj<sup>6,18</sup>, and S. S. Brown<sup>1,3</sup> 


<sup>1</sup>Chemical Sciences Division, NOAA Earth System Research Laboratory, Boulder, CO, USA, <sup>2</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA, <sup>3</sup>Department of Chemistry, University of Colorado Boulder, Boulder, CO, USA, <sup>4</sup>Now at the Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Nova Scotia, Canada, <sup>5</sup>Department of Chemistry, University of York, York, UK, <sup>6</sup>Department of Atmospheric Sciences, University of Utah, Salt Lake City, UT, USA, <sup>7</sup>Jacobs Technology, Inc., Durham, NC, USA, <sup>8</sup>Now at California Air Resources Board, Sacramento, CA, USA, <sup>9</sup>Department of Atmospheric Sciences, University of Washington, Seattle, WA, USA, <sup>10</sup>Now at ARM Aerial Facility, Pacific Northwest National Laboratory, Richland, WA, USA, <sup>11</sup>Office of Research and Development, Environmental Protection Agency, Durham, NC, USA, <sup>12</sup>Department of Soil, Water, and Climate, University of Minnesota, Twin Cities, St. Paul, MN, USA, <sup>13</sup>Department of Chemistry, University of Toronto, Toronto, Ontario, Canada, <sup>14</sup>Now at Department of Chemistry, York University, Toronto, Ontario, Canada, <sup>15</sup>NOAA Pacific Marine Environmental Laboratory, Seattle, WA, USA, <sup>16</sup>Now at the Institut für Ionenphysik und Angewandte Physik, Universität Innsbruck, Innsbruck, Austria, <sup>17</sup>Now at the Institute for Environmental and Climate Research, Jinan University, Guangzhou, China, <sup>18</sup>Division of Air Quality, Utah Department of Environmental Quality, Salt Lake City, UT, USA

**Abstract** Wintertime ammonium nitrate aerosol pollution is a severe air quality issue affecting both developed and rapidly urbanizing regions from Europe to East Asia. In the United States, it is acute in western basins subject to inversions that confine pollutants near the surface. Measurements and modeling of a wintertime pollution episode in Salt Lake Valley, Utah, demonstrate that ammonium nitrate is closely related to photochemical ozone through a common parameter, total odd oxygen,  $O_{x,total}$ . We show that the traditional nitrogen oxide and volatile organic compound ( $NO_x$ -VOC) framework for evaluating ozone mitigation strategies also applies to ammonium nitrate. Despite being nitrate-limited, ammonium nitrate aerosol pollution in Salt Lake Valley is responsive to VOCs control and, counterintuitively, not initially responsive to  $NO_x$  control. We demonstrate simultaneous nitrate limitation and  $NO_x$  saturation and suggest this phenomenon may be general. This finding may identify an unrecognized control strategy to address a global public health issue in regions with severe winter aerosol pollution.

**Plain Language Summary** Particulate matter (PM) is dangerous to human health and impacts visibility and climate. In the United States, Europe, and Asia, PM is severe in urban areas in the winter when ammonium nitrate,  $NH_4NO_3$ , comprises an appreciable fraction of the total PM mass. A key control strategy is to reduce emissions of the limiting reagent. Using measurements from a recent field campaign in the Salt Lake Valley, Utah, which experiences high PM levels in winter, we demonstrate that emission control strategies can be evaluated using the same framework commonly used to control ozone, another common pollutant that occurs at high levels in urban areas in the summer. We show that initial control of the  $NO_x$  precursor is ineffective at reducing  $NH_4NO_3$  aerosol in the Salt Lake Valley, while initial control of volatile organic compounds, which are not a direct precursor for either nitrate or ammonium, is effective due to their influence on oxidation cycles. This finding differs from many mitigation strategies in the western United States and may also be relevant to other regions in Europe and Asia which experience high wintertime PM.

## 1. Introduction

Air quality in most of the United States has been steadily improving for decades due to emissions control technologies and regulations (Hand et al., 2014). Ozone ( $O_3$ ), which is a health hazard and greenhouse gas in the troposphere, has decreased markedly (Cooper et al., 2014) and is well understood in terms of its photochemical formation mechanism from nitrogen oxides ( $NO_x = NO + NO_2$ ) and volatile organic compounds (VOCs; Lin et al., 1988; Seinfeld, 1989). In contrast, many urban areas, such as the Salt Lake Valley (SLV) and San Joaquin Valley in the western United States, the Po Valley in Europe, and Beijing, China, in east Asia, experience severe wintertime pollution in the form of particulate matter smaller than 2.5 microns in diameter ( $PM_{2.5}$ ; Green et al., 2015; Figure 1). The SLV in northern Utah regularly experiences episodes of  $PM_{2.5}$  pollution (Silcox et al., 2012), exacerbated by the onset of persistent cold air pools (PCAPs; high-pressure systems that cap colder air within the basin), which prevent the vertical mixing of pollutants away from the surface (Whiteman et al., 2014). The trapped gas-phase pollutants, including  $NO_x$  and VOCs from anthropogenic sources, undergo complex chemical oxidation reactions, forming nitric acid ( $HNO_3$ ) that reacts with gas-phase ammonia ( $NH_3$ ) to form ammonium nitrate aerosol ( $NH_4NO_3$ ), which comprises ~75% of the  $PM_{2.5}$  composition in the SLV during pollution episodes (Baasandorj et al., 2017; Franchin et al., 2018; Kuprov et al., 2014).




The SLV exceeds the 24-hr U.S. National Ambient Air Quality Standard for  $PM_{2.5}$  ( $35 \mu g/m^3$ ) on an average of 18 days/year (Whiteman et al., 2014), representing an important public health issue (Beard et al., 2012). A better understanding of the chemical processes that lead to the formation of ammonium nitrate and its precursors in the SLV and other polluted areas across the world is required to employ more effective control strategies.

Previous studies have demonstrated that during PCAP-induced pollution events, the valleys in the Salt Lake region typically have an excess of ammonia (Franchin et al., 2018; Kelly et al., 2013; Kuprov et al., 2014), and that the cold temperatures and high relative humidity drive the reversible reaction (R1) to form particulate ammonium nitrate (abbreviated  $pNO_3^-$ ; Blanchard et al., 2000; Franchin et al., 2018; Weber et al., 2016). The recent analysis of Franchin et al. (2018) demonstrates that the observed ratio between gas-phase nitric acid and total nitrate,  $HNO_3/(HNO_3 + pNO_3^-)$ , has a median value of <8% during PCAP episodes, and the calculated aerosol pH curve is insensitive to small changes in pH (supporting information Figure S1). Furthermore, Franchin et al. (2018) show that the initial response of total aerosol mass to reductions in total nitrate ( $HNO_3 + pNO_3^-$ ) is linear but substantially less than linear in its initial response to reductions in total reduced nitrogen ( $NH_3 + NH_4^+$ ). Therefore, the formation rate of  $pNO_3^-$  is mainly limited by the chemical formation rate of nitric acid ( $HNO_3$ ) from oxidation of  $NO_x$ , which is emitted from fossil fuel combustion. This  $HNO_3$  limitation has led typical western U.S. control strategies to focus on emission reductions of  $NO_x$  rather than  $NH_3$  (Franchin et al., 2018; Kelly et al., 2018; Pusede et al., 2016). We demonstrate in this study that ammonium nitrate aerosol pollution may be treated in the same manner as  $O_3$  pollution through a common parameter,  $O_{x,total}$ , and may be most effectively reduced in the SLV by initially controlling VOC emissions rather than controlling emissions of  $NH_3$  or  $NO_x$ .

## 2. Methods

### 2.1. Observations of Wintertime Pollution Episodes in Northern Utah

The Utah Winter Fine Particulate Study (UWFPS) was a 4-week field campaign in the SLV region during the winter of 2016–2017 designed to observe the chemical processes that form ammonium nitrate aerosol during PCAPs (Baasandorj et al., 2018; Bares et al., 2018; Franchin et al., 2018). Ground sites made continuous surface-level measurements of trace gases (including  $O_3$ ,  $NO_x$ , and VOCs) and the chemical composition of  $PM_{2.5}$ , while a Twin Otter light aircraft made periodic measurements aloft with a similar payload. This analysis primarily uses observations from the main SLV site at the University of Utah (UU), with auxiliary measurements from the Hawthorne site on the SLV floor and the Logan site in the nearby Cache Valley (Figure S2). Two major PCAP episodes occurred during the campaign, in which the 24-hr  $PM_{2.5}$  concentrations surpassed the National Ambient Air Quality Standard in at least one of the three major valleys for a



**Figure 1.** Seasonal trends in  $PM_{2.5}$  nonattainment in three regions with high ammonium nitrate aerosol content. (a) Yearly profiles of  $PM_{2.5}$  measured in the Salt Lake Valley, United States, between 1999 and 2017, with the average shown in red. Virtually all exceedances of the U.S. National Ambient Air Quality Standard ( $35 \mu\text{g}/\text{m}^3$ ; dashed line) occur during the winter. (b) The fraction of winter and summer days in nonattainment of local air quality standards for  $PM_{2.5}$  in the Salt Lake Valley (EPA, 2018), Ispra, Italy in the Po Valley (EMEP, 2018), and Beijing, China (DOS, 2018; see Text S1); these three regions are noted for their poor wintertime air quality and high ammonium nitrate aerosol mass loading. In the Salt Lake Valley and Po Valley, summertime exceedances are rare, while wintertime exceedances are persistent. In China, summertime exceedances have decreased in recent years, while wintertime exceedances remain frequent.  $PM_{2.5}$  = particulate matter smaller than 2.5 microns in diameter.

parameter,  $O_{x,\text{total}}$ , which is equal to the sum of all gas- and particulate-phase species that contain an odd oxygen (O or  $O_3$ ).

$$O_{x,\text{total}} = O_3 + NO_2 + 2 \times NO_3 + 3 \times N_2O_5 + ClNO_2 + \text{peroxyacyl nitrates} + 2 \times \text{alkyl nitrates} + OH + 1.5 \times (HNO_3 + pNO_3^-), \quad (R2)$$

where the stoichiometric coefficients indicate the number of odd oxygen species it contains and are derived in Text S2.  $NH_3$  is reduced nitrogen and is therefore not a component of  $O_{x,\text{total}}$ . The coefficient for the sum of gas-phase  $HNO_3$  and  $pNO_3^-$  is denoted here as 1.5 but may be as high as 2, depending on the mechanism that formed it, and therefore,  $O_{x,\text{total}}$  as defined here should be considered a lower bound.  $O_{x,\text{total}}$  may be more broadly defined than (R2) if one considers the products of the reactions of OH,  $O_3$ , and  $NO_3$  with VOCs, for example, but for this analysis, we focus only on the propagation of odd oxygen through nitrogen chemistry.

The nighttime mechanism results in no net change in  $O_{x,\text{total}}$ , as those reactions represent only a repartitioning of its components. The daytime mechanism, on the other hand, generates one  $NO_2$  molecule for each cycle without loss of an  $O_3$  molecule and therefore represents an increase in  $O_{x,\text{total}}$  before terminating in  $HNO_3$  or alkyl nitrate production. Transport of air masses with background concentrations of  $O_3$  and negligible concentrations of the other  $O_{x,\text{total}}$  components into the SLV would deplete the other  $O_{x,\text{total}}$

total of 8 days. This analysis investigates the mechanisms for the  $PM_{2.5}$  production that lead to the exceedance during one of the most intense pollution episode of that winter, which occurred between 27 January and 4 February 2017 (Figure S2). These results are compared to observations of trace gas species from the recent Uintah Basin Winter Ozone Study, where a major PCAP in January 2013 led to elevated levels of  $O_3$  (Ahmadov et al., 2015; Edwards et al., 2014). Measurement techniques for both campaigns are described in detail in Text S1 (Lee et al., 2014; Liao et al., 2017; Wild et al., 2014; Wild et al., 2016).

## 2.2. $O_{x,\text{total}}$ as a Parameter for Describing both $HNO_3$ and $O_3$ Formation

There are two mechanisms for  $HNO_3$  formation: daytime production from the gas-phase reaction of  $OH + NO_2$  and nighttime production from  $N_2O_5$  uptake onto aerosol particles (Jones & Seinfeld, 1983). The full mechanistic details are listed in Text S2 and summarized here. In the daytime, VOCs are photolyzed or oxidized by the hydroxyl radical (OH) to form  $HO_2$  or  $RO_2$  radicals, collectively termed  $HO_x$  radicals ( $HO_x = OH + HO_2 + RO_2$ ). Radical propagation reactions with NO generate an  $NO_2$  molecule with each iteration, while the dominant radical termination reaction of OH with  $NO_2$  generates  $HNO_3$ . The overall  $HNO_3$  production efficiency of this cycle depends on (1) the availability of  $HO_x$  radicals, which are generally less abundant in winter than in summer, and (2) the number of iterations each  $HO_x$  radical makes prior to termination (i.e., chain length), which is determined by the relative abundance of VOC and  $NO_x$  (Kleinman, 2005; Lin et al., 1988). During the day,  $NO_2$  rapidly interconverts with NO and  $O_3$ , and therefore, the same  $HO_x$  cycle that produces  $HNO_3$  is also the dominant mechanism for  $O_3$  production. At night, photolytic production of radicals and  $O_3$  halts, and NO titrates the  $O_3$  to  $NO_2$ , which then oxidizes to form  $N_2O_5$ , which further reacts heterogeneously with aerosol to form  $HNO_3$  and  $ClNO_2$  (Kelly et al., 2018).

The parameter  $O_x (=O_3 + NO_2)$  has often been used to quantify the net photochemical formation of  $O_3$  beyond its rapid daytime interconversion with  $NO_2$  (Liu, 1977; Wood et al., 2009). Here we introduce an analogous

components by their stoichiometric coefficients ( $\geq 1$ ) for every  $O_3$  molecule added, leading to a net decrease of  $O_{x,total}$ . Dry deposition of individual species to the surface is also a net loss process. Therefore, growth in  $O_{x,total}$  beyond its background level in unpolluted air is a measure of the role that photochemistry plays in the net production of  $HNO_3 + pNO_3^-$  during the PCAP.


### 2.3. Photochemical Box Modeling

The Dynamically Simple Model for Atmospheric Chemical Complexity (DSMACC) (Emmerson & Evans, 2009), a photochemical zero-dimensional box model, was used to characterize the growth of  $O_{x,total}$  in the SLV during one of the most severe PCAPs sampled between 27 January and 4 February 2017. DSMACC utilizes the Master Chemical Mechanism (v3.3.1; Jenkin et al., 2015), a near-explicit mechanism to characterize the degradation of 143 VOCs by photochemistry and oxidation, using >15,000 reactions and >3,500 intermediate species. A subset of the Master Chemical Mechanism was used here, consisting of the 58 VOCs for which we had experimental constraints for the starting conditions. The Tropospheric Ultraviolet and Visible Radiation Model (v5.2; Madronich et al., 1998) was used to calculate the photolysis rate constants in the chemical mechanism. A zero-dimensional model treats the chemistry as a single system evolving in time and does not consider meteorology or boundary layer dynamics, but the relative simplicity of these models allows the use of a highly detailed chemical mechanism. The stagnant air and constrained meteorology in these valleys make this an appropriate technique for these conditions (Edwards et al., 2014). Further model details are described in Text S3 (de Gouw et al., 2017; Kurtenbach et al., 2001; McDuffie, 2018; Wesely & Hicks, 2000; Wild et al., 2017; Zhang et al., 2012).

## 3. Results


Figure 2a shows the most abundant components of  $O_{x,total}$  observed at the UU sampling site during the UWFPS campaign. As the PCAP progressed,  $O_{x,total}$  increased steadily at a rate of approximately 5.5 ppbv/day, and its composition quickly became dominated by  $NO_2$  and  $pNO_3^-$ . A similar trend is observed in Twin Otter vertical profiles (Figure 2b). These observations of  $O_{x,total}$  growth and partitioning are compared to a recent study in the nearby Uintah Basin that also experiences frequent wintertime PCAPs (Ahmadov et al., 2015). In contrast to SLV, the sparsely populated Uintah Basin in northeastern Utah experiences high wintertime levels of  $O_3$  (UBWOS, 2014) due to photochemical oxidation of VOCs emitted in abundance from local oil and natural gas operations in a relatively  $NO_x$ -poor environment (Edwards et al., 2014). Figure 2c shows a similar growth rate of  $O_{x,total}$  in the Uintah Basin in during a PCAP in January 2013, but unlike the SLV,  $O_{x,total}$  in the Uintah Basin is partitioned nearly entirely to  $O_3$ . The SLV is densely populated and contains no oil and natural gas wells, though it does have significant agricultural and industrial sectors, and  $NO_x$  and VOC emission profiles typical of U.S. urban areas (Figure S3a). The closely related  $O_{x,total}$  growth in Figure 2 is reflective of the same photochemical mechanisms playing important roles in both valleys (Kleeman et al., 2005; Meng et al., 1997; Nguyen & Dabdub, 2002), and the widely divergent partitioning of its components to either  $pNO_3^-$  in the SLV or  $O_3$  in the Uintah Basin derives from their starkly different emission profiles (Figure S3b). The common parameter  $O_{x,total}$  can therefore be used to evaluate mitigation strategies for controlling both criteria pollutants, as demonstrated below.

The production of tropospheric  $O_3$ , frequently the major component of  $O_{x,total}$ , is often evaluated in terms of its nonlinear sensitivity to  $NO_x$  and VOC emissions (Lin et al., 1988; Milford et al., 1989; Seinfeld, 1989; Trainer et al., 1993). Because small changes in the availability of  $NO_x$  can change the relative branching of the  $HO_x$  radical reaction pathways, the  $O_3$  production rate and yield may either increase or decrease with  $NO_x$  availability, depending on the VOC concentration and composition (Lin et al., 1988), which often requires detailed modeling to characterize fully. Edwards et al. (2014) used a zero-dimensional box model to show that  $O_3$  production in the Uintah Basin was close to its maximum efficiency with respect to  $NO_x$  at the prevailing VOC concentrations in winter. Here we develop an analogous model to explain the growth of  $O_{x,total}$  in the SLV. We modified the DSMACC box model to split into two concurrent boxes at night to model the residual layer (RL) and nocturnal boundary layer (NBL) separately, to more accurately model the vertical structure of the PCAP, as shown in Figure 3a.




**Figure 2.**  $O_{x,total}$  measured during the Utah Winter Fine Particulate Study 2017 and Uintah Basin Winter Ozone Study 2013 campaigns. (a)  $O_{x,total}$  measured at the University of Utah ground site in the Salt Lake Valley during the UWFPS campaign with the major measured components of  $O_{x,total}$  in filled colors.  $O_3$  and  $NO_2$  were the only components of  $O_{x,total}$  directly measured at University of Utah, so  $pNO_3^-$  was calculated from the total particulate matter smaller than 2.5 microns in diameter measurement. The other gas-phase components were estimated to be minor contributors. When the persistent cold air pool (PCAP) begins on 26 January, a steady increase in  $O_{x,total}$  is observed, along with a change in its partitioning, with  $NO_2$  and  $pNO_3^-$  dominating and  $O_3$  being depleted. (b) A vertical profile of the boundary layer on 28 January (denoted by blue circle in (a) from the Twin Otter. The height of the boundary layer is approximately 800-m above ground level, and  $O_{x,total}$  remains at 50 ppbv above that height and grows well in excess of that budget beneath it. (c) The  $O_{x,total}$  measured during the 2013 Uintah Basin Winter Ozone Study campaign in Uintah Basin, with some missing data due to instrument downtime. The PCAP begins on 29 January 2013 and persists for 9 days. In stark contrast to the Salt Lake Valley,  $O_3$  represents approximately 85% of the  $O_{x,total}$  throughout the PCAP in the Uintah Basin.

Figure 3b shows the base case model compared with the observations. The base case was selected based on daytime agreement between modeled and observed  $O_{x,total}$ , VOCs, and  $NO_x$ , as detailed in Text S3. The model did not attempt to replicate the partitioning of  $HNO_3$  between the gas and aerosol phase and simply treated the two as the lumped sum  $HNO_3 + pNO_3^-$ . Franchin et al. (2018) showed this partitioning to be >92% to the aerosol phase during UWFPS PCAP events. The UU site is located at slightly higher elevation



**Figure 3.** Results of the photochemical box model of  $O_{x,total}$  growth. (a) An idealized representation of the daily evolution of the pollution layer as implemented in the model. The gray dashed line shows the top of the capping inversion layer. The black line shows the nighttime division of the layer into a NBL and a RL. During the day, weak solar heating convectively mixes the two layers (black dashed lines). The elevation of the three ground sites above the SLV floor and the Twin Otter's cruising altitude range are on the right. (b) Comparison between model prediction and the UU observations of  $O_{x,total}$ . Emission rates were tuned such that the daytime (yellow rectangles) agreement between the modeled and observed  $O_{x,total}$  was optimized. The location of the UU site on the valley wall is consistent with the nighttime measurement lying between the prediction for the RL and the NBL. SLV = Salt Lake Valley; UU = University of Utah.



**Figure 4.** Isopleths of O<sub>3</sub>, O<sub>x,total</sub>, and HNO<sub>3</sub> + pNO<sub>3</sub><sup>-</sup> formation. (a) Isopleths of the predicted maximum daytime O<sub>3</sub>, O<sub>x,total</sub>, and HNO<sub>3</sub> + pNO<sub>3</sub><sup>-</sup> on the final day of the model in the Salt Lake Valley, as a function of normalized NO<sub>x</sub> and VOC emissions. The black circles indicate the base case, and the contours show the predicted maximum concentration of each species as the normalized emissions are varied between 0 and 2. (b) Cuts through each isopleth at the base case, with normalized NO<sub>x</sub> and VOC emissions of 1. VOC = volatile organic compound.

on the side of the valley wall and may sample from either the NBL or RL at night at different times during the PCAP (see Text S3.1). Additionally, that site may be subject to upslope and downslope canyon flows along the basin walls (Baasandorj et al., 2017). Agreement between the model and UU observations was therefore only expected during daytime hours when the pollution layer is mixed. The split box model successfully captures the overall growth of O<sub>x,total</sub> over the four model days, accurately predicts that there is O<sub>3</sub> depletion in the NBL due to dilution and reaction with NO<sub>2</sub> but not in the RL, and reproduces the observed partitioning of O<sub>x,total</sub> (Figure S4).

The input emission rates may be adjusted from the model base case to determine the effect of various emission control strategies. Isopleths determined in this way are often used to predict the effect of emission controls on O<sub>3</sub> production (Kleinman, 2005; Seinfeld, 1989). VOC controls on PM<sub>2.5</sub> formation have been considered in the past in California (Kleeman et al., 2005; Nguyen & Dabdub, 2002; Pun & Seigneur, 2001). However, VOC controls in the context of the chemical equivalence of O<sub>3</sub> and pNO<sub>3</sub><sup>-</sup> in terms of O<sub>x,total</sub> during weak photochemical environments, such as the SLV, have not been considered. Calculated isopleths of O<sub>3</sub>, O<sub>x,total</sub>, and the sum of HNO<sub>3</sub> + pNO<sub>3</sub><sup>-</sup> are shown in Figure 4. They indicate that the SLV is in a NO<sub>x</sub>-saturated regime, where initial decreases in NO<sub>x</sub> emissions will increase O<sub>3</sub>, O<sub>x,total</sub>, and HNO<sub>3</sub> + pNO<sub>3</sub><sup>-</sup>, absent concurrent VOC reductions.

NO<sub>x</sub> saturation in the SLV is attributed to the modeled average HO<sub>x</sub> chain length or the number of times a HO<sub>x</sub> radical undergoes a propagation reaction cycle generating NO<sub>2</sub> before terminating (Jeffries & Tonnesen, 1994; Mao et al., 2010). The model predicts this value to be 1.07 in the SLV (Figure S5), so HO<sub>x</sub> is near its peak efficiency for HNO<sub>3</sub> production, as over 99% of HO<sub>x</sub> radicals terminate in HNO<sub>3</sub>, rather than an alkyl nitrate or HO<sub>x</sub> self-reaction. This observation is also consistent with the low observed aerosol-phase organic nitrates (Franchin et al., 2018). NO<sub>x</sub> serves to quench the HO<sub>x</sub>-NO<sub>x</sub> cycle, and therefore, reductions in NO<sub>x</sub> increase the HO<sub>x</sub> chain length and drive more efficient production of O<sub>3</sub> and HNO<sub>3</sub>. In the Uintah Basin, on the other hand, the average chain length is 4.3, indicating that HO<sub>x</sub> radicals are highly efficient in generating O<sub>3</sub> before termination largely via self-reaction. The ratio of NO<sub>x</sub> to VOCs (Figure S3b) is a key parameter that dictates the ultimate fate of radicals: High NO<sub>x</sub>/VOCs yields high HNO<sub>3</sub> in the SLV, whereas low NO<sub>x</sub>/VOC yields high O<sub>3</sub> in Uintah Basin.

It is often assumed that when pNO<sub>3</sub><sup>-</sup> production is HNO<sub>3</sub>-limited (as opposed to NH<sub>3</sub>-limited), HNO<sub>3</sub> production must be NO<sub>x</sub>-limited (Guo et al., 2018; Wen et al., 2018), and previous studies in California's San

Joaquin Valley have found that >50% NO<sub>x</sub> reductions would reduce pNO<sub>3</sub><sup>-</sup> (Kelly et al., 2018; Pusede et al., 2016). However, Figure 4 demonstrates that the SLV is HNO<sub>3</sub>-limited and NO<sub>x</sub>-saturated simultaneously (see also Pun & Seigneur, 2001). Our finding implies a counterintuitive control strategy to initially reduce VOC emissions could be more effective in controlling ammonium nitrate aerosol exceedances in the NO<sub>x</sub>-rich SLV, though a combination of NO<sub>x</sub> and VOC control will eventually be necessary for reduction below U.S. air quality standards. The model can also elucidate the relative roles of the daytime and nighttime chemical mechanisms for forming HNO<sub>3</sub>. We derive a 43% contribution to total HNO<sub>3</sub> formation by nighttime N<sub>2</sub>O<sub>5</sub> uptake on aerosol (Figure S6), with nighttime HNO<sub>3</sub> production occurring primarily in the RL, where there are no NO<sub>x</sub> emissions.

#### 4. Conclusions

O<sub>x,total</sub> is a parameter that describes both O<sub>3</sub> and pNO<sub>3</sub><sup>-</sup> production, demonstrating that they are different outcomes of the same chemical cycle, driven by different NO<sub>x</sub>/VOC emission ratios, and may be considered chemically equivalent. We have therefore modeled the predicted response of O<sub>x,total</sub> to changes in NO<sub>x</sub> and VOC emissions in the same manner that O<sub>3</sub> production is normally considered. Although this conceptual model is simplified, its conclusions are robust with respect to the underlying assumptions, as detailed in sensitivity tests in Text S4 (Li et al., 2014; Stemmler et al., 2006; Zhou et al., 2011). We find that this framework is an effective way of describing the response of ammonium nitrate aerosol to emission controls in an urban wintertime environment that is sensitive to HNO<sub>3</sub>. Furthermore, a more general three-component analysis which considers NO<sub>x</sub>, VOCs, and NH<sub>3</sub> could describe ammonium nitrate formation under any conditions in the context of O<sub>x,total</sub>. We find that the SLV is likely to be well within a NO<sub>x</sub>-saturated regime, and therefore, control strategies that initially focus on VOCs rather than NO<sub>x</sub> or NH<sub>3</sub> would be most effective, contrary to the view that NO<sub>x</sub> reductions are most effective in HNO<sub>3</sub> (rather than NH<sub>3</sub>) limited systems. The current understanding of U.S. VOC emissions and the cost and feasibility of their reduction continues to evolve (McDonald et al., 2018). Further detailed evaluation of the unique and changing biogenic and anthropogenic VOC profiles in the SLV and other valleys would be required to formulate an effective VOC emission control strategy.

Recent work has noted that summertime O<sub>3</sub> and PM<sub>2.5</sub> in the United States have steadily declined (Cooper et al., 2014; Hand et al., 2014) in response to emissions controls, but wintertime PM<sub>2.5</sub>, nitrate aerosol in particular, has not decreased as strongly (Green et al., 2015; Shah et al., 2018). Our analysis further suggests that these more gradual decreases in western U.S. PM<sub>2.5</sub> may be attributable to recent reductions in VOC rather than NO<sub>x</sub> emissions and that NO<sub>x</sub> reductions may in fact have had the counterintuitive effect of slowing the rate of decrease of PM<sub>2.5</sub> in areas dominated by ammonium nitrate. This analysis may also be relevant to both Europe and East Asia (Figure 1), where nitrate is increasingly a major component of wintertime PM<sub>2.5</sub> (Xu et al., 2018), NO<sub>x</sub> concentrations are high (Guo et al., 2014; Wang et al., 2017, 2018; Wen et al., 2018), and ammonia is typically in excess (Guo et al., 2018). Mitigation strategies have normally focused on NO<sub>x</sub> control (Guo et al., 2018), though recent work has indicated that PM<sub>2.5</sub> in northern China has a negative sensitivity to NO<sub>x</sub> (Zhao et al., 2017). Chinese emission inventories show recent reductions in NO<sub>x</sub> but increase in VOCs nationwide, a trend that could serve to exacerbate the severity of nitrate aerosol (Wang et al., 2017). Although the NO<sub>x</sub> and VOC emission profiles in European and Chinese cities may differ substantially from valleys in northern Utah, this framework for modeling aerosol suggests new and promising avenues for improved control strategies.

#### References

- Ahmadov, R., McKeen, S., Trainer, M., Banta, R., Brewer, A., Brown, S., et al. (2015). Understanding high wintertime ozone pollution events in an oil- and natural gas-producing region of the western US. *Atmospheric Chemistry and Physics*, 15(1), 411–429. <https://doi.org/10.5194/acp-15-411-2015>
- Baasandorj, M., Brown, S. S., Hoch, S., Crosman, E., Long, R., Silva, P., et al. (2018). 2017 Utah Winter Fine Particulate Study Final Report. Retrieved from <https://www.esrl.noaa.gov/csd/groups/csd7/measurements/2017uwfyps/finalreport.pdf>
- Baasandorj, M., Hoch, S. W., Bares, R., Lin, J. C., Brown, S. S., Millet, D. B., et al. (2017). Coupling between chemical and meteorological processes under persistent cold-air pool conditions: Evolution of wintertime PM<sub>2.5</sub> pollution events and N<sub>2</sub>O<sub>5</sub> observations in Utah's Salt Lake Valley. *Environmental Science & Technology*, 51(11), 5941–5950. <https://doi.org/10.1021/acs.est.6b06603>

#### Acknowledgments

The authors thank NOAA Aircraft Operations, particularly Jason Clark, Rob Mitchell, Rob Militec, and Lindsay Norman, and the Utah Division of Air Quality, particularly Chris Pennell, Brock LeBaron, and Patrick Barrickman. The authors acknowledge Gail Tonnesen, Stuart McKeen, Sebastian Hoch, Eric Crosman, Michael Trainer, and David Parrish for helpful discussions. This is NOAA PMEL contribution number 4819. The views expressed in this article are those of the authors and do not necessarily represent the views or policies of the U.S. Environmental Protection Agency. This study was partially supported by Utah State Contract 170856. M. B., J. C. L., and S. S. B. initiated and supervised the UWFPs study. C. C. W., E. E. M., R. B., K. S. D., W. P. D., D. L. F., A. F., L. G., R. L., A. M. M., D. B. M., A. M., J. G. M., T. P. R., J. A. T., L. C. V., A. R. W., M. B., and S. S. B. collected and curated the UWFPs data. J. A. dG., P. K. Q., J. M. R., P. R. V., R. J. W., C. W., B. Y., and S. S. B. collected and curated the UBWOS data. J. B. G. and B. H. L. curated and validated the UWFPs data. C. C. W., E. E. M., P. M. E., and S. S. B. developed the split box model. C. C. W. and S. S. B. wrote the manuscript. All authors contributed to and edited the manuscript. Authors declare no competing interests. All raw data can be accessed on the UWFPs website (<https://www.esrl.noaa.gov/csd/groups/csd7/measurements/2017uwfyps/>). The DSMACC box model may be downloaded for free online (<https://github.com/barronh/DSMACC>).



- Bares, R., Lin, J. C., Hoch, S. W., Baasandorj, M., Mendoza, D. L., Fasoli, B., et al. (2018). The wintertime covariation of CO<sub>2</sub> and criteria pollutants in an urban valley of the Western United States. *Journal of Geophysical Research: Atmospheres*, *123*, 2684–2703. <https://doi.org/10.1002/2017JD027917>
- Beard, J. D., Beck, C., Graham, R., Packham, S. C., Traphagan, M., Giles, R. T., & Morgan, J. G. (2012). Winter temperature inversions and emergency department visits from asthma in Salt Lake County, Utah, 2003–2008. *Environmental Health Perspectives*, *120*(10), 1385–1390. <https://doi.org/10.1289/ehp.1104349>
- Blanchard, C. L., Roth, P. M., Tanenbaum, S. J., Ziman, S. D., & Seinfeld, J. H. (2000). The use of ambient measurements to identify which precursor species limit aerosol nitrate formation. *Journal of the Air & Waste Management Association*, *50*(12), 2073–2084. <https://doi.org/10.1080/10473289.2000.10464239>
- Cooper, O. R., Parrish, D. D., Ziemke, J., Balashov, N. V., Cupeiro, M., Galbally, I. E., et al. (2014). Global distribution and trends of tropospheric ozone: An observation-based review. *Elementa: Science of the Anthropocene*, *2*(29). <https://doi.org/10.12952/journal.elementa.000029>
- de Gouw, J. A., Gilman, J. B., Kim, S. W., Lerner, B. M., Isaacman-VanWertz, G., McDonald, B. C., et al. (2017). Chemistry of volatile organic compounds in the Los Angeles basin: Nighttime removal of alkenes and determination of emission ratios. *Journal of Geophysical Research: Atmospheres*, *122*, 11,843–11,861. <https://doi.org/10.1002/2017JD027459>
- DOS (2018). US Department of State. Mission China Air Quality Monitoring Program Archive. Retrieved from <https://www.stateair.net/web/historical/1/1.html>, Accessed September 5, 2018.
- Edwards, P. M., Brown, S. S., Roberts, J. M., Ahmadov, R., Banta, R. M., deGouw, J. A., et al. (2014). High winter ozone pollution from carbonyl photolysis in an oil and gas basin. *Nature*, *514*(7522), 351–354. <https://doi.org/10.1038/nature13767>
- EMEP (2018). European Monitoring and Evaluation Programme. Norwegian Institute for Air Research EBAS [internet database]. Retrieved from <https://ebas.nilu.no/Default.aspx>, Accessed December 11, 2018.
- Emmerson, K. M., & Evans, M. J. (2009). Comparison of tropospheric gas-phase chemistry schemes for use within global models. *Atmospheric Chemistry and Physics*, *9*(5), 1831–1845. <https://doi.org/10.5194/acp-9-1831-2009>
- EPA (2018). US Environmental Protection Agency. Air Quality System Data Mart [internet database]. Retrieved from <https://www.epa.gov/ttn/airs/aqsdatamart>, Accessed September 5, 2018.
- Franchin, A., Fibiger, D. L., Goldberger, L., McDuffie, E. E., Moravek, A., Womack, C. C., et al. (2018). Airborne and ground-based observations of ammonium nitrate dominated aerosols in a shallow boundary layer during intense winter pollution episodes in northern Utah. *Atmospheric Chemistry and Physics*, *18*(23), 17,259–17,276. <https://doi.org/10.5194/acp-18-17259-2018>
- Green, M. C., Chow, J. C., Watson, J. G., Dick, K., & Inouye, D. (2015). Effects of snow cover and atmospheric stability on winter PM<sub>2.5</sub> concentrations in Western U.S. valleys. *Journal of Applied Meteorology and Climatology*, *54*(6), 1191–1201. <https://doi.org/10.1175/jamcd-14-0191.1>
- Guo, H., Otjes, R., Schlag, P., Kiendler-Scharr, A., Nenes, A., & Weber, R. J. (2018). Effectiveness of ammonia reduction on control of fine particle nitrate. *Atmospheric Chemistry and Physics*, *18*(16), 12,241–12,256. <https://doi.org/10.5194/acp-18-12241-2018>
- Guo, S., Hu, M., Zamora, M. L., Peng, J., Shang, D., Zheng, J., et al. (2014). Elucidating severe urban haze formation in China. *Proceedings of the National Academy of Sciences*, *111*(49), 17,373–17,378. <https://doi.org/10.1073/pnas.1419604111>
- Hand, J. L., Schichtel, B. A., Malm, W. C., Copeland, S., Molenaar, J. V., Frank, N., & Pitchford, M. (2014). Widespread reductions in haze across the United States from the early 1990s through 2011. *Atmospheric Environment*, *94*, 671–679. <https://doi.org/10.1016/j.atmosenv.2014.05.062>
- Jeffries, H. E., & Tonnesen, S. (1994). A comparison of two photochemical reaction mechanisms using mass balance and process analysis. *Atmospheric Environment*, *28*(18), 2991–3003. [https://doi.org/10.1016/1352-2310\(94\)90345-X](https://doi.org/10.1016/1352-2310(94)90345-X)
- Jenkin, M. E., Young, J. C., & Rickard, A. R. (2015). The MCM v3.3.1 degradation scheme for isoprene. *Atmospheric Chemistry and Physics*, *15*(20), 11,433–11,459. <https://doi.org/10.5194/acp-15-11433-2015>
- Jones, C. L., & Seinfeld, J. H. (1983). The oxidation of NO<sub>2</sub> to nitrate—Day and night. *Atmospheric Environment*, *17*(11), 2370–2373. [https://doi.org/10.1016/0004-6981\(83\)90239-1](https://doi.org/10.1016/0004-6981(83)90239-1)
- Kelly, J. T., Parworth, C. L., Zhang, Q., Miller, D. J., Sun, K., Zondlo, M. A., et al. (2018). Modeling NH<sub>4</sub>NO<sub>3</sub> over the San Joaquin Valley during the 2013 DISCOVER-AQ Campaign. *Journal of Geophysical Research: Atmospheres*, *123*, 4727–4745. <https://doi.org/10.1029/2018JD028290>
- Kelly, K. E., Kotchenruther, R., Kuprov, R., & Silcox, G. D. (2013). Receptor model source attributions for Utah's Salt Lake City airshed and the impacts of wintertime secondary ammonium nitrate and ammonium chloride aerosol. *Journal of the Air & Waste Management Association*, *63*(5), 575–590. <https://doi.org/10.1080/10962247.2013.774819>
- Kleeman, M. J., Ying, Q., & Kaduwela, A. (2005). Control strategies for the reduction of airborne particulate nitrate in California's San Joaquin Valley. *Atmospheric Environment*, *39*(29), 5325–5341. <https://doi.org/10.1016/j.atmosenv.2005.05.044>
- Kleinman, L. I. (2005). The dependence of tropospheric ozone production rate on ozone precursors. *Atmospheric Environment*, *39*(3), 575–586. <https://doi.org/10.1016/j.atmosenv.2004.08.047>
- Kuprov, R., Eatough, D. J., Cruickshank, T., Olson, N., Cropper, P. M., & Hansen, J. C. (2014). Composition and secondary formation of fine particulate matter in the Salt Lake Valley: Winter 2009. *Journal of the Air & Waste Management Association*, *64*(8), 957–969. <https://doi.org/10.1080/10962247.2014.903878>
- Kurtenbach, R., Becker, K. H., Gomes, J. A. G., Kleffmann, J., Lörzer, J. C., Spittler, M., et al. (2001). Investigations of emissions and heterogeneous formation of HONO in a road traffic tunnel. *Atmospheric Environment*, *35*(20), 3385–3394. [https://doi.org/10.1016/S1352-2310\(01\)00138-8](https://doi.org/10.1016/S1352-2310(01)00138-8)
- Lee, B. H., Lopez-Hilfiker, F. D., Mohr, C., Kurtén, T., Worsnop, D. R., & Thornton, J. A. (2014). An iodide-adduct high-resolution time-of-flight chemical-ionization mass spectrometer: Application to atmospheric inorganic and organic compounds. *Environmental Science & Technology*, *48*(11), 6309–6317. <https://doi.org/10.1021/es500362a>
- Li, X., Rohrer, F., Hofzumahaus, A., Brauers, T., Haseler, R., Bohn, B., et al. (2014). Missing gas-phase source of HONO inferred from zeppelin measurements in the troposphere. *Science*, *344*(6181), 292–296. <https://doi.org/10.1126/science.1248999>
- Liao, J., Brock, C. A., Murphy, D. M., Sueper, D. T., Welti, A., & Middlebrook, A. M. (2017). Single-particle measurements of bouncing particles and in situ collection efficiency from an airborne aerosol mass spectrometer (AMS) with light-scattering detection. *Atmospheric Measurement Techniques*, *10*(10), 3801–3820. <https://doi.org/10.5194/amt-10-3801-2017>
- Lin, X., Trainer, M., & Liu, S. C. (1988). On the nonlinearity of the tropospheric ozone production. *Journal of Geophysical Research*, *93*(D12), 15,879–15,888. <https://doi.org/10.1029/JD093iD12p15879>
- Liu, S. C. (1977). Possible effects on tropospheric O<sub>3</sub> and OH due to NO emissions. *Geophysical Research Letters*, *4*(8), 325–328. <https://doi.org/10.1029/GL004i008p00325>

- Madronich, S., McKenzie, R. L., Björn, L. O., & Caldwell, M. M. (1998). Changes in biologically active ultraviolet radiation reaching the Earth's surface. *Journal of Photochemistry and Photobiology B: Biology*, *46*(1-3), 5–19. [https://doi.org/10.1016/S1011-1344\(98\)00182-1](https://doi.org/10.1016/S1011-1344(98)00182-1)
- Mao, J., Ren, X., Chen, S., Brune, W. H., Chen, Z., Martinez, M., et al. (2010). Atmospheric oxidation capacity in the summer of Houston 2006: Comparison with summer measurements in other metropolitan studies. *Atmospheric Environment*, *44*(33), 4107–4115. <https://doi.org/10.1016/j.atmosenv.2009.01.013>
- McDonald, B. C., de Gouw, J. A., Gilman, J. B., Jathar, S. H., Akherati, A., Cappa, C. D., et al. (2018). Volatile chemical products emerging as largest petrochemical source of urban organic emissions. *Science*, *359*(6377), 760–764. <https://doi.org/10.1126/science.aaq0524>
- McDuffie, E. (2018). Measurements and modeling of nitrogen oxides: Tropospheric transformations during summer and winter in polluted regions across the U.S., Doctoral dissertation thesis, University of Colorado, Boulder.
- Meng, Z., Dabdub, D., & Seinfeld, J. H. (1997). Chemical coupling between atmospheric ozone and particulate matter. *Science*, *277*(5322), 116–119. <https://doi.org/10.1126/science.277.5322.116>
- Milford, J. B., Russell, A. G., & McRae, G. J. (1989). A new approach to photochemical pollution control: Implications of spatial patterns in pollutant responses to reductions in nitrogen oxides and reactive organic gas emissions. *Environmental Science & Technology*, *23*(10), 1290–1301. <https://doi.org/10.1021/es00068a017>
- Nguyen, K., & Dabdub, D. (2002). NO<sub>x</sub> and VOC control and its effects on the formation of aerosols. *Aerosol Science and Technology*, *36*(5), 560–572. <https://doi.org/10.1080/02786820252883801>
- Pun, B. K., & Seigneur, C. (2001). Sensitivity of particulate matter nitrate formation to precursor emissions in the California San Joaquin Valley. *Environmental Science & Technology*, *35*(14), 2979–2987. <https://doi.org/10.1021/es0018973>
- Pusede, S. E., Duffey, K. C., Shusterman, A. A., Saleh, A., Laughner, J. L., Wooldridge, P. J., et al. (2016). On the effectiveness of nitrogen oxide reductions as a control over ammonium nitrate aerosol. *Atmospheric Chemistry and Physics*, *16*(4), 2575–2596. <https://doi.org/10.5194/acp-16-2575-2016>
- Seinfeld, J. H. (1989). Urban air pollution: State of the science. *Science*, *243*(4892), 745–752. <https://doi.org/10.1126/science.243.4892.745>
- Shah, V., Jaeglé, L., Thornton, J. A., Lopez-Hilfiker, F. D., Lee, B. H., Schroder, J. C., et al. (2018). Chemical feedbacks weaken the wintertime response of particulate sulfate and nitrate to emissions reductions over the eastern United States. *Proceedings of the National Academy of Sciences*, *115*(32), 8110–8115. <https://doi.org/10.1073/pnas.1803295115>
- Silcox, G. D., Kelly, K. E., Crosman, E. T., Whiteman, C. D., & Allen, B. L. (2012). Wintertime PM<sub>2.5</sub> concentrations during persistent, multi-day cold-air pools in a mountain valley. *Atmospheric Environment*, *46*, 17–24. <https://doi.org/10.1016/j.atmosenv.2011.10.041>
- Stemmler, K., Ammann, M., Donders, C., Kleffmann, J., & George, C. (2006). Photosensitized reduction of nitrogen dioxide on humic acid as a source of nitrous acid. *Nature*, *440*(7081), 195–198. <https://doi.org/10.1038/nature04603>
- Trainer, M., Parrish, D. D., Buhr, M. P., Norton, R. B., Fehsenfeld, F. C., Anlauf, K. G., et al. (1993). Correlation of ozone with NO<sub>y</sub> in photochemically aged air. *Journal of Geophysical Research*, *98*(D2), 2917–2925. <https://doi.org/10.1029/92JD01910>
- UBWOS (2014). Uinta Basin 2013 Winter Ozone Study: Final Report. Retrieved from <https://deq.utah.gov/legacy/destinations/u/uinta-basin/ozone/strategies/studies/UBOS-2013.htm>
- Wang, H., Lu, K., Chen, X., Zhu, Q., Wu, Z., Wu, Y., & Sun, K. (2018). Fast particulate nitrate formation via N<sub>2</sub>O<sub>5</sub> uptake aloft in winter in Beijing. *Atmospheric Chemistry and Physics*, *18*(14), 10,483–10,495. <https://doi.org/10.5194/acp-18-10483-2018>
- Wang, J., Zhao, B., Wang, S., Yang, F., Xing, J., Morawska, L., et al. (2017). Particulate matter pollution over China and the effects of control policies. *Science of the Total Environment*, *584*–585, 426–447. <https://doi.org/10.1016/j.scitotenv.2017.01.027>
- Weber, R. J., Guo, H., Russell, A. G., & Nenes, A. (2016). High aerosol acidity despite declining atmospheric sulfate concentrations over the past 15 years. *Nature Geoscience*, *9*(4), 282–285. <https://doi.org/10.1038/ngeo2665>
- Wen, L., Xue, L., Wang, X., Xu, C., Chen, T., Yang, L., et al. (2018). Summertime fine particulate nitrate pollution in the North China Plain: Increasing trends, formation mechanisms and implications for control policy. *Atmospheric Chemistry and Physics*, *18*(15), 11,261–11,275. <https://doi.org/10.5194/acp-18-11261-2018>
- Wesely, M. L., & Hicks, B. B. (2000). A review of the current status of knowledge on dry deposition. *Atmospheric Environment*, *34*(12), 2261–2282. [https://doi.org/10.1016/S1352-2310\(99\)00467-7](https://doi.org/10.1016/S1352-2310(99)00467-7)
- Whiteman, C. D., Hoch, S. W., Horel, J. D., & Charland, A. (2014). Relationship between particulate air pollution and meteorological variables in Utah's Salt Lake Valley. *Atmospheric Environment*, *94*, 742–753. <https://doi.org/10.1016/j.atmosenv.2014.06.012>
- Wild, R. J., Dubé, W. P., Aikin, K. C., Eilerman, S. J., Neuman, J. A., Peischl, J., et al. (2017). On-road measurements of vehicle NO<sub>2</sub>/NO<sub>x</sub> emission ratios in Denver, Colorado, USA. *Atmospheric Environment*, *148*, 182–189. <https://doi.org/10.1016/j.atmosenv.2016.10.039>
- Wild, R. J., Edwards, P. M., Bates, T. S., Cohen, R. C., de Gouw, J. A., Dubé, W. P., et al. (2016). Reactive nitrogen partitioning and its relationship to winter ozone events in Utah. *Atmospheric Chemistry and Physics*, *16*(2), 573–583. <https://doi.org/10.5194/acp-16-573-2016>
- Wild, R. J., Edwards, P. M., Dubé, W. P., Baumann, K., Edgerton, E. S., Quinn, P. K., et al. (2014). A measurement of total reactive nitrogen, NO<sub>y</sub>, together with NO<sub>2</sub>, NO, and O<sub>3</sub> via cavity ring-down spectroscopy. *Environmental Science & Technology*, *48*(16), 9609–9615. <https://doi.org/10.1021/es501896w>
- Wood, E. C., Herndon, S. C., Onasch, T. B., Kroll, J. H., Canagaratna, M. R., Kolb, C. E., et al. (2009). A case study of ozone production, nitrogen oxides, and the radical budget in Mexico City. *Atmospheric Chemistry and Physics*, *9*(7), 2499–2516. <https://doi.org/10.5194/acp-9-2499-2009>
- Xu, W., Sun, Y., Wang, Q., Zhao, J., Wang, J., Ge, X., et al. (2018). Changes in aerosol chemistry from 2014 to 2016 in winter in Beijing: Insights from high resolution aerosol mass spectrometry. *Journal of Geophysical Research: Atmospheres*, *124*, 1132–1147. <https://doi.org/10.1029/2018JD029245>
- Zhang, L., Jacob, D. J., Knipping, E. M., Kumar, N., Munger, J. W., Carouge, C. C., et al. (2012). Nitrogen deposition to the United States: Distribution, sources, and processes. *Atmospheric Chemistry and Physics*, *12*(10), 4539–4554. <https://doi.org/10.5194/acp-12-4539-2012>
- Zhao, B., Wu, W., Wang, S., Xing, J., Chang, X., Liou, K. N., et al. (2017). A modeling study of the nonlinear response of fine particles to air pollutant emissions in the Beijing–Tianjin–Hebei region. *Atmospheric Chemistry and Physics*, *17*(19), 12,031–12,050. <https://doi.org/10.5194/acp-17-12031-2017>
- Zhou, X., Zhang, N., TerAvest, M., Tang, D., Hou, J., Bertman, S., et al. (2011). Nitric acid photolysis on forest canopy surface as a source for tropospheric nitrous acid. *Nature Geoscience*, *4*(7), 440–443. <https://doi.org/10.1038/ngeo1164>