

COMPARATIVE MODEL FOR CLASSIFICATION OF FOREST DEGRADATION

AHMED A MEHDAWI

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (Remote Sensing)

Faculty of Geoinformation and Real Estate
Universiti Teknologi Malaysia

MARCH 2015

Status Declaration Letter

Librarian

MARCH 2015

Perpustakaan Sultanah Zanariah

UTM, Skudai

Johor

Sir,

CLASSIFICATION OF THESIS AS CONFIDENTIAL


COMPARATIVE MODEL FOR CLASSIFICATION OF FOREST
DEGRADATION, Ahmed A Mehdawi, and Baharin bin Ahmad.

Please be informed that the above mentioned thesis entitled "**COMPARATIVE
MODEL FOR CLASSIFICATION OF FOREST DEGRADATION**" be
classified as CONFIDENTIAL for a period of five (5) years from the date of this
letter. The reason for the classifications are:

- (i) The authors plan to publish more journal articles based on the thesis.
- (ii) The authors had developed a conceptual model which needs to be implement in
real model nearly.

Thank you.

Sincerely yours,



ASSOC. PROF. DR. BAHARIN BIN AHMAD

Department of Geoinformation and Real Estate

Phone: +(6)075530873 (Office)

This thesis is dedicated to my beloved family, my lovely wife who paid the price during my postgraduate studies.

ACKNOWLEDGEMENT

First and foremost, all the praises and gratitude are due to Allah SWT for bestowing His mercy and blessings, which enabled me to complete this thesis. May peace and blessings be upon holy prophet Muhammad SAW, his family and his companions.

I would like to express my deep appreciation and respect to my supervisor, Assoc. Prof. Dr. Baharin Bin Ahmad, for all his guidance, help, suggestions, cooperation, understanding, patience, support and encouragement throughout my entire research period despite his immeasurable commitments.

My profound gratitude also goes to my parents, Colonel Abdul Jalil Mehdawi and Hajjah Salwa Abdul Ghani for their unfinished support all my life. Their unalloyed love kept me going, even during the most challenging moments; very genuine appreciation goes to my beloved father whom always gave me motivation and courage to look on the bright side every time I felt unmotivated, whom that never let me down and whom I respect the most in my heart. I am also indebted to my father and mother in law Alhaji Abdullah Mohd Razi and Hajjah Hasemat Binti Daud as well as my aunties- Salimh, Intesar, Fthi, Salwa Ahmed for their support and prayers all the time. To my siblings- Mohammed, Amu Abdul Rahman, Saleh, Mohand, Mahmud, Zoza and Hartini you are most appreciated. I will never be able to thank my wife, Amu Abdul Jalil, enough for the inspirational and motivational role she played during the course of my Ph.D. Your love and caring are my strength. Finally, I am grateful to all friends, especially Dr Abdullah Algahtani, Abdul Wahid Almrhon, Dr Aomar Lawal Dano, Mohd Hayyi Mat Zin colleagues, and relatives who have supported me in one way or the other during the course of my stay here in Malaysia. I implore Allah (SWT), The Almighty to reward you all abundantly.

ABSTRACT

The challenges of forest degradation together with its related effects have attracted research from diverse disciplines, resulting in different definitions of the concept. However, according to a number of researchers, the central element of this issue is human intrusion that destroys the state of the environment. Therefore, the focus of this research is to develop a comparative model using a large amount of multi-spectral remote sensing data, such as IKONOS, QUICKBIRD, SPOT, WORLDVIEW-1, Terra-SARX, and fused data to detect forest degradation in Cameron Highlands. The output of this method in line with the performance measurement model. In order to identify the best data, fused data and technique to be employed. Eleven techniques have been used to develop a Comparative technique by applying them on fifteen sets of data. The output of the Comparative technique was used to feed the performance measurement model in order to enhance the accuracy of each classification technique. Moreover, a Performance Measurement Model has been used to verify the results of the Comparative technique; and, these outputs have been validated using the reflectance library. In addition, the conceptual hybrid model proposed in this research will give the opportunity for researchers to establish a fully automatic intelligent model for future work. The results of this research have demonstrated the Neural Network (NN) to be the best Intelligent Technique (IT) with a 0.912 of the Kappa coefficient and 96% of the overall accuracy, Mahalanobis had a 0.795 of the Kappa coefficient and 88% of the overall accuracy and the Maximum likelihood (ML) had a 0.598 of the Kappa coefficient and 72% of the overall accuracy from the best fused image used in this research, which was represented by fusing the IKONOS image with the QUICKBIRD image as finally employed in the Comparative technique for improving the detectability of forest change.

ABSTRAK

Cabaran dalam menangani degradasi hutan dan kesannya telah menarik pelbagai penyelidikan dari pelbagai bidang, mengakibatkan definisi terhadap konsep yang berbeza. Walaubagaimanapun, menurut sebahagian penyelidik, elemen utama dalam isu ini ialah pencerobohan manusia yang telah memusnahkan keadaan alam sekitar. Oleh itu, fokus penyelidikan ini ialah untuk membangunkan model perbandingan menggunakan sejumlah besar data remote sensing multi spektrum seperti IKONOS, QUICKBIRD, SPOT, WORLDVIEW-1, Terra-SARX dan data gabungan untuk mengesan degradasi hutan di Cameron Highlands. Hasil dari kaedah ini sejajar dengan model pengukuran prestasi. Ini akan membantu dalam mengenalpasti data yang terbaik, data gabungan dan teknik yang akan digunakan. Sebelas teknik telah digunakan untuk membangunkan Teknik Perbandingan dengan mengaplikasikannya kepada lima belas set data. Hasil dari Teknik Perbandingan telah digunakan untuk membekalkan model pengurusan prestasi supaya dapat memperbaiki ketepatan setiap klasifikasi teknik. Bahkan, model pengurusan prestasi telah digunakan untuk mengenal pasti hasil dari Teknik perbandingan, dan hasil ini telah disahkan menggunakan perpustakaan pantulan. Sebagai tambahan, model konsep hibrid yang dicadangkan dalam penyelidikan ini akan memberi peluang kepada para penyelidik untuk mewujudkan model pintar automatik untuk kerja masa hadapan. Hasil dari kajian ini telah menunjukkan Rangkaian saraf (NN) adalah Teknik Pintar (IT) terbaik dengan 0.912 bagi Kapa Koefisien dan 96% bagi ketepatan keseluruhan. Manakala Mahalanobis memperoleh 0.795 Kapa Koefisien dan 88% bagi ketepatan keseluruhan. *Maximum likelihood* (ML) memperoleh 0.598 bagi Kapa Koefisien dan 72% bagi ketepatan keseluruhan daripada imej gabungan yang terbaik dalam kajian ini, yang diwakili dengan menggabungkan imej IKONOS dengan imej QUICKBIRD yang akhirnya digunakan dalam Teknik Perbandingan untuk memperbaiki kebolehsesan degradasi hutan.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xii
	LIST OF FIGURES	xiii
	LIST OF ABBREVIATIONS	xvi
	LIST OF APPENDICES	xvix
<u>1</u>	<u>INTRODUCTION</u>	1
	1.1 Background	1
	1.2 Problem Statements	4
	1.3 Study Area	5
	1.4 Research Objectives	9
	1.5 Research Questions	9
	1.6 Scopes of the Research	10
	1.7 Significance of Research	10
	1.8 Thesis Organization and Flow Chart	12
	1.9 Summary of the Chapter	14
<u>2</u>	LITERATURE REVIEW	15
	2.1 Introduction	15

2.2 Forests	16
2.2.1 Tropical Dry Forests	17
2.2.2 Forest in Southeast Asia	17
2.2.3 Forest in Malaysia	18
2.2.3.1 Cameron Highland as unique Biodiversity	18
2.3 Forest Degradation	19
2.3.1 The Difficulty of Forest Degradation	21
2.3.2 Explanations of Forest and Forest Degradation	22
2.3.3 Reasons of Forest Degradation	25
2.3.4 Displays of Forest Degradation	25
2.4 Remote Sensing	27
2.4.1 Monitoring Forest by Satellite Image	27
2.4.2 Optical and Radar Remote Sensing System	32
2.4.3 Optical Remote Sensing System	35
2.4.4 Active Radar Remote Sensing	36
2.5 Difference amongst Satellite Images	37
2.6 The Impact of Resolution	39
2.6.1 Radiometric Resolution	40
2.6.2 Spectral Resolution	41
2.6.3 Spatial Resolution	42
2.6.4 Temporal Resolution	43
2.7 Supporting Technologies to Remote Sensing Data	44
2.8 Practical Difficulties of Land and Forest Cover Monitoring in the Tropics	47
2.8.1 Frequent Cloud Cover	47
2.8.2 Atmospheric Effects	48
2.8.3 Mosaicking Artifacts of Multispectral Datasets	49
2.9 Practical Difficulties due to lacking Data Stability	50
2.9.1 Pre-processing	51
2.9.2 Atmospheric Correction Methods	53
2.9.3 Orthorectification	55
2.10 Data Fusion	56
2.10.1 Multi-Source Data Fusion	60

2.10.2 Fusion for Post Classification	62
2.10.3 Fusion for Change Detection	64
2.11 Classification	66
2.11.1 Choice of a Classification Scheme and Training Model	66
2.11.2 Remote-Sensing Classification Procedure	67
2.12 Explanations of Classification	68
2.12.1 Unsupervised Classification Approachs	68
2.12.1.1 Pixel-Oriented Techniques	69
2.12.2 Supervised Classification	71
2.13 Techniques used in supervised classification	73
2.14 Techniques Used In Unsupervised Classification	86
2.15 Particle Swarm Optimization (PSO) Algorithm	92
2.16 Selection of a Classification System and Training Samples	93
2.17 Advanced classification approaches	95
2.18 Hybrid Image Classification	96
2.19 Performance Measurement Model	97
2.20 Classification Accuracy Assessment	97
2.20.1 Classification Techniques and Forest Degedation	99
2.20.2 Classification Techniques for Forest Degradation in Tropical Region within Asia	107
2.20.3 Modelling of Forest Degradation	111
2.20.4 Modelling Requirement	111
2.20.5 Types of models	112
2.21 Summary of the Chapter	115
3	
METHODOLOGY AND DATA PREPARATION	117
3.1 Introduction	117
3.2 Research Design and Methods	119
3.3 Datasets	122
3.3.1 Satellite Data	122
3.3.1.1 IKONOS Image	124
3.3.1.2 SPOT Image	125

3.3.1.3 QuickBird Image	128
3.3.1.4 Worldview-1	129
3.3.2 TerraSAR-X	130
3.4 Ancillary Data	132
3.5 Preprocessing stage	134
3.5.1 Image Registration	135
3.5.2 Image Orthorectification	136
3.5.3 Atmospheric Correction (AC)	137
3.5.4 Cloud Masking	140
3.5.5 Mosaicking	142
3.6 Post processing	142
3.6.1 Fusion	144
3.7 Comparative Technique	146
3.7.1 Supervised Classification Methods	150
3.7.2 Unsupervised classification	151
3.8 Performance Measurement Model	153
3.9 Validation	156
3.10 Summary of the Chapter	157
4	
RESULT AND DISCUSSION	158
4.1 Introduction	158
4.2 The Fusion Techniques	159
4.2.1 IKONOS Fused with QuickBird, SPOT, WorldView, and Terra SAR	160
4.2.2 QuickBird Fused with SPOT, Terra SAR, and Worldview	162
4.2.3 SPOT Satellite Image Fusion with Terra SAR and Worldview	162
4.3 Comparative Technique	163
4.3.1 Techniques used in Supervised Classification	163
4.3.1.1 Neural Network	164
4.3.1.2 Mahalanobis Distance Classification	165
4.3.1.3 Maximum likelihood	166
4.3.1.4 Spectral Information Divergence (SID)	167

	4.3.1.5 Fuzzy logic	168
	4.3.1.6 Parallel piped Classification	169
	4.3.1.7 Minimum Distance Classification	170
	4.3.1.8 Spectral Angle Mapper (SAM) Method	170
	4.3.1.9 Binary Encoding Classification	171
	4.3.2 Unsupervised Classification Methods	171
	4.3.2.1 K-Means Classification	172
	4.3.2.2 ISODATA Classification	173
	4.4 Overall Accuracy	174
	4.5 Performance Measurement Model (PMM)	175
	4.6 Validation Techniques	179
	4.7 Summary of the Chapter	182
5	CONCLUSION	184
	5.1 Introduction	184
	5.1.1 A restatement of the research objectives	184
	5.1.2 Actualisation of the research objectives	185
	5.2 Summary of the Chapter	187
	<u>REFERENCES</u>	188
	Appendix A-R	211-267

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	Multiband Regions of Multispectral Sensors	30
2.2	QB and Ikonos Criteria	31
2.3	Differences between Passive and Active Sensor	34
2.4	Fusion Technique Equations	59
2.5	Summary of the Possible Ways of Classifying Models	113
3.1	The Coordinate of Covered Area	122
3.2	Specification of topography map	133
3.3	Deliberate fused image	145
3.4	Equations of the Classification Techniques	147
4.1	Overall Accuracy and Kappa Coefficient of the Classification Techniques used	174
4.2	Average Accuracy of the Eleven Techniques Applied for Quickbird and its Fused Images1	177
4.3	Minimum Squared Error of the Eleven Techniques Applied for Quickbird and its Fused Images	177

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
1.1	Map of the study area	6
1.2	The location of Cameron Highlands, located between Perak and Pahang	8
1.3	Thesis Organizational Flow Diagram	13
2.1	Centroid model	26
2.2	Different sensors; (a) Passive Sensor, (b) Active Sensor	33
2.3	Fundamental method of the remote sensing	35
2.4	Atmosphere transmissions	36
2.5	Shows the components of Spectral Reflectance	40
2.6	Diagramme of a Hyper-spectral image	42
2.7	Shows the pixel size concept on the image	43
2.8	Shows the orthorectification from the viewing direction	55
2.9	Processing level of data fusion	57
2.10	Levels of fusion	61
2.11	The threshold logic unit	75
2.12	Neural network processing element	76
2.13	Setting the parallel pipe boundaries by inspecting Class histograms in each band	77
2.14	A set of two-dimensional parallelpipe	78
2.15	Classification associated with connected data displaying areas of inseparability	79
2.16	(a) Representing pixel vectors by their angles, (b) segmenting the spectral space by angle	82
2.17	(a) A simple binary decision tree, and (b) multi-class decisions using binary classifiers in parallel, followed by decision logic	84
2.18	(a) Representing pixel vectors by their angles, (b) segmenting the spectral space by angle	87

2.19	The general structure of fuzzy image processing	90
3.1	Flowchart of the Research Activities	118
3.2	Phases of the General Framework of the Study	121
3.3	Satellite Dataset	123
3.4	IKONOS image shows the study area	124
3.5	Outline of the main steps of the pre-process in IKONOS	125
3.6	SPOT -5 image with some information	126
3.7	Preprocessing for SPOT image	127
3.8	QuickBird Image for study area in 2007	128
3.9	Worldview-1pan	129
3.10	Terra SAR image for study area	131
3.11	District and sub-district maps of the study area	133
3.12	Flowchart of pre-processing as employed in the study	134
3.13	Steps of Image Registration	136
3.14	The main key of FLAASH procedures	138
3.15	The original IK and QB image with cloud impact	140
3.16	IK and QB after masking	141
3.17	Flowchart of the Post Classification	143
3.18	Flowchart of the Comparative Technique	149
3.19	Flowchart of the Supervised Classification	151
3.20	Flowchart of the Unsupervised Classification	153
3.21	Performance measures on three different levels	154
3.22	Draws the general strategy of the PMM	155
3.23	The validation steps of this research	156
4.1	IKONOS fused images	160
4.2	Neural Network classification	164
4.3	Mahalanobis distance classification	166
4.4	Maximum Likelihood Classification	167
4.5	The Performance Measurement Model of the Neural Network Classification	176
4.6	The Performance Measurement Model of the Mahalanobis Classification	178
4.7	Validations of QB, SPOT, and IKONOS based on the Spectral Library	180

4.8	Validation based on the Spectral Reflectance Map	181
4.9	Validation based on the Spectral Profile from the Neural Network Classified Image	182

LIST OF ABBREVIATIONS

AC	-	Atmospheric Correction
AI	-	Artificial Intelligence
AID	-	Automatic Interaction Detection
AIRSAR	-	Airborne Synthetic Aperture Radar
ANN	-	Artificial Neural Networks
ARSM	-	“Malaysia Remote Sensing Agency”
AVHRR	-	Advanced Very High Resolution Radiometer
BP	-	Back Propagation
BRDF	-	Bidirectional reflectance distribution function
CHAID	-	Chi-squared Automatic Interaction Detector
CM	-	Comparative Model
DEM	-	Digital Elevation Models
DTC	-	Decision Tree Classifications
EAs	-	Evolutionary Algorithms
ENVI	-	Environment for Visualizing Images
EO	-	Earth Observation
FFNN	-	Feed Forward Neural Network
FLAASH	-	Fast Line-of-sight Atmospheric Analysis of Spectral Hyper
FLS	-	Fuzzy Logic System
FNN	-	Fuzzy Nearest Neighbour
GA	-	Genetic Algorithms
GCP	-	Ground Control Points
GIS	-	Geographic Information System
GPS	-	Global Position System
GSI	-	Ground Sampling Interval
HH	-	Horizontal transmit and horizontal receive polarizations

HRG	-	High Resolution Geometrical
HRVIR	-	High-Resolution Visible and Infrared
HSV	-	Hue-Saturation-Value
HV	-	Horizontal transmit and vertical receive polarizations
ID3	-	Induction of Decision Tree
HIS	-	Intensity-Hue-Saturation
IK	-	IKONOS
IR	-	Infrared
ISODATA	-	Iterative Self- Organizing Data Analysis Technique
IUCN	-	International Union for Conservation of Nature
KNN	-	k Nearest Neighbor
LAI	-	Leaf Area Index
LULC	-	Land Use and Land Cover
MD	-	Minimum Distance
MLC	-	Maximum Likelihood Classifier
MODTRAN	-	Moderate Resolution Atmospheric Transmission
MS	-	Multispectral
NASA	-	National Aeronautics and Space Administration
NDVI	-	Normalized Difference Vegetation Index
NIR	-	Near Infrared
NN	-	Neural Network
OAA	-	One Against All
OAo	-	One Against One
PAN	-	Panchromatic
PCA	-	Principal Component Analysis
PE	-	Processing Elements
PMM	-	Performance Measurement Model
PNN	-	Probabilistic Neural Networks
PRFs	-	Permanent Reserved Forests
PSO	-	Particle Swarm Optimization
QB	-	Quick Bird
RADAR	-	Radio Detection and Ranging
RBF	-	Radial Basis Function

RMSE	-	Root Mean Square Error
RPC	-	Rational Polynomial Coefficients
ROI	-	Region Of Interest
SAM	-	Spectral Angel Map
SAR	-	Synthetic Aperture Radar
SI	-	Swarm Intelligent
SID	-	Spectral Information Divergence
SM	-	Stripmap mode
SP	-	SPOT
SPOT	-	Satellite Pour "Observation" de la Terre
SVMs	-	Support Vector Machines
TGO	-	Trimble geometrics
TLU	-	Threshold Logic Unit
V	-	Visible
VH	-	Vertical Horizontal
VV	-	Vertical Vertical
WV	-	World View

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Conceptual Hybrid Model	211
B	Reflectance and Spectral Library	217
C	GPS coordinate points	227
D	Quickbird Fused with SPOT, TerraSAR, and Worldview	229
E	Quickbird Fused with SPOT, TerraSAR, and Worldview	230
F	Mahalanobis Distance Classification Map of IKONOS Fused Images	231
G	Maximum Likelihood Classification Map of IKONOS Fused Images	233
H	Spectral Information Divergence Classification Map of IKONOS Fused Images	236
I	Fuzzy Logic Classification Map of IKONOS Fused Images	239
J	Parallelepiped Classification Map of IKONOS-Fused Images	242
K	Spectral Angle Mapper Classification Map of IKONOS Fused Images	245
L	Binary Encoding Classification Map of IKONOS Fused Images	248
M		251
	Applying K-Means Classification	
N	IsoData Classification Map of IKONOS Fused Images	255
O	PMM Input to Matlab	261
P	PMM Results	262

Q	Reflectance Library Coordinate	266
R	Filedwork Pictures	267

CHAPTER 1

INTRODUCTION

1.1 Background

Worldwide, forests are very much an important element of the Earth's ecology. Forests play a very critical role in the air composition that living things breathe, provision of environmental diversity, protection of the soil from erosion, and sustaining the water cycle (Bonan, 2008). For instance, forest products are no doubt the main source of living for billions of people and the entire world. Additionally, they minimise greenhouse gas emissions into the atmosphere through the means of carbon repossession. In 2008, Ollingera et al. highlighted that the role forests play in reducing climate change is above expectation. Regrettably, forest cover degradation has become part and parcel of human behaviour in the past decades. The degradation includes, amongst others, cutting down trees, conversion to plantation or cropland as well as man-made or natural catastrophes (Potter et al., 2003). Even though natural catastrophes play a major role in forest degradation, a huge percentage of deforestation is as a result of direct human action in an attempt to increase economic status, and social as well as demographic forces. Anyway, the effects of deforestation can be very big and last for decades (Kumar, 2011).

Remote sensing has a potential role as a source of information in sustainable management around the world. This potential is largely determined by the unique characteristics of remote sensing data which provide synoptic, repetitive, and quantitative and spatially explicit functions (Franklin, 2001). Remote sensing, taking into account the above -mentioned meaning, is a very powerful and promising tool in providing independent data, which can also be used to evaluate forest change (Lucas et al., 2002; Lu et al., 2005).

In the developed countries, remote sensing data helps immensely in development projects, such as creating the appropriate action plans for forest monitoring and forecasting purposes. Changes in Land Use and Land Cover (LULC) as well as forest changes can be detected by utilising remote sensing as the main tool to monitor the local, regional as well as international resources. Remote sensing provides the basis for a better understanding of the relationships between human and natural phenomena (Vahidi et al., 2013).

Remote sensing studies on image classification have become a hotspot topic nowadays by researchers. The remote sensing community considers classification results as benchmarks for several environmental and socio-economic applications. Researchers and practitioners have successfully developed advanced classification strategies and techniques to improve the classification accuracy and reliability (Foody, 1996; Gong et al., 1999; Stuckens et al., 2000; Franklin et al., 2002; Pal and Mather, 2003; Gallego, 2004; Brekke and Solberg, 2005; and Feng et al., 2007). However, the classification of data obtained from the remote sensing technique into a thematic map remains a challenge due to many factors, such as the complexity of the landscape in a given study area, selected remote sensing data, image processing and classification strategies. These factors may cause an incorrect classification.

Digital image classification is primarily based on several algorithms that have the ability to provide an automatic system to successfully determine similarities and distinguish different surfaces. The degree of identification achieved by these algorithms includes an efficient utilisation of the remote sensing data. Accordingly, various algorithms have been developed in an attempt to classify vegetation, forest, Land Use and Land Cover (LULC) etc. In this research, a Comparative Model (CM) was developed to be able to enhance the detectability of forest change. Moreover, a Performance Measurement Model (PMM) was equally employed in this study to check the accuracy of the classification outputs.

Previous researches on image classification as well as the latest reviews of classification methods for tropical region are scarce (Tso and Mather, 2001; Landgrebe, 2003). Based on recent findings, classification algorithms and techniques for forest areas require an intensive review. This will be very valuable for guiding or selecting a suitable classification method for a particular study, as revealed in this research.

The use of Artificial Intelligence (AI) in environmental modelling with recognition of its potential acceptance has increased. AI imitates the human perception, learning and thinking in order to solve complex problems (Chen et al., 2008). This research is based on artificial neural networks, fuzzy models, swarm intelligence, machine learning and hybrid systems in order to improve the Comparative technique used in this research.

Data retrieval from manual and semi-automatic strategies has been greatly affected by the interpreters. But the automatic extraction, so far, depends on the performance of the algorithm as well as the content of the information obtained from the image (Al- Dossary and Marfurt, 2007). Particle swarm optimisation (PSO) is one of the AI techniques improving the performance of Artificial Neural Networks (ANNs). ANNs are used to develop a land use map or other environmental variables,

which present with a membership value of 0 to 1. This depends on the degree of closeness of the class for each class used in the training .

1.2 Problem Statements

Despite the growing global awareness of the importance of the conservation of tropical rainforests, the available methods to classify humid tropical forests are not sufficient. Deforestation, degradation and loss of biodiversity would eventually cause negative effects in the livelihoods of people who depend on the forest resources. As Komlos (2008) stated, it is well known that rapid development modifications can significantly degrade the surrounding environment. These development activities manifest as land modifications, such as the conversion of wetlands to settlements and alterations in the land cover in the area. As a result, major changes in the vegetation cover and surface water occur with related implications for productivity and sustainable development in the area. Moreover, these activities result in more occurrences of severe land degradation. It is necessary to seek a detailed understanding of the linkages between the manifestations of land degradation and human settlement intensification through the identification and areal quantification of land modifications associated with land degradation.

Tropical forest regions have suffered from forest degradation, which as a result has brought a major concern amongst the local communities. Particularly, Cameron Highlands have suffered a lot from landslides, erosion, and so on. Previously, there has been no study conducted in the study area with regards to forest degradation detection to monitor forest degradation. To achieve the foregoing, comparative techniques should be the appropriate technique. There is a lack of large original and fused satellite data usage from the previous studies in obtaining a suitable database to be used for the Comparative technique.

Remote sensing classification techniques are faced with difficulties of integrating statistical data classification. The successful use of Artificial Neural Networks (ANNs) with regards to remote sensing has really been encouraged by many previous researchers (Brekke and Solberg, 2005; and Sadly and Faisal, 2011). However, the reliability of the outputs from ANN needs more investigation. The performance measurement of the classification output is the main key to enhance the accuracy of the overall classification. This is another important area that needs further investigation.

Swarm intelligent algorithms, such as PSO, have just recently been highlighted in the theoretical datasets utilising binary or maybe even multiclass strategies. More basic classifier methods just produce an excellent outcome across the evaluation of datasets. In addition, no research projects or scientific studies have recently ended up with records on the subject of employing population-based research strategies to optimise the feed forward neural network, FFNN. Development of a new mathematical model is necessary to map highly located forest cover change, such as in Cameron Highlands.

1.3 Study Area

The study area was in the mainland of Cameron Highlands (4° 28' N) (101° 23' E), Pahang, Malaysia. It is located on the main range of Peninsular Malaysia as shown in Figure 1.1 It covers a total area of 71 000 715 km² (Fortuin, 2006). Generally, the terrain is mountainous and strongly dissected with 10–35° slopes. More than 66 percent of the land has a gradient of more than 20°. Cameron Highlands is about 715 km² in area settled between roughly 900 and 1800 m and surrounded by forested peaks rising to 2032 m. The Malaysian lowlands are heavily disturbed, so upland forests like those of Cameron Highlands are an important refuge

for biodiversity. Cameron Highlands is significantly cooler than Malaysia's lowlands, with a mean daily minimum of 14.8°C, a mean daily maximum of 21.1°C, which suits temperate crops. The rainfall averages 2660 mm yr⁻¹, humidity is high and there is no marked dry season.




Figure 1.1 Map of the study area (Ismail et al., 2012)

In peninsular Malaysia, Cameron Highlands is a tourist resort and it is referred to as the 'Green Bowl' which is the second most important state for growing vegetables (mostly cabbage, tomatoes, and leafy vegetables) and is also important for tea, flowers and fruit. In addition, the Cameron Highlands Catchment area is a source of water supply to many areas of Peninsular Malaysia. Sg. Telom, Sg. Bertam and Sg. Lemoi are the three main rivers of the Cameron Highlands Catchment, which drains the northern, middle and southern sections of the highlands (Makalahmad et al., 2008). Cameron Highlands is located approximately 200 km north of Kuala Lumpur on the east side of the border between the states of Perak and Pahang in Malaysia as shown in Figure 3.4. At about 1,500 meters above sea level, it is the highest area on the mainland stretching along a plateau set. Cameron Highlands was

named after William Cameron, a British Government surveyor who discovered it in 1885 whilst on a mapping expedition. However, he failed to mark his discovery and it was not until 1925 that Sir George Maxwell recorded Cameron's discovery and decided to develop it as a hill resort. (Tenaga Nasional Berhad Research, 2009). Cameron Highlands covers from Brincang town, down to Habu. The location of the study area is shown precisely in the data acquisition section. The study area consists of the Mentigi forest reserve, which is one of the gazetted forest reserves in Malaysia. This study area includes various types of land uses, such as forest, urban, tea and vegetable. The choice of this area was due to encroachment issues by the vegetable farmers and illegal logging.




Figure 1.2 The location of Cameron Highlands, located between Perak and Pahang (Malaysian Journal of Environmental Management (Barrow et al., 2005))

1.4 Research Objectives

The general aim of the research has been to develop a Comparative technique for classification of forest degradation. The research has the following sub-objectives:

1. To assess the reliability of optical and radar remote sensing data for monitoring the current situation of forest degradation.
2. To develop a Performance Measurement Model (PMM) that can assist to examine as well as improve the efficiency of the remote sensing data.
3. To establish a new conceptual classifier model using hybrid PSO methods with FFNN.

1.5 Research Questions

The following questions were addressed by this study in order to categorise and map the procedures of forest change:

1. What is the efficiency of using optical, radar, and fused remote sensing data in monitoring and assessing forest degradation?
2. Are the results obtained from the Performance Measurement Model in line with the results obtained from the Comparative Model?
3. How does the hybrid technique help in detecting forest degradation?

1.6 Scope of the Research

In achieving the study's objectives, the scope of the study was designed to carry out a research case study along with the classification framework within satellite remote sensing data to determine suitable strategies coupled with obtaining a better understanding for the purpose of establishing a Comparative technique for Classification of Forest Degradation.

1. In order to develop and test the methodology, the study area has been selected in Malaysia where the preservation of forest process is in practice.
2. The case study area was classified by optical, radar and fused images by using a Comparative model. The results have been compared and evaluated with the traditional classification methods, such as parallel pipe, minimum distance, and so on.
3. The research has mainly emphasized the utilisation of artificial neural networks trained with multispectral values to forecast forest degradation. The advantage of the proposed technique is that it needs very few variables and very little facts.
4. Sensitivity analysis/validation has been carried out in order to check the model's efficiency using a confusion matrix.

1.7 Significance of Research

Considering the current availability of satellite images, the variety of Passive and Active sensors provides the ideal possibility of effective tropical forest observation. The initial strategy is going to enhance the Cameron Highlands with data which are readily available from Optical sensors such as IKONOS, QUICKBIRD, SPOT, WORLDVIEW, and TERRASAR-X data. Therefore, these varieties will also provide significant methods which have more tendency of discovering the forest degradation. This kind of multi-sensor strategy has already been developed in this research for beneficial achievement specifically in the situation where the multi-sensor data can be merged together to get more data which is known as image fusion. This will generate more data and enhance the outcome provided by the Comparative technique. Definitely, this research will be useful to several industries to obtain information and facts when dealing with a particular research location.


Moreover, the Comparative Classification Model used in this study including various supervised and unsupervised classification techniques applied on a large amount of optical and radar data of the humid forest has never been published going by the several past studies as in (Kumar, 2006; Shaoqing and Lu, 2008; Rozenstein and Karnieli, 2011; Veetil and Zanardi, 2012; Kalra et al., 2013; Janardhana and Venugopala, 2014; and Mallinis et al., 2014). All the above mentioned studies did not use a large amount of data or many techniques to enhance the classification output. Due to the shortage of specific information and facts, there were serious complications which did not even allow multispectral satellite images to be used. This research has efficiently established a Comparative Model which includes various kinds of data as believed to play the role of benchmark information and facts for forest degradation detection in different humid tropical rainforest and mountainous regions. On the subject of employing population-based research strategies to optimise a feed forward neural network, FFNN, as stated in the problem statement, a conceptual model has been developed. The development of this

conceptual intelligent model based on a hybrid of AI and swarm intelligent techniques will help researchers and decision makers in enhancing the outputs from the classification maps. Hopefully, it can come into reality in the near future.

1.8 Thesis Organisation and Flow Chart

The organisation of this thesis is shown in Figure 1.1. In chapter 1, general information on the research, research background, problem description, research aim and objectives, research questions, scope of the research, significance of the research and general organisation of this thesis have been discussed. Chapter 2 contains detailed information on forest changes, locally and globally, and a literature review on forest change detection using remote sensing and Artificial Intelligent techniques. Moreover, the components of the Comparative Model (CM) are discussed. The Performance Measurement Model (PMM) is introduced. Specific techniques of forest change detection utilised in this thesis are discussed and the rationale for selecting this set of techniques is offered.

In chapter 3, the research methodology is discussed. The processes of data acquisition, derivation, and spatial classification of the forest change detection are thoroughly explained. The classification techniques used in this research, the CM and PMM mathematical models are discussed in detail. Moreover, in chapter 4, the results from the fusion, CM and PMM are presented and discussed. The various classification techniques are tested and discussed; and the best classification techniques are identified. The validation of the CM was performed using the PMM and reflectance library. Chapter 5 discusses the main findings including the contribution to the body of knowledge, recommendations, and future research direction.






Figure 1.3 Thesis Organizational Flow Diagram

1.9 Summary of the Chapter

The alarming and persistent occurrence of forest degradation, which affects the forest in Cameron Highlands, has caused tremendous damages and distortions to human lives, economics, and properties. Since forest degradation takes place across natural landforms that possess some specific geomorphology and geological makeup, there is an ultimate need to thoroughly examine the study area in order to investigate and analyse the nature of forest degradation. One has yet to see any published article that has used the Comparative Model for the classification of forest degradation challenges in Cameron Highlands. This is detrimental to the populace and their lives as a whole because a more effective tool ought to be used in order to minimise to the maximum the alarming forest degradation epidemic in the area.

REFERENCES

- Abdolrassoul, S. Mahiny, B.Turner J, (2007).A Comparison of Four Common Atmospheric Correction Methods. Photogrammetric Engineering and Remote Sensing Vol. 73, No. 4, 361–368.
- Achard, F., Eva, H.D., Stibig, H.-J., Mayaux, P., Gallego, J., Richards, T. and Malingreau, J.-P. (2002). Determination of deforestation rates of the world's humid tropical forests. *Science* 297: 999-1002.
- Adhikari, R., Agrawal, K., R. (2013) Hybridization of Artificial Neural Network and Particle Swarm Optimization Methods for Time Series Forecasting.
- Akgün, A. E., A Hüsnü Turk, N (2004). Comparing Different Satellite Image Classification Methods: An Application In Ayvalık District, Western Turkey. XXth International Congress for Photogrammetry and Remote Sensing, Istanbul, Turkey.
- Al-Ahmadi, F. S., Hames, A.S. – Comparison of Four Classification Methods to Extract Land Use and Land Cover from Raw Satellite Images for Some Remote Arid Areas, Kingdom of Saudi Arabia, *Earth Sci.*, Vol. 20 No.1, pp: 167-191, (2009).
- Al-Dossary, S., and K. J. Marfurt, (2007), Lineament-preserving smoothing: *Geophysics*, 72, P1-P8.
- Allen, D.M.(2005). Cameron Highlands Malaysia's Coolest Hill Resort.Sunspot Productions Malaysia.
- Alshennawy, A., Ayman A (2009). "Edge detection in digital images using fuzzy logic technique."*World Academy of science, engineering and technology* **51**: 178-186.
- Althausen, J.D., (2002), What remote sensing system should be used to collect the data? In J.D. Bossler, J.R. Jensen, R.B. McMaster and C. Rizos (Eds),

- Manual of Geospatial Science and Technology, pp. 276–297 (New York: Taylor and Francis).
- Arijit Laha and J. Das. (2003). "Fuzzy Rules and Evidence Theory for Satellite Image Analysis", In proceedings of International Conference on Advances in Pattern Recognition.
- Armenteras, D., Rudas, G. Rodriguez, N. Sua, S., and Romero, M. (2004). Pattern and cause of deforestation in Colombian Amazon. *Indicator*, 6: 353-368.
- Asner, G.P., Carlson, K.M., Martin, R.E. (2005). Substrate age and precipitation effects on Hawaiian forest canopies from space imaging spectroscopy. *Remote sensing of environment*, 98:457-467.
- Asner, G.P., Clark, J.K., Mascaro, J., Galindo Garcia, G.A., Chadwick, K.D., Navarrete Encinales, D.A., Paez-Acosta, G., Cabrera Montenegro, E., Kennedy-Bowdoin, T., Duque, A., Balaji, A., von Hildebrand, P., Maatoug, L., Phillips Bernal, J.F., Yepes Quintero, A.P., Knapp, D.E., Garcia Davila, M.C., Jacobson, J., Ordonez, M.F., (2012). High-resolution mapping of forest carbon stocks in the Colombian Amazon. *Biogeosciences* 9, 2683–2696.
- Awrangjeb, M., Ravanbakhsh, M., and Fraser, C. S. (2010). Automatic detection of residential buildings using LIDAR data and multispectral imagery. *ISPRS Journal of Photogrammetry and Remote Sensing*, 65(5):457-467.
- Baatz, M., Heynen, M., Hofmann, P., Lingenfelder, I., Mimler, M., Schäpe, A., Weber, M. and Willhauck, G. (2000). *eCognition user guide*. Definiens imaging GmbH, München, Germany.
- Bandyopadhyay, S., and Maulik, U. (2002). "Genetic clustering for automatic evolution of clusters and application to image classification," *Pattern Recognition*, vol. 35, no. 6, pp. 1197–1208
- Barnsley, M.J., (1999), *Digital remote sensing data and their characteristics*.
- Barrow, C.J.1, Clifton, J.2, Chan, N.W.3 & Tan, Y.L. (2005). Sustainable Development in the Cameron Highlands, Malaysia. *Malaysian Journal of Environmental Management* 6 (2005): 41 – 57.
- Baugh, W.M. and Groeneveld, D.P. (2006). Broadband vegetation index performance evaluated for a low-cover environment. *International Journal of Remote Sensing* 27, (21):4715–4730.

- Berlanga Robles, C. A., and A. Ruiz Luna. (2002). Land use mapping and change detection in the coastal zone of northwest Mexico using remote sensing techniques. *Journal of Coastal Research* 18:514-522.
- Binaghi, E., Brivio, P. A., Ghezzi, P., and Rampini, A. (1999). A fuzzy set accuracy assessment of soft classification. *Pattern Recognition Letters*, 20, pp. 935–948.
- Bischof, H., Schneider, W. and Pinz, A. J. (1992) Multispectral classification of Landsat-images using neural networks. *IEEE Transactions on Geo Science and Remote Sensing*, 30, 482-489.
- Bloom, A. L., Fielding, E. J. and Fu, X. (1988). A demonstration of stereophotogrammetry with combined SIR-B and Landsat TM images. *International Journal of Remote Sensing* 9, 1023- 38.
- Bonan GB (2008) Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science* 320:1444–1449.
- Bork, E. W., Su, J. G., (2007). Integrating LIDAR data and multispectral imagery for enhanced classification of rangeland vegetation: A meta analysis. *Remote Sensing of Environment*, 111(2007), pp. 11–24.
- Bowen, M. E., C. A. McAlpine, A. P. N. House, and G. C. Smith. (2007). Regrowth forests on abandoned agricultural land: A review of their habitat values for recovering forest fauna. *Biological Conservation* 140:273w-296.
- Boyd, D. and F. Danson (2005). "Satellite remote sensing of forest resources: three decades of research development." *Progress in Physical Geography* 29(1): 1-26.
- Brandt Tso, Paul Mather, (2009), *Classification Methods For Remotely Sensed Data*, Taylor & Francis Group, LLC.
- Brekke, C and Solberg, A. H. S. (2005). "Oil spill detection by satellite remote sensing," *Remote Sens. Environ.*, vol. 95, no. 1, pp. 1–13, Mar.
- Brisco, B. and Brown, R. J. (1995). Multi date SAR/TM synergism for crop classification in western Canada," *Photogrammetric Engineering and Remote Sensing*, Vol. 61, No. 7, 1009{1014.
- Broich, M., Hansen, M. C., Potapov, P. V., Stehman, S. V. (2011). Time-series analysis of multiresolution optical imagery for quantifying forest cover loss in Sumatra and Kalimantan, Indonesia. *J. Appl. Earth Obs. Geoinform.* 13, 277–291.

- Bryan, J.E., Shearman, P.L., Asner, G.P., Knapp, D.E., Aoro, G., Lokes, B., (2013). Extreme differences in forest degradation in Borneo: comparing practices in Sarawak, Sabah, and Brunei. *PLoS ONE* 8, e69679. <http://dx.doi.org/10.1371/journal.pone.0069679>.
- Bucki, M., Cuypers, D., Mayaux, P., Achard, F., Estreguil, C., Grassi, G. (2012). Assessing REDD+performance of countries with low monitoring capacities: the matrix approach. *Environ. Res. Lett.* 7, 014031 (13pp) <http://dx.doi.org/10.1088/1748-9326/7/1/014031>.
- California Scientific, (2007). "BrainMaker Neural Network Application Examples."
- Canada Center for Remote Sensing (2008). Introduction to RADAR remote sensing.
- Carlson, K.M., Curran, L.M., Asner, G.P., McDonald Pittman, A., Trigg, S.N., Adeney, J.M., (2013). Carbon emissions from forest conversion by Kalimantan oil palm plantations. *Nature Clim. Change* 3, 283–287.
- Carmen, G. (2008). Optical And Radar Remote Sensing Applied To Agricultural Areas In Europe. Doctoral thesis. Universitat de València. Departament de Física Aplicada.
- Chambers JQ, Asner GP, Morton DC, Anderson LO, Saatchi SS, Espírito-Santo FDB (2007) Regional ecosystem structure and function: ecological insights from remote sensing of tropical forests. *Trends Ecol Evol* 22:414–423.
- Chang, C.I., (1999). Spectral information divergence for hyperspectral image analysis. *Geoscience and Remote Sensing Symposium* 1, 509–511.
- Chanussot, J., Benediktsson, J.A. and Fauvel, M. (2006). Classification of remote sensing images from urban areas using a fuzzy possibility model. *IEEE Geoscience and Remote Sensing Letters* .3, 40–4.
- Chen, D. and STOW, D.A., (2002). The effect of training strategies on supervised classification at different spatial resolution. *Photogrammetric Engineering and Remote Sensing*, 68, pp. 1155–1162.
- Cheng, L. B., Wenxing (2014). "Remote sensing image classification based on optimized support vector machine." *TELKOMNIKA Indonesian Journal of Electrical Engineering* **12**(2): 1037-1045.
- Chuvieco, E. and Huete, A. (2010). *Fundamental of satellite remote sensing*, Taylor and Francis Group, New York.

- Cihlar J., (2000), Land cover mapping of large areas from satellites: status and research priorities. *International Journal of Remote Sensing*, 21, pp. 1093–1114.
- Cihlar,J., Xiao, Q., Chen, J., Beaubien, J., Fung, K. And Latifovic, R., (1998),Classification by progressive generalization: a new automated methodology for remote sensing multispectral data. *International Journal of Remote Sensing*, 19, pp.2685–2704.
- Cingolani, A.M., Renison, D., Zak, M.R. And Cabido, M.R., (2004), Mappingvegetation in a heterogeneous mountain rangeland using Landsat data: an alternative method to define and classify land-cover units. *Remote Sensing of Environment*, 92, pp. 84–97.
- Clark, J, Zajkowski, T, Lannom, K.(2004). Remote sensing imagery support for burned area emergency response teams on (2003) southern California wildfires.
- Clark, M., Roberts, D. A. and Clark, D. B. (2005). Hyper spectral discrimination of tropical rain forest tree species at leaf to crown scales, *Remote Sensing of Environment*, 96(3- 4), 375-398.
- Cohen, W.B. and Goward, S.N. (2004). Landsat role in ecological applications of remote sensing.
- Congalton , R.G., (1991), A review of assessing the accuracy of classification of remotely sensed data. *Remote Sensing of Environment*, 37, pp. 35–46.
- Congalton, R.G. and Green, K., (1993), A practical look at the sources of confusion in error matrix generation. *Photogrammetric Engineering and Remote Sensing*, 59, pp. 641–644.
- Congalton, R.G. and Green, K., (1999), *Assessing the Accuracy of Remotely Sensed Data: Principles and practices*. Boca Raton, London, New York: Lewis Publishers.
- Coppin, P. R., and Bauer, M. E., (1994).Processing of multitemporal Landsat TM imagery to optimise extraction of forest cover change features.*IEEE Transactions on Geoscience and Remote Sensing*, 32, 918–927.
- Corbera, E., Estrada, M., Brown, K., (2010). Reducing greenhouse gas emissions from deforestation and forest degradation in developing countries: revisiting the assumptions. *Climatic Change* 100, 355–388.

- Curran, P.J., Dungan, J.L., and Gholz, H.L. (1992). Seasonal LAI in slash pine estimated with Landsat TM. *Remote Sensing of Environment* 39: 3 – 13.
- Dahl T.E. (2004). Remote Sensing as a Tool for Monitoring Wetland Habitat Change. *Monitoring Science and Technology Symposium: Unifying knowledge for sustainability in the Western Hemisphere*; 2004 September 20-24; Denver, CO. Proceedings RMRS-P-000. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Dai, X and Khorram, S. (1998). "Effects of image misregistration on the accuracy of remotely data base prepared as part of a coordinated research program on carbon fluxes in the tropics. *Chemosphere*, **29**, 5.1015-1062.
- Dai, X.L., and Khorram, S., (1999), Remotely Sensed Change detection based on Artificial Neural Networks. *Photogrammetric Engineering & Remote Sensing*, 65(10), pages 1187-1194.
- Davidar, P., Sahoo, S., Mammen, P.C., Acharya, P., Puyravaud, J.P., Arjunan, M., Garrigues, J.P. and Roessingh, K. (2010). Assessing the extent and causes of forest degradation in India: Where do we stand? *Biological Conservation* 143: 2937-2944.
- DeFries R. S. and Chan J. (2000). Multiple Criteria for Evaluating Machine Learning Algorithms for Land Cover Classification from Satellite Data. *REMOTE SENS. ENVIRON.* 74:503–515 (2000)
- Dewan, A. M. and Yamaguchi, Y. (2009). Land use and land cover change in greater Dhaka, Bangladesh: Using remote sensing to promote sustainable urbanization. *Applied Geography*, 29 (3): 390-401.
- Ding, H., Wang, R. C., Wu, J. P., Zhou, B., Shi, Z., & Ding, L. X. (2007). Quantifying land use change in Zhejiang coastal region, China using multi-temporal Landsat TM/ETM+ images. *Pedosphere*, 17, 712–720.
- Dirzo, R. and Raven, P., (2003). Global state of biodiversity and loss. *Annu. Rev. Environ. Resour.*, 28, pp.137–167.
- Dominguez, S., Avouac, J-P, and Michel, R. (2003). "Horizontal coseismic deformation of the 1999 Chi-Chi earthquake measured from SPOT satellite images; implications for the seismic cycle along the western foothills of central Taiwan," *Journal of Geophysical Research*, vol. 108, no.B2, 2003.

- Droj, Gabriela (2007), The Applicability Of Fuzzy Theory In Remote Sensing Image Classification, *STUDIA University Babes-Bolyai, INFORMATICA*, Volume LII, Number 1.
- Du H., C.-I.Chang, H. Ren, C-C Chang, J. O. Jensen, and F. M. D'Amico (2004) "New hyperspectral discrimination measure for spectral characterization", *Optical Engineering*, 43, no 8, 1777–1786.
- Echeverria, C., Coomes, D. A., Hall, M., and Newton, A. C. (2008). Spatially explicit models to analyze forest loss and fragmentation between 1976 and 2020 in southern Chile. *Ecological Modelling*, 212(3e4), 439e449.
- Egan, W. G. (2003). *Optical remote sensing: science and technology*, CRC Press.
- Ehrlich D and C. Bielski. Texture based change detection of built-up on spot panchromatic imagery using pca. In *Joint Urban Remote Sensing Event*, (2011).
- Englhart, S., Keuck, V., Siegert, F., (2011). Aboveground biomass retrieval in tropical forests—The potential of combined X- and L-band SAR data use. *Remote Sens. Environ.* 115, 1260–1271.
- Envisat-Asar and Irs P6 Liss-Iii Satellite Data." A Case Study Over Tropical Moist Deciduous Forested Regions Of Karnataka, India." Vol. Xxxvii. Part B6b: 329-334.
- Estes, J.E. and Loveland, T.R., (1999), Characteristics, sources, and management of remotely-sensed data. In P. Longley,
- European Space Agency. (2009) *History of Europe in space. 50 years of earth observation.*
- FAO, (2000). *On Definitions of Forest and Forest Change. Forest Resources Assessment Programme. Working Paper No. 33.* Rome, Italy.
- FAO. (2009). *Low Greenhouse Gas Agriculture: Mitigation and Adaptation Potential of Sustainable Farming Systems.*
- FAO (2010). *Global Forest Assessment 2010.* FAO forestry paper, 163. Food and Agriculture Organization (FAO) of the United Nations (UN), Rome.
- Felde, G. W., Anderson, G. P., Adler-Golden, S. M., Matthew, M. W., and Berk, A., (2003). Analysis of Hyperion Data with the FLAASH Atmospheric Correction Algorithms and Technologies for Multispectral, Hyperspectral, and Ultraspectral Imagery IX. *SPIE Aerosense Conference*, Orlando. 21-25 April 2003.

- Feng, Y.L., Wang, J.F., Sang, W.G. (2007) Irradiance acclimation, capture ability, and efficiency in invasive and non-invasive alien plant species. *Photosynthetica* 45: 245-253.
- Finn, J.T., (1993), Use of the average mutual information index in evaluating classification error and consistency. *International Journal of Geographical Information Systems*, 7, pp. 349–366.
- Foody, G.M., (1996), Approaches for the production and evaluation of fuzzy land coverclassification from remotely-sensed data. *International Journal of Remote Sensing*, 17,pp. 1317–1340.
- Foody, G.M., Palubinsakas, G., Lucas, R.M., Curran, P.J., and Honzak, M., (1996), Identifying Terrestrial Carbon Sinks: Classification of Successional Stages in Regenerating Tropical Forest from Landsat TM Data, *Remote Sens. Environ.*, 55:205-216.
- Foody, G.M., (2003). Remote sensing of tropical forest environments: towards the monitoring of Forest, Wildlife and related Threats on the Island of Borneo. WWF Germany.
- Fortuin, R. (2006). Soil Erosion in Cameron Highlands, an Erosion Rate Study of a Highland Area. Saxion University Deventer. fragmentation and its impact on species diversity: an analysis using remote sensing and GIS. *Biology and Conservation*, 14, 1681-1698.
- Franke, J., Navratil, P., Keuck, V., Peterson, K., Siegert, F., (2012). Monitoring fire and selective logging activities in tropical peat swamp forests. *IEEE J. Sel.Top. Appl. Earth Obs. Remote Sens.* 5, 1811–1820.
- Franklin, S.E. (2001). Remote Sensing for Sustainable Forest Management. Lewis Publishers. Boca Raton.
- Franklin, S.E., Peddle, D.R., Dechka, J.A. And Stenhouse, G.B., (2002), Evidential reasoning with Landsat TM, DEM and GIS data for land cover classification In support of grizzly bear habitat mapping. *International Journal of Remote Sensing*, 23,pp. 4633–4652.
- Franklin SE, Wulder MA, Gerylo GR. (2001). Texture analysis of IKONOS panchromatic imagery for Douglas-fir forest age class separability in British Columbia. *International Journal of Remote Sensing* 22: 2627–2632.
- Friedl, M. A., and Brodley, C. E., (1997). Decision tree classification of land cover from remotely sensed data. *Remote Sensing of Environment*, 61, 399–409.

- Fuller, D. O. (2006). "Tropical forest monitoring and remote sensing: A new era of transparency in forest governance?" *Singapore Journal of Tropical Geography* **27**(1): 15-29.
- Funk, M. W. (2006). *Image Registration in Airborne Remote Sensing*, University of North Carolina at Charlotte.
- Gallego, D. , Canovas, F., Esteve, M.A. & Galian, J. (2004). Descriptive biogeography of *Tomicus* (Coleoptera: Scolytidae) in Spain. *Journal of Biogeography*, 31: 2011-2024.
- Gamba, P. and Chanussot, J. (2008).Foreword to the special issue on data fusion.*IEEE Transaction on Geo Science and Remote Sensing* 46, 1283-88.
- Gao, F., Masek, J., Schwaller, M. and Hall, F. (2006). On the blending of the Landsat and MODIS surface reflectance: predicting daily Landsat surface reflectance. *IEEE Transactions on Geoscience and Remote Sensing* 44, 2207-18.
- Gardner, T.A., Ribeiro-Júnior, M.A., Barlow, J., Cristina, T., Ávila-Pires, S., Hoogmoed, M.S. and Peres, C.A. (2007) The Value of primary, secondary and plantation forests for a Neotropical herpetofauna. *Conservation Biology*, **21**, 775-787.
- Geist HJ, Lambin EF (2001) Proximate causes and underlying driving forces of tropical deforestation. *Bioscience* 52:143–150.
- Gibson, P. and Power, C.(2000). *Introductory remote sensing: digital image processing and applications*. Routledge, London.
- Giridhar, M. V. and. Viswanadh, G. K. (2008).Evaluation of watershed parameters using RS and GIS.*Earth and Space Conference: Proceedings of the 11th Aerospace Division International Conference on Engineering, Science, Construction, and Operations in Challenging Environments* 323.
- Goetz, S., A. Baccini, N. Laporte, T. Johns, W. Walker, J. Kellndorfer, R. Houghton, and M. Sun. (2009). "Mapping and Monitoring Carbon Stocks with Satellite Observations: A Comparison of Methods." *Carbon Balance and Management* 4 (2): 1–7.
- Gong, P., G.S. Biging, S.M. Lee, X. Mei, Y. Sheng, R. Pu, B. Xu, K. Schwarz, and M. Mostafa, (1999). Photo ecometrics for forest inventory, *Geographic Information Science*, 5:9–14.
- Gonzalez, R. C., and Woods, R. E., (1992).*Digital Image Processing*.

- Gopal, S. and Woodcock, C., (1994), Theory and methods for accuracy assessment of thematic maps using fuzzy sets. *Photogrammetric Engineering and Remote Sensing*, 60, pp. 181–188.
- Guariguata, M.R., Nasi, R., Kanninen, M., (2009). Forest degradation: it is not a matter of new definitions. *Conservation Letters* 2, 286–287.
- Gunarso Petrus, Manjela Eko Hartoyo, Fahmuddin Agus and Timothy J. Killeen .(2013). Oil palm and land use change in Indonesia, Malaysia and Papua New Guinea .Reports from the Technical Panels of the 2nd Greenhouse Gas Working Group of the Roundtable on Sustainable Palm Oil (RSPO).
- Günlü, A., Zeki, E.B., Kadioğulları, I.A. and Altun, L. (2008). Forest site classification using Landsat 7 ETM data: A case study of Maçka-Ormanüstü forest, Turkey. *Environment Monitoring and Assessment*, DOI 10.1007/s10661-008-0252-3.
- Haala ,N and Brenner, C.(1999). Extraction of buildings and trees in urban environments. *ISPRS Journal of Photogrammetry and Remote Sensing*,54:130|137, 1999.
- Healey, S.P., Cohen, W.B., Zhiqiang, Y., Krankina, O.N.(2005). Comparison of tasseled-cap-based landsat data structures for use in forest disturbance detection. *Remote Sens. Environ.* 7, 301–310.
- Hall, G.F., D.E. Strelbel and Sellers, P.J. (1988). Linking knowledge among spatial scales: vegetation, atmosphere climate and remote sensing. *Landscape Eco*, 2: 3-22.
- Häme, T., Kilpi, J., Ahola, H.A., Rauste, Y., Antropov, O., Rautiainen, M., Sirro, L., Bounpone, S. (2013). Improved mapping of tropical forests with optical and SAR imagery, Part I: forest cover and accuracy assessment using multi-resolution data. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* 6, 74–91.
- Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R., Kommareddy, A., Egorov, A., Chini, L., Justice, C.O., Townshend, J.R.G.(2013). High-resolution global maps of 21st-century forest cover change. *Science* 342, 850–853.
- Hardin, P.J. and Shumway, J.M., (1997), Statistical significance and normalized confusion matrices. *Photogrammetric Engineering and Remote Sensing*, 63, pp. 735–740.

- Harvey, K.R, Hill, J.E .(2001). Vegetation mapping of a tropical freshwater swamp in the Northern Territory, Australia: a comparison of aerial photography, Landsat TM and SPOT satellite imagery. *Remote Sens Environ* 22:2911–2925.
- Hegarar-Masclé, S.L., Bloch, I. and Vidal-Madjar, D.(1998).Introduction of neighborhood information in evidence theory and application to data fusion of radar and optical images with partial cloud cover. *Pattern Recognition* 31, 1811-23.
- Hepner, G. F., Logan, T., Ritter, N. and Bryant, N. (1990) Artificial neural network classification using a minimal training set: comparison to conventional supervised classification. *Photogrammetric Engineering and Remote Sensing*, 56,469-473.
- Herold, M., Román-Cuesta, R.M., Hirata, Y., Van Laake, P., Asner, G., Souza, C., Avitabile, V., Skutsch, M., MacDicken, K., (2011). Options for monitoring and estimating historical carbon emissions from forest degradation in the context of REDD+. *Carbon Balance and Management* 6 (13).
- Herold, M., Skutch, M., (2011). Monitoring, reporting and verification for national REDD+ programmes: two proposals. *Environmental Research Letters* 6 , <http://dx.doi.org/10.1088/1748-9326/6/1/014002>.
- Herren HR, Binns P, Najam A, Khan S, Halberg N, Cim Li Ching, Treyer S, Lal R, Atapattu S, Kodituwakku D, Tamo M, Neuenschwander P, Fernandez M, Bassi A, Kassam A, De Fraiture C, Kingamkono R, Perfecto Y, Khan ZR .(2011). *UNEP Green Economy Report: Agriculture*. Geneva (forthcoming).
- Hett, C., Castella, J.-C., Heinemann, A., Messerli, P., Pfund, J.L.(2012). A landscape mosaics approach for characterizing swidden systems from a REDD+ perspective. *Appl. Geogr.* 32, 608–618.
- Holmgren., P. (2008). Role of satellite remote sensing in REDD .MRV Working Paper.
- Hu Z., J. W. Lee, K. Chandran, S. Kim, K. Sharma, A. C. Brotto, and S. K. Khanal (2013). Nitrogen transformations in intensive aquaculture system and its implication to climate change through nitrous oxide emission. *Bioresource Technology* 130, 314–320. doi: 10.1016/j.biortech.2012.12.033, ISSN: 09608524.

- Huang, C., Goward, S.N., Masek, J.G., Thomas, N., Zhu, Z., Vogelmann, J.E., (2010). An automated approach for reconstructing recent forest disturbance history using dense Landsat time series stacks. *Remote Sens. Environ.* 114, 183–198.
- Hubert-Moy, L., Cotonnec, A., Le Du, L., Chardin, A. And Perez, P., (2001), A comparison of parametric classification procedures of remotely sensed data applied on different landscape units. *Remote Sensing of Environment*, 75, pp. 174–187.
- Idrisi Klimanjaro, (2004), *Guide to GIS and Image Processing Volume 2*, Idrisi Production, Clark Labs, USA, pp.57-82
- Ismail, M, H., Othman, K., C., Abd Malek, A., I and Abdullah, A., S. (2012). *Land Use Trends Analysis Using SPOT-5 Images and Its Effect on the Landscape of Cameron Highland, Malaysia*
- Jablonski, D. A. (2010). *NDVI and panchromatic image correlation using texture analysis*, DTIC Document.
- Jahne, B. (1991). *Digital Image Processing*, New York: Springer-Verlag.
- Jarvis, C. H., and Stuart, N. (1996). "The sensitivity of a neural network for classifying remotely sensed imagery." *Computers & Geosciences*, 22(9), 959-967.
- Jensen, J. R. (1996) *Introductory Digital Image Processing - A Remote Sensing Perspective*. (London: Prentice Hall).
- Jensen, J. R., Qiu, F., and Ji, M. (1999). "Predictive modelling of coniferous forest age using statistical and artificial neural network approaches applied to remote sensor data." *International Journal of Remote Sensing*, 20(14), 2805-2822.
- Jensen, J.R. and Cowen, D.C., (1999), *Remote sensing of urban/suburban infrastructure and socioeconomic attributes*. *Photogrammetric Engineering and Remote Sensing*, 65, pp.611–622.
- Jensen, J.R. (2000). *Remote Sensing Of The Environment: An Earth Resource Perspective*, Prentice-Hall, Upper Saddle River, NJ.
- Jensen, J. R. (2005). *Introductory Digital Image Processing- A remote sensing perspective*, Pearson.

- Ji, C. Y. (2000). "Land-Use Classification of Remotely Sensed data Using Kohonen Self-Organizing Feature Map Neural Networks." *Photogrammetric Engineering & Remote Sensing*, 66(12), 1451-1460.
- Joseph S (2008) Assessment of landcover dynamics and its conservation implications in tropical forests of Western Ghats. In: India student conference on conservation science, Cambridge University, UK.
- Joshi, C., De Leeuw, J., Skidmore, A.K., van Duren, I.C., van Oosten, H.(2006). Remotely sensed estimation of forest canopy density: a comparison of the performance of four methods. *Int. J. Appl. Earth Observ. Geoinf.* 8, 84–95.
- Joyce, K. E. B., Stella E Samsonov, Sergey V McNeill, Stephen J Glassey, Phil J (2009). "A review of the status of satellite remote sensing and image processing techniques for mapping natural hazards and disasters." *Progress in Physical Geography* **33**(2): 183-207.
- Jubanski, J., Ballhorn, U., Kronseder, K., Franke, J., Siegert, F., (2013). Detection of large above-ground biomass variability in lowland forest ecosystems by airborne LiDAR. *Biogeosciences* 10, 3917–3930.
- Kandrika, S., and Roy, P. S. (2008). Land use land cover classification of Orissa using multitemporal IRS-P6 AWiFS data: A decision tree approach. *International Journal of Applied Earth Observation and Geoinformation*, 10(2), 186–193.
- Kaur, J. K., Kirandeep (2012). "Remote Image Classification Using Particle Swarm Optimization."
- Kavzoglu, (2001), An Investigation of the design and use offered-forward artificial neural networks in the classification of remote sensed images.
- Kennedy, J. and Eberhart, R. (1995). Particle swarm optimization, *Proceeding. IEEE International Conference on Neural Networks (ICNN)*, Nov./Dec., Australia, Pages 1942–1948.
- Kennedy, R.E., Yang, Z., Cohen, W.B., (2010). Detecting trends in forest disturbance and recovery using yearly Landsat time series: 1. LandTrendr — temporal segmentation algorithms. *Remote Sens. Environ.* 114, 2897–2910.
- Key, J., Yang, P., Baum, B. and Nasiri, S. (2002). Parameterization of shortwave ice cloud optical properties for various particle habits. *Journal of Geophysical Research*, 107: 4181-4191.

- Khoi, D. D., & Murayama, Y. (2011). Modeling deforestation using a neural network Markov model. In Y. Murayama, & R. B. Thapa (Eds.), *Spatial analysis and modeling in geographical transformation process: GIS-based applications* (pp. 169-190). Dordrecht: Springer.
- Kim, J. R. and Muller, J. P. (2002). 3D reconstruction from very high resolution satellite stereo and its application to object identification. In ISPRS, Ottawa, Canada, 2002. July 8-12.
- Knorn, J., Kuemmerle, T., Radeloff, V. C., Szabo, A., Mindrescu, M., Keeton, W. S. (2012). Forest restitution and protected area effectiveness in post-socialist Romania. *Biological Conservation*, 146, 204–212.
- Komlos, L. (2008). Examination of land use/land cover changes in the vicinity of Playa del Carmen, Mexico and their impact on groundwater quality. MA thesis. Fayetteville: University of Arkansas, Department of Geography.
- Kovacs, J.M., Wang, J.F., Flores-Verdugo, F. (2005). Mapping mangrove leaf area index at the species level using IKONOS and LAI-2000 sensors for the Agua Brava Lagoon, Mexican Pacific. *Estuar Coast Shelf Sci* 62:377–384.
- Kronseder, K., Ballhorn, U., Böhm, V., Siegert, F., (2012). Above ground biomass estimation across forest types at different degradation levels in central Kalimantan using LiDAR data. *Int. J. Appl. Earth Observ. Geoinf.* 18, 37–48
- Kumar, U. (2006). Comparative Evaluation of the Algorithms for Land Cover Mapping using Hyperspectral Data. Master Thesis submitted to the International Institute for Geoinformation Science and Earth Observation in partial fulfillment of the requirements.
- Kumar, J., (2011). Mapping and Analysis of Land Use/Land Cover of Kanpur City Using Remote Sensing and GIS Technique, 2006. *Transaction Institute of Indian Geographers*. 33 (1), 44-53.
- Kumar, D., S. Borah and U. Shankar, (2010). Monitoring forest cover change using remote sensing in Amchang Wildlife Sanctuary, Assam, India. Communicated data.
- Kuplich, T.M., (2006). Classifying regenerating forest stages in Amazonia using remotely sensed images and a neural network. *Forest Ecology and Management*, 234(1-3): 1-9.
- Landgrebe, D.A., (2003), *Signal Theory Methods in Multispectral Remote Sensing* (Hoboken, NJ: John Wiley and Sons).

- Lambin, E.F., (1999). Monitoring forest degradation in tropical regions by remote sensing: some methodological issues. *Global Ecology and Biogeography* 8 (3/4) 191–198.
- Lambin, E.; Geist, H. & Lepers, E., (2003). Dynamics of land-use and land-cover change in tropical regions. *Annual Review of Environment and Resources* 28, pp. 205-241.
- Langner, A. (2009). *Monitoring Tropical Forest Degradation and Deforestation in Borneo, Southeast Asia*.
- Langner, A., Samejima, H., Ong, R.C., Titin, J., Kitayama, K., (2012). Integration of carbon conservation into sustainable forest management using high resolution satellite imagery: a case study in Sabah, Malaysian Borneo. *Int. J. Appl. Earth Observ. Geoinf.* 18, 305–312.
- Laurina, G. Vaglio Gaia,*, Veraldo Liesenberg, Qi Chend, Leila Guerriero, Fabio Del Frate, Antonio Bartolini, David Coomes, Beccy Wileborec, Jeremy Lindsell, Riccardo Valentini, (2013). Optical and SAR sensor synergies for forest and land cover mapping in a tropical site in West Africa. *International Journal of Applied Earth Observation and Geoinformation* 21 (2013) 7–16.
- Lee, D., Lee, K., and Lee, S. (2008). Fusion of LiDAR and imagery for reliable building extraction. *Photogrammetric Engineering and Remote Sensing, Journal of the American Society for Photogrammetry and Remote Sensing*, 74(2):215–225.
- Lefsky, M.A. and Cohen, W.B., (2003), Selection of remotely sensed data. In M.A. Wulder and S.E. Franklin (Eds), *Remote Sensing of Forest Environments: Concepts and case studies*, pp. 13–46, Boston: Kluwer Academic Publishers.
- Leong, Y.K. (1992.) *Conservation and Development of Cameron Highlands*. Hill Development Proceedings of the Seminar, pp. 20-32.
- Leprince, S., Barbot, S., Ayoub, F. and Avouac, J. (2007). Automatic and precise orthorectification, coregistration, and subpixel correlation of satellite images, application to ground deformation measurements. *IEEE Transaction Geoscience and Remote Sensing* 45, 1529–58.
- Levin, N. (1999). *Fundamentals of Remote Sensing*.
- Liang, W., D. Hoja, M. Schmitt, and U. Stilla. Comparative study of change detection for reconstruction monitoring based on very high resolution optical data. In

- U. Stilla, P. Gamba, C. Juergens, and D. Maktav, editors, IEEE GRSS and ISPRS Joint Urban Remote Sensing Event, (2011).
- Li L, Ustin SL, Lay M (2005) Application of multiple end member spectral mixture analysis (MESMA) to AVIRIS imagery for coastal salt marsh mapping: a case study in China Camp, CA, USA. *Int J Remote Sens* 26:5193–5207.
- Li, Z. F., Jefferson M (2011). "Integrating Mahalanobis typicality's with a neural network for rubber distribution mapping." *Remote Sensing Letters* 2(2): 157-166.
- Lillesand, M. T., Kiefer, W. R. and Chipman, N, J. (2008). *Remote Sensing And Image Interpretation* (6th ed). John Wiley and Sons, Inc, New York.
- Lillesand, T.M. and Kiefer, R.W. (2000) *Remote Sensing and Digital Image Interpretation*, Wiley, New York, 724 p.
- Liu, J., M. Liu, D. Zhuang, Z. Zhang and X. Deng. (2003). Study on spatial pattern of land-use change in China during 1995-2000. *Sci. China Ser. D Earth Sci.*, 46: 373-378.
- Lu, D. Weng, Q (2007). "A survey of image classification methods and techniques for improving classification performance." *International Journal of Remote Sensing* 28(5): 823-870.
- Lu D, M. Batistella, and E. Moran, "Satellite estimation of aboveground biomass and impacts of forest stand structure," *Photogrammetric Engineering and Remote Sensing*, vol. 71, no. 8, pp. 967–974, (2005). View at Scopus.
- Lucas, R. M., Honzak, M., Do Amaral, I., Curran, P. J., and Foody, G. M., (2002), Forest regeneration on abandoned clearances in central Amazonia. *International Journal of Remote Sensing*, 23, 965–988.
- Lucas. C., M. , Schöngart, J. , Sheikh., P., Wittmann, F., Piedade, M, McGrath, D.(2014). Effects of land-use and hydroperiod on aboveground biomass and productivity of secondary Amazonian floodplain forests.
- Lund, H.G., (2002). When is a forest not a forest? *Journal of Forestry* 100, 21–28.
- Lund, H. G. (coord.), (2007). *Definitions of Forest, Deforestation, Afforestation, and Reforestation*. [Online] Gainesville, VA: Forest Information Services. Available at: [http:// home.comcast.net/gyde/DEFpaper.htm](http://home.comcast.net/gyde/DEFpaper.htm) (accessed: 02.05.2013).
- Maeda, E. E., Almeida, C. M., Ximenes, A. C., Formaggio, A. R., Shimabukuro, Y. E., & Pellikka, P. (2011). Dynamic modeling of forest conversion: simulation

- of past and future scenarios of rural activities expansion in the fringes of the Xingu National Park, Brazilian Amazon. *International Journal of Applied Earth Observation and Geoinformation*, 13(3), 435e446.
- Maikhuri, R.K., Rao, K.S. and Semwal, R.L. (2001). Changing scenario of Himalayan agroecosystem: Loss of agrobiodiversity, an indicator of environmental change in central Himalaya, India. *The Environmentalist* 20:23-39.
- Makalahmad, A., Eisakhani, M. and Isa, M.H., (2008). Developing MIKE-11 Model for Water Quality Simulation in Bertam River, Cameron Highlands. In: *International Conference on Construction and Building Technology – ICCBT 2008*, 16th – 20th June 2008, Kuala Lumpur, Malaysia. 8 pp.
- Makhfi, P. (2007). "Introduction to Knowledge Modeling."
- Martinez Morales, R., Idol, T. and Friday, J.B. (2011). Assessment of Acacia koa forest health across environmental gradients in Hawai‘i using fine resolution remote sensing and GIS, *Sensors*, 11, 5677-5694; doi:10.3390/s110605677.
- Masek, J.G., Huang, C., Wolfe, R., Cohen, W., Hall, F., Kutler, J., Nelson, P., (2008). North American forest disturbance mapped from a decadal landsat record. *Remote Sens. Environ.* 112, 2914–2926.
- Maselli, F., Conese, C. And Petkov, L., (1994), Use of probability entropy for the estimation and graphical representation of the accuracy of maximum likelihood classifications.
- Masser, I. (2001)."Managing our urban future: the role of remote sensing and geographic information systems."*Habitat International*25(4): 503-512.
- Mather, P. M. (2004). *Computer processing of remotely sensed images*. (3rd ed), John Wiley and Sons, Ltd.
- Matricardi, E.A.T., Skole, D.L., Pedlowski, M.A., Chomentowski, W., Fernandes, L.C., (2010). Assessment of tropical forest degradation by selective logging and fire using landsat imagery. *Remote Sens. Environ.* 114, 1117–1129.
- May, A.M.B, Pinder, J.E, Kroh, G.C .(1997). A comprison of LANDSAT Thematic Mapper and SPOT multi-spectral imagery for the classification of shrub and meadow vegetation in Northern California, USA.*Int J Remote Sens* 18:3719–3728.
- McCulloch, W. S., and Pitts, W. (1943)."A logical calculus of the ideas immanent in nervous activity."*Bulletin of Mathematical Biology*, 5, 115-133.

- MelGani F, L Bruzzone (2004) Classification of Hyperspectral Remote Sensing Images with Support Vector Machines. *IEEE Transactions on Geoscience and Remote Sensing*; 42(8): 1778-1790.
- Messerli, P., Heinimann, A., Epprecht, M., (2009). Finding homogeneity in heterogeneity - a new approach to quantifying landscape mosaics developed for Lao PDR. *Hum. Ecol.* 37, 291–304.
- Miettinen, J., Shi C., Tan W.J. & Liew S.C. (2007) Land cover map of insular Southeast Asia in 250m spatial resolution. *Remote Sensing Letters*, 3, 11-20.
- Miettinen, J., Liew, S.C., (2010). Status of peatland degradation and development in Sumatra and Kalimantan. *AMBIO* 39, 394–401.
- Miettinen, J., Stibig ,H-J., Achard, F .(2014).Remote sensing of forest degradation in Southeast Asia—Aiming for a regional view through 5–30 m satellite data .*Global Ecology and Conservation* 2 (2014) 24–36.
- Millington, A.C., X.M. Velez-Liendo and A.V. Bradley, (2003). Scale dependence in multi temporal mapping of forest fragmentation in Bolivia: implications for explaining temporal trends in landscape ecology and applications to biodiversity conservation. *ISPRS J. Photogram*, 57: 289-299.
- Ministry of Forestry (MoF), (2011). Digital Land Cover Map 2009. Ministry of Forestry, Jakarta, Indonesia (unpublished).
- Mitchard, E.T.A., Saatchi, S.S., Woodhouse, I.H., Nangendo, G., Ribeiro, N.S., Williams, M., Ryan, C.M., Lewis, S.L., Feldpausch, T.R., Meir, P., (2009). Using satellite radar backscatter to predict above-ground woody biomass: a consistent relationship across four different African landscapes. *Geophys. Res. Res.*
- Mitchard, E.T.A., Saatchi, S.S., Lewis, S.L., Feldpausch, T.R., Woodhouse, I.H., Sonké, B., Rowland, C., Meir, P., (2011). Measuring biomass changes due to woody encroachment and deforestation/degradation in a forest–savanna boundary region of central Africa using multi-temporal L-band radar backscatter. *Remote Sens. Environ.* 115, 2861–2873.
- Mohammed, J. (2013). "Land use and cover change assessment using Remote Sensing and GIS: Dohuk City, Kurdistan, Iraq (1998-2011)." *International Journal of Geomatics & Geosciences* 3 (3).
- Mon, M.S., Mizoue, N., Htun, N.Z., Kajisa, T., Yoshida, S.(2012). Estimating forest canopy density of tropical mixed deciduous vegetation using Landsat data: a

- comparison of three classification approaches. *Int. J. Remote Sens.* 33, 1042–1057.
- Morel, A.C., Saatchi, S.S., Malhi, Y., Berry, N.J., Banin, L., Burslem, D., Nilus, R., Ong, R.C., (2011). Estimating aboveground biomass in forest and oil palm plantation in Sabah, Malaysian Borneo using ALOS PALSAR data. *Forest Ecol. Manag.* 262, 1786–1798.
- Morgan, J. N. and J. A. Sonquist (1963). Problems in the analysis of survey data, and a proposal. *Journal of the American Statistical Association*, 58, 415-434.
- Morgan, J. N. and R. C. Messenger (1973). THAID a sequential analysis program for analysis of nominal scale dependent variables. Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor.
- Margono, B.A., Potapov, P.V., Turubanova, S., Stolle, F., Hansen, M.C., (2014). Primary forest cover loss in Indonesia over 2000–2012.. (advanced online publication).
- Margono, B.A., Turubanova, S., Zhuravleva, I., Potapov, P., Tyukavina, A., Baccini, A., Goetz, S., Hansen, M.C., 2012. Mapping and monitoring deforestation and forest degradation in Sumatra (Indonesia) using Landsat time series data sets from 1990 to 2010. *Environ. Res.*
- Mulligan, M. and Wainwright, J., Modelling and Model Building, in Wainwright J. and Mulligan M. (eds.), *Environmental Modelling: Finding Simplicity in Complexity*, West Sussex, England, UK: John Wiley & Sons Ltd., 7-73, (2004)
- Murdiyarso, D., van Noordwijk, M., Puntodewo, A., Widayati, A., Lusiana, B., (2008). District scale prioritization for A/R CDM project activities in Indonesia in line with sustainable development objectives. *Agriculture, Ecosystems & Environment* 126 (1–2) 59–66.
- Narimani, R. N., Ahmad (2013). "Classification credit dataset using particle swarm optimization and probabilistic neural network models based on the dynamic decay learning algorithm."
- Negron-Juarez, R.I., Chambers, J.Q., Marra, D.M., Ribeiro, G.H.P.M., Rifai, S.W., Higuchi, N., Roberts, D., (2011). Detection of subpixel treefall gaps with landsat imagery in Central Amazon forests. *Remote Sens. Environ.* 115, 3322–3328.

- Ngamabou, R. S. (2006). "Evaluating the efficacy of remote sensing techniques in monitoring forest cover and forest cover change in the Mount Cameroon region."
- Nizalapur, V. (2008). "Land Cover Classification Using Multi-Source Data Fusion.
- Ollingera S. V. , A. D. Richardsona, M. E. Martina, D. Y. Hollingerb, S. E. Froelkinga, P. B. Reichc , L. C. Plourdea, G. G. Katuld, J. W. Mungere, R. Orend, M.-L. Smithb,f, K. T. Paw Ug, P. V. Bolstadc , B. D. Cookc , M. C. Daya, T. A. Martinh, R. K. Monsoni , and H. P. Schmid . (2008) .Canopy nitrogen, carbon assimilation, and albedo in temperate and boreal forests: Functional.
- Omran M. (2005) Particle Swarm Optimization Methods For Pattern Recognition And Image Processing, PhD Thesis, University of Pretoria.
- Orekan ,V.,O.,A .(2007) Implementation of the local land-use and land-cover change model CLUE-s for Central Benin by using socio-economic and remote sensing data .PhD thesis . University of Bonn
- Otukei, J.R. and Blaschke, T. (2010). Land cover change assessment using decision trees, support vector machines and maximum likelihood classification algorithms. *International Journal of Applied Earth Observation and Geoinformation*,12, S27-S31.
- Ouchi, K. (2013). "Recent Trend and Advance of Synthetic Aperture Radar with Selected Topics." *Remote Sensing* 5(2): 716-807.
- Pais-Barbosa, J. (2007), *Hidroformas e hidromorfologias costeiras locais* (in Portuguese). PhD Thesis, Faculty of Engineering of the University of Porto, 2 Vol.
- Pal, M., and Mather, P. M. (2003). An assessment of the effectiveness of decision tree methods for land cov classification. *Remote Sensing of Environment*, 86, 554–565.
- Panta, M., Kim, K., Joshi, C., (2008). Temporal mapping of deforestation and forest degradation in Nepal: applications to forest conservation. *Forest Ecol. Manag.* 256, 1587–1595.
- Pengra BW, Johnston CA, Loveland, T.R .(2007). Mapping an invasive plant, *Phragmites australis*, in coastal wetlands using the EO-1 Hyperion hyperspectral sensor. *Remote Sens Environ* 108:74–81.

- Penuelas J, Blanchard B, Blanchard A. (1993). Assessing community type, plant biomass, pigment composition and photosynthetic efficiency of aquatic vegetation from spectral reflectance. *Remote Sens Environ* 46:110–118.
- Phinn, S.R., (1998), A framework for selecting appropriate remotely sensed data dimensions for environmental monitoring and management. *International Journal of Remote Sensing*, 19, pp. 3457–3463.
- Phinn, S.R., Menges, C., Hill, G.J.E. And Stanford, M., (2000), Optimizing remotely sensed solutions for monitoring, modeling, and managing coastal environments. *Remote Sensing of Environment*, 73, pp. 117–132.
- Pithon, S., Jubelin, G., Guitet, S., Gond, V., (2013). A statistical method for detecting logging-related canopy gaps using high-resolution optical remote sensing. *Int. J. Remote Sens.* 34, 700–711.
- Pohl, C. and van Genderen, J.L. (1998). *Multisensor Image Fusion In Remote Sensing: Concepts*.
- Potapov, P., Yaroshenko, A., Turubanova, S., Dubinin, M., Laestadius, L., Thies, C., Aksenov, D., Egorov, A., Yesipova, Y., Glushkov, I., Karpachevskiy, M., Kostikova, A., Manisha, A., Tsybikova, E., Zhuravleva, I. (2008). Mapping the world's intact forest landscapes by remote sensing. *Ecol. Soc.* 13, 51. 16p.
- Potter, C., Tan, P., Steinbach, M., Klooster, S., Kumar, V., Myneni, R., and Genovese, V. (2003). Major disturbance events in terrestrial ecosystems detected using global satellite data sets. *Global Change Biology* 9, 7, 1005–1021. ACM.
- Powell, R.L., Matzke, N., De Souza Jr, C., Clark, M., Numata, I., Hess, L.L. And Roberts, D.A., (2004), Sources of error in accuracy assessment of thematic land-cover maps in the Brazilian Amazon. *Remote Sensing of Environment*, 90, pp. 221–234.
- Prabhu, S. T., D (2013). "Efficient Technique For The Classification Of Satellite Images Using Fuzzy Rule Classifier." *Journal of Theoretical & Applied Information Technology* **53** (3).
- Proisy C, Coueron P, Fromard F. (2007). Predicting and mapping mangrove biomass from canopy grain analysis using Fourier-based textural ordination of IKONOS images. *Remote Sens Environ* 109:379–392.

- Pu, R., N. M. Kelly, Q. Chen, and Gong, P. (2008). Spectroscopic determination of health levels of Coast Live Oak (*Quercus agrifolia*) leaves. *Geocarto International*, 23(1), 3- 20.
- Putz, F.E., Redford, K.H., (2010). The importance of defining ‘forest’: tropical forest degradation, deforestation, long-term phase shifts, and further transitions. *Biotropica* 42 (1) 10–20.
- Puyravaud, J. (2007) Forest degradation in the Western Ghats biodiversity hotspot: resource collection, livelihood concerns and sustainability. *Current Science*, Vol. 93(11), pp. 1573 - 1578.
- Qin, J., Burks, T.F., Ritenour, M.A., Bonn, W.G., (2009). Detection of citrus canker using hyper spectral reflectance imaging with spectral information divergence. *Journal of Food Engineering* 93 (2), 183–191.
- Qiu, F., and J.R. Jensen, (2004). Opening the black box of neural networks for remote sensing image classification, *International Journal of Remote Sensing*, 25:1749–1768.
- Quinlan, J. R. (1975). *Machine Learning*, Volume 1(1) University of Sydney York: Lewis Publishers.
- Quinlan, J. R. (1993). *C4.5: Programs For Machine Learning*. San Mateo, California: Morgan Kauffmann Publishers.
- Radke, R. J., Andra, S., O.Al-Kofahi and Roysam, B., (2005). Image detection algorithms: a systematic survey. *IEEE Trans. Image Processing*, 14(3), pp. 294–307.
- Ramachandra TV, Kumar U (2004). Geographic Resources Decision Support System for land use, land cover dynamics analysis. *Proceedings of the FOSS/GRASS Users Conference held in Bangkok, Thailand*.
- Ramankutty, N., A.T. Evan, C. Monfreda, and J.A. Foley. (2008). Farming the planet: 1. Geographic distribution of global agricultural lands in the year 2000. *Global Biogeochemical Cycles* 22: 1-19.
- Ranchin, T. and Wald, L. (2000). Fusion of high spatial and spectral resolution images: The ARSIS concept and its implementation. *Photogrammetric Engineering and Remote Sensing* 45, 49-61.
- Rao, A. R., and Jain, R., (1998). Knowledge Representation and Control in Computer Vision Systems. *IEEE Expert System and their Applications*, pp 65-79.

- Rauste, Y., (2005). Multi-temporal JERS SAR data in boreal forest biomass mapping. *Remote sensing of Environment*, 97(2): 263-275. Relations and potential climate feedbacks.
- Rendong L, Jiyuan ,L .(2004). Estimating wetland vegetation biomass in the Poyang Lake of central China from Landsat ETM data. *IEEE Trans Geosci Remote Sen* 4:4590–4593 (IGARSS apos).
- Repaka, S.R. And Truax, D.D. (2004): Comparing spectral and object based approaches for classification and transportation feature extraction from high resolution multi-spectral imagery. *Proceedings of the ASPRS 2004 Annual Conference, Denver, USA. May 23-28.*
- Reshitnyk, L. Y. (2013). *Acoustic and Satellite Remote Sensing Of Shallow Near shore Marine Habitats In The Gwaii Haanas National Marine Conservation Area, University of Victoria.*
- Reyes C., Taylor F., Martínez E. y López F. (2006). Geo-Cybernetics: A new Avenue of Research in Geomatics, *Cartographica*, 41, Issue 1. DOI. 10.3138/C034-6P5T-W322- 1G72.
- Richards, J. A. (2013). *Remote sensing digital image analysis: an introduction, Springer.*
- Richards, J. A. and X. Jia, (2006). *Remote Sensing Digital Image Analysis (4th ed.): An Introduction. Verlag, Berlin, Heidelberg.*
- Richards, J. A. and X. Jia, (2006). *Remote Sensing Digital Image Analysis, An Introduction, Springer, Fourth Edition.*
- Rignot, E., Salas, W.A. and Skole, D.L., (1997). Mapping deforestation and secondary growth in Rondonia, Brazil, using imaging radar and thematic mapper data. *Remote Sensing of Environment*, 59(2): 167-179.
- Rikimaru, A., Roy, P.S., Miyatake, S. (2002). Tropical forest cover density mapping. *Trop. Ecol.* 43, 39–47.
- Rindfuss, R., Entwisle, B., Walsh, S. J., Prasartkul, P., Sawangdee, Y., Crawford, T. W. and Reade, T. (2002). Continuous and discrete: where they have met in Nang Rong, Thailand. In: Walsh, S. J. and Crews-Meyer, K. (eds.). *Linking people, place and policy: A GIS science approach, Kluwer Academic Press, Boston.*

- Riitters, K., Vogt, P., Soille, P., Estreguil, C., (2009). Landscape patterns from mathematical morphology on maps with contagion. *Landscape Ecology* 24, 699–709.
- Roberts, D.A., Green, R.O. and Adams, J.B. (1997). Temporal and spatial patterns in vegetation and atmospheric properties from AVIRIS. *Remote sensing of environment*, 62: 223 – 240.
- Roberts, J. V. A., J, Ahmed, FB (2008). "Medium resolution image fusion, does it enhance forest structure assessment".
- Robila, S. (2005). An investigation of spectral metrics in hyper-spectral image pre-processing for classification. *Geospatial goes global: from your neighborhood to the whole planet. ASPRS Annual Conference, Baltimore, Maryland.*
- Rogers, R.H., and Wood, L. (1990). The history and status of merging multiple sensor data.
- Romijn E, Ainembabazi JH, Wijaya A, Herold M, Angelsen A, Verchot L, Murdiyarso D (2013) Exploring different forest definitions and their impact on developing REDD+ reference emission levels: a case study for Indonesia. *Environmental Science and Policy*, 33, 246–259.
- Rosso PH, Ustin SL, Hastings A (2005) Mapping marshland vegetation of San Francisco Bay, California, using hyperspectral data. *Int J Remote Sens* 26:5169–5191.
- Rottensteiner .F, Trinder .J, Clode .S, and Kubik .K.(2003). Building detection using LIDAR data and multispectral images. In *APRS*, volume 2, pages 673{682, Sydney, 2003.
- Roughgarden, J., S.W. Running and P.A. Matson, (1991). What does remote sensing do for ecology? *Ecology*, 72: 1981-1982.
- Rudel TK (2006) Shrinking tropical forests, human agents of change, and conservation policy. *Conserv Biol* 20:1604–1609.
- Russell G. Congalton, (1991) A Review of Assessing the Accuracy of Classification of Remotely Sensed Data, *Remote Sens, Environ.* 37:35- 46.
- Ryan, C.M., Hill, T., Woollen, E., Ghee, C., Mitchard, E., Cassells, G., Grace, J., Woodhouse, I.H., Williams, M., 2012. Quantifying small-scale deforestation and forest degradation in African woodlands using radar imagery. *Global Change Biol.* 18, 243–257.

- Sadly, M and Faisal, Y.(2011). Application of a Hybrid Neural Network Model for Multispectral Remotely Sensed Image Classification in the Belopa Area South Sulawesi of Indonesia . ISBN: 978-979-1421-11-9.
- Sahar, L. (2009). "Using remote-sensing and GIS technology for automated building extraction."
- Sands, R.(2005). Forestry in a Global Context.CABI Publishing, Wallingford, UK.
- Sasaki, N., Putz, F.E., (2009). Critical need for new definitions of forest and forest degradation in global climate change agreements: policy perspective. *Conservation Letters* 2, 226–232.
- Schiewe J. (2002): Segmentation of High-Resolution Remotely Sensed Data – Concepts,
- Schmidt KS, Skidmore AK .(2003). Spectral discrimination of vegetation types in a coastal wetland. *Remote Sens Environ* 85:92–108.
- Schowengerdt, R. A. (2007). *Remote Sensing, Models And Methods For Image Processing*. (3rd ed), Elsevier Inc. New York, USA.
- Schowengerdt,R.A. (1997) *Remote Sensing, Models And Methods For Image Processing*, Second edn. Academic Press, London.
- Selvalakshmi, S. Jayakumar, V.S. Ramachandran.(2014). Are the sensitive zones degrading? A modelling approach using GIS and remote sensing.
- Shaikh M, Green D, Cross H (2001) A remote sensing approach to determine environmental flow for wetlands of lower Darling River, New South Wales, Australia. *Int J Remote Sens* 22:1737–1751.
- Shalaby, A., and Tateishi, R., (2007). Remote sensing and GIS for mapping and monitoring land cover and land-use changes in the Northwestern coastal zone of Egypt. *Applied Geography*, 27, 28–41.
- Sharma, M. G., Rashmi Kumar, Deepak Kapoor, Rajiv (2011). "Efficacious approach for satellite image classification."*Journal of Electrical and Electronics Engineering Research* 3(8): 143-150.
- Sharma, R. G., Aniruddha Joshi, PK (2013). "Decision tree approach for classification of remotely sensed satellite data using open source support."
- Shearman, P.L., Ash, J., Mackey, B., Bryan, J.E., Lokes, B., (2009). Forest conversion and degradation in Papua New Guinea 1972–2002. *Biotropica* 41, 379–390.

- Shi, Y. (2013). "A Remote Sensing and GIS-based Wetland Analysis In Canaan Valley, West Virginia."
- Shijo Joseph • M. S. R. Murthy • A. P. Thomas .(2011).The progress on remote sensing technology in identifying tropical forest degradation: a synthesis of the present knowledge and future perspectives.
- Sneath,P. H. A., And R. R. Sokal. (1973). Numerical taxonomy. W. H. Freeman, San Francisco.
- Simard, M. G., G De Saatchi, S Mayaux, P (2002)."Mapping tropical coastal vegetation using JERS-1 and ERS-1 radar data with a decision tree classifier."International Journal of Remote Sensing**23**(7): 1461-1474.
- Sneath, P. H. A. and Sokal, R. R. (1973) Numerical Taxonomy.(San Francisco:W.H. Freeman).
- Singh, Saurabh, Gupta, Abhinav, and Efros, Alexei A. Unsupervised discovery of mid-level discriminative patches. In ECCV, (2012).
- Soares-Filho, B. S., Rodrigues, H., & Follador, M. (2013). A hybrid analytical-heuristic method for calibrating land-use change models. Environmental Modelling & Software, 43, 80e87. <http://dx.doi.org/10.1016/j.envsoft.2013.01.010>.
- Sohn . Gand Dowman .I.(2007). Data fusion of high-resolution satellite imagery and LiDAR data for automatic building extraction. ISPRS Journal of Photogrammetry and Remote Sensing, 62(1):43{63, May (2007).
- Soudani, K., Francois, C., Maire, G. L., Dantec, V.L. and Dufrene, E. (2006).Comparative analysis of Ikonos, Spot, and ETM data for leaf area index estimation in temperate coniferous and deciduous forest stands. Remote Sensing of Environment, 102, 161– 175.
- Souza Jr., C., Firestone, L., Silva, M.L., Roberts, D.A., (2003). Mapping forest degradation in the Eastern Amazon from SPOT 4 through spectral mixture models. Remote Sens. Environ. 87, 494–506.
- StatSoft, I. (2003). Neural Networks.
- Stehman, S.V., (2004), A critical evaluation of the normalized error matrix in map accuracy assessment. Photogrammetric Engineering and Remote Sensing, 70, pp. 743–751.

- Stuckens, J., Coppin, P. R., & Bauer, M. E. (2000). Integrating contextual information with per-pixel classification for improved land cover classification. *Remote Sensing of Environment*, 71, 282–296.
- Tehrany, M. S., B. Pradhan, et al. (2013). "A comparative assessment between object and pixel-based classification approaches for land use/land cover mapping using SPOT 5 imagery." *Geocarto International* (ahead-of-print): 1-19.
- Thapa, R. B., & Murayama, Y. (2012). Scenario based urban growth allocation in Kathmandu Valley, Nepal. *Landscape and Urban Planning*, 105(1e2), 140e148.
- Thapa ,B .R., Shimada, M., Watanabe, M., Motohka,T., Shiraishi ,T. (2013). The tropical forest in south east Asia: Monitoring and scenario modeling using synthetic aperture radar data . *Applied Geography* 41 (2013) 168e178.
- Toriman, M., Karim, O., Mokhtar, M., Gazim, M., and Abdullah, M. (2010). Use of Info Work RS in Modeling the Impact of Urbanization on Sediment Yield in Cameron Highlands, Malaysia. *Nature and Science*.
- Tottrup, C. and Rasmussen, M. S. (2004). Mapping long-term changes in savannah crop productivity in Senegal through trend analysis of time series of remote sensing data. *Agriculture.Ecosystems and Environment*, 103(3): 545-560.
- Treitz, P., and Rogan, J. (2004). Remote sensing for mapping and monitoring land-cover and land-use change—an introduction. *Progress in Planning*, 61, 269–279.
- Tso, B. and P.M. Mather, (2001). *Classification Methods for Remotely Sensed Data*. Taylor and Francis, London.
- Turner, W., S. Spector, N. Gardiner, M. Fladeland, E. Sterling, and M. Steininger.(2003). Remote sensing for biodiversity science and conservation, *TRENDS in Ecology and Evolution*, 18(3): 306-314.
- Vahidi,H., Monabbati, e. (2013).contextual image classification approach for monitoring of agricultural land cover by support vector machines and markov random fields .
- Vaiphasa C, Ongsomwang S, Vaiphasa T, Skidmore ,AK. (2005). Tropical mangrove species discrimination using hyper spectral data: a laboratory study. *Estuary Coast Shelf Sci* 65:371–379.
- Van Der Werf, G.R., Dempewolf, J., Trigg, S.N., Randerson, J.T., Kasibhatia, P.S., Giglio, L., Murdiyarsa, D., Peters, W., Morton, D.C., Collatz, G.J., Dolman,

- A.J.& DeFries, R.S. (2008). Climate regulations of fire emissions and deforestation in equatorial Asia.
- Van Noordwijk, M., Minang, P.A., (2009). If we cannot define it, we cannot save it. ASB Policybriefs, 15 Available at: <http://www.asb.cgiar.org/pdfwebdocs/ASBPB15.pdf> (accessed: 17.04.13).
- Venkateswaran, C. J. V., R Saravanan, AM (2013). "A Fuzzy Based Approach to Classify Remotely Sensed Images." *International Journal of Engineering and Technology*.
- Verchot, L.V., Zomer, R.b., Van Straaten, O., Muys, B., (2007). Implications of country-level decisions on the specification of crown cover in the definition of forests for land area eligible for afforestation and reforestation activities in the CDM. *Climatic Change* 81 (3–4) 415–430.
- Vögtle T., Schilling K.-J., (1995): Wissensbasierte Extraktion von Siedlungsbereichen in der Satellitenbildanalyse, *ZPF* (63) 5, 199-207.
- Walter, V., (2004). Object-based evaluation of LIDAR and multispectral data for automatic change detection in GIS databases, *IAPRS*, Vol. 35, Part B2, pp.723-728.
- Waltz. E (2001).*The Principles And Practice Of Image And Spatial Data Fusion*. In *Multisensory Data Fusion*.CRC Press, D. Hall and J. Llinas edition, 2001.
- Wardle DA, Walker LR, Bardgett RD (2004) Ecosystem properties and forest decline in contrasting long-term chronosequences. *Science* 305:509–513.
- Wenbo .W, Jing .Y, and Tingjun K. K.(2008) Study of remote sensing image fusion and its application in image classification. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*,37:1141{1146, 2008.
- Weng, Q. (2010). *Remote Sensing Of Impervious Surfaces*, CRC Press.
- Wijedasa, L.S., Sloan, S., Michelakis, D.G., Clements, G.R., (2012). Overcoming limitations with landsat imagery for mapping of peat swamp forests in Sundaland. *Remote Sens.* 4, 2595–2618.
- Wilkie, D. S. ,and Finn, J. T. (1996). *Remote sensing imagery for natural resources monitoring: A guide for first-time users*. New York: Columbia University Press.
- Wilson, J. P. and Fotheringham S. A. (2008).*The Hand Book Of Geographic Information Science*. Blackwell publishing, Victoria, Australia.

- Woodcock, C. E. and Strahler, A. H. (1987) The factor of scale in remote sensing. *Remote Sensing of Environment*, 21, 311-332.
- Woodcock, C.E. And Gopal, S., (2000), Fuzzy set theory and thematic maps: accuracy assessment and area estimation. *International Journal of Geographic Information Science*, 14, pp. 153–172.
- Wright, S. J. and Muller-Landau, H. C. (2006).The future of tropical forest species.*Biotropica* 38, 3: 287–3011. doi:10.1111/j.1744- 7429.2006.00154.x
- Xu, H. (2007). Extraction of Urban Built-up Land Features from Landsat Imagery Using a Thematicoriented Index Combination Technique. *Photogrammetric Engineering and Remote Sensing*, 73, 12:1381–1391.
- Xu, R. a. W., Donald (2005)."Survey of clustering algorithms." *Neural Networks, IEEE Transactions on* **16**(3): 645-678.
- Yang, J. G., Peng Fu, Rong Zhang, Minghua Chen, Jingming Liang, Shunlin Xu, Bing Shi, Jiancheng Dickinson, Robert (2013). "The role of satellite remote sensing in climate change studies." *Nature Climate Change* **3**(10): 875-883.
- Yiming Y (1994). *Rules Based Fuzzy Logic Inference*. *IEEE Int. Conf. Syst. Man Cybernet.*, 1: 465-470.
- Zakaria .H. E. (2010). *Integration of Remote Sensing and GIS in Studying Vegetation Trends and Conditions in the Gum Arabic Belt in North Kordofan, Sudan* .Dissertation of Doctor of Natural Science. Technical University of Dresden.
- Zhang, J.W., Oliver, W.W. and Busse, M.D. (2006). Growth and development of ponderosa pine on sites of contrasting productivities: relative importance of stand density and shrub competition effects. *Can. J. For. Res.* 36, 2426–2438.
- Zhang, J.W., Powers, R.F. and Skinner, C.N. (2010). To manage or not to manage: the role of silviculture in sequestering carbon in the specter of climate change. In *Integrated Management of Carbon Sequestration and Biomass Utilization Opportunities in a Changing Climate*. T.B. Jain, R.T. Graham and J. Sandquist (tech. eds.), pp. 95–110. *Proceedings of the 2009 National Silviculture Workshop*, June 15–18, 2009; Boise, Idaho. RMRS-P-61.
- Zhou, G., Baysal, O., Kaye, J., Habib, S., Wang, C., (2004).Concept design of future intelligent.