

IMPROVEMENT OF CYCLODEXTRIN GLUCANOTRANFERASE
EXCRETION AND CELL VIABILITY OF RECOMBINANT
Escherichia coli IMMOBILIZED ON HOLLOW
FIBER MEMBRANE

ROHAIDA BINTI CHE MAN

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (Bioprocess Engineering)

Faculty of Chemical and Energy Engineering
Universiti Teknologi Malaysia

MARCH 2016

To my beloved parents (Hj. Che Man bin Mamat and Hjh. Hasmah binti Mohd Noh), my husband (Saiful Aflah bin Abdol Karim), daughter (Nur Alya Safiah), son (Muhammad Aiman Rafiqin), sisters (Fauziah, Rohaniah, Rohaniza and Rohaina) and brothers (Mohd Hatta, Mohd Lutffi, Mohd Asri, Mohd Nizam, Muhammad and Mohd Hafizullah). I dedicated this work in sincere gratitude for their patience, love and support.

ACKNOWLEDGEMENT

Bismillahirrahmanirahim....

In the name of Allah, The Most Gracious, The Most Merciful. Praise is to Allah S.W.T by whose grace and blessing I receive guidance in completing this study. Thanks for His greatest love and blessing.

First of all, I wish to convey my deepest appreciation and sincere thanks to my supervisor co-supervisor, Prof. Dr. Rosli bin Md Illias and Prof. Dr. Ahmad Fauzi bin Ismail for the advice, guidance and criticisms throughout this study. I'm very much indebted to them.

I gratefully acknowledge Universiti Malaysia Pahang for providing the scholarship during my study.

I would like to express my deepest gratitude to my research associate in genetic laboratory (Yan, Has, Bai, Abbas, Shalyda, Iza, Atul, Intan, Aisyah, Ling, Amal, Hazlin, Faiz, Joyce, Dr Tim, Dr Aizi, Dr Anuar, Sammy, Kimi, Dayah, Dilin, Ummu, Joanne, and Yeng) for their help and friendship during study.

I wish to show my appreciation to Mr Yaakop, Mr Jefri, Mr Razis and Mr Bee Chee for their help during the experimental set up and invaluable assistance. Without their help, the labwork might not be complete successfully.

Last but not least, I want to extend my gratitude to my parents, husband, daughter, son, sisters and brothers for their love and support through the years.

ABSTRACT

The excretion of a recombinant enzyme into culture medium presents significant advantages over cytoplasmic expression. However, cell lysis is one of the major drawbacks during the excretion of recombinant enzyme when using *Escherichia coli* (*E. coli*) as a host. Cell immobilization is a promising solution for the enhancement of enzyme excretion with reduction of cell lysis. In the present study, a recombinant *E. coli* was immobilized using hollow fiber membrane to improve the enzyme excretion, cell viability and plasmid stability. The effects of different polymers of hollow fiber membrane and culture conditions on the cyclodextrin glucanotransferase (CGTase) excretion, cell lysis and plasmid stability of immobilized *E. coli* were investigated. The immobilized cells on a polyvinylidene fluoride polymer exhibited a 2.0–4.5-fold increase in the CGTase excretion with 18–95% reduction of cell lysis and over 100% increment of plasmid stability compared to the free cells. The CGTase excretion was successfully optimized by response surface methodology. Under the optimized conditions [25 °C of post- induction temperature, 0.011 mM isopropyl β -D-1-thiogalactopyronoside and pH 8.8], the CGTase excretion was 3.8-fold higher with 80% reduction of cell lysis compared to the value before optimization process. The use of low tryptone concentration (5 g/l) reduced the occurrence of cell lysis (90% reduction) and increased the plasmid stability (86% increment) without significant change in CGTase excretion in comparison with initial tryptone concentration (20 g/l). This approach (5 g/l) produced an approximately two times higher CGTase excretion (compared with 20 g/l during) recycle process. The membrane bioreactor also showed 2.5-fold increase in the CGTase excretion (473×10^3 U/ml) with 75% reduction of cell lysis compared to shake flask culture (190×10^3 U/ml of CGTase activity). Hence, the immobilization of *E. coli* on hollow fiber membrane proved to be valuable for the excretion of recombinant proteins in *E. coli* with high cell stability.

ABSTRAK

Perembesan enzim rekombinan ke dalam media kultur adalah pendekatan yang lebih baik berbanding pengungkapan sitoplasmik. Walau bagaimanapun, lisis sel adalah salah satu masalah utama dalam perembesan enzim rekombinan apabila menggunakan *Escherichia coli* (*E. coli*) sebagai perumah. Imobilisasi sel adalah penyelesaian yang baik untuk peningkatan perembesan enzim dengan pengurangan kadar lisis sel. Dalam kajian ini, *E. coli* rekombinan telah diimobilisasikan menggunakan polimer membran gentian berongga bertujuan untuk meningkatkan perembesan enzim, bilangan sel hidup dan kestabilan plasmid. Kesan polimer membran gentian berongga yang berbeza dan keadaan pertumbuhan untuk perembesan siklodextrin glukano transferase (CGTase), kadar lisis sel dan kestabilan plasmid bagi sel imobilisasi telah dikaji. Sel imobilisasi pada poliviniliden fluorida polimer mempamerkan 2.0-4.5 kali ganda peningkatan dalam perembesan CGTase dengan 18-95% pengurangan kadar lisis sel dan peningkatan kestabilan plasmid melebihi 100% berbanding dengan sel bebas. Perembesan CGTase berjaya dioptimumkan dengan menggunakan kaedah gerak balas permukaan. Dengan menggunakan keadaan optimum [25 °C suhu induksi, 0.011 mM isopropil β -D-1-thiogalaktopiranosida dan pH 8.8], perembesan CGTase adalah 3.8 kali ganda tinggi dengan pengurangan kadar lisis sel sebanyak 80% berbanding dengan nilai sebelum proses pengoptimuman. Penggunaan kepekatan tripton yang rendah (5 g/l) mengurangkan kadar lisis sel dengan 90% pengurangan dan meningkatkan kestabilan plasmid (86% peningkatan) tanpa perubahan perembesan CGTase yang ketara berbanding dengan kepekatan tripton yang asal (20 g/l). Pendekatan ini (5 g/l) membuktikan penghasilan perembesan CGTase dengan kira-kira dua kali ganda berbanding 20 g/l sepanjang proses berulang. Bioreaktor membran juga menunjukkan peningkatan perembesan CGTase (473×10^3 U/ml) sebanyak 2.5 kali ganda dengan pengurangan lisis sel sebanyak 75% berbanding kelalang kon (190×10^3 U/ml aktiviti CGTase). Oleh itu, sel *E. coli* imobilisasi pada membran gentian berongga berguna untuk tujuan perembesan protein rekombinan dengan kadar sel hidup yang tinggi.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xiii
	LIST OF FIGURES	xvi
	LIST OF SYMBOLS	xxxii
	LIST OF ABBREVIATIONS	xxxiii
	LIST OF APPENDICES	xxxv
1	INTRODUCTION	1
	1.1 Introduction	1
	1.2 Objectives of the study	4
	1.3 Scopes of the study	4
	1.4 Problem statement and novelties of the study	5
2	LITERATURE REVIEW	6
	2.1 Cyclodextrin glucanotransferase (CGTase)	6

2.2	Recombinant protein production	9
2.3	System for producing recombinant protein	10
2.3.1	<i>Escherichia coli</i> (<i>E. coli</i>)	10
2.3.2	Yeasts	11
2.3.3	Filamentous fungi	12
2.3.4	Mammalian cells	12
2.4	<i>Escherichia coli</i> as host	13
2.4.1	Cytoplasmic expression	16
2.4.2	Periplasmic secretion	17
2.4.3	Excretion (extracellular secretion)	18
2.5	Problem encountered in excretion of recombinant protein expression	20
2.6	Cell immobilization	22
2.7	Techniques of cell immobilization	24
2.7.1	Immobilization on solid support surfaces	24
2.7.2	Entrapment within porous matrix	26
2.7.3	Cell flocculation (aggregation)	28
2.7.4	Mechanical containment behind a barrier	31
	2.7.4.1 Cell immobilization using membrane	31
	2.7.4.2 Cell immobilization using microcapsules	35
2.8	Factors affecting the cell immobilization system	37
2.8.1	Techniques of cell immobilization	38
2.8.2	Culture conditions	39
	2.8.2.1 Medium	39
	2.8.2.2 Inducer concentration	40

	2.8.2.3 Temperature	40
	2.8.2.4 Time	41
	2.8.2.5 Agitation rate	42
	2.8.2.6 pH	43
3	MATERIALS AND METHODS	44
3.1	Strategies for improvement of CGTase excretion with high cell viability of immobilized <i>Escherichia coli</i>	44
3.2	<i>Escherichia coli</i> strains and plasmid selection	46
3.3	Chemicals and reagents	47
3.4	Preparation of bacterial glycerol stock	47
3.5	Preparation of competent cell	47
3.6	Plasmid DNA isolation (Ehrt and Schnappinger, 2003)	48
3.7	Transformation into <i>E. coli</i> cell	49
3.8	Cell immobilization	49
3.9	Expression study on CGTase excretion and cell viability of immobilized cell	50
3.9.1	Screening of the cultural conditions using one factor at one time method (OFAT)	50
3.9.1.1	Polymer membrane selection	50
3.9.1.2	Expression medium selection	51
3.9.1.3	Contact time for cell immobilization	51
3.9.1.4	Inducer concentration	52
3.9.1.5	Post induction temperature	52
3.9.1.6	Post induction time	52

3.9.1.7	Agitation rate	53
3.9.1.8	pH	53
3.9.2	Experimental design on CGTase excretion and cell viability of immobilized cell	53
3.9.2.1	Screening of the cultural conditions by using full factorial design (FFD)	54
3.9.2.2	Optimization of the cultural conditions by using response surface methodology (RSM)	56
3.9.3	Formulation of expression medium (effect of tryptone)	58
3.9.4	Effect of cross linkers and reusability of immobilized cells	59
3.9.5	Membrane bioreactor	60
3.10	Preparation of enzymes	65
3.11	Analytical methods	65
3.11.1	CGTase activity	65
3.11.2	Xylanase activity	65
3.11.3	β -Mannosidase activity	66
3.11.4	β -Galactosidase activity (cell lysis)	66
3.11.5	Plasmid stability	67
3.11.6	Cell concentration	67
3.11.7	Sodium dodecyl sulfate polyacrylamide gel electrophoresis	68
3.11.8	Field emission scanning electron microscope (FESEM)	69
3.11.9	Nutrients content	70
3.11.10	Cell concentration	70

4	RESULT AND DISCUSSION	71
4.1	Screening of culture conditions on CGTase excretion and cell viability of immobilized cell using one factor at one time method (OFAT)	71
4.1.1	Effect of types of polymer membrane on CGTase excretion and cell viability of immobilized cell	72
4.1.2	Effect of expression medium on CGTase excretion and cell viability of immobilized cell	78
4.1.3	Effect of contact time for cell immobilization	84
4.1.4	Effect of IPTG concentration	90
4.1.5	Effect of post induction temperature	95
4.1.6	Effect of post induction time	100
4.1.7	Effect of agitation rate	109
4.1.8	Effect of pH	117
4.2	Screening of culture conditions affecting on CGTase excretion and cell viability of immobilized cell using design of experiment (full factorial design)	122
4.3	Optimization of culture conditions on CGTase excretion and cell viability of immobilized cell using response surface methodology (central composite design)	128
4.4	Medium formulation for CGTase excretion and cell viability of immobilized cell	141
4.4.1	Kinetics evaluation of immobilized and free cell fermentation	154
4.5	Reusability of immobilized cell	167

4.6	High-level excretion of CGTase in membrane bioreactor	179
4.6.1	Kinetics evaluation of immobilized and free cell in membrane bioreactor	187
4.6.2	Reusability of immobilized cell in membrane bioreactor	194
5	CONCLUSION AND RECOMMENDATIONS	198
5.1	Conclusion	198
5.2	Recommendations	199
5.2.1	Excretion of different recombinant protein using cell immobilization system	199
5.2.2	Effect of inoculum size	200
5.2.3	Addition of permeabilizer	200
5.2.4	Cultivation mode of membrane bioreactor	201
5.2.5	Effect of dissolved oxygen in the membrane bioreactor	201
	REFERENCES	202
	Appendices A-C	226-243

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	Applications of CD within different industries	8
2.2	Relative merits of different compartments for gene expression in <i>E. coli</i>	15
2.3	The support material for cell immobilization through surface attachment	26
2.4	Entrapment of cell within porous matrix	28
2.5	The flocculating agents for cell immobilization	30
2.6	Cell immobilization using a membrane as a barrier	32
2.7	Cell immobilization in microcapsules	36
2.8	Factors affecting the cell immobilization system	37
3.1	Host strains and plasmid	46
3.2	Thickness and porosity of hollow fiber membrane for each polymer	50
3.3	Independent variables and the levels of the screening design	54
3.4	Experimental design of 2^5 full factorial design	55
3.5	Actual and coded values of the design variables for the optimization process	56
3.6	Experimental design of the central composite design	57
4.1	Comparison of CGTase expression from various microorganisms	93
4.2	Nutrient content in the expression medium	101
4.3	The excretion of xylanase and mannosidase in recombinant	

	<i>E. coli</i> immobilized on the hollow fiber membrane	107
4.4	Comparison of immobilization matrices and free cells reported in the literature showing productivity for product formation	108
4.5	Diffusion of molecules in immobilization matrix	115
4.6	Experimental design and results of full factorial design	122
4.7	Regression analysis of the full factorial design for CGTase activity	125
4.8	Regression analysis of the full factorial design for β -galactosidase activity	126
4.9	Statistical analysis for CGTase activity and β -galactosidase activity	127
4.10	Experimental design and results of the central composite design	129
4.11	ANOVA for response surface quadratic model for CGTase excretion	130
4.12	ANOVA for response surface quadratic model for β -galactosidase activity	131
4.13	Summary of the optimized cultural conditions for the CGTase activity and β -galactosidase activity	140
4.14	The CGTase excretion, cell viability, plasmid stability and cell concentration of immobilized and free cell in medium containing 20 g/l and 5 g/l of tryptone	154
4.15	Growth kinetics of immobilized and free cell using medium containing 20 g/l and 5 g/l of tryptone	160
4.16	Kinetics parameter values of product formation for immobilized and free cell	165
4.17	Cumulative CGTase activity and β -galactosidase activity of immobilized cells using different cross-linkers during seven cycles	173
4.18	Comparison of average productivity obtained from reusability of immobilized cell reported in the literature	178

4.19	CGTase excretion, cell lysis and plasmid stability of immobilized and free cell in shake flask and bioreactor after 24 h of post induction time	181
4.20	Comparison of CGTase activity using different types of reactor	184
4.21	Growth kinetics of the immobilized and free cell	190
4.22	Kinetics parameters values of product formation for immobilized and free cell	193
4.23	Cumulative CGTase activity and β -galactosidase activity of immobilized and free cells in the membrane bioreactor during seven cycles	195
4.24	Comparison of average productivity obtained from reusability of immobilized cell using different types of reactor	197

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Cyclodextrins formation under catalytic reaction of CGTase	7
2.2	Chemical structure of α -, β - and γ -cyclodextrin (CD), consisting of 6, 7 or 8 glucose units, respectively	7
2.3	Schematic diagram of membrane cell composition in <i>E. coli</i> . LPS, lipopolysaccharide; PL, phospholipids; IM, inner membrane; OM, outer membrane	14
2.4	Cleavage of the signal peptide by signal peptidase. A, D, F, Y, E and G is one of the translocation machinery found in <i>E. coli</i> . Once the preprotein is translocated across the membrane, the signal peptide is cleaved off by the signal peptidase	18
2.5	Basic methods of cell immobilization	23
2.6	Hollow fiber membrane (A); Cross section area of hollow fiber membrane (B)	33
3.1	Research design for the improvement of CGTase excretion and cell viability of immobilized cell	45
3.2	Experimental apparatus for CGTase excretion by the immobilized <i>E. coli</i> . T, P and pH represent the temperature, pressure and pH sensor, respectively	62
3.3	Dimension of membrane bioreactor	63

3.4	Schematic diagram of membrane bioreactor	64
4.1	Effect of polymer membranes on cell concentration of immobilized and free cell. The expression medium used was LB. The expression conditions for immobilized and free cell system were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The PVDF+ NMP polymer showed the highest cell concentration compared to other polymers	73
4.2	Effect of the polymer membranes on enzyme activity of immobilized and free cell. The expression medium used was LB. The expression conditions for immobilized and free cell system were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The PVDF+NMP polymer showed the highest CGTase excretion while the PES+PEG polymer showed the lowest cell lysis	74
4.3	Porosity of hollow fiber membrane for each polymer	76
4.4	Effect of the polymer membranes on plasmid stability of immobilized and free cell. The expression medium used was LB. The expression conditions for immobilized and free cell system were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The PEI+NMP polymer showed the highest plasmid stability	77
4.5	Effect of expression medium on enzyme activity of immobilized (on PVDF polymer) and free cell. The expression conditions for immobilized and free cell system were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The SOB was used as expression medium for free cell. The 2xYT medium showed the highest CGTase excretion and cell lysis	79

- 4.6 Effect of expression medium on cell concentration of immobilized (on PVDF polymer) and free cell. The expression conditions for immobilized and free cell system were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The SOB was used as expression medium for free cell. The 2xYT medium showed the highest cell concentration compared to other mediums 80
- 4.7 SDS-PAGE analysis of CGTase excretion of immobilized cell using different types of expression medium at 24 h of post induction time. 1: Marker. 2: LB. 3: SOB. 4: TB. 5: 2xYT. 6: M9. The arrow shows the CGTase with a molecular weight of approx. 75 kDa. The expression conditions of immobilized cell were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate and pH 7. The sample for SDS-PAGE analysis was collected after 24 h of post induction time 82
- 4.8 Effect of expression medium on plasmid stability of immobilized (on PVDF polymer) and free cell. The expression conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The SOB was used as expression medium for free cell. The M9 medium showed the highest plasmid stability 84
- 4.9 Effect of contact time on cell immobilization process. The polymer membrane and medium used were PVDF and SOB, respectively. The constant value for each variable was 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate and pH 7. The cell concentration of immobilized cell was measured at every time interval (6, 12, 18, 24 and 30 h). The highest cells that immobilized on the hollow fiber membrane were detected at 30 h 85

- 4.10 Effect of contact time for cell immobilization on enzyme activity. The polymer membrane and medium used were PVDF and SOB, respectively. The expression conditions of immobilized cell were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The 18 h of contact time showed the highest CGTase activity with low occurrence of cell lysis 87
- 4.11 Effect of contact time for cell immobilization on plasmid stability. The polymer membrane and medium used were PVDF and SOB, respectively. The expression conditions of immobilized cell were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The 18 h of contact time showed the highest plasmid stability 88
- 4.12 Growth profile of recombinant *E. coli* cultivated in LB medium. The contact time of 18 h and 30 h were categorized as stationary and death phase, respectively 89
- 4.13 Effect of inducer concentration on enzyme activity of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for the immobilized and free cells system were as follows: 30 °C, 200 rpm, 24 h of post induction time and pH 7. The IPTG with concentration of 0.01 mM was used in the free cell system. The highest excretion of CGTase with low occurrence of cell lysis was detected at 0.01 mM of IPTG 91
- 4.14 Effect of inducer concentration on plasmid stability of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cells were as follows: 30 °C, 200 rpm, 24 h of post induction time

- and pH 7. The IPTG with concentration of 0.01 mM was used in the free cell system. The high plasmid stability of immobilized cell was obtained when using low concentration of IPTG (0.005 and 0.01 mM) 92
- 4.15 Effect of inducer concentration on cell concentration of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as follows: 30 °C, 200 rpm, 24 h of post induction time and pH 7. The IPTG with concentration of 0.01 mM was used in the free cell system 94
- 4.16 Effect of post induction temperature on cell concentration of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions of immobilized and free cells were as follows: 0.01 mM IPTG, 200 rpm, 24 h of post induction time and pH 7. The post induction temperature for free cell was 30°C. The cell concentration of immobilized cell was increased as the post induction temperature increased 96
- 4.17 Effect of post induction temperature on enzyme activity of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 200 rpm, 24 h of post induction time and pH 7. The post induction temperature for free cell was 30°C. The temperature of 30°C showed the highest CGTase activity with low occurrence of cell lysis 98
- 4.18 Effect of post induction temperature on plasmid of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression

- conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 200 rpm, 24 h of post induction time and pH 7. The post induction temperature for free cell was 30°C. The plasmid stability was lower at higher temperature 99
- 4.19 Effect of post induction time on enzyme activity of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate and pH 7. The post induction time for free cell was 24 h. The high CGTase excretion with low occurrence of cell lysis was detected at 24 h of post induction time 101
- 4.20 Effect of post induction time on cell concentration of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 30 °C, 200 rpm of agitation rate, 24 h of post induction time and pH 7. The post induction time for free cell was 24 h 105
- 4.21 (A) SDS-PAGE analysis of recombinant CGTase excretion at different post induction time. (B) Western blot analysis of recombinant CGTase excretion at different post induction time corresponding to the SDS-PAGE analysis. Lane M, protein size marker; lane 1, recombinant CGTase at 0 h post induction time (control); lane 2, recombinant CGTase at 8 h post induction time; lane 3, recombinant CGTase at 24 h post induction time; lane 4, recombinant CGTase at 48 h post induction time; lane 5, recombinant CGTase at 72 h post induction time 103

- 4.22 Field emission scanning electronic microscopy (FESEM) showing (A) cross-section structure of hollow fiber membrane(uninoculated); (B) recombinant *E. coli* immobilized on hollow fiber membrane; and (C) ruptured and lysed recombinant *E. coli* cells after 24 h of post induction time (highlighted in circle) 104
- 4.23 Effect of post induction time on plasmid stability of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cells were as follows: 0.01 m M IPTG, 30 °C, 200 rpm of agitation rate and pH 7. The post induction time for free cell was 24 h. The plasmid stability of immobilized cell decreased at longer post induction time (48 and 72 h) 105
- 4.24 Effect of agitation rate on enzyme activity of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 30 °C, 24 h of post induction time and pH 7. The agitation rate used for free cell was 200 rpm. The highest CGTase excretion with low cell lysis was observed at 200 rpm of agitation rate 110
- 4.25 Effect of agitation rate on cell concentration of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 30 °C, 24 h of post induction time and pH 7. The agitation rate used for free cell was 200 rpm. The highest cell concentration of immobilized cell was obtained at 200 rpm of agitation rate 111
- 4.26 Effect of agitation rate on plasmid stability of immobilized (on PVDF polymer) and free cell. The

- expression medium used was SOB. The expression conditions of immobilized and free cell were as follows: 0.01 mM IPTG, 30 °C, 24 h of post induction time and pH 7. The agitation rate used for free cell was 200 rpm. The high plasmid stability of immobilized cell was observed at low agitation rate 114
- 4.27 Effect of medium pH on enzyme activity of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 30 °C, 24 h of post induction time and 200 rpm of agitation rate. The medium pH used for free cell was pH 9. The buffer solutions of different pHs were phosphate-citrate buffer (pH 5 and pH 6), Sorensen's phosphate buffer (pH 7 and pH 8) and glycine-NaOH buffer (pH 9 and pH 10). The highest CGTase activity with low cell lysis was detected at pH 9 118
- 4.28 Effect of medium pH on plasmid stability of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as follows: 0.01 mM IPTG, 30 °C, 24 h of post induction time and 200 rpm of agitation rate. The medium pH used for free cell was pH 9. The buffer solutions of different pHs were phosphate-citrate buffer (pH 5 and pH 6), Sorensen's phosphate buffer (pH 7 and pH 8) and glycine-NaOH buffer (pH 9 and pH 10). The high plasmid stability was obtained at alkaline condition 119
- 4.29 Effect of medium pH on cell concentration of immobilized (on PVDF polymer) and free cell. The expression medium used was SOB. The expression conditions for immobilized and free cell were as

- follows: 0.01 mM IPTG, 30 °C, 24 h of post induction time and 200 rpm of agitation rate. The medium pH used for free cell was pH 9. The buffer solutions of different pHs were phosphate-citrate buffer (pH 5 and pH 6), Sorensen's phosphate buffer (pH 7 and pH 8) and glycine-NaOH buffer (pH 9 and pH 10) 120
- 4.30 Response surface plot of CGTase excretion: concentration of IPTG vs. pH with constant level of post induction temperature (25°C). The CGTase activity of immobilized cell was measured after 24 h of post induction time. The optimum pH and IPTG for the highest CGTase excretion of immobilized cell were pH 9 and 0.01 mM, respectively 134
- 4.31 Response surface plot of CGTase excretion: post induction temperature vs. pH with constant level of IPTG concentration (0.01 mM). The CGTase activity of immobilized cell was measured after 24 h of post induction time. The optimum post induction temperature and pH for the highest CGTase excretion of immobilized cell were 25 °C and pH 9, respectively 135
- 4.32 Response surface plot of β -galactosidase activity: IPTG vs. post induction temperature with the constant level of pH (pH9). The β -galactosidase activity of immobilized cell was measured after 24 h of post induction time. The optimum IPTG and post induction temperature for the lowest occurrence of cell lysis were 0.005 mM and 22°C, respectively 137
- 4.33 Response surface plot of β -galactosidase activity: post induction temperature vs. pH with constant level of IPTG (0.01 mM). The β -galactosidase activity of immobilized cell was measured after 24 h of post induction time. The optimum post

- induction temperature and pH for the lowest occurrence of cell lysis were 22°C and pH 9, respectively 138
- 4.34 Effect of tryptone concentration on enzyme activity of immobilized cell. The polymer membrane used was PVDF. The expression was conducted using the optimized cultural conditions (25 °C, 0.011 mM IPTG and pH 8.8). The enzyme activity was measured after 24 h of post induction time. The tryptone concentration of 20 g/l was used as a control. The high CGTase excretion with low occurrence of cell lysis was observed at 5 g/l of tryptone 143
- 4.35 Effect of tryptone concentration on the plasmid stability of immobilized cell. The polymer membrane used was PVDF. The expression was conducted using optimized cultural conditions (25 °C, 0.011 mM IPTG and pH 8.8). The plasmid stability was measured after 24 h of post induction time. The tryptone concentration of 20 g/l was used as a control. High plasmid stability was obtained at low tryptone concentration 144
- 4.36 Effect of tryptone concentration on cell concentration of the immobilized cell. The polymer membrane used was PVDF. The expression was conducted using optimized cultural conditions (25 °C, 0.011 mM IPTG and pH 8.8). The cell concentration was measured after 24 h of post induction time. The tryptone concentration of 20 g/l was used as a control. The cell concentration of the immobilized cell increased as the tryptone concentration increased 145
- 4.37 The effect of tryptone concentration on enzyme activity of immobilized (on PVDF polymer) and free cell using different concentrations of tryptone, 20 g/l

- and 5 g/l. Tryptone concentration of 20 g/l was used as a control. The expression was conducted using the optimized cultural conditions (25 °C, 0.011 mM IPTG and pH 8.8). The enzyme activity was measured at selected time interval 147
- 4.38 The effect of tryptone concentration on plasmid stability of immobilized (on PVDF polymer) and free cell using different concentrations of tryptone, 20 g/l and 5 g/l. Tryptone concentration of 20 g/l was used as a control. The expression was conducted using the optimized cultural conditions (25 °C, 0.011 mM IPTG and pH 8.8). The plasmid stability was measured at selected time interval 148
- 4.39 The effect of tryptone concentration on β -galactosidase activity (cell lysis indicator) of immobilized (on PVDF polymer) and free cell using different concentrations of tryptone, 20 g/l and 5 g/l. The concentration of 20 g/l was used as a control. The expression was conducted using the optimized cultural conditions (25 °C, 0.011 mM IPTG and pH 8.8). The β -galactosidase activity was measured at selected time interval 149
- 4.40 The structure of the immobilized cell in medium containing 20 g/l of tryptone. The polymer membrane used was PVDF. The expression conditions were 25 °C of post induction temperature, 0.011 mM IPTG and pH 8.8. The picture shows more cells were ruptured and lysed 150
- 4.41 The structure of the immobilized cell in medium containing 5 g/l of tryptone. The polymer membrane used was PVDF. The expression conditions were 25 °C of post induction temperature, 0.011 mM IPTG and pH 8.8. The picture shows the cells were not ruptured 151

- 4.42 (A) The effect of tryptone concentration on cell concentration of immobilized (on PVDF polymer) and free cell using different concentration of tryptone, 20 g/l and 5 g/l; (B) The enlarged scale for cell concentration of the free cell from (A). The concentration of 20 g/l tryptone was used as a control. The expression was conducted using the optimized cultural conditions (25 °C, 0.011 mM IPTG and pH 8.8). The cell concentration of immobilized and free cell was measured at selected time interval 152
- 4.43 Growth kinetics of the immobilized cell using medium containing 20 g/l of tryptone. The cell was immobilized on PVDF polymer. The growth conditions for the immobilized cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (0.019) represents the value of μ_{\max} for the immobilized cell 157
- 4.44 Growth kinetics of the immobilized cell using medium containing 5 g/l of tryptone. The cell was immobilized on PVDF polymer. The growth conditions for the immobilized cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (0.012) represents the value of μ_{\max} for the immobilized cell 158
- 4.45 Growth kinetics of the free cell using medium containing 20 g/l of tryptone. The growth conditions for the free cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (0.168) represents the value of μ_{\max} for the free cell 159
- 4.46 Growth kinetics of the free cell using medium containing 5 g/l of tryptone. The growth conditions for the free cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (0.132) represents the value of μ_{\max} for the free cell 160

- 4.47 Kinetics for product formation (CGTase excretion) of immobilized cell using medium containing 20 g/l of tryptone. The polymer membrane used was PVDF. The growth conditions for the immobilized cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (29.02) represents the value of α for the immobilized cell 162
- 4.48 Kinetics for product formation (CGTase excretion) of immobilized cell using medium containing 5 g/l of tryptone. The polymer membrane used was PVDF. The growth conditions for the immobilized cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (33.15) represents the value of α for the immobilized cell 163
- 4.49 Kinetics for product formation of free cell using medium containing 20 g/l of tryptone. The growth conditions for the free cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graphs (2.974) represents the value of α for the free cell 164
- 4.50 Kinetics for product formation of free cell using medium containing 5 g/l of tryptone. The growth conditions for the free cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (4.178) represents the value of α for the free cell 165
- 4.51 Effect of cross-linkers (GA, PLL and PEI) concentration on CGTase excretion of the immobilized cell. The polymer membrane used was PVDF. The expression of immobilized cell using treated membrane was conducted at 25 °C, 0.011 mM IPTG and pH 8.8. The enzyme activity was measured after 24 h of post induction time 168

- 4.52 Effect of cross-linkers (GA, PLL and PEI) concentration on β -galactosidase activity (indicator of cell lysis) of the immobilized cell. The polymer membrane used was PVDF. The expression of immobilized cell using treated membrane was conducted at 25 °C, 0.011 mM IPTG and pH 8.8. The enzyme activity was measured after 24 h of post induction time 169
- 4.53 Comparison of CGTase excretion between immobilized cell (on non-treated membrane and treated membrane) and free cell by repeated fermentation in medium containing 20 g/l of tryptone. The hollow fiber membrane was treated with 0.3% of glutaraldehyde, poly-L-lysine and polyethylenimine. The non-treated hollow fiber membrane was used as a control. The conditions for expression were 0.011 mM IPTG, 25 °C and pH 8.8. Enzyme activity was measured for every cycle (24 h of post induction time) until 7 cycles 172
- 4.54 Comparison of CGTase excretion between immobilized cell (on non-treated membrane and treated membrane) and free cell by repeated fermentation in medium containing 5 g/l of tryptone. The hollow fiber membrane was treated with 0.3% of glutaraldehyde, poly-L-lysine and polyethylenimine. The non-treated hollow fiber membrane was used as control. The conditions for growth and expression were 0.011 mM IPTG, 25 °C and pH 8.8. Enzyme activity was measured for every cycle (24 h of post induction time) until 7 cycles 175
- 4.55 Comparison of cell lysis between immobilized cell (on non-treated membrane and treated membrane) and free cell by repeated fermentation in medium containing 20 g/l of tryptone. The hollow fiber membrane was treated with 0.3% of glutaraldehyde,

- poly-L-lysine and polyethylenimine. The non-treated hollow fiber membrane was used as a control. The conditions for expression were 0.011 mM IPTG, 25 °C and pH 8.8. Enzyme activity was measured for every cycle (24 h of post induction time) until 7 cycles 176
- 4.56 Comparison of cell lysis between immobilized cell (on non-treated membrane and treated membrane) and free cell by repeated fermentation in medium containing 5 g/l of tryptone. The hollow fiber membrane was treated with 0.3% of glutaraldehyde, poly-L-lysine and polyethylenimine. The non-treated hollow fiber membrane was used as a control. The conditions for expression were 0.011 mM IPTG, 25 °C and pH 8.8. Enzyme activity was measured for every cycle (24 h of post induction time) until 7 cycles 177
- 4.57 Profiles of the immobilized and free cell on CGTase excretion in the membrane bioreactor. The expression was conducted using the optimized conditions (0.011 mM IPTG, 25 °C and pH 8.8). The CGTase activity was measured at selected time interval 185
- 4.58 Profiles of the immobilized and free cell on cell lysis in the membrane bioreactor. The expression was conducted using the optimized conditions (0.011 mM IPTG, 25 °C and pH 8.8). The β -galactosidase activity was measured at selected time interval 186
- 4.59 Growth kinetics of the immobilized cell in the membrane bioreactor. The cell was immobilized on PVDF polymer. The growth conditions of immobilized cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (0.012) represents the value of μ_{\max} for the immobilized cell 189

- 4.60 Growth kinetics of the free cell in the membrane bioreactor. The growth conditions of free cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (0.149) represents the value of μ_{\max} for the free cell 190
- 4.61 Kinetics for product formation (CGTase excretion) of immobilized cell in the membrane bioreactor. The polymer membrane used was PVDF. The growth conditions of immobilized cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (1.317) represents the value of α for the immobilized cell 192
- 4.62 Kinetics for product formation (CGTase excretion) of free cell in the membrane bioreactor. The growth conditions of free cell were 25 °C, 0.011 mM IPTG and pH 8.8. The slope of the graph (0.004) represents the value of α for the free cell 193
- 4.63 Comparison of CGTase excretion between immobilized and free cell by repeated fermentation in the membrane bioreactor. The conditions for expression were 0.011 mM IPTG, 25°C and pH 8.8. Enzyme activity was measured for every cycle (24 h of post induction time) until 7 cycles 195
- 4.64 Comparison of cell lysis between immobilized and free cell by repeated fermentation in the membrane bioreactor. The conditions for expression were 0.011 mM IPTG, 25°C and pH 8.8. Enzyme activity was measured for every cycle (24 h of post induction time) until 7 cycles 196

LIST OF SYMBOLS

g	-	gram
h	-	hour
l	-	liter
min	-	minute
ml	-	milliliter
sec	-	second
v/v	-	volume solute per volume solution
w/v	-	weight solute per volume solution

LIST OF ABBREVIATIONS

A_{600}	-	absorbance at wavelength 600 nm
ANOVA	-	analysis of variance
BRP	-	bacteriocin release protein
CCD	-	central composite design
CD	-	cyclodextrin
CGTase	-	cyclodextrin glucanotransferase
Da, kDa	-	dalton, kilodalton
DNA	-	deoxyribonucleic acid
<i>E. coli</i>	-	<i>Escherichia coli</i>
EDTA	-	ethylenediaminetetra-acetate
FESEM	-	field emission scanning electron microscopy
FFD	-	full factorial design
HCl	-	hydrochloric acid
IM	-	inner membrane
IPTG	-	isopropyl β -D-1-thiogalactopyranoside
<i>lac</i>	-	lactose
LB	-	luria bertani
LPS	-	lipopolysaccharide
MgCl	-	magnesium chloride
MW	-	molecular weight
NMP	-	1-methyl-2-pyrrolidon
OD	-	optical density
OFAT	-	one factor at one time
OM	-	outer membrane

ONPG	-	<i>ortho</i> -nitrophenyl- β -galactoside
PEG	-	polyethylene glycol
PES	-	polyethersulfone
rpm	-	revolution per minutes
RSM	-	response surface methodology
SDS-PAGE	-	sodium dodecyl sulfate polyacrylamide gel electrophoresis
Sp.	-	species

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A1	Medium and buffers preparation	228
A2	Antibiotic and inducer	232
A3	Transformation and storage solution (TSS) reagent	232
A4	SDS-PAGE buffer and solutions	233
A5	Calculation of CGTase activity	235
A6	Calculation of β -galactosidase activity	237
A7	Calculation for the volume of free cell in shake flask	238
B1	Results for screening process of cultural conditions using FFD	239
B2	Results for optimization process of cultural conditions using RSM	242
C	Publications and awards	244

CHAPTER 1

INTRODUCTION

1.1 Introduction

Protein excretion is important, as it enables active and stable enzyme production, which is essential for successful biocatalysis. In recombinant protein expression, excreted proteins are preferable to intracellular proteins production. Excretory production of recombinant proteins provides several advantages over intracellular protein production, including increased protein stability and solubility, correct formation of disulfide bonds and facilitation of downstream processing (Mergulhao *et al.*, 2005). However, studies have shown that excretion of enzymes in *Escherichia coli* (*E. coli*) may cause cell lysis because of pressure build-up through overproduction of the expressed recombinant protein (Fu *et al.*, 2005). Thus, the production of recombinant proteins by excretion without (or with less) cell lysis caused by overexpression of the recombinant protein in the host cell is difficult. Various attempts have been made to reduce cell lysis resulting from the excretion of proteins, such addition of glycine or Triton-X to the medium (Jang *et al.*, 1999), using bacteriocin release protein (BRP) (Fu *et al.*, 2003) and cultivating cells using favorable conditions (Shokri *et al.*, 2003).

Cell immobilization for the excretion of enzymes has various advantages over the conventional process using a free cell system. These advantages include the repeated and expanded use of cells, preservation of plasmid-bearing cells, simplicity of performing a continuous process for extended periods of time, decreased risk of contamination, protection of the cells from environmental stresses and easier downstream processing (Jamuna and Ramakrishna, 1992, Mamo and Gessesse, 1997, Moriwaki *et al.*, 2007). Using cell immobilization techniques, increased protein excretion and reduction in cell lysis can be achieved. Various techniques for cell immobilization, such as adsorption on surfaces (Mazzer *et al.*, 2008), covalent bonding to carriers (Atanasova *et al.*, 2009), entrapment in a polymer gel (Kunamneni *et al.*, 2007) and self-aggregation (Yang *et al.*, 2006), have been used to obtain cells with high levels of enzyme production and stability (Chen *et al.*, 2004). Therefore, the choice of immobilization technique and the mechanical properties of the matrix are significant factors affecting the enzyme activity and long-term stability of biocatalysts obtained from immobilized cells.

Calcium alginate, glass beads, polyacrylamide gel, alginate, silanized magnetite, agarose, polyurethane foam and carrageenan are the most commonly used matrices for the immobilization of cells. However, the use of alginate for gel entrapment is insufficient because the solidification of the gel requires the use of calcium chloride, which reduces the pH of the mixture and brings a negative effect on cell growth (Atanasova *et al.*, 2009). Furthermore, covalent binding to a matrix has the major disadvantages of high cost and low yield because of exposure of the cells to toxic reagents and rigorous reaction conditions. The development of cell immobilization techniques using hollow fiber membranes has attracted considerable interest because of the high surface-to-volume ratio that can be obtained with this type of membrane (Wang *et al.*, 1991). Several studies have been conducted using hollow fiber membranes as a matrix for the production of β -lactamase (Inloes *et al.*, 1983a), ammonia (Wang *et al.*, 1991), ethanol (Mehaia and Cheryan, 1984, Shukla *et al.*, 1989) and ethylene (Lloyd and Bunch, 1996).

In the present study, a hollow fiber membrane was employed as a matrix for cell immobilization to increase the excretion of cyclodextrin glucanotransferase (CGTase) with a low occurrence of cell lysis. *E. coli* were immobilized via adsorption and entrapment within the polymeric matrix based on the high porosity of the membrane. The major benefit of immobilization through adsorption is direct contact between nutrients and the matrix. Moreover, the limitation of diffusion imposed by cell immobilization through entrapment in a hollow fiber membrane is less severe because of the presence of liquid medium in the intra-capillary space of the membrane. In addition to being readily available and inexpensive, hollow fibers were selected as the matrix because of several other desirable properties, such as high mechanical strength, operational durability and lack of toxicity. Another advantage of hollow fiber membranes is the free exchange of nutrients and metabolic products, given the short diffusion distance between cells and the liquid medium (Bunch, 1988).

Most studies for CGTase production thus far have focused on attachment to cotton (Kriaa *et al.*, 2012); entrapment in different gels, such as polyacrylamide, agar, alginate and polyvinyl alcohol-cryogel gels (Abdel-Naby *et al.*, 2000, Martins *et al.*, 2003, Kunamneni *et al.*, 2007, Mazzer *et al.*, 2008); or covalent linking to a flat sheet membrane (Atanasova *et al.*, 2009). To the best of our knowledge, there have been no studies on the excretion of CGTase by *E. coli* that have been immobilized through a combination of adsorption and entrapment of the cells using a hollow fiber membrane. The results presented here suggest that immobilized cell is a promising method for CGTase excretion with less occurrences of cell lysis.

1.2 Objectives of the study

1. To improve CGTase excretion of immobilized recombinant *E. coli* on the hollow fiber membrane in comparison with free cell.
2. To increase cell viability with high plasmid stability of immobilized recombinant *E. coli* on the hollow fiber membrane.

1.3 Scopes of the study

Scopes of this study are:

- a) Screening the effect of the culture conditions (polymer membrane, expression medium, time for cell immobilization, IPTG, post induction temperature, post induction time, agitation rate and pH) on enhancement of CGTase excretion and cell viability of the immobilized cell using one factor at one time method (OFAT).
- b) Determination of significant factors of culture conditions (IPTG, post induction temperature, post induction time, agitation rate and pH) on the immobilized cell for the increment of CGTase excretion and cell viability using full factorial design (FFD).
- c) Optimization of the culture conditions (IPTG, post induction temperature and pH) of the immobilized cell for the excretion of recombinant CGTase and reduction of cell lysis using response surface methodology (RSM).
- d) Determination of the effect of expression medium formulation and cross-linkers on the immobilized cell for the improvement of CGTase excretion with less occurrence of cell lysis.
- e) Production of high-level CGTase excretion of immobilized cell in the membrane bioreactor.

1.4 Problem statement and novelties of the study

In recent years, there has been great interest in the production of heterologous proteins by utilizing recombinant DNA technology. Some of these heterologous proteins have already been produced at the industrial level. However, the recombinant protein products accumulate in the microorganisms especially when using *Escherichia coli* as a host. Therefore, the excretion of these proteins to the external medium would be attractive with regard to further downstream processing and the stability of the protein product. However, the excretion of heterologous proteins remains problematic. In *E. coli*, excretion always correlates with non-specific leakage and cell lysis due to pressure build-up in the periplasmic space by high level production of the recombinant protein. Thus, it becomes a challenge when the recombinant protein needs to be excreted with low cell lysis. Using the cell immobilization method, increased protein excretion with reduction in cell lysis can be achieved.

There are three novelties of the study. The novelties of the study are as follows:

- a) The cell immobilization method improves the excretion of recombinant protein with increase in cell viability and plasmid stability as compared to the conventional fermentation (free cell).
- b) Cell immobilization on hollow fiber membrane involves adsorption technique with three interactions (two electrostatic interactions and one hydrophobic interaction).
- c) High CGTase excretion with low cell lysis of immobilized cell using low tryptone concentration and non-treated hollow fiber membrane in reusability process.

REFERENCES

- Abdel-Naby, M. A., Reyad, R. M. & Abdel-Fattah, A. F. (2000). Biosynthesis of cyclodextrin glucosyltransferase by immobilized *Bacillus amyloliquefaciens* in batch and continuous cultures. *Biochemical Engineering Journal*, 5, 1-9.
- Adinarayana, K., Jyothi, B. & Ellaiah, P. (2005). Production of alkaline protease with immobilized cells of *Bacillus subtilis* PE-11 in various matrices by entrapment technique. *AAPS PharmSciTech*, 6, E391-E397.
- Altshuler, G. L., Dziejulski, D. M., Soweck, J. A. & Belfort, G. (1986). Continuous hybridoma growth and monoclonal antibody production in hollow fiber reactors–separators. *Biotechnology and Bioengineering*, 28, 646-658.
- Anal, A. K. & Singh, H. (2007). Recent advances in microencapsulation of probiotics for industrial applications and targeted delivery. *Trends in Food Science & Technology*, 18, 240-251.
- Andersson, H. & von Heijne, G. (1991). A 30-residue-long" export initiation domain" adjacent to the signal sequence is critical for protein translocation across the inner membrane of *Escherichia coli*. *Proceedings of the National Academy of Sciences*, 88, 9751-9754.
- Andries, M., Van Beveren, P., Goffing, O., Rajotte, P. & Masschelein, C. (2000). Results on semi-industrial continuous top fermentation with the Meura-Delta immobilised yeast fermenter. *Brauwelt International*, 134-136.
- Angkawidjaja, C., Kuwahara, K., Omori, K., Koga, Y., Takano, K. & Kanaya, S. (2006). Extracellular secretion of *Escherichia coli* alkaline phosphatase with a C-terminal tag by type I secretion system: purification and biochemical characterization. *Protein Engineering Design and Selection*, 19, 337-343.

- Anisha, G. & Prema, P. (2008). Cell immobilization technique for the enhanced production of α -galactosidase by *Streptomyces griseolalbus*. *Bioresource Technology*, 99, 3325-3330.
- Atanasova, N., Kitayska, T., Yankov, D., Safarikova, M. & Tonkova, A. (2009). Cyclodextrin glucanotransferase production by cell biocatalysts of alkaliphilic bacilli. *Biochemical Engineering Journal*, 46, 278-285.
- Ates, S., Dingil, N., Bayraktar, E. & Mehmetoglu, U. (2002). Enhancement of citric acid production by immobilized and freely suspended *Aspergillus niger* using silicone oil. *Process Biochemistry*, 38, 433-436.
- Avci, A. (2012). Purification and characterization of a thermostable cyclodextrin glycosyltransferase from *Thermoanaerobacter* sp. P4. *African Journal of Biotechnology*, 11, 10407-10415.
- Barany, S. & Szepesszentgyörgyi, A. (2004). Flocculation of cellular suspensions by polyelectrolytes. *Advances in Colloid and Interface Science*, 111, 117-129.
- Bečka, S., Škrob, F., Plháčková, K., Kujan, P., Holler, P. & Kyslík, P. (2003). Cross-linked cell aggregates of *Trigonopsis variabilis*: D-amino acid oxidase catalyst for oxidation of cephalosporin C. *Biotechnology Letters*, 25, 227-233.
- Birch, R. M. & Walker, G. M. (2000). Influence of magnesium ions on heat shock and ethanol stress responses of *Saccharomyces cerevisiae*. *Enzyme and Microbial Technology*, 26, 678-687.
- Bisbee, C. (1993). Current perspectives on manufacturing and scaleup of biopharmaceuticals. *Genetic Engineering News*, 13, 8-10.
- Bisht, D., Yadav, S. K. & Darmwal, N. S. (2013). Optimization of immobilization conditions by conventional and statistical strategies for alkaline lipase production by *Pseudomonas aeruginosa* mutant cells: scale-up at bench-scale bioreactor level. *Turkish Journal of Biology*, 37, 392-404.
- Biwer, A., Antranikian, G. & Heinzle, E. (2002). Enzymatic production of cyclodextrins. *Applied Microbiology and Biotechnology*, 59, 609-617.
- Börner, R. A., Zaushitsyna, O., Berillo, D., Scaccia, N., Mattiasson, B. & Kirsebom, H. (2014). Immobilization of *Clostridium acetobutylicum* DSM 792 as macroporous aggregates through cryogelation for butanol production. *Process Biochemistry*, 49, 10-18.

- Bučko, M., Schenk Mayerová, A., Gemeiner, P., Vikartovská, A., Mihovilovič, M. D. & Lacík, I. (2011). Continuous testing system for Baeyer–Villiger biooxidation using recombinant *Escherichia coli* expressing cyclohexanone monooxygenase encapsulated in polyelectrolyte complex capsules. *Enzyme and Microbial Technology*, 49, 284-288.
- Bunch, A. W. (1988). The uses and future potential of microbial hollow-fibre bioreactors. *Journal of Microbiological Methods*, 8, 103-119.
- Busscher, H. J. & Weerkamp, A. H. (1987). Specific and non-specific interactions in bacterial adhesion to solid substrata. *FEMS Microbiology Letters*, 46, 165-173.
- Caunt, P., Impoolsup, A. & Greenfield, P. (1988). Stability of recombinant plasmids in yeast. *Journal of Biotechnology*, 8, 173-192.
- Chandel, A. K., Lakshmi Narasu, M., Chandrasekhar, G., Manikyam, A. & Venkateswar Rao, L. (2009). Use of *Saccharum spontaneum* (wild sugarcane) as biomaterial for cell immobilization and modulated ethanol production by thermotolerant *Saccharomyces cerevisiae* VS3. *Bioresource Technology*, 100, 2404-2410.
- Chen, J., Xu, Y., Xin, J., Li, S., Xia, C. & Cui, J. (2004). Efficient immobilization of whole cells of *Methylobionas* sp. strain GYJ3 by sol–gel entrapment. *Journal of Molecular Catalysis B-Enzymatic*, 30, 167-172.
- Chen, X. A., Xu, Z. N., Cen, P. L. & Wong, W. (2006). Enhanced plasmid stability and production of hEGF by immobilized recombinant *E. coli* JM101. *Biochemical Engineering Journal*, 28, 215-219.
- Chien, C. S., Lee, W. C. & Lin, T. J. (2001). Immobilization of *Aspergillus japonicus* by entrapping cells in gluten for production of fructooligosaccharides. *Enzyme and Microbial Technology*, 29, 252-257.
- Choi, J. & Lee, S. (2004). Secretory and extracellular production of recombinant proteins using *Escherichia coli*. *Applied Microbiology and Biotechnology*, 64, 625-635.
- Choi, J. H., Keum, K. C. & Lee, S. Y. (2006). Production of recombinant proteins by high cell density culture of *Escherichia coli*. *Chemical Engineering Science*, 61, 876-885.

- Chou, C. P. (2007). Engineering cell physiology to enhance recombinant protein production in *Escherichia coli*. *Applied Microbiology and Biotechnology*, 76, 521-532.
- Chu, Y. F., Hsu, C. H., Soma, P. K. & Lo, Y. M. (2009). Immobilization of bioluminescent *Escherichia coli* cells using natural and artificial fibers treated with polyethyleneimine. *Bioresource Technology*, 100, 3167-3174.
- Corona-González, R. I., Miramontes-Murillo, R., Arriola-Guevara, E., Guatemala-Morales, G., Toriz, G. & Pelayo-Ortiz, C. (2014). Immobilization of *Actinobacillus succinogenes* by adhesion or entrapment for the production of succinic acid. *Bioresource Technology*, 164, 113-118.
- Czaczyk, K. & Myszka, K. (2007). Biosynthesis of extracellular polymeric substances (EPS) and its role in microbial biofilm formation. *Polish Journal of Environmental Studies*, 16, 799.
- D'souza, S. & Kamath, N. (1988). Cloth bioreactor containing yeast cells immobilized on cotton cloth using polythylenimine. *Applied Microbiology and Biotechnology*, 29, 136-140.
- D'Souza, S. & Melo, J. (1991). A method for the preparation of coimmobilizates by adhesion using polyethyleneimine. *Enzyme and Microbiology Technology*, 13, 508-511.
- Danner, H., Madzingaidzo, L., Thomasser, C., Neureiter, M. & Braun, R. (2002). Thermophilic production of lactic acid using integrated membrane bioreactor systems coupled with monopolar electro dialysis. *Applied Microbiology and Biotechnology*, 59, 160-169.
- De León, A., Hernández, V., Galindo, E. & Ramí, O. T. (2003). Effects of dissolved oxygen tension on the production of recombinant penicillin acylase in *Escherichia coli*. *Enzyme and Microbial Technology*, 33, 689-697.
- de Souza, E. A., Rossi, D. M. & Ayub, M. A. Z. (2014). Bioconversion of residual glycerol from biodiesel synthesis into 1, 3-propanediol using immobilized cells of *Klebsiella pneumoniae* BLh-1. *Renewable Energy*, 72, 253-257.
- Derman, A. I., Prinz, W. A., Belin, D. & Beckwith, J. (1993). Mutations that allow disulfide bond formation in the cytoplasm of *Escherichia coli*. *Science*, 262, 1744-1747.

- Diaz Ricci, J. C. & Hernández, M. E. (2000). Plasmid effects on *Escherichia coli* metabolism. *Critical Reviews in Biotechnology*, 20, 79-108.
- Dincbas, V., Hortacsu, A. & Camurdan, A. (1993). Plasmid stability in immobilized mixed cultures of recombinant *Escherichia coli*. *Biotechnology Progress*, 9, 218-220.
- Ding, R., Li, Z., Chen, S., Wu, D., Wu, J. & Chen, J. (2010). Enhanced secretion of recombinant α -cyclodextrin glucosyltransferase from *E. coli* by medium additives. *Process Biochemistry*, 45, 880-886.
- Donlan, R. M. & Costerton, J. W. (2002). Biofilms: survival mechanisms of clinically relevant microorganisms. *Clinical Microbiology Reviews*, 15, 167-193.
- Doran, P. M. & Bailey, J. E. (1986). Effects of immobilization on growth, fermentation properties, and macromolecular composition of *Saccharomyces cerevisiae* attached to gelatin. *Biotechnology and Bioengineering*, 28, 73-87.
- Ehrt, S. & Schnappinger, D. (2003). Isolation of plasmids from *E. coli* by alkaline lysis. *E. coli Plasmid Vectors*. Springer.
- Einsfeldt, K., Júnior, J. B. S., Argondizzo, A. P. C., Medeiros, M. A., Alves, T. L. M., Almeida, R. V. & Larentis, A. L. (2011). Cloning and expression of protease ClpP from *Streptococcus pneumoniae* in *Escherichia coli*: Study of the influence of kanamycin and IPTG concentration on cell growth, recombinant protein production and plasmid stability. *Vaccine*, 29, 7136-7143.
- El-Mansi, E. M. T., Bryce, C. F., Demain, A. L. & Allman, A. R. *Fermentation Microbiology and Biotechnology*, CRC press. 2011
- Fahey, R. C., Hunt, J. S. & Windham, G. C. (1977). On the cysteine and cystine content of proteins. *Journal of Molecular Evolution*, 10, 155-160.
- Fang, N., Zhong, C.-Q., Liang, X., Tang, X.-F. & Tang, B. (2010). Improvement of extracellular production of a thermophilic subtilase expressed in *Escherichia coli* by random mutagenesis of its N-terminal propeptide. *Applied Microbiology and Biotechnology*, 85, 1473-1481.
- Fischer, D., Bieber, T., Li, Y., Elsässer, H.-P. & Kissel, T. (1999). A novel non-viral vector for DNA delivery based on low molecular weight, branched

- polyethylenimine: effect of molecular weight on transfection efficiency and cytotoxicity. *Pharmaceutical Research*, 16, 1273-1279.
- Fletcher, M. (1977). The effects of culture concentration and age, time and temperature on bacterial attachment to polystyrene. *Canadian Journal of Microbiology*, 23, 1-6.
- Fu, Z., Hamid, S. B. A., Razak, C. N. A., Basri, M., Salleh, A. B. & Rahman, R. N. Z. A. (2003). Secretory expression in *Escherichia coli* and single-step purification of a heat-stable alkaline protease. *Protein Expression and Purification*, 28, 63-68.
- Fu, Z., Ng, K., Lam, T. & Wong, W. (2005). Cell death caused by hyper-expression of a secretory exoglucanase in *Escherichia coli*. *Protein Expression and Purification*, 42, 67-77.
- Fuzi, S. F. Z. M., Razali, F., Jahim, J. M., Rahman, R. A. & Illias, R. M. (2014). Simplified feeding strategies for the fed-batch cultivation of *Kluyveromyces lactis* GG799 for enhanced recombinant xylanase production. *Bioprocess and Biosystems Engineering*, 37, 1887-1898.
- Gadgil, M., Kapur, V. & Hu, W. S. (2005). Transcriptional response of *Escherichia coli* to temperature shift. *Biotechnology Progress*, 21, 689-699.
- Ganatsios, V., Koutinas, A. A., Bekatorou, A., Kanellaki, M. & Nigam, P. (2014). Promotion of maltose fermentation at extremely low temperatures using a cryotolerant *Saccharomyces cerevisiae* strain immobilised on porous cellulosic material. *Enzyme and Microbial Technology*.
- Gawande, B. & Patkar, A. (1999). Application of factorial designs for optimization of cyclodextrin glycosyltransferase production from *Klebsiella pneumoniae pneumoniae* AS-22. *Biotechnology and Bioengineering*, 64, 168-173.
- Gawande, P. V. & Kamat, M. Y. (1998). Immobilization of *Aspergillus* sp. on nylon bolting cloth for production of xylanase. *Journal of Fermentation and Bioengineering*, 86, 243-246.
- Geesey, G. G., Wigglesworth-Cooksey, B. & Cooksey, K. (2000). Influence of calcium and other cations on surface adhesion of bacteria and diatoms: a review. *Biofouling*, 15, 195-205.

- Glick, B. R. (1995). Metabolic load and heterologous gene expression. *Biotechnology Advances*, 13, 247-261.
- Göksungur, Y. & Güvenç, U. (1997). Batch and continuous production of lactic acid from beet molasses by *Lactobacillus delbrueckii* IFO 3202. *Journal of Chemical Technology and Biotechnology*, 69, 399-404.
- Haaland, P. D. *Experimental design in biotechnology*, CRC press. 1989
- Hamdy, M., Kim, K. & Rudtke, C. (1990). Continuous ethanol production by yeast immobilized on to channeled alumina beads. *Biomass*, 21, 189-206.
- Hausner, M. & Wuertz, S. (1999). High rates of conjugation in bacterial biofilms as determined by quantitative in situ analysis. *Applied and Environmental Microbiology*, 65, 3710-3713.
- Helander, I. M., Alakomi, H.-L., Latva-Kala, K. & Koski, P. (1997). Polyethyleneimine is an effective permeabilizer of gram-negative bacteria. *Microbiology*, 143, 3193-3199.
- Hiemstra, H., Dijkhuizen, L. & Harder, W. (1983). Diffusion of oxygen in alginate gels related to the kinetics of methanol oxidation by immobilized *Hansenula polymorpha* cells. *European Journal of Applied Microbiology and Biotechnology* 18, 189-196.
- Hikita, C. & Mizushima, S. (1992). Effects of total hydrophobicity and length of the hydrophobic domain of a signal peptide on in vitro translocation efficiency. *Journal of Biological Chemistry*, 267, 4882-4888.
- Hoffmann, F. & Rinas, U. (2001). Plasmid amplification in *Escherichia coli* after temperature upshift is impaired by induction of recombinant protein synthesis. *Biotechnology Letters*, 23, 1819-1825.
- Hsu, C. H., Chu, Y. F., Argin-Soysal, S., Hahm, T. S. & Lo, Y. M. (2004). Effects of surface characteristics and xanthan polymers on the immobilization of *Xanthomonas campestris* to fibrous matrices. *Journal of Food Science*, 69, E441-E448.
- Hu, C. K., Bai, F. W. & An, L. J. (2003). Enhancing ethanol tolerance of a self-flocculating fusant of *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae* by Mg^{2+} via reduction in plasma membrane permeability. *Biotechnology Letters*, 25, 1191-1194.

- Hu, Y., Tang, T., Yang, W. & Zhou, H. (2009). Bioconversion of phenylpyruvic acid to L-phenylalanine by mixed-gel immobilization of *Escherichia coli* EP8-10. *Process Biochemistry*, 44, 142-145.
- Huang, J., Dhulster, P., Thomas, D. & Barbotin, J.-N. (1990). Agitation rate effects on plasmid stability in immobilized and free-cell continuous cultures of recombinant *E. coli*. *Enzyme and Microbial Technology*, 12, 933-939.
- Hucík, M., Bučko, M., Gemeiner, P., Štefuca, V., Vikartovská, A., Mihovilovič, M. D., Rudroff, F., Iqbal, N., Chorvát Jr, D. & Lacík, I. (2010). Encapsulation of recombinant *E. coli* expressing cyclopentanone monooxygenase in polyelectrolyte complex capsules for Baeyer–Villiger biooxidation of 8-oxabicyclo [3.2. 1] oct-6-en-3-one. *Biotechnology letters*, 32, 675-680.
- Hughes, J., Ramsden, D. & Symes, K. (1990). The flocculation of bacteria using cationic synthetic flocculants and chitosan. *Biotechnology Techniques*, 4, 55-60.
- Idris, A. & Suzana, W. (2006). Effect of sodium alginate concentration, bead diameter, initial pH and temperature on lactic acid production from pineapple waste using immobilized *Lactobacillus delbrueckii*. *Process Biochemistry*, 41, 1117-1123.
- Inloes, D. S., Smith, W. J., Taylor, D. P., Cohen, S. N., Michaels, A. S. & Robertson, C. R. (1983a). Hollow-fiber membrane bioreactors using immobilized *E. coli* for protein synthesis. *Biotechnology and Bioengineering*, 25, 2653-2681.
- Inloes, D. S., Taylor, D. P., Cohen, S. N., Michaels, A. S. & Robertson, C. R. (1983b). Ethanol production by *Saccharomyces cerevisiae* immobilized in hollow-fiber membrane bioreactors. *Applied and Environmental Microbiology*, 46, 264-278.
- Ismail, N. F., Hamdan, S., Mahadi, N. M., Murad, A. M. A., Rabu, A., Bakar, F. D. A., Klappa, P. & Illias, R. M. (2011). A mutant L-asparaginase II signal peptide improves the secretion of recombinant cyclodextrin glucanotransferase and the viability of *Escherichia coli*. *Biotechnology Letters*, 33, 999-1005.
- Isoai, A., Kimura, H., Reichert, A., Schörgendorfer, K., Nikaido, K., Tohda, H., Giga-Hama, Y., Mutoh, N. & Kumagai, H. (2002). Production of D-amino

- acid oxidase (DAO) of *Trigonopsis variabilis* in *Schizosaccharomyces pombe* and the characterization of biocatalysts prepared with recombinant cells. *Biotechnology and Bioengineering*, 80, 22-32.
- Jamuna, R. & Ramakrishna, S. (1992). Continuous synthesis of thermostable α -amylase by *Bacillus* cells immobilized in calcium alginate. *Enzyme and Microbial Technology*, 14, 36-41.
- Jang, K., Seo, J., Song, K., Kim, C. & Rhee, S. (1999). Extracellular secretion of levansucrase from *Zymomonas mobilis* in *Escherichia coli*. *Bioprocess Engineering*, 21, 453-458.
- Jeong, K. J. & Lee, S. Y. (2000). Secretory production of human leptin in *Escherichia coli*. *Biotechnology and Bioengineering*, 67, 398-407.
- Jin, Y. L. & Speers, R. A. (1998). Flocculation of *Saccharomyces cerevisiae*. *Food Research International*, 31, 421-440.
- Jonet, M. A., Mahadi, N. M., Murad, A. M. A., Rabu, A., Bakar, F. D. A., Rahim, R. A., Low, K. O. & Illias, R. M. (2012). Optimization of a heterologous signal peptide by site-directed mutagenesis for improved secretion of recombinant proteins in *Escherichia coli*. *Journal of Molecular Microbiology and Biotechnology*, 22, 48-58.
- Jong, W. S., Saurí, A. & Luirink, J. (2010). Extracellular production of recombinant proteins using bacterial autotransporters. *Current Opinion in Biotechnology*, 21, 646-652.
- Jouenne, T., Bonato, H., Mignot, L. & Junter, G.-A. (1993). Cell immobilization in composite agar layer microporous membrane structures: growth kinetics of gel-entrapped cultures and cell leakage limitation by a microporous membrane. *Applied Microbiology and Biotechnology*, 38, 478-481.
- Junter, G. A. & Jouenne, T. (2004). Immobilized viable microbial cells: from the process to the proteome... or the cart before the horse. *Biotechnology Advances*, 22, 633-658.
- Kanellaki, M., Bekatorou, A. & Koutinas, A. A. (2014). Low-temperature production of wine, beer and distillates using cold-adapted yeasts. *Cold-adapted Yeasts*. Springer.

- Karatan, E. & Watnick, P. (2009). Signals, regulatory networks, and materials that build and break bacterial biofilms. *Microbiology and Molecular Biology Reviews*, 73, 310-347.
- Kawaguti, H. Y., Buzzato, M. F., Orsi, D. C., Suzuki, G. T. & Sato, H. H. (2006). Effect of the additives polyethylenimine and glutaraldehyde on the immobilization of *Erwinia* sp. D12 cells in calcium alginate for isomaltulose production. *Process Biochemistry*, 41, 2035-2040.
- Khushoo, A., Pal, Y., Singh, B. N. & Mukherjee, K. (2004). Extracellular expression and single step purification of recombinant *Escherichia coli* l-asparaginase II. *Protein Expression and Purification*, 38, 29-36.
- Kilonzo, P., Margaritis, A. & Bergougnou, M. (2009). Airlift-driven fibrous-bed bioreactor for continuous production of glucoamylase using immobilized recombinant yeast cells. *Journal of Biotechnology*, 143, 60-68.
- Kilonzo, P., Margaritis, A. & Bergougnou, M. (2011). Effects of surface treatment and process parameters on immobilization of recombinant yeast cells by adsorption to fibrous matrices. *Bioresource Technology*, 102, 3662-3672.
- Kim, S.-G., Kweon, D.-H., Lee, D.-H., Park, Y.-C. & Seo, J.-H. (2005). Coexpression of folding accessory proteins for production of active cyclodextrin glycosyltransferase of *Bacillus macerans* in recombinant *Escherichia coli*. *Protein Expression and Purification*, 41, 426-432.
- Kjelleberg, S., Humphrey, B. A. & Marshall, K. C. (1982). Effect of interfaces on small, starved marine bacteria. *Applied and Environmental Microbiology*, 43, 1166-1172.
- Korber, D. R., Lawrence, J. R. & Caldwell, D. E. (1994). Effect of motility on surface colonization and reproductive success of *Pseudomonas fluorescens* in dual-dilution continuous culture and batch culture systems. *Applied and Environmental Microbiology*, 60, 1421-1429.
- Kosseva, M. R., Panesar, P. S., Kaur, G. & Kennedy, J. F. (2009). Use of immobilised biocatalysts in the processing of cheese whey. *International Journal of Biological Macromolecules*, 45, 437-447.

- Kourkoutas, Y., Bekatorou, A., Banat, I. M., Marchant, R. & Koutinas, A. (2004). Immobilization technologies and support materials suitable in alcohol beverages production: a review. *Food Microbiology*, 21, 377-397.
- Krapf, M. E. M., Lartiges, B. S., Merlin, C., Francius, G., Ghanbaja, J. & Duval, J. F. (2012). Polyethyleneimine-mediated flocculation of *Shewanella oneidensis* MR-1: Impacts of cell surface appendage and polymer concentration. *Water Research*, 46, 1838-1846.
- Krastanov, A. (1997). Continuous sucrose hydrolysis by yeast cells immobilized to wool. *Applied Microbiology and Biotechnology*, 47, 476-481.
- Kriaa, M., Ayadi, D. Z., Jemli, S., Sahnoun, M., Bejar, S. & Kammoun, R. (2012). Improvement of cyclodextrin glycosyltransferase (CGTase) production by recombinant *Escherichia coli* pAD26 immobilized on the cotton. *Biologia*, 67, 1049-1055.
- Kubitschek, H. (1968). Constancy of uptake during the cell cycle in *Escherichia coli*. *Biophysical Journal*, 8, 1401.
- Küenzi, M. T. & Fiechter, A. (1969). Changes in carbohydrate composition and trehalase-activity during the budding cycle of *Saccharomyces cerevisiae*. *Archiv für Mikrobiologie*, 64, 396-407.
- Kunamneni, A., Prabhakar, T., Jyothi, B. & Ellaiah, P. (2007). Investigation of continuous cyclodextrin glucanotransferase production by the alginate-immobilized cells of alkalophilic *Bacillus* sp. in an airlift reactor. *Enzyme and Microbial Technology*, 40, 1538-1542.
- Kurland, C. & Dong, H. (1996). Bacterial growth inhibition by overproduction of protein. *Molecular Microbiology*, 21, 1-4.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227, 680-685.
- Lahooti, S. & Sefton, M. V. (2000). Effect of an immobilization matrix and capsule membrane permeability on the viability of encapsulated HEK cells. *Biomaterials*, 21, 987-995.
- Lavallie, E. R., DiBlasio, E. A., Kovacic, S., Grant, K. L., Schendel, P. F. & McCoy, J. M. (1993). A thioredoxin gene fusion expression system that circumvents inclusion body formation in the *E. coli* cytoplasm. *Bio/Technology*, 11, 187-93.

- Lee, S. Y., Chang, H. N., Um, Y. S. & Hong, S. H. (1998). Bacteriorhodopsin production by cell recycle culture of halobacterium halobium. *Biotechnology Letters*, 20, 763-765.
- Lee, T., Ahn, J. & Ryu, D. D. (1983). Performance of an immobilized yeast reactor system for ethanol production. *Enzyme and Microbial Technology*, 5, 41-45.
- Li, Z. F., Li, B., Liu, Z. G., Wang, M., Gu, Z. B., Du, G. C., Wu, J. & Chen, J. (2009). Calcium leads to further increase in glycine-enhanced extracellular secretion of recombinant α -cyclodextrin glycosyltransferase in *Escherichia coli*. *Journal of Agricultural and Food Chemistry*, 57, 6231-6237.
- Liu, C., Wang, K., Jiang, J.-H., Liu, W.-J. & Wang, J.-Y. (2015). A novel bioflocculant produced by a salt-tolerant, alkaliphilic and biofilm-forming strain *Bacillus agaradhaerens* C9 and its application in harvesting *Chlorella minutissima* UTEX2341. *Biochemical Engineering Journal*, 93, 166-172.
- Liu, L., Liu, S. & Tan, X. (2010). Zirconia microbial hollow fibre bioreactor for *Escherichia coli* culture. *Ceramics International*, 36, 2087-2093.
- Lloyd, J. R. & Bunch, A. W. (1996). The physiological state of an ethylenogenic *Escherichia coli* immobilized in hollow-fiber bioreactors. *Enzyme and Microbial Technology*, 18, 113-120.
- Lo, P. K., Hassan, O., Ahmad, A., Mahadi, N. M. & Illias, R. M. (2007). Excretory over-expression of *Bacillus* sp. G1 cyclodextrin gluconotransferase (CGTase) in *Escherichia coli*: Optimization of the cultivation conditions by response surface methodology. *Enzyme and Microbial Technology*, 40, 1256-1263.
- Low, K. O., Mahadi, N. M., Rahim, R. A., Rabu, A., Bakar, F. D. A., Murad, A. M. A. & Illias, R. M. (2011). An effective extracellular protein secretion by an ABC transporter system in *Escherichia coli*: statistical modeling and optimization of cyclodextrin gluconotransferase secretory production. *Journal of Industrial Microbiology and Biotechnology*, 38, 1587-1597.
- Luedeking, R. & Piret, E. L. (1959). A kinetic study of the lactic acid fermentation. Batch process at controlled pH. *Journal of Biochemical and Microbiology*, 1, 393-412.
- Luirink, J., Watanabe, T., Wu, H., Stegehuis, F., De Graaf, F. & Oudega, B. (1987). Modification, processing, and subcellular localization in *Escherichia coli* of

- the pCloDF13-encoded bacteriocin release protein fused to the mature portion of beta-lactamase. *Journal of Bacteriology*, 169, 2245-2250.
- Luo, W., Liu, Y., Zhu, X., Zhao, W., Huang, L., Cai, J., Xu, Z. & Cen, P. (2011). Cloning and characterization of purine nucleoside phosphorylase in *Escherichia coli* and subsequent ribavirin biosynthesis using immobilized recombinant cells. *Enzyme and Microbial Technology*, 48, 438-444.
- Lusta, K. A., Chung, I. K., Sul, I. W., Park, H. S. & Shin, D. I. (2000). Immobilization of fungus *Aspergillus* sp. by a novel cryogel technique for production of extracellular hydrolytic enzymes. *Process Biochemistry*, 35, 1177-1182.
- Ma, T., Yang, S.-T. & Kniss, D. A. (1999). Development of an in vitro human placenta model by the cultivation of human trophoblasts in a fiber-based bioreactor system. *Tissue Engineering*, 5, 91-102.
- Mah, T. F. C. & O'Toole, G. A. (2001). Mechanisms of biofilm resistance to antimicrobial agents. *Trends in Microbiology*, 9, 34-39.
- Mahat, M. K., Illias, R. M., Rahman, R. A., Rashid, N. A. A., Mahmood, N. A. N., Hassan, O., Aziz, S. A. & Kamaruddin, K. (2004). Production of cyclodextrin glucanotransferase (CGTase) from alkalophilic *Bacillus* sp. TS1-1: media optimization using experimental design. *Enzyme and Microbial Technology*, 35, 467-473.
- Makrides, S. C. (1996). Strategies for achieving high-level expression of genes in *Escherichia coli*. *Microbiological Reviews*, 60, 512-538.
- Mamo, G. & Gessesse, A. (1997). Thermostable amylase production by immobilized thermophilic *Bacillus* sp. *Biotechnology Techniques*, 11, 447-450.
- Man, R. C., Ismail, A. F., Ghazali, N. F., Fuzi, S. F. Z. M. & Illias, R. M. (2015). Effects of the immobilization of recombinant *Escherichia coli* on cyclodextrin glucanotransferase (CGTase) excretion and cell viability. *Biochemical Engineering Journal*.
- Marshall, K. C. (1985). Mechanisms of bacterial adhesion at solid-water interfaces. *Bacterial Adhesion*, 133-161.
- Martins, R. F., Plieva, F. M., Santos, A. & Hatti-Kaul, R. (2003). Integrated immobilized cell reactor-adsorption system for β -cyclodextrin production: a

- model study using PVA-cryogel entrapped *Bacillus agaradhaerens* cells. *Biotechnology Letters*, 25, 1537-1543.
- Mathew, A. K., Crook, M., Chaney, K. & Humphries, A. C. (2013). Comparison of entrapment and biofilm mode of immobilisation for bioethanol production from oilseed rape straw using *Saccharomyces cerevisiae* cells. *Biomass & Bioenergy*, 52, 1-7.
- Matos, C. F., Branston, S. D., Albinia, A., Dhanoya, A., Freedman, R. B., Keshavarz-Moore, E. & Robinson, C. (2012). High-yield export of a native heterologous protein to the periplasm by the tat translocation pathway in *Escherichia coli*. *Biotechnology and Bioengineering*, 109, 2533-2542.
- Matsui, T., Sato, H., Sato, S., Mukataka, S. & Takahashi, J. (1990). Effects of nutritional conditions on plasmid stability and production of tryptophan synthase by a recombinant *Escherichia coli*. *Agricultural and Biological Chemistry*, 54, 619-624.
- Mazzer, C., Ferreira, L. R., Rodella, J. R. T., Moriwaki, C. & Matioli, G. (2008). Cyclodextrin production by *Bacillus firmus* strain 37 immobilized on inorganic matrices and alginate gel. *Biochemical Engineering Journal*, 41, 79-86.
- Mehaia, M. A. & Cheryan, M. (1984). Ethanol production in a hollow fiber bioreactor using *Saccharomyces cerevisiae*. *Applied Microbiology and Biotechnology*, 20, 100-104.
- Mergulhao, F., Summers, D. & Monteiro, G. (2005). Recombinant protein secretion in *Escherichia coli*. *Biotechnology Advances*, 23, 177-202.
- Mill, P. (1964). The nature of the interactions between flocculent cells in the flocculation of *Saccharomyces cerevisiae*. *Journal of General Microbiology*, 35, 61-68.
- Moriwaki, C., Mangolim, C. S., Ruiz, G. B., de Moraes, G. R., Baesso, M. L. & Matioli, G. (2014). Biosynthesis of CGTase by immobilized alkalophilic bacilli and crystallization of beta-cyclodextrin: Effective techniques to investigate cell immobilization and the production of cyclodextrins. *Biochemical Engineering Journal*, 83, 22-32.

- Moriwaki, C., Pelissari, F. M., Gonçalves, R. A. C., Gonçalves, J. E. & Matioli, G. (2007). Immobilization of *Bacillus firmus* strain 37 in inorganic matrix for cyclodextrin production. *Journal of Molecular Catalysis B-Enzymatic*, 49, 1-7.
- Mundra, P., Desai, K. & Lele, S. (2007). Application of response surface methodology to cell immobilization for the production of palatinose. *Bioresource Technology*, 98, 2892-2896.
- Munton, T. & Russell, A. (1973). Effect of glutaraldehyde on cell viability, triphenyltetrazolium reduction, oxygen uptake, and β -galactosidase activity in *Escherichia coli*. *Applied Microbiology*, 26, 508-511.
- Naim, R. & Ismail, A. F. (2013). Effect of polymer concentration on the structure and performance of PEI hollow fiber membrane contactor for CO₂ stripping. *Journal of Hazardous Materials*, 250, 354-361.
- Nam, S. W., Park, H. Y., Kim, J. H., Seo, J. H., Han, N. S. & Kim, B. W. (2001). Expression of *Bacillus macerans* cyclodextrin glucanotransferase gene in *Saccharomyces cerevisiae*. *Biotechnology Letters*, 23, 727-730.
- Narayanan, N., Khan, M. & Chou, C. P. (2010). Enhancing functional expression of heterologous lipase B in *Escherichia coli* by extracellular secretion. *Journal of Industrial Microbiology & Biotechnology*, 37, 349-361.
- Nedović, V. & Willaert, R. *Applications of cell immobilisation biotechnology*, Springer. 2006
- Neubauer, P., Lin, H. & Mathiszik, B. (2003). Metabolic load of recombinant protein production: inhibition of cellular capacities for glucose uptake and respiration after induction of a heterologous gene in *Escherichia coli*. *Biotechnology and Bioengineering*, 83, 53-64.
- Nguyen, A. L. & Luong, J. H. T. (1986). Diffusion in κ -carrageenan gel beads. *Biotechnology and Bioengineering*, 28, 1261-1267.
- Ni, Y. & Chen, R. (2009). Extracellular recombinant protein production from *Escherichia coli*. *Biotechnology Letters*, 31, 1661-1670.
- Norde, W. & Lyklema, J. (1989). Protein adsorption and bacterial adhesion to solid surfaces: a colloid-chemical approach. *Colloid Surface*, 38, 1-13.

- Novotny, J., Ganju, R. K., Smiley, S. T., Hussey, R. E., Luther, M. A., Recny, M. A., Siliciano, R. F. & Reinherz, E. L. (1991). A soluble, single-chain T-cell receptor fragment endowed with antigen-combining properties. *Proceedings of the National Academy of Sciences*, 88, 8646-8650.
- Obradovic, B., Nedović, V. A., Bugarski, B., Willaert, R. G. & Vunjak-Novakovic, G. (2004). Immobilised cell bioreactors. *Fundamentals of cell immobilisation biotechnology*. Springer.
- Ogbonna, J. C., Amano, Y. & Nakamura, K. (1989). Elucidation of optimum conditions for immobilization of viable cells by using calcium alginate. *Journal of Fermentation and Bioengineering*, 67, 92-96.
- Ong, R. M., Goh, K. M., Mahadi, N. M., Hassan, O., Rahman, R. N. Z. R. A. & Illias, R. M. (2008). Cloning, extracellular expression and characterization of a predominant β -CGTase from *Bacillus* sp. G1 in *E. coli*. *Journal of Industrial Microbiology and Biotechnology*, 35, 1705-1714.
- Onoda, T., Enokizono, J., Kaya, H., Oshima, A., Freestone, P. & Norris, V. (2000). Effects of calcium and calcium chelators on growth and morphology of *Escherichia coli* L-form NC-7. *Journal of Bacteriology*, 182, 1419-1422.
- Paetzel, M., Dalbey, R. E. & Strynadka, N. C. (2000). The structure and mechanism of bacterial type I signal peptidases: a novel antibiotic target. *Pharmacology & Therapeutics*, 87, 27-49.
- Pai, O., Banoth, L., Ghosh, S., Chisti, Y. & Banerjee, U. C. (2014). Biotransformation of 3-cyanopyridine to nicotinic acid by free and immobilized cells of recombinant *Escherichia coli*. *Process Biochemistry*, 49, 655-659.
- Pan, K. L., Hsiao, H. C., Weng, C. L., Wu, M. S. & Chou, C. P. (2003). Roles of DegP in prevention of protein misfolding in the periplasm upon overexpression of penicillin acylase in *Escherichia coli*. *Journal of Bacteriology*, 185, 3020-3030.
- Park, J. & Chang, H. (2000). Microencapsulation of microbial cells. *Biotechnology Advances*, 18, 303-319.

- Park, S. L., Kwon, M. J., Kim, S. K. & Nam, S. W. (2004). GroEL/ES chaperone and low culture temperature synergistically enhanced the soluble expression of CGTase in *E. coli*. *Journal of Microbiology and Biotechnology*, 14, 216-219.
- Petsch, D. & Anspach, F. B. (2000). Endotoxin removal from protein solutions. *Journal of Biotechnology*, 76, 97-119.
- Pilkington, P., Margaritis, A., Mensour, N. & Russell, I. (1998). Fundamentals of immobilised yeast cells for continuous beer fermentation: a review. *Journal of the Institute of Brewing*, 104, 19-31.
- Raafat, A. I., Araby, E. & Lotfy, S. (2012). Enhancement of fibrinolytic enzyme production from *Bacillus subtilis* via immobilization process onto radiation synthesized starch/dimethylaminoethyl methacrylate hydrogel. *Carbohydrate Polymers*, 87, 1369-1374.
- Ramírez, D. M. & Bentley, W. E. (1993). Enhancement of recombinant protein synthesis and stability via coordinated amino acid addition. *Biotechnology and Bioengineering*, 41, 557-565.
- Rehn, G., Grey, C., Branney, C. & Adlercreutz, P. (2013). Chitosan flocculation: An effective method for immobilization of *E. coli* for biocatalytic processes. *Journal of Biotechnology*, 165, 138-144.
- Rehn, G., Grey, C., Branney, C., Lindberg, L. & Adlercreutz, P. (2012). Activity and stability of different immobilized preparations of recombinant *E. coli* cells containing ω -transaminase. *Process Biochemistry*, 47, 1129-1134.
- Reinikainen, P., Korpela, K., Nissinen, V., Olkku, J., Söderlund, H. & Markkanen, P. (1989). *Escherichia coli* plasmid production in fermenter. *Biotechnology and Bioengineering*, 33, 386-393.
- Roble, N., Ogbonna, J. & Tanaka, H. (2003). A novel circulating loop bioreactor with cells immobilized in loofa (*Luffa cylindrica*) sponge for the bioconversion of raw cassava starch to ethanol. *Applied Microbiology and Biotechnology*, 60, 671-678.
- Robledo-Ortíz, J., Ramírez-Arreola, D., Gomez, C., González-Reynoso, O. & González-Núñez, R. (2010). Bacterial immobilization by adhesion onto agave-fiber/polymer foamed composites. *Bioresource Technology*, 101, 1293-1299.

- Rochex, A., Lecouturier, D., Pezron, I. & Lebeault, J.-M. (2004). Adhesion of a *Pseudomonas putida* strain isolated from a paper machine to cellulose fibres. *Applied Microbiology and Biotechnology*, 65, 727-733.
- Rols, J. L. & Goma, G. (1989). Enhancement of oxygen transfer rates in fermentation using oxygen-vectors. *Biotechnology Advances*, 7, 1-14.
- Rozkov, A., Avignone-Rossa, C., Ertl, P., Jones, P., O'Kennedy, R., Smith, J., Dale, J. & Bushell, M. (2004). Characterization of the metabolic burden on *Escherichia coli* DH1 cells imposed by the presence of a plasmid containing a gene therapy sequence. *Biotechnology and Bioengineering*, 88, 909-915.
- Safarikova, M., Atanasova, N., Ivanova, V., Weyda, F. & Tonkova, A. (2007). Cyclodextrin glucanotransferase synthesis by semicontinuous cultivation of magnetic biocatalysts from cells of *Bacillus circulans* ATCC 21783. *Process Biochemistry*, 42, 1454-1459.
- Saswathi, N., Sheela, R., Jamuna, R. & Ramakrishna, S. (1995). Synthesis of cyclodextrin glycosyl transferase by immobilized cells of *Bacillus circulans*. *Bioprocess Engineering*, 12, 283-286.
- Sato, K. & Toda, K. (1983). Oxygen uptake rate of immobilized growing *Candida lipolytica*. *Journal of Fermentation Technology*, 61, 239-245.
- Sayadi, S., Nasri, M., Berry, F., Barbotin, J. N. & Thomas, D. (1987). Effect of temperature on the stability of plasmid pTG201 and productivity of xylE gene product in recombinant *Escherichia coli*: development of a two-stage chemostat with free and immobilized cells. *Journal of General Microbiology*, 133, 1901-1908.
- Schein, C. H. & Noteborn, M. H. (1988). Formation of soluble recombinant proteins in *Escherichia coli* is favored by lower growth temperature. *Nature Biotechnology*, 6, 291-294.
- Senthuran, A., Senthuran, V., Mattiasson, B. & Kaul, R. (1997). Lactic acid fermentation in a recycle batch reactor using immobilized *Lactobacillus casei*. *Biotechnology and Bioengineering*, 55, 841-853.
- Shi, X., Karkut, T., Chamankhah, M., Alting-Mees, M., Hemmingsen, S. M. & Hegedus, D. (2003). Optimal conditions for the expression of a single-chain

- antibody (scFv) gene in *Pichia pastoris*. *Protein Expression and Purification*, 28, 321-330.
- Shin, H. D. & Chen, R. R. (2008). Extracellular recombinant protein production from an *Escherichia coli* lpp deletion mutant. *Biotechnology and Bioengineering*, 101, 1288-1296.
- Shokri, A., Sandén, A. & Larsson, G. (2002). Growth rate-dependent changes in *Escherichia coli* membrane structure and protein leakage. *Applied Microbiology and Biotechnology*, 58, 386-392.
- Shokri, A., Sandén, A. & Larsson, G. (2003). Cell and process design for targeting of recombinant protein into the culture medium of *Escherichia coli*. *Applied Microbiology and Biotechnology*, 60, 654-664.
- Shreve, G. S. & Vogel, T. M. (1993). Comparison of substrate utilization and growth kinetics between immobilized and suspended *Pseudomonas* cells. *Biotechnology and Bioengineering*, 41, 370-379.
- Shukla, R., Kang, W. & Sirkar, K. (1989). Novel hollow fiber immobilization techniques for whole cells and advanced bioreactors. *Applied Biochemistry and Biotechnology*, 20, 571-586.
- Shuler, M. L. & Kargi, F. *Bioprocess engineering*, Prentice Hall New York. 2002
- Sian, H. K., Said, M., Hassan, O., Kamaruddin, K., Ismail, A. F., Rahman, R. A., Mahmood, N. A. N. & Illias, R. M. (2005). Purification and characterization of cyclodextrin glucanotransferase from alkalophilic *Bacillus* sp. G1. *Process Biochemistry*, 40, 1101-1111.
- Silva, F., Passarinha, L., Sousa, F., Queiroz, J. A. & Domingues, F. C. (2009). Influence of growth conditions on plasmid DNA production. *Journal of Microbiology and Biotechnology*, 19, 1408-1414.
- Sletta, H., Nedal, A., Aune, T., Hellebust, H., Hakvåg, S., Aune, R., Ellingsen, T., Valla, S. & Brautaset, T. (2004). Broad-host-range plasmid pJB658 can be used for industrial-level production of a secreted host-toxic single-chain antibody fragment in *Escherichia coli*. *Applied and Environmental Microbiology*, 70, 7033-7039.
- Sleytr, U. B. (1997). I. Basic and applied S-layer research: an overview. *FEMS Microbiology Reviews*, 20, 5-12.

- Smet, C., Van Derlinden, E., Mertens, L., Noriega, E. & Van Impe, J. F. (2015). Effect of cell immobilization on the growth dynamics of *Salmonella Typhimurium* and *Escherichia coli* at suboptimal temperatures. *International Journal of Food Microbiology*, 208, 75-83.
- Song, B. & Leff, L. G. (2006). Influence of magnesium ions on biofilm formation by *Pseudomonas fluorescens*. *Microbiological Research*, 161, 355-361.
- Steinberger, R. & Holden, P. (2004). Macromolecular composition of unsaturated *Pseudomonas aeruginosa* biofilms with time and carbon source. *Biofilms*, 1, 37-47.
- Stoodley, P., Cargo, R., Rupp, C., Wilson, S. & Klapper, I. (2002). Biofilm material properties as related to shear-induced deformation and detachment phenomena. *Journal of Industrial Microbiology and Biotechnology*, 29, 361-367.
- Sunitha, K., Kim, Y.-O., Lee, J.-K. & Oh, T.-K. (2000). Statistical optimization of seed and induction conditions to enhance phytase production by recombinant *Escherichia coli*. *Biochemical Engineering Journal*, 5, 51-56.
- Sutherland, I. W. (2001a). Biofilm exopolysaccharides: a strong and sticky framework. *Microbiology*, 147, 3-9.
- Sutherland, I. W. (2001b). Microbial polysaccharides from Gram-negative bacteria. *International Dairy Journal*, 11, 663-674.
- Swamy, K. & Goldberg, A. (1982). Subcellular distribution of various proteases in *Escherichia coli*. *Journal of Bacteriology*, 149, 1027-1033.
- Swartz, J. R. (1996). *Escherichia coli* recombinant DNA technology. *Escherichia coli and Salmonella: cellular and molecular biology*, 2nd ed. ASM Press, Washington, DC, 1693-1711.
- Tachibana, Y., Kuramura, A., Shirasaka, N., Suzuki, Y., Yamamoto, T., Fujiwara, S., Takagi, M. & Imanaka, T. (1999). Purification and characterization of an extremely thermostable cyclomaltodextrin glucanotransferase from a newly isolated hyperthermophilic archaeon, a *Thermococcus* sp. *Applied and Environmental Microbiology*, 65, 1991-1997.
- Takagi, M., Shiwaku, K., Inoue, T., Shirakawa, Y., Sawa, Y., Matsuda, H. & Yoshida, T. (2003). Hydrodynamically stable adhesion of endothelial cells

- onto a polypropylene hollow fiber membrane by modification with adhesive protein. *Journal of Artificial Organs*, 6, 222-226.
- Talmadge, K. & Gilbert, W. (1982). Cellular location affects protein stability in *Escherichia coli*. *Proceedings of the National Academy of Sciences*, 79, 1830-1833.
- Terpe, K. (2006). Overview of bacterial expression systems for heterologous protein production: from molecular and biochemical fundamentals to commercial systems. *Applied Microbiology and Biotechnology*, 72, 211-222.
- Toda, K. & Shoda, M. (1975). Sucrose inversion by immobilized yeast cells in a complete mixing reactor. *Biotechnology and Bioengineering*, 17, 481-497.
- Tokuda, H. (2009). Biogenesis of outer membranes in Gram-negative bacteria. *Bioscience, Biotechnology, and Biochemistry*, 73, 465-473.
- Tripathi, A., Sami, H., Jain, S. R., Vilorio-Cols, M., Zhuravleva, N., Nilsson, G., Jungvid, H. & Kumar, A. (2010). Improved bio-catalytic conversion by novel immobilization process using cryogel beads to increase solvent production. *Enzyme and Microbial Technology*, 47, 44-51.
- Urbance, S. E., Pometto III, A. L., DiSpirito, A. A. & Denli, Y. (2004). Evaluation of succinic acid continuous and repeat-batch biofilm fermentation by *Actinobacillus succinogenes* using plastic composite support bioreactors. *Applied Microbiology and Biotechnology*, 65, 664-670.
- van der Wal, F. J., Luirink, J. & Oudega, B. (1995). Bacteriocin release proteins: mode of action, structure, and biotechnological application. *FEMS Microbiology Reviews*, 17, 381-399.
- van Dijl, J. M., de Jong, A., Smith, H., Bron, S. & Venema, G. (1991). Signal peptidase I overproduction results in increased efficiencies of export and maturation of hybrid secretory proteins in *Escherichia coli*. *Molecular and General Genetics MGG*, 227, 40-48.
- Van Loosdrecht, M., Lyklema, J., Norde, W., Schraa, G. & Zehnder, A. (1987). Electrophoretic mobility and hydrophobicity as a measured to predict the initial steps of bacterial adhesion. *Applied and Environmental Microbiology*, 53, 1898-1901.

- Van Loosdrecht, M., Lyklema, J., Norde, W. & Zehnder, A. (1990). Influence of interfaces on microbial activity. *Microbiological Reviews*, 54, 75-87.
- Vassileva, A., Beschkov, V., Ivanova, V. & Tonkova, A. (2005). Continuous cyclodextrin glucanotransferase production by free and immobilized cells of *Bacillus circulans* ATCC 21783 in bioreactors. *Process Biochemistry*, 40, 3290-3295.
- Vassileva, A., Burhan, N., Beschkov, V., Spasova, D., Radoevska, S., Ivanova, V. & Tonkova, A. (2003). Cyclodextrin glucanotransferase production by free and agar gel immobilized cells of *Bacillus circulans* ATCC 21783. *Process Biochemistry*, 38, 1585-1591.
- Velaithan, V., Chin, S. C., Yusoff, K., Illias, R. M. & Rahim, R. A. (2014). Novel synthetic signal peptides for the periplasmic secretion of green fluorescent protein in *Escherichia coli*. *Annals of Microbiology*, 64, 543-550.
- Verdoes, J. C., Punt, P. & Van Den Hondel, C. (1995). Molecular genetic strain improvement for the overproduction of fungal proteins by filamentous fungi. *Applied Microbiology and Biotechnology*, 43, 195-205.
- Villaverde, A. & Carrió, M. M. (2003). Protein aggregation in recombinant bacteria: biological role of inclusion bodies. *Biotechnology Letters*, 25, 1385-1395.
- Vinogradov, A. M., Hugo Schmidt, V., Tuthill, G. F. & Bohannon, G. W. (2004). Damping and electromechanical energy losses in the piezoelectric polymer PVDF. *Mechanics of Materials*, 36, 1007-1016.
- von Weymarn, N., Kiviharju, K. & Leisola, M. (2002). High-level production of D-mannitol with membrane cell-recycle bioreactor. *Journal of Industrial Microbiology and Biotechnology*, 29, 44-49.
- Vučurović, V. M. & Razmovski, R. N. (2012). Sugar beet pulp as support for *Saccharomyces cerevisiae* immobilization in bioethanol production. *Industrial Crops and Products*, 39, 128-134.
- Walls, E. L. & Gainer, J. L. (1991). Increased protein productivity from immobilized recombinant yeast. *Biotechnology and Bioengineering*, 37, 1029-1036.
- Wang, J. Y. & Chao, Y. P. (2006). Immobilization of cells with surface-displayed chitin-binding domain. *Applied and Environmental Microbiology*, 72, 927-931.

- Wang, S. C., Jin, M. R. & Hall, D. O. (1991). Immobilization of *Anabaena azollae* in hollow fibre photobioreactors for ammonia production. *Bioresource Technology*, 38, 85-90.
- Wang, Z., Le, G., Shi, Y. & Węgrzyn, G. (2001). Medium design for plasmid DNA production based on stoichiometric model. *Process Biochemistry*, 36, 1085-1093.
- Watanabe, I., Miyata, N., Ando, A., Shiroma, R., Tokuyasu, K. & Nakamura, T. (2012). Ethanol production by repeated-batch simultaneous saccharification and fermentation (SSF) of alkali-treated rice straw using immobilized *Saccharomyces cerevisiae* cells. *Bioresource Technology*, 123, 695-698.
- Wróbel, B. & Węgrzyn, G. (1997). Amplification of pSC101 replicons in *Escherichia coli* during amino acid limitation. *Journal of Biotechnology*, 58, 205-208.
- Wu, T., Malinverni, J., Ruiz, N., Kim, S., Silhavy, T. J. & Kahne, D. (2005). Identification of a multicomponent complex required for outer membrane biogenesis in *Escherichia coli*. *Cell*, 121, 235-245.
- Xu, Z. N., Shen, W. H., Chen, H. & Cen, P. L. (2005). Effects of medium composition on the production of plasmid DNA vector potentially for human gene therapy. *Journal of Zhejiang University- Science B*, 6, 396.
- Yang, P., Teo, W. K. & Ting, Y. P. (2006). Design and performance study of a novel immobilized hollow fiber membrane bioreactor. *Bioresource Technology*, 97, 39-46.
- Ying, L., Kang, E. & Neoh, K. (2002). Covalent immobilization of glucose oxidase on microporous membranes prepared from poly (vinylidene fluoride) with grafted poly (acrylic acid) side chains. *Journal of Membrane Science*, 208, 361-374.
- Yoon, S. H., Kim, S. K. & Kim, J. F. (2010). Secretory production of recombinant proteins in *Escherichia coli*. *Recent Patents on Biotechnology*, 4, 23-29.
- Young, V. R. & Marchini, J. S. (1990). Mechanisms and nutritional significance of metabolic responses to altered intakes of protein and amino acids, with reference to nutritional adaptation in humans. *The American Journal of Clinical Nutrition*, 51, 270-289.

- Yu, J., Yue, G., Zhong, J., Zhang, X. & Tan, T. (2010). Immobilization of *Saccharomyces cerevisiae* to modified bagasse for ethanol production. *Renewable Energy*, 35, 1130-1134.
- Zaushitsyna, O., Berillo, D., Kirsebom, H. & Mattiasson, B. (2014). Cryostructured and crosslinked viable cells forming monoliths suitable for bioreactor applications. *Topics in Catalysis*, 57, 339-348.
- Zheng, C., Zhao, Z., Li, Y., Wang, L. & Su, Z. (2011). Effect of IPTG amount on apo-and holo-forms of glycerophosphate oxidase expressed in *Escherichia coli*. *Protein Expression and Purification*, 75, 133-137.
- Zhinan, X., Gang, L., Peilin, C. & Wong, W. (2000). Factors influencing excretive production of human epidermal growth factor (hEGF) with recombinant *Escherichia coli* K12 system. *Bioprocess Engineering*, 23, 669-674.