

HIGH SPATIAL RESOLUTION AND HYPERSPECTRAL REMOTE SENSING  
FOR MAPPING VEGETATION SPECIES IN TROPICAL RAINFOREST

ALVIN LAU MENG SHIN

A thesis submitted in fulfilment of the  
requirements for the award of the degree of  
Doctor of Philosophy (Remote Sensing)

Faculty of Geoinformation Science and Engineering  
Universiti Teknologi Malaysia

DECEMBER 2009

For my wife and my family.

## ACKNOWLEDGEMENTS

I would like to express my heartiest thanks and profound gratitude to my supervisor, Prof. Dr. Mazlan bin Hashim for his untiring guidance, invaluable assistance, endless encouragement and strong support throughout the period of my research. My sincere thanks to Prof. Dr. Toshinori Okuda, my external supervisor, for giving me many useful advices during my research and throughout the period of my six months' attachment in the Graduate School of Integrated Art and Sciences, Hiroshima University, Japan.

I must also express my heartfelt gratitude to Prof. Dr. Shatri Mansor, Prof. Dr. Mohd Ibrahim Seeni Mohd and Assoc. Prof. Dr. Ab Latif Ibrahim (who kindly sat on my viva voce panels) as well as Assoc. Prof. Dr. Md. Noor Kamaruddin (chairman of the viva voce panel). They have all given me many kind and useful advice during the viva voce for the improvement of my research.

I also wish to express my deepest appreciation to the staff of the Forest Research Institute Malaysia, who have been helpful in providing me with census data and assistance. I am truly indebted to the Universiti Teknologi Malaysia for the award of the scholarship and the study leave granted to me, thus providing me such a golden opportunity to pursue my post-graduate studies.

Last but not least, I wish to thank in particular my parents and my wife for their unselfish and unfailing love, tender care, affectionate encouragement and constant prayers. To all those whose names are too many to be mentioned here but have rendered me their assistance in many wonderful ways during my research and thesis writing, I also most sincerely acknowledge my thanks and gratitude.

## ABSTRACT

The focus of this study is on vegetation species mapping using high spatial resolution IKONOS-2 and digital Color Infrared (CIR) Aerial Photos (spatial resolution 4 m for IKONOS-2 and 20 cm for CIR) and Hyperion Hyperspectral data (spectral resolution 10 nm) in Pasoh Forest Reserve, Negeri Sembilan. Spatial and spectral separability in distinguishing vegetation species were investigated prior to vegetation species mapping to provide optimal vegetation species discrimination. A total of 88 selected vegetation species and common timber groups of the dominant family Dipterocarpaceae with diameter at breast height more than 30 cm were used in this study, where trees spectra were collected by both in situ and laboratory measurements of foliar samples. The trees spectra were analysed using first and second order derivative analysis together with scatter matrix plot based on multi-objective optimization algorithm to identify the best separability and sensitive wavelength portions for vegetation species mapping. In high spatial resolution data mapping, both IKONOS-2 and CIR data were classified by supervised classification approach using maximum likelihood and neural network classifiers, while the Hyperion data was classified by spectral angle mapper and linear mixture modeling. Results of this study indicate that only a total of ten common timber group of dominant Dipterocarpaceae genus were able to be recognized at significant divergence. Both high spatial resolution data (IKONOS-2 and CIR) gave very good classification accuracy of more than 83%. The classified hyperspectral data at 30 m spatial resolution gave a classification accuracy of 65%, hence confirming that spatial resolution is more sensitive in identification of tree genus. However, for species mapping, both high spatial and spectral remotely sensed data used are marginally less sensitive than at genus level.

## ABSTRAK

Kajian ini memfokuskan pemetaan spesies tumbuhan dengan menggunakan data resolusi spatial yang tinggi IKONOS-2 dan foto udara berdigit inframerah berwarna (CIR) (resolusi spatial 4 m bagi data IKONOS-2 dan 20 cm untuk CIR) dan data *Hyperion Hyperspectral* (resolusi spektral 10 nm) di Hutan Simpanan Pasoh, Negeri Sembilan. Untuk memberi perbezaan spesies tumbuhan yang optimum, keupayaan pemisahan spatial dan spektral dikaji terlebih dahulu sebelum pemetaan spesies tumbuhan dijalankan. Sejumlah 88 jenis spesies tumbuhan terpilih dan tumbuhan dalam kumpulan balak umum untuk keluarga dominan *Dipterocarpaceae* yang mempunyai diameter pada ketinggian paras dada lebih daripada 30 cm telah digunakan di mana spektra pokok dikumpulkan dengan penyampelan folia lapangan dan makmal. Untuk mengenalpasti pemisahan bahagian panjang gelombang yang baik dan sensitif dalam pemetaan spesies tumbuhan, spektra pokok dianalisis dengan analisa derivatif pertama dan kedua bersama dengan plot matrik serakan berasaskan algoritma optimikasi multi-objektif. Dalam pemetaan data resolusi spatial tinggi, data IKONOS-2 dan CIR dikelaskan dengan pendekatan pengkelasan berpenyelia menggunakan pengkelas kemungkinan maksimum dan rangkaian saraf manakala data *Hyperion* dikelaskan dengan pemeta spektra bersudut dan model campuran berkadar langsung. Keputusan kajian ini menunjukkan bahawa hanya sejumlah sepuluh jenis kumpulan balak umum dalam keluarga dominan *Dipterocarpaceae* dapat dikenali dengan nyata perbezaannya pada peringkat genus. Kedua-dua data resolusi spatial yang tinggi (IKONOS-2 dan CIR) memberi ketepatan pengkelasan yang sangat baik, iaitu melebihi 83%. Data *hyperspectral* yang telah dikelaskan pada resolusi spatial 30 m memberi ketepatan pengkelasan 65%, maka disahkan bahawa resolusi spatial adalah lebih sensitif dalam mengenalpasti genus pokok. Walau bagaimanapun, bagi pemetaan spesies, kedua-dua data remote sensing yang mempunyai resolusi spatial dan spektral yang tinggi adalah kurang sensitif jika berbanding dengan pemetaan tumbuhan pada peringkat genus.

## TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	<b>DECLARATION</b>	ii
	<b>DEDICATION</b>	iii
	<b>ACKNOWLEDGEMENTS</b>	iv
	<b>ABSTRACT</b>	v
	<b>ABSTRAK</b>	vi
	<b>TABLE OF CONTENTS</b>	vii
	<b>LIST OF TABLES</b>	xi
	<b>LIST OF FIGURES</b>	xiii
	<b>LIST OF ABBREVIATIONS</b>	xix
	<b>LIST OF APPENDICES</b>	xxi
<b>1</b>	<b>INTRODUCTION</b>	1
	1.1 Introduction	1
	1.2 Importance of Vegetation Study	3
	1.3 The Use of Remote Sensing Analysis in Vegetation Studies	5
	1.4 Problem Statement	7
	1.5 Objectives of the Study	11
	1.6 Scopes of the Study	12
	1.7 Significance of Research	13
	1.8 Study Area	14
	1.9 Thesis Outline	17

<b>2</b>	<b>LITERATURE REVIEW</b>	<b>18</b>
2.1	Introduction	18
2.2	High Spatial Remote Sensing and Its Applications	18
2.3	Hyperspectral Remote Sensing and Its Applications	20
2.4	Remote Sensing in Vegetation Mapping	24
	2.4.1 Spectral Characteristic of Vegetation	25
	2.4.2 Spatial Characteristic of Vegetation	28
	2.4.3 Modeling Canopy Reflectance	31
	2.4.4 Temporal Characteristic of Vegetation	32
2.5	Inference on Previous Studies	34
2.6	Summary	42
<b>3</b>	<b>DATA PREPARATION AND PRE-PROCESSING</b>	<b>47</b>
3.1	Introduction	47
3.2	Data Acquisition	49
	3.2.1 Remotely Sensed Data	49
	3.2.1.1 High Spatial Remotely Sensed Data	49
	3.2.1.2 Hyperspectral Remotely Sensed Data	56
	3.2.1.3 Spectral Radiometer Data	58
	3.2.2 Ancillary Data	58
3.3	Data Pre-processing	59
	3.3.1 Radiometric Calibration using Empirical Line Calibration Method	59
	3.3.2 Geometric Correction and Data Mosaicking	64
	3.3.2.1 Geometric Rectification of Hyperion Hyperspectral Data	65
	3.3.2.2 Data Mosaicking	67
	3.3.3 Clouds and Shadow Effects Masking	70
3.4	Image Enhancement of Hyperion Hyperspectral Data	71
	3.4.1 Minimum Noise Fraction	72

3.4.2	Pixel Purity Index	78
3.5	Census Data Plot of Study Area	81
3.5.1	Establish DBH to Crown Size Relationship	82
3.5.2	Derivative of Crown Data	85
3.6	Analysis and Discussion	88
3.7	Summary	90
<b>4</b>	<b>CREATION AND ANALYSIS OF SPECTRAL LIBRARY</b>	<b>91</b>
4.1	Introduction	91
4.2	Role of Spectral Library in Hyperspectral Analysis	91
4.3	Creation of Spectral Library	92
4.3.1	Conceptual Design of Spectral Library	95
4.3.2	Creation of Spectral Library by Field Campaign	96
4.3.3	Creation of Spectral Library by Laboratory Analyses	99
4.3.4	Compiling of Spectral Library	103
4.4	Spectral Analysis and Discussion	107
4.4.1	Spectral Separability and Sensitivity Analyses	112
4.4.1.1	Spectral Separability by First and Second Order Derivative Analysis	112
4.4.1.2	Spectral Separability by Combination of Three or Four Wavebands	115
4.4.1.3	Spectral Separability by Scatter Matrix Plot Analyses	116
4.4.2	Compatibility Test of Spectral Library	125
4.4.3	Spatial Sensitivity Analysis	128
4.5	Summary	131



<b>5</b>	<b>VEGETATION SPECIES MAPPING</b>	132
5.1	Introduction	132
5.2	Vegetation Mapping using High Spatial Data	132
5.2.1	Vegetation Species Mapping using Maximum Likelihood Classifier	133
5.2.2	Vegetation Species Mapping using Neural Network Classifier	139
5.3	Vegetation Mapping using Hyperspectral Data	145
5.3.1	Feature Extraction from Hyperspectral Data	145
5.3.1.1	Band Selection	146
5.3.1.2	Feature Extraction	147
5.3.2	Spectral Angle Mapper	149
5.3.3	Linear Mixture Modelling	153
5.4	Classification Accuracy Assessment	157
5.5	Analysis and Discussion	161
5.6	Summary	164
<b>6</b>	<b>CONCLUSIONS AND RECOMMENDATIONS</b>	165
6.1	Conclusions	165
6.2	Recommendations	168
	<b>REFERENCES</b>	169
	Appendices A-B	190-211

## LIST OF TABLES

<b>TABLE NO.</b>	<b>TITLE</b>	<b>PAGE</b>
1.1	The most important families for 50-ha plot of Pasoh Forest Reserve	15
2.1	Summary of some vegetation and related studies done using remote sensing techniques.	42
3.1	Specification of IKONOS-2 Data Used.	50
3.2	Detailed Specifications of the Z/I Imaging's DMC.	52
3.3	Specification of ancillary data i.e. topographic maps used in this study.	58
3.4	Summary of geometric correction control points used in Hyperion data.	66
3.5	Eigenvalues for MNF bands.	75
3.6	DBH and mean crown area used to develop DBH-crown size relationship for eight common species at genus level.	84
4.1	Specifications of Spectroradiometer used in this study.	96
4.2	List of spectra collected from Pasoh Forest Reserve, PFR according to family, genus and species.	98
4.3	List of spectra collected from 50-ha plot of Pasoh Forest Reserve during second field campaign.	101
4.4	List of spectra collected from secondary forest, FRIM during third field campaign.	102
4.5	Divergence analysis results of the scatter plots at species and genus level.	119

4.6	Lists of Species used in compatibility test of spectral library.	125
4.7	Paired Samples Statistics of Spectra collected from Pasoh Forest Reserve and FRIM, Kepong.	126
4.8	Paired samples correlations of selected vegetation spectra collected from Pasoh Forest Reserve and FRIM, Kepong	127
4.9	Paired sample test of selected vegetation spectra collected from Pasoh Forest Reserve and FRIM, Kepong.	127
5.1	Information extraction and classification accuracy with different techniques used for high spatial resolution data.	159
5.2	Information extraction and classification accuracy with different techniques used for Hyperion hyperspectral data.	160

## LIST OF FIGURES

FIGURES NO.	TITLE	PAGE
1.1	Study area; (a) location map, (b) the raw IKONOS satellite image of the study area and (c) the detailed plan of Pasoh Forest Reserve.	16
2.1	The hyperspectral concept.	21
2.2	Spectral reflectance characteristic of a healthy, green vegetation for the wavelength interval 0.4-2.6 $\mu\text{m}$ .	26
3.1	Data processing involved four main parts: (1) data acquisition and preprocessing; (2) spectral separability and sensitivity analysis, (3) building and analysis of spectral library; and (4) vegetation species mapping.	48
3.2	(a) and (b) Raw IKONOS-2 data, (c) Mosaiced IKONOS data with which (d) the subset of the study area are shown with clouds and its shadows being masked in black color.	51
3.3	(a) optics frame, (b) mounting cameras on optics frame, (c) camera sensors, (d) outlook view of camera and (e) camera mounted on optics frame and (f) DMC installed in airplane ready for data acquisition.	53
3.4	Flight index map for DMC mission over Pasoh Forest Reserve.	54
3.5	Raw CIR data of study area- 50-ha plot of Pasoh Forest Reserve in which Infra-red, Red, and Green bands are loaded in RGB.	55

3.6	(a) Raw Hyperion hyperspectral data, (b) subset of Pasoh Forest Reserve (c) 50-ha plot of Pasoh Forest Reserve in which bands 40, 31, 13 loaded in R, G, B.	57
3.7	Spectra extracted from reference and same target for (a) dark area and (b) light area from Hyperion image.	60
3.8	Regression plot for reference and image spectra.	61
3.9	Image (a) before radiometric calibration and (b) after radiometric calibration using Empirical Line Calibration Method in which bands 40, 31, 13 were loaded in R, G, B.	62
3.10	Spectra plots before and after radiometric calibration for (a) plantation a, (b) plantation b, (c) forest a and (d) forest b extracted from Hyperion image using empirical radiometric calibration method.	63
3.11	Ground control points location selected from Hyperion data.	65
3.12	Geometrically corrected Hyperion data in which bands 40, 30, 13 are loaded in RGB.	67
3.13	Two sets of georeferenced IKONOS-2 images loaded for mosaicking (scene po14395_0000 loaded in green frame and scene po140395_0100 loaded in red frame).	68
3.14	Two sets of IKONOS-2 data successfully mosaicked in which bands 3, 2, 1 are loaded in RGB.	69
3.15	Mask built to IKONOS-2 data, in which the clouds and shadow effects were being displayed in white color.	70
3.16	IKONOS-2 image after masking out the clouds and shadow areas.	71
3.17	The MNF process employed in this study.	74
3.18	MNF eigenvalues plot.	75

3.19	The first six MNF Bands generated from MNF Transformation for Hyperion data.	76
3.20	The PPI algorithm works by projecting points in the data set onto random skewers.	79
3.21	Pixel Purity Index plot up to 100 000 iterations.	80
3.22	PPI result with white pixels representing pure pixels.	81
3.23	The longest axis of crown and the longest axis perpendicular to the first axis were measured and the means were calculated to give the crown size value in diameter.	83
3.24	Scatter plot and equation with coefficient of the determination for the relationship between DBH and crown size in Pasoh 50-ha plot.	85
3.25	Census plot of Pasoh 50-ha plot at genus level for tree with DBH more than 30 cm.	86
3.26	Census plot has been converted into raster form in various spatial resolutions to test its sensitivity in spatial variation.	87
4.1	Methods employed for building the spectral library.	94
4.2	The data model of Spectral Library System of featured vegetation species.	95
4.3	Field Spectral sampling configuration used in the study.	97
4.4	Spectral sampling configuration used in the study.	99
4.5	Welcoming screen of spectral library.	103
4.6	Vegetation family selection screen of the spectral library; in this demonstration, vegetation family <i>DIPTEROCARPACEAE</i> is selected.	104
4.7	Vegetation genus selection screen of the spectral library; in this example, vegetation genus <i>Shorea</i> was selected.	104

4.8	Vegetation species selection screen of the spectral library; in this example, vegetation species <i>Multiflora</i> was selected.	105
4.9	Detailed spectral library plot for <i>Shorea Multiflora</i> .	106
4.10	Spectral plot of all 37 vegetation species of Pasoh Forest Reserve acquired during field campaigns.	108
4.11	Spectra Plots of common timber species of Pasoh Forest Reserve acquired during field campaigns.	109
4.12	Typical spectra plots of vegetation under various sampling condition, to simulate natural sampling environment analyses.	110
4.13	Spectra plots of <i>Xerosperrum noronhi</i> showing that mossy leaves gives higher reflectance values.	111
4.14	First order derivative (1/nm) of spectral reflectance plot.	113
4.15	Second order derivative (1/nm) of spectral reflectance plot.	114
4.16	Second order derivative (1/nm) of spectral reflectance plot for genus <i>Shorea</i> .	114
4.17	The separability of genus <i>Shorea</i> analyzed by choosing (a) three and (b) four bands combination.	115
4.18	Spectra plot for common timber groups of the dominant family Diterocarpaceae with selected wavelength portion, namely blue-green, green-peak, yellow-edge, red-well and near infrared.	117
4.19	(a) Spectral matrix plot of common timber species of the dominant family Diterocarpaceae of Pasoh Forest Reserve at blue-green portions of wavelengths and (b) closer zoom in of scatter matrix plot gives a clearer view of good separability wavelengths.	120

4.20	(a) Spectral matrix plot of common timber groups of the dominant family Dipterocarpaceae of Pasoh Forest Reserve at blue-green portions of wavelengths and (b) closer zoom in of scatter matrix plot gives a clearer view of good separability wavelengths.	122
4.21	Spectra plot of selected vegetation species collected from Pasoh Forest Reserve and FRIM.	126
4.22	Spectra plot at leaf-scale for various vegetation species	129
4.23	Leaf-scale spectra showing unhealthy leaves having higher reflectance at visible and near infrared portion of wavelength.	129
4.24	Leaf-scale vs. pixel-scale spectra plot for <i>ELATTA sp.</i>	130
4.25	Spectra plot of <i>ELATTA sp.</i> at various scales.	130
5.1	Classification result of IKONOS-2 data for selected vegetation species of Pasoh 50-ha plot (PFR) using Maximum Likelihood Classifier.	135
5.2	Classification result of IKONOS-2 data for common timber groups of the dominant family Dipterocarpaceae in the Pasoh 50-ha plot (PFR) using Maximum Likelihood Classifier.	136
5.3	Classification result of CIR data for selected vegetation species of Pasoh 50-ha plot (PFR) using Maximum Likelihood Classifier.	137
5.4	Classification result of CIR data for common timber groups of the dominant family Dipterocarpaceae in the Pasoh 50-ha plot (PFR) using Maximum Likelihood Classifier.	138
5.5	Classification result of IKONOS-2 data for selected vegetation species of Pasoh 50-ha plot (PFR) using Neural Network Classifier.	141



5.6	Classification result of IKONOS-2 data for common timber groups of the dominant family Dipterocarpaceae in the Pasoh 50-ha plot (PFR) using Neural Network Classifier.	142
5.7	Classification result of CIR data for selected vegetation species of Pasoh 50-ha plot (PFR) using Neural Network Classifier.	143
5.8	Classification result of CIR data for common timber groups of the dominant family Dipterocarpaceae in the Pasoh 50-ha plot (PFR) using Neural Network Classifier.	144
5.9	20 vegetation classes successfully extracted from Hyperion Image using Linear Discriminant Analysis.	148
5.10	Two dimensional example of the Spectral Angle Mapper.	149
5.11	SAM classified Hyperion images for vegetation genus classes with (a) 0.03, (b) 0.04, (c) 0.05, (d) 0.06, (e) 0.07 and (f) 0.08 spectral angles (radian).	151
5.12	Results of the unmixing analysis for endmembers (a) <i>Dacryodes</i> , (b) <i>Dipterocarpus</i> , (c) <i>Neobalanocarpus</i> , (d) <i>Shorea</i> , (e) <i>Drypetes</i> and (f) <i>Eugenia</i> extracted from Hyperion image of 50-ha plot of PFR.	155
5.13	Classification results from IKONOS-2 image showing many pixels (highlighted area) nearby cloud coverage have been wrongly classified as Shorea class and contribute to lower accuracy if compared with CIR data.	161
5.14	Classification results of Pasoh Forest Reserve 50-ha plot for mapping selected vegetation species done by (a) Maximum Likelihood Classifier and (b) Neural Network Classifier, in which more unclassified area were shown by Neural Network Classifier.	162

## LIST OF ABBREVIATIONS

APAR	-	Absorbed Photosynthetically Active Radiation
ASMS	-	Airborne Sensor Management System
BRDF	-	Bidirectional Reflectance Distribution Function
CASI	-	Compact Airborne Spectrographic Imager
CCD	-	Charge-Coupled Device
CIR	-	Colour Infrared
DBH	-	Diameter at Breast Height
DC	-	Dark Current
DMC	-	Digital Mapping Camera
DN	-	Digital Number
ETM+	-	Enhanced Thematic Mapper Plus
FOV	-	Field-of-View
FRIM	-	Forest Research Institute Malaysia
GCP	-	Ground Control Point
GMT	-	Greenwich Mean Time
GPS	-	Global Positioning System
GRABS	-	Greenness Above Bare Soil
HRV	-	High Resolution Visible
II	-	Infrared Index
IFOV	-	Instantaneous Field-of-View
IR	-	Infrared
JERS	-	Japanese Earth Resources Satellite
JPL	-	Jet Propulsion Lab
LAI	-	Leaf Area Index
LDA	-	Linear Discriminant Analysis
LMM	-	Linear Mixture Modelling
MESSR	-	Multispectral Electronic Self-Scanning Radiometer

MIR	-	Middle Infrared
MNF	-	Minimum Noise Fraction
MOS	-	Marine Observation Satellite
MSI	-	Moisture Stress Index
MSS	-	Multispectral Scanner
NASA	-	National Aeronautics and Space Administration
NIES	-	National Institute of Environmental Study
NIR	-	Near Infrared
NNC	-	Neural Network Classifier
OIF	-	Optimum Index Factor
PCA	-	Principal Components Analysis
PPI	-	Pixel Purity Index
PFR	-	Pasoh Forest Reserve
RMSE	-	Root Mean Square Error
ROI	-	Region of Interest
RWC	-	Relative Water Content
SAM	-	Spectral Angle Mapper
S/N	-	Signal-to-Noise
SAIL	-	Scattering by Arbitrarily Inclined Leaves
SAR	-	Synthetic Aperture Radar
SPOT	-	Satellite Pour l'Observation de la Terre
SWIR	-	Short-wave Infrared
TM	-	Thematic Mapper
USGS	-	United States Geological Survey
UTM	-	Universal Transverse Mercator
VNIR	-	Visible Near Infrared

**LIST OF APPENDICES**

<b>APPENDIX</b>	<b>TITLE</b>	<b>PAGE</b>
A	Spectral Matrix Plot of Common Timber Species of the Dominant Family Dipterocarpaceae of Pasoh Forest Reserve	190
B	Spectral Matrix Plot of Common Timber Groups of the Dominant Family Dipterocarpaceae of Pasoh Forest Reserve	201

## **CHAPTER 1**

### **INTRODUCTION**

#### **1.1 Introduction**

Vegetation can be defined as plant life of a region (i.e. plant community), which refers to land cover provided by plants without specific reference to any structure, spatial extent, or any other specific botanical or geographical characteristics (Jensen, 2000; Austin and Heyligers, 1991). Vegetation is broader than the term flora which refers exclusively to species composition (Laubenfels, 1975). Vegetation is known to have a strong influence on land-atmosphere interactions, and major changes in land cover, associated primarily with deforestation, have been shown to have significant impacts on the local climate (Fuentes et al., 2006). Vegetation is critical in local and global energy balances which helps in regulating the flow of numerous biogeochemical cycles, including water, carbon, and nitrogen.

Most people have an understanding that vegetation refers to forest, plantation and grassland in which these terms conjure up an image of what such vegetation looks like. However, ecologists discriminate vegetation structure at a much more detailed level, whereby the vegetation structure is determined by an interacting combination of environmental and historical factors, as well as species composition (Kabat et al., 2004). In this study, the term vegetation is thus confined to the forest vegetation that exists in the study area.

Vegetation studies over the forested area have become a great interest of ecologists and scientists due to its importance in maintaining a sustainable biological diversity in this complex ecosystem (Clark et al., 2005). Over 30 percent of the Earth's surface is covered with forests, a community of plants, animals, and many other micro-organisms with vegetation dominating the largest organism found on this planet. Out of this 30 percent, 6.4 percent is covered by tropical rainforest. The existing of tropical rainforest may not be so significant by looking at the percentage of total forests exist on Earth but it maintain a large proportion of the world's biological diversity (Thomas et al., 2004; Whitmore, 1990). Tropical rainforest has become the most notable storehouses of biological diversity on land, accumulating two-thirds of known terrestrial species and protects the largest share of threatened species. The type of forest in a given area depends on many elements, including the climate, soil, water source, rainfall patterns, seed sources and human influence. Forest can also be defined as a biological system with distinctively big numbers of interrelationships of the living part of the environment (such as plants, animals and micro-organisms) to each other and to the non-living, inorganic or abiotic parts (e.g. soil, climate, water, organic debris, rocks) (Maarel, 2004).

The tropical rain forest of Malaysia is a highly complex ecosystem which is rich and varied in plant and animal life. The forest maintains the environment stability of the country and is a store house of plant and animal species in such a way that their richness and diversity are considered as the centre of origin and diversity of many present-day as well as future crop plants (Chin and Lai, 1993).

Vegetation studies, particularly the forests inventory and management, including species richness mapping have always remained as an issue of concern by all parties over the world (Revilla, 1994). It plays an important role in the economic development of a country, generating much government revenues, especially in terms of foreign exchange earnings, such as the development of local wood-based and related industries and employment. The complex ecological relationships involving forests could allow humans to benefit from them in a variety of ways (e.g. processing the wood from trees, adopting nutrition from animals, making the forest

into a recreation park, as a medicinal source and so on) (WWF, 2005). The importance of vegetation studies are further discussed in next section.

## **1.2 Importance of Vegetation Studies**

Vegetation is one of the most important components in the biosphere. Vegetation being the primary producer and capable of photosynthesis, its growth, maintenance and development are sensitive to the environment. Therefore, it provides evidence or clues for us to understand other processes of the ecosystem, such as climate changes, carbon cycling, ecosystem evolution, and human-nature interaction. Understanding the vegetation community (including its species richness and evenness), alternations in vegetation phenological (growth) cycles, and modifications in plant physiology and morphology also provide us with valuable insight into the changing of climatic, geologic, and physiographic characteristics of an area (Jensen, 2000).

The importance of vegetation study is getting more significant with the establishment of the Kyoto Protocol. The Kyoto Protocol is a protocol to the United Nations Framework Convention on Climate Change (UNFCCC or FCCC) and an international environmental treaty produced at the United Nations Conference on Environment and Development (UNCED) which is informally known as the Earth Summit held in Rio de Janeiro, Brazil from 3–14 June 1992. During the convention, the member countries (and Malaysia is one of the member countries) who signed the treaty agreed and intended to achieve the "stabilization of greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system" (UNFCCC, 2005). The Kyoto Protocol establishes legally binding commitments for the reduction of four greenhouse gases (carbon dioxide, methane, nitrous oxide, sulfur hexafluoride). Vegetation plays an important role in reducing the emission of greenhouse gases and the circulation of carbon (Smith, 2003). Warmer global temperatures which linked directly to greenhouse gas emission may alter tree growth rates, recruitment and

mortality, thereby creating new assemblages of tree as global temperature increase and extinction of some vegetation species (Laurance et al., 2004).

Some observations made have concluded that 80 percent of the nutrients in the tropical ecosystem are in the vegetation and this should always be carefully weighed in the design/formulation of forest management system in the tropics (McGinley and Finegan, 2003; Mendoza et al., 1999; Prabhu et al., 1996). This situation is more serious in cases where there is a danger of accelerated soil erosion resulting in the loss of most nutrients in the soil, or, in the event where the original vegetation has to give way to plantations (Zhang et al., 2003 and Revilla, 1994). Tropical rainforest biodiversity is endangered by large scale (e.g., 10-500 ha) deforestation and logging activities (Achard et al., 2002). Some vegetation species are very sensitive to environmental changes. For example, wetland vegetation species can be a good indicator of the environmental changes once the anticipated biophysical parameters are determined. The importance of vegetation in biodiversity has led to the concerns for forest inventories that include type of forest inventories involving precision of estimates, accuracy of estimates, control of re-enumeration, systematic sampling and monitoring of forest change.

Malaysia is fully aware of the need for effective forest management and conservation, not only to ensure a sustained supply of timber but also to maintain environmental stability, providing sanctuary for wildlife and to serve as an invaluable storehouse of genetic resources useful for the improvement of its indigenous tree species, agricultural crops and live stocks (Razak et al., 2002; Appanah, 1999; Salleh and Musa, 1994). From individual crown to landscape scales, vegetation in tropical rainforest has a dominant role in maintaining its rich environment sustainability. Furthermore, vegetation of the tropical rainforest represents a major pool of terrestrial carbon. In this instance, remote sensing can be used as an effective tool to monitor, perform inventory mapping and detecting the sustainable environment.

The growing needs to conserve and provide accurate information on forest biophysical parameters implies the relative importance of different vegetation species,



their occurrence and distribution to be studied. Knowledge of vegetation species composition and diversity is therefore the most useful tool in evaluating overall species diversity and endemism in tropical forest, and it is critical for conservation planning. Subsequently, there is a need to formulate sound management policies and guidelines for the conservation of these species (Ashton, 1990). In order to meet this need, remote sensing technology has been widely used in vegetation and its related studies due to its advantages in wide coverage, cost effective and availability of multi temporal archived data.

### **1.3 The Use of Remote Sensing Analysis in Vegetation Studies**

According to Yamada (1997), the obstacles faced by foresters and ecologists in studying the tropical rain forest can be categorized into three broad categories, namely: 1) human factors, 2) natural factors and 3) biological factors. Among these factors, one of the main problems in studying the tropical rain forest is accessibility. This can be further explained as the understanding, monitoring and inventory mapping of tropical rain forests are being influenced by a lack of spatially and temporally extensive information on vegetation composition, species richness and structure. Due to expensive costs and inaccessibility (as mentioned earlier), most available data only comes from relatively small field plots with infrequent re-sampling intervals. With the existing plots, it is difficult to scale-up the field data to the landscape, regional or global scales needed in sustainability analysis of ecosystem in tropical rainforest (Tuomisto et al., 2003).

Since the launching of Landsat in 1972, remote sensing has become a powerful means in providing data from which updated forest cover and related information can be obtained. Passive remote sensing sensors provide multi-scale, good spatial and temporal measurement of radiance from tropical rainforest canopies which can be linked to species composition and richness mapping (Foody et al., 2006; Nagendra, 2001). However, remote sensing studies in tropical rainforest have only focused on the mapping of general forest cover classes for calculating the rate and extent of regional deforestation and forest fragmentation relied upon medium spatial

resolution imagery from multispectral spaceborne sensors due to the reasonable costs and acceptable accuracy (Roberts et al., 2002; Steininger et al., 2001). Remote sensing technology helps to resolve the accessibility problem as mentioned earlier in mapping the forest vegetation species and its related studies with reasonable costs.

In vegetation studies that employ remote sensing techniques, two widely used data are satellite and airborne colour infrared (CIR) images (Gould, 2000; Madden et al., 1999; Welch, 1996). The first and second generation satellite data (i.e. the first generation of satellite data refer to Landsat MSS and second generation of satellite data refer to Landsat TM, SPOT HRV and other multispectral sensors with broad bands spectral resolution ) mostly do not have adequate spatial resolution to differentiate between detailed ground information. The airborne CIR images were then used to complement the satellite data. Both of these data complement each other when studying large areas where airborne data can be very costly to acquire, and are mostly used when detailed studies involving smaller areas are of interest. However, the complexity of the vegetation species composition in a tropical rainforest due to high tree diversity and both natural and human disturbances results in complex radiance signals that are difficult to discriminate using broad spectral and spatial resolution sensors. In order to delineate vegetation properly, the recent high spatial and hyperspectral remote sensing data therefore meet the needs in terms of both spatial and spectral resolution.

A new generation of high spatial resolution multispectral sensor (less than 4 meters) permitted the mapping of individual tree crown at species level as a group of image pixels (Gougeon & Leckie, 2003). Such advancement in remote sensing technology could greatly improve multi-scale forest classifications of vegetation species richness, habitat and disturbance history mapping. With the temporal data, individual tree crown analyses may also provide a means to systematically monitor changes in some special valuable species (e.g. common timber species) due to logging and climate change. (Clark et al., 2004; Read et al., 2003 and Nagendra, 2001).

With the advancement in remote sensing technology, the development of sensor has a great achievement in developing hand-held, airborne and spaceborne hyperspectral optical sensors which allow the spectral measurement over 50 narrow, continuous bands spanning the visible (400 nm to 700 nm), near infrared (700-1327) and two short wave infrared (1467-1771 nm and 1994-2435 nm respectively) regions of electromagnetic spectrum (Asner, 1998). The automated vegetation classification and species identification from tropical rainforest are now possible with hyperspectral imagery which is fine enough in terms of spectral wise. However, the accuracy may vary from different spatial resolution hyperspectral data and feature extraction techniques used, which will be discussed in later chapters.

In Malaysia, the use of remote sensing techniques in mapping the tropical rain forest has been explored since 1990s (Nuraznin and Hashim, 2007; Okuda et al., 2004; Okuda et al., 2003; Hashim et al., 2002; Hashim et al, 1999; Khali et al., 1993, Radzali, et al., 1992, Sawada et al., 1991). Series of researches had been carried out by Forest Research Institute Malaysia (FRIM) with the aid of remote sensing technology to extract forest and vegetation information. Khali et al. (1993) had started a research at the North Selangor peat swamp forest to assess the forest condition after logging by using remote sensing techniques. Landsat-TM data were used and results showed that the peat swamp forest could be mapped using remote sensing analysis and the area was differentiated into different classes of damage severity and the area extent was calculated. (Khali et al, 1994). Because of the high demand for assessing and monitoring vegetation change, vegetation observation has long been an important application of remote sensing.

#### **1.4 Problem Statement**

Remote sensing is an important tool for measuring global biodiversity in forest ecosystems. Remote sensing techniques offer to deliver structural information about forest stands such as the nature of the canopy surface, the layering within the canopy and even individual tree identification (Innes and Koch, 1998). In this study, this information is being linked with ecological species information derived from

ground sampling to give estimates of species richness and distribution over much larger scales than previously available.

Moderate spatial resolution systems such as Landsat TM are able to provide valuable information on macro-level deforestation and fragmentation. It is used to make measurements of ecosystem properties and correlate these properties with 'invisible' properties such as biodiversity and species distribution that serve as key indicator of forest sustainable management. Since eight years ago, high resolution remote sensing system with a spatial resolution finer than 4 m and spectral resolution better than 10 nm are available commercially in the market. It provides detailed information and shows great potential in providing information for many applications especially in vegetation and its related studies. With the existence of the new system, the ability to detect every single tree species, canopy structure and even soil properties is now possible (Johnson, 2002; Newman et al. 1998).

However, the increase in spatial and spectral resolutions raises two common issues for effective extraction of information in either high spatial or spectral resolutions data respectively. In the case of high spatial resolution, the inherent effectiveness of spatial information have somehow ensured 'busyness' or variation in a 'class of target' hence preprocessing such as texture analysis need to be carried out prior to information extraction (Schowengerdt, 2007). On the other hand, the high spectral resolution such as found in hyperspectral data often face "Rayleigh phenomenon" and problem in selecting optimum bands from the voluminous data set. "Rayleigh phenomenon" refers to the fraction of light scattered are sensitive with the narrow bands and sometimes give false information in interpreting the target object (Borengasser et al., 2007). In selecting the optimum band for selected application, the voluminous spectral resolution data will have a lot of redundancy. A total of about 200 narrow bands may be available in hyperspectral data but only small portion of narrow wavebands are useful (for example in case of vegetation species recognition).

It appears that the advancement of high resolution system which provides high spatial and hyperspectral data is having a dilemma - as the imagery is so

detailed that individual pixels often lose their representation of the landscape. Consequently, a traditional pixel-based process becomes incapable due to insufficient consideration of the rich amount of information on spatial and spectral association. Vegetation information extraction, especially species mapping from high spatial resolution imagery and hyperspectral remote sensing analysis is one of the most important research topics in the field but not much effort has gone into this topic so far, especially in tropical rainforest of this region.

Identification of individual crown species was traditionally performed by using visual interpretation method from high spatial resolution aerial photographs which are taken from film camera. New digital forms of high spatial resolution imagery from airborne and spaceborne multispectral sensors (i.e. Digital Color Infrared Aerial photo and IKONOS-2 in this study) have stimulated the development of automated techniques for individual crown species detection, delineation and subsequent measurement of related information (Chubey et al. 2006). Automated individual crown species identification from micro to macro (which can be further explained as leaf to crown scale) algorithms have been optimized for species mapping in some tropical forests, and it is not clear how these algorithms will perform in tropical rainforest in Malaysia with high species diversity and complex canopies. Consequently, this study is undertaken to further examine the algorithms and techniques available for micro to macro level vegetation species mapping and to suggest the algorithm and processing flow which would be suitable for use in tropical rainforest region with some modifications of algorithm by selecting the suitable spectral regions which were identified and tested in this study for the purpose of species richness and forest health mapping. With all the selected algorithms and techniques examined and tested using high spatial and hyperspectral remote sensing analysis at micro level vegetation mapping, the challenge lies on how to implement the best algorithm chosen for the macro-level analysis in order to enable large scale biodiversity mapping so as to make full use of the algorithm for vegetation species mapping and its related analysis.

There are some pre-processing requirements that should be first addressed when applying the hyperspectral remote sensing analysis in vegetation species

mapping. These include the creation of a spectral library and the calibration of hyperspectral data. A spectral library is needed to compare the spectra recorded by the sensor to the spectra collected in the field work as a reference to find out the endmember (known as class in multispectral analysis). Most of the spectral libraries available created by Jet Propulsion Lab, (JPL) and United States Geological Survey, (USGS) are more focused on geological study such as mineral exploration. Recently, USGS has started developing spectral libraries for common vegetation types. In Malaysia, a complete set of hyperspectral libraries has yet to be developed. Therefore, to study Malaysian vegetations using hyperspectral data, a new set of spectral libraries is needed. These libraries should be complete because different regions of study may need a different spectral library to increase the accuracy of the study. From the spectral library created, series of analysis need to be carried out to study the separability of different species available in study area to make sure the species identification at micro level (leaf scale) is possible and further scale-up the species mapping to macro level (entire study area of this study).

With the doubts and problems discussed, it is worthwhile to carry out the study and the detailed objectives and scope of this study are defined in the next two sections.

## 1.5 Objectives of the Study

The overall aim of this study is to perform vegetation species mapping in tropical rainforest using high spatial resolution and hyperspectral remote sensing analysis. In order to achieve the overall aim, the following sets of specific objectives are formulated:

- 1 To create a spectral library for tropical rainforest and investigate the spectral variation among tropical rainforest vegetation species, thereby permitting spectral-based species discrimination;
- 2 To identify the spectral regions and spatial scale which provide optimal vegetation species discrimination by the analysis of spatial and spectral separability and variability in distinguishing vegetation species using high spatial and hyperspectral remote sensing analysis;
- 3 To examine and identify the existing techniques in vegetation species mapping of tropical rainforest and develop an analytical procedure to perform species mapping of tropical rainforest using high spatial resolution and hyperspectral remote sensing analysis;
- 4 To perform accuracy assessment on results obtained in objectives (1), (2) and (3) as compared to baseline information by using statistical approaches.

## 1.6 Scopes of the Study

The scopes of the study are as follows:

1. Vegetation species mapping in this study is confined to the identification of different types of vegetation, which means counting of the number of different type of plants in the study area by mapping the vegetation community in species, genus or family. The vegetation species mapping in this study is further confined to big families (at least 100 trees in each genus that existed in the study area with diameter at breast height, DBH > 30 cm). Trees with DBH larger than 30 cm were used so as to ensure that the top layer of the canopy can be analysed by using remotely sensed data. In addition to this, common timber species available in the study area will also be identified.
2. The high spatial resolution remotely sensed data used in the study is confined to multispectral IKONOS-2 data with 4 m spatial resolution and digital color infrared aerial photo by Z/I Imaging's Digital Mapping Camera with 20 cm spatial resolution. Hyperspectral remotely sensed data used in this study are confined to Hyperion (provides 220 continuous spectral bands ranging in wavelength from 0.4  $\mu\text{m}$  to 2.4  $\mu\text{m}$  with spatial resolution of 30 m). More details will be given in Chapter 3 on Data Acquisition, Preprocessing and Data Enhancement.
3. The study area is confined to tropical rainforest located in Pasoh Forest Reserve (PFR) 50 ha plot which contains 335,240 trees in 814 species, 290 genus and 78 families. On a per hectare basis, the species diversity of trees at Pasoh is comparable with those recorded anywhere in the world species-rich tropical rain forest similar to many seasonal forests of Malaysia. The availability and continuous census used as ground truth is the important selection factor apart from the scientific documents on tree species mapping undertaken.



## 1.7 Significance of Research

The monitoring, conservation and management of tropical rainforests has become a great challenge due to lack of spatially and temporally extensive information on tree floristic composition and vegetation species mapping (Foody et al., 2006 and Tuomisto et al., 2003). The vegetation species distribution map can be an important tool for understanding the extent and pattern of old growth forests, predicting rare plant habitat, modelling the spread of invasive species, and modelling how vegetation might change under various climate change scenarios (Clark et al., 2005). The introducing of high spatial resolution and hyperspectral remote sensing provided a good solution in vegetation species mapping which was done earlier in relatively small field plots with infrequent re-sampling intervals but involved prohibitive costs and inaccessibility problems. Thus, this study is carried out to perform vegetation species mapping of tropical rainforest using high spatial and hyperspectral remotely sensed data. Furthermore such a study have not being reported for tropical rainforest species or even at genus level due to the remoteness, dense, complex multi-storey canopies.

Spectral library of tropical rainforest species even for common timber species have not been reported to be available. Lack of this basic information, have cause major set-back for not enabling hyperspectral data to be classified optimally at full capacity to derive absolute forest information. Worst still, even the fine resolution hyperspectral airborne data is only being relatively classified due to lack of the spectral library. This study have successfully created spectral library which consists of common timber species of the dominant family Dipterocarpaceae; thereby would serve as baseline information for vegetation species identification in any tropical rainforest of this region by using hyperspectral remote sensing analysis.

This study has also placed contribution to the assessment of the spatial and spectral sensitivity of high spatial resolution and hyperspectral remotely sensed data in mapping the vegetation species in tropical rainforest. In particular whole electromagnetic spectrum from 400-1200 nm have been explored and had successfully identified the wavelengths as well as spectral band for identification of

all common timber groups of the dominant family Dipterocarpaceae. This information is very vital for any mission and design of future sensing systems including satellite for forestry and related vegetation studies in tropical rainforest.

Apart from the above, this thesis has also significantly contribute to new knowledge on understanding of the vegetation species mapping and determine the advantages and limitations of high spatial and hyperspectral remote sensing analysis in vegetation species mapping in tropical rainforest environment. This research thus, produces a series of accurate vegetation distribution maps of vegetation in genus and species level for Pasoh Forest Reserve.

## **1.8 Study Area**

The area of this study is confined to Pasoh Forest Reserve which located at 2° 55' N latitude and 102° 18' E longitude, about 8 km from the town of Simpang Pertang, Negeri Sembilan, approximately 70 km southeast of Kuala Lumpur (Figure 1.1). The Pasoh Forest Reserve comprises of low hills and alluvium rising to a granite ridge along its eastern border. A buffer zone of 700 ha within the western and southern margins was logged in the 1950s. The study area was further confined to 50-ha plot 1 km long by 0.5 km wide of Pasoh Forest Reserve. The plot was established between 1985 and 1988 for research purposes. The first detailed descriptions of the forest botany and stand structure were published in 1990 (Kochummen et al., 1990; Manokaran and LaFrankie 1990). The enumeration included all free standing trees and shrubs > 30 cm diameter-at-breast-height (DBH) excluding climbers. According to the details by which the plot was first surveyed, the 50-ha plot contains 335 240 trees in 814 species, 290 genus and 78 families. In 2000, another detailed survey was carried out and there were 338 924 trees in 818 species in 295 genus and 81 families. The number of trees has increased and the variety of the vegetation community in the 50-ha plot of Pasoh Forest Reserve is enough to represent any other tropical rain forest available within this region (Okuda et al., 2003).


The 30 species of Dipterocarpaceae family dominated the 50-ha plot, accounting for 27.3% of the total basal area (Manokaran and LaFrankie, 1990). The Euphorbiaceae with 85 species was the richest family in the plot and had the highest number of trees in the plot with 13.4% of total tree numbers. Shorea was the most important genus in the 5-ha research plot in terms of tree number (20 960 trees, 6.2 % of all trees). Shorea was the fifth most diverse genus in the plot with 14 species (1.7% of all species). As for the common timber groups, the family Dipterocarpaceae once again dominated the plot with 10 common genus. Red Meranti was the biggest genus under Shorea with a total of 13 401 trees (43.36% of total basal area) follow by Balau under same genus (shorea) with 6842 trees (22.13%) (Davies et al., 2003). More details can be obtained in Table 1.1.

The selection of the study area was based on: (1) the variety of vegetation types and (2) the availability of airborne and spaceborne high spatial and hyperspectral data over the study area. As mentioned earlier, the variety of the vegetation species that exist in the 50-ha plot of Pasoh Forest Reserve is enough to represent any other tropical rain forest available within this region and suitable to be selected as study area due to its completeness of secondary data, for example the complete census plot of the 50-ha plot of Pasoh Forest Reserve. It is not easy to acquire both high spatial resolution and hyperspectral remote sensing data for same tropical rainforest with ready and complete set of secondary data. Large number of biological researches had been carried out at Pasoh Forest Reserve since 1970s which also helps when performing the literature search.

**Table 1.1:** The most important families for 50-ha plot of Pasoh Forest Reserve.

Family	Basal Area (m <sup>2</sup> )	Family	Total number of Trees	Family	Species number
Dipterocarpaceae	453.21	Euphorbiaceae	45436	Euphorbiaceae	85
Fabaceae	141.47	Dipterocarpaceae	31178	Lauraceae	49
Euphorbiaceae	120.46	Annoaceae	24752	Myrtaceae	48
Burseraceae	100.91	Rubiaceae	20506	Rubiaceae	47
Myrtaceae	56.96	Burseraceae	17701	Annoaceae	42

*Source: Davies et al., (2003)*



**Figure 1.1:** Study area; (a) location map, (b) the raw IKONOS satellite image of the study area and (c) the detailed plan of Pasoh Forest Reserve.

## 1.9 Thesis Outline

This thesis comprises of six chapters. Chapter 1 explains the research background of the study and gives the problem statement, objectives and scope of the study. The review of the high spatial and hyperspectral remote sensing techniques, together with its applications and works previously done using remote sensing techniques for vegetation and its related studies are addressed in Chapter 2. The literature review on vegetation mapping, especially on vegetation health mapping and species richness mapping (identification of different vegetation species), are also addressed in this chapter. In Chapter 3, preparation of two sets of high spatial and hyperspectral remote sensing data and pre-processing employed in the study are discussed. Data preparation includes the collection of field and image spectra for spectral library. All the pre-processing works including radiometric calibration, geometric correction and data mosaic, data masking, and image enhancements (Minimum Noise Fraction and Pixel Purity Index), which need to be done prior to feature extraction and data classification of vegetation species mapping, are also presented in this chapter. Chapter 4 presents the flow for creating a spectral library for different vegetation species available in the study area, which will be used later in the hyperspectral data processing. The sensitivity of high spatial resolution data and hyperspectral remote sensing data used in this study were also analysed in the same chapter. The feature extraction process and data classification, which are carried out using different approaches (i.e. Band Selection Feature Extraction, Spectral Angle Mapper, and Linear Mixture Modelling for hyperspectral data) on hyperspectral data (neural network classifier and maximum likelihood for high spatial data) are discussed in Chapter 5. Results and analyses of the feature extraction and classification of the vegetation are also presented and discussed in this chapter. Conclusions of the research and recommendations for future work are given in Chapter 6.

## REFERENCES

- Aardt, J. A. N. and Wynne, R. H. (2007). Examining Pine Spectral Separability using Hyperspectral Data from an Airborne Sensor: An Extension of Field-based Results. *International Journal of Remote Sensing*. Vol. 28 (2), pp. 431-436.
- Achard, F., Eva, H. D., Stibig, H. J., Mayaux, P., Gallego, J. and Richards, T. (2002). Determination of Deforestation Rates of the World's Humid Tropical Forests. *Science*. Vol. 297(5583), pp. 999– 1002.
- Asner, G. P. (1998). Biophysical and Biochemical Sources of Variability in Canopy Reflectance. *Remote Sensing of Environment*. Vol. 64(3), pp. 234– 253.
- Appanah, S., Ismail Harun, P.F. Chong and M. Kleine (1999). *Internal Assessment Procedures for Sustainable Forest Management in Malaysia –Consultancy Report to the Forest certification Project (GTZ)*. Kuala Lumpur, Malaysia.
- Aschbacher, J., Ofren, R. S., Delsol, J. P., Suselo, T. B., Vibulsresth, S. and Charrupat, T. (1995). An Integrated Comparative Approach to Mangrove Vegetation Mapping using Remote Sensing and GIS Technologies: Preliminary Results. *Hydrobiologia*. Vol. 295, pp. 285-294.
- Ashton, P. (1990). Plant and Conservation in Tropical Asia. *International Conference on Conservation of Tropical Biodiversity in Harmony with Nature*. 12-16 June, 1990.

- ASTER Spectral Library (2008). *ASTER Digital Spectral Library*. Retrieved on 7 June 2008, from <http://asterweb.jpl.nasa.gov/>.
- Atkinson, P. M. (1993). The Effect of Spatial Resolution on the Experimental Variogram of Airborne MSS Imagery. *International Journal of Remote Sensing*, Vol. 14, pp. 1005–1011.
- Atkinson, P. M., Dunn, R., & Harrison, A. R. (1996). Measurement Error in Reflectance Data and its Implications for Regularizing the Variogram. *International Journal of Remote Sensing*. Vol. 17, pp. 3735–3750.
- Austin, M. P. and Heyligers, P. C. (1991). Vegetation Survey Design, a New Approach: Gradsect sampling. In Nature Conservation: Cost Effective Biological Surveys and Data Analysis. *Proceedings of CONCOM Workshop eds. C.R. Margules and Austin, M.P.*, pp. 31-36.
- Beaudemin, M. and Fung, K. B. (2001). On Statistical Band Selection for Image Visualisation. *Photogrammetric Engineering and Remote Sensing*. Vol. 67(5), pp. 571-574.
- Bernstein, R., (1983). Image Geometry and Rectification. *Manual of Remote Sensing*. R. N. Colwell, ed., Bethesda, MD: American Society of Photogrammetry, 1, pp. 875-881.
- Boardman J.W. (1989). Inversion of Imaging Spectrometry Data Using Singular Value Decomposition. *Proceedings, IGARSS '89, 12th Canadian Symposium on Remote Sensing*. Vol. 4, pp. 2069-2072.
- Boardman J. W., and Kruse, F. A. (1994). Automated Spectral Analysis: A Geologic Example Using AVIRIS Data. North Grapevine Mountains, Nevada. *Proceedings, Tenth Thematic Conference on Geologic Remote Sensing*. Ann Arbor, MI: Environmental Research Institute of Michigan.

- Boardman, J. W., Kruse, F. A. and Green, R. O. (1995). Mapping Target Signatures via Partial Unmixing of AVIRIS Data. In *Summaries, Fifth JPL Airborne Earth Science Workshop*. JPL Publication 95-1, Vol. 1.
- Borengasser, M., Hungate, W. S. and Watkins, R. (2007). *Hyperspectral Remote Sensing Principles and Applications*. CRC Press, Taylor & Francis Group.
- Brown, L. and G. Borstad (1999). Mapping Wetland Breeding Habitats in the Frazier River Delta, Backscatter. *Association of Marine Remote Sensing*. Vol. 10(1), pp. 8-11.
- Buckingham R., Staenz K, and Hollinger A (2002). Review of Canadian Airborne and Space Activities in Hyperspectral Remote Sensing, *Canadian Aeronautics and Space Journal*. Vol. 48, No. 1, March 2002 Vol. 48, No. 1.
- Campbell, J.B. (1996). *Introduction to Remote Sensing*. Second edition. New York: The Guildford Press.
- Castro, E., K. L., Sa´nchez-Azofeifa, G. A. and Caelli, T. (2004). Discrimination of Lianas and Trees with Leaf-level Hyperspectral Data. *Remote Sensing of Environment*. Vol. 90, pp. 353– 372.
- Center for the Study of Earth from Space (CSES) (1992). *SIPS User’s Guide, The Spectral Image Processing System*. University of Colorado, Boulder. Vol. 1.1, p. 74.
- Chen, S. S., Fang; L. G., Liu, Q. H., Chen, L. F., Tong, Q. X. (2005). The Design and Development of Spectral Library of Featured Crops of South China. *IEEE Transactions on Geoscience and Remote Sensing*. Volume 2(25-29), 4 pp.
- Chin. T. Y and Lai F.S. (1993). Challenges in Planning and Managing Forest Catchments in Peninsular Malaysia. *International Symposium on Management of Rivers for the Future*, 16-18 Nov 1993, Kuala Lumpur.



- Chubey, M. S., Franklin, S. E., & Wulder, M. A. (2006). Object-oriented Analysis of IKONOS-2 Imagery for Extraction of Forest Inventory Parameters. *Photogrammetric Engineering and Remote Sensing*. Vol. 72, pp. 383–394.
- Clark, D. B., Read, J. M., Clark, M. L., Murillo Cruz, A., Fallas Dotti, M., and Clark, D. A. (2004). Application of 1-m and 4-m Resolution Satellite Data to Studies of Tree Demography, Stand Structure and Land-use Classification in Tropical Rain Forest Landscapes. *Ecological Applications*. Vol. 14(1), pp. 61–74.
- Clark, R.N., Swayze, G.A., Wise, R., Livo, E., Hoefen, T., Kokaly, R., Sutley, S.J. (2007), USGS Digital Spectral Library SPLIB06A. *U.S. Geological Survey, Digital Data Series 231*.
- Clark, M. L., Roberts, D. A., Clark, D. B. (2005). Hyperspectral Discrimination of Tropical Rain Forest Tree Species at Leaf to Crown Scales. *Remote Sensing of Environment*. Vol. 96, pp. 375 -398.
- Clark, R. N., Swayze, G. A., Gallagher, A., King, T. V. and Calvin, W. M. (1993). The U. S. Geological Survey, Digital Spectral Library 0.2 to 3 $\mu$ m. *U. S. Geological Survey Open File Report*. pp. 93-592.
- Clark, R. N., King, T. V. V., Klejwa, M., and Swayze, G. A. (1990). High Spectral Resolution Spectroscopy of Minerals. *Journal of Geophysical Research*. Vol. 95, No. B8, pp. 12653 - 12680.
- Cleveland, William and Marylyn M. G. (1988). *Dynamic Graphics for Statistics*, Wadsworth & Brooks.
- Cochrane, M. A. (2000). Using Vegetation Reflectance Variability for Species Level Classification of Hyperspectral Data. *International Journal of Remote Sensing*, Vol. 21(10), pp. 2075– 2087.

- Conel, J. E., Green, R.O., Vane, G., Bruegge, C. J., Alley, R.E., and Curtiss, B., J., (1987). Airborne Imaging Spectrometer-2: Radiometric Spectral Characteristic and Comparison of Ways to Compensate For the Atmosphere. *Proceedings of SPIE*. Vol. 834, pp. 140-157.
- Congalton, R., K. Birch, R. Jones, and J. Schriever (2002). Evaluating Remotely Sensed Techniques for Mapping Riparian Vegetation. *Computers and Electronics in Agriculture*. Vol. 37. pp. 113-126.
- de Kok R., Schneider T., & Ammer U. (1999). Object-oriented Classification and Applications in the Alpine Forest Environment. *International Archives of Photogrammetry and Remote Sensing*. Vol. 32, Part 7-4-3 W6, 3-4 June 1999, Valladolid, Spain.
- Danson, F. M. (1998). Teaching the Physical Principles of Vegetation Canopy Reflectance using SAIL Model. *Photogrammetric Engineering & Remote Sensing*. Vol 64(8), pp. 8010-812.
- Daughtry, C.S.T. (2001). Discriminating Crop Residues from Soil by Short-Wave Infrared Reflectance. *Agronomy Journal*. Vol. 93, pp. 125-131.
- Davies S. J., Supardi, N Md. N, LaFrankie J. V. and Ashton P. S. (2003). The Trees of Pasoh Forest: Stand Structure and Floristic Composition of the 50-ha Forest Research Plot, Pasoh. *Ecology of Lowland Rain Forest in Southeast Asia*. Springer pp. 35-50.
- Debinski, D. M., and Humphrey, P. S. (1997). An Integrated Approach to Biological Diversity Assessment. *Natural Areas Journal*. Vol. 17, pp. 355-365.
- Duchene, J. and Leclercq, S., (1988). An Optimal Transformation for Discriminant and Principal Component Analysis. *IEEE Transactions on Pattern Analysis and Machine Intelligence*. Vol. 10(6), pp. 978-983.

- EO-1-NASA Web (2008). *Earth Observation Mission 1*. Retrieved on 15 August, 2008, from <http://eo1.gsfc.nasa.gov/>.
- ESA Earthnet (2007). *IKONOS-22 Third Party Mission*. Retrieved on 17 March, 2007, from <http://earth.esa.int/object/index.cfm?fobjectid=5097>.
- Foody, G.M. and Cutler, M.E.J. (2006). Mapping the Species Richness and Composition of Tropical Forests from Remotely Sensed Data with Neural Networks. *Ecological Modelling*. Vol. 195, pp. 37-42.
- Forsyth, F., G. Borstad, W. Horniak, and L. Brown (1998). Prince Rupert Intertidal Habitat Inventory Project. *Unpublished Report to Prince Rupert Port Corporation, the Canadian Department of Fisheries and Oceans, and the City of Prince Rupert*.
- Franklin, S. E. (1994). Discrimination of Subalpine Forest Species and Canopy Density Using Digital CASI, SPOT PLA, and Landsat TM Data. *Photogrammetric Engineering and Remote Sensing*. Vol. 60, pp. 1233–1241.
- Frohn, R. C. (1998). *Remote Sensing for Landscape Ecology*. Boca Raton, FL: Lewis Publishers, 99 pp.
- Fuentes, D. A., Gamon, J. A., Cheng, Y. F., Helen C., Claudio, Qiu, H. L., Mao, Z. Y. Sims, D. A., Rahman, A. F., Oechel, W., Luo, H. Y. (2006). Mapping Carbon and Water Vapor Fluxes in a Chaparral Ecosystem Using Vegetation Indices Derived from AVIRIS. *Remote Sensing of Environment*. Volume 103, Issue 3, pp. 312-323.
- Fung, T., Ma, F. Y. and Siu, W. L. (1998). Hyperspectral Data Analysis for Subtropical Tree Species Recognition. *Symposium Proceedings, IGARSS '98, Sensing and Managing the Environment*, Vol. 3, pp. 1298– 1300.

- Gat, N., Subramanian, S., Barhen, J., Toomarian, N. (1997). Spectral Imaging Applications: Remote Sensing, Environmental Monitoring, Medicine, Military Operations, Factory Automation, and Manufacturing. *Proceedings of SPIE*. Vol. 2962, pp. 63-77.
- Gates, D. M., Keegan, J. J., Schleter, J. C. and Weidner, V. R. (1965), Spectral Properties of Plants, *Applied Optics*. Vol. 4(1), pp. 11-20.
- Gausman, H. W. (1985). Plant Leaf Optical Properties in Visible and Nearinfrared Light. Lubbock, Texas' Texas Tech Press.
- Gaussmann, H. W., Allen, W. A. and Cardenas, R. (1969). Reflectance of Cotton Leaves and their Structure. *Remote Sensing of Environment*. Vol. 1, pp. 110-22.
- Goel, N. S. (1988). Modes of Vegetation Canopy Reflectance and their Use in Estimation of Biophysical Parameters from Reflectance Data. *Remote Sensing Review*. Vol. 4, pp. 1-212.
- Goetz, S. J., Wright, R., Smith, A. J., Zinecker, E., & Schaub, E. (2003). IKONOS Imagery for Resource Management: Tree Cover, Impervious Surfaces and Riparian Buffer Analyses in the Mid-Atlantic Region. *Remote Sensing of Environment*. Vol. 88, pp. 195–208.
- Goetz, A. F. H., and Srivastava, V. (1985). Mineralogical Mapping in the Cuprite Mining District, Nevada. *Proceedings of the Airborne Imaging Spectrometer Data Analysis Workshop*. JPL Publication 85-41, Jet Propulsion Laboratory, Pasadena, CA, pp. 22-29.
- Gong P., Pu, R. and Yu, B. (1997). Conifer Species Recognition: An Exploratory Analysis of In Situ Hyperspectral Data. *Remote Sensing of Environment*. Vol. 62, pp. 189-200

- Gougeon, F. A., & Leckie, D. G. (2003). Forest Information Extraction from High Spatial Resolution Images using an Individual Tree Crown Approach. *Information Report BC-X-396*. Victoria, British Columbia, Canadian Pacific Forestry Centre, Canadian Forest Service.
- Gould, W. (2000). Remote Sensing of Vegetation, Plant Species Richness, and Regional Biodiversity Hotspots. *Ecological Applications*, Vol. 10, No. 6, pp. 1861-187
- Goward, S. N., Huemmrich, K. F. and Waring, R. H. (1994). Visible-near Infrared Spectral Reflectance of Landscape Components in Western Oregon. *Remote Sensing of Environment*. Vol. 47, pp. 190-203.
- Grant, L. (1987). Diffuse and Specular Characteristics of Leaf Reflectance. *Remote Sensing of Environment*. Vol. 22, pp. 309– 322.
- Green, A. A., Berman, M., Switzer, P, and Craig, M. D. (1988). A Transformation for Ordering Multispectral Data in terms of Image Quality with Implications for Noise Removal. *IEEE Transactions on Geoscience and Remote Sensing*. Vol. 26, No. 1, pp. 65-74.
- Green E.P., C.D. Clark, A.J. Edwards and A.C. Ellis (1998). Remote Sensing Techniques for Mangrove Mapping. *International Journal of Remote Sensing*. Vol. 19, No. 5, pp. 935-956.
- Gregory, P. A., Martin, R. E. (2008). Spectral and Chemical Analysis of Tropical Forests: Scaling from Leaf to Canopy Levels. *Remote Sensing of Environment*. Volume 112, Issue 10, pp. 3958-3970.
- Guerric, M., François, C., Soudani, K., Berveiller, D., Pontailier, J., Bréda, N., Genet, H., Davi, H., Dufrêne, E. (2008). Calibration and Validation of Hyperspectral Indices for the Estimation of Broadleaved Forest Leaf Chlorophyll Content, Leaf Mass per Area, Leaf Area Index and Leaf Canopy Biomass. *Remote Sensing of Environment*. Volume 112, Issue 10, pp. 3846-3864.

- Hashim, M., Yoshida, K., Adachi, N., Okuda, T., and Bonkik, M. (2002). Applying the Geographical Information System to Evaluate Land Use Options: Oil Palm Plantation versus Tropical Forests in the Pasoh Forest Region. *Annual Meeting of NIES-FFPRI-FRIM Joint Research Project- Toward the Sustainable Management of Tropical Rainforest*. January 2002. Tsukuba, Japan.
- Hashim, M., Wan Kadir, W. H. and Lee, K. Y. (1999). Global Rainforest Mapping Activities in Malaysia (GRFM): Radar Remote Sensing for Forest Survey and Biomass Indicators. *Principal Investigator Report for Contract Research in GRFM*. Remote Sensing Centre of Japan, National Aerospace Administration of Japan (NASDA).
- Hay, G. J., Castilla, G., Wulder, M., & Ruiz, J. R. (2005). An Automated Object Oriented Approach for the Multiscale Image Segmentation of Forest Scenes. *International Journal of Applied Earth Observation and Geoinformation*. Vol. 7, pp. 339–359.
- Heiden, U., Segl, K., Roessner, S. and Kaufmann, H. (2007). Determination of Robust Spectral Features for Identification of Urban Surface Materials in Hyperspectral Remote Sensing Data, *Remote Sensing of Environment*. Vol. 111, pp. 537-552
- Huete, A. and Justice, C. (1999). Modis Vegetation Index (MOD 13) Algorithm Theoretical Basis Documents, Greenbelt: NASA Goddard Space Center. Retrieved from <http://modarch.gsfc.nasa.gov/MODIS/LAND/#vegetation-indicies>, 129 pp.
- Innes, J. L. and Koch, B. (1998). Forest Biodiversity and its Assessment by Remote Sensing. *Global Ecology and Biogeography*. Vol 7(6), pp. 397-419.
- Jacquemoud, S., Baret, F., Andrieu, B., Danson, F. M. And Jagard, K. W. (1995). Extraction of Vegetation Biophysical Parameters by Inversion of PROSPECT+ SAIL Models on Sugar Beet Canopy Reflectance Data: Application to TM and AVIRIS Sensors. *Remote Sensing of Environment*. Vol. 52, pp. 163-172.

- Jensen, J.R. (2000). *Remote Sensing of the Environment: An Earth Resource Perspective*. (First Edition). United States of America: Prentice Hall.
- Jensen J. R. (1999). *Introductory Digital Image Processing: A Remote Sensing Perspective*. Second Edition. United States of America: Prentice Hall.
- Johansen, K., Coops, N. C., Gergel, S. E., Stange, Y. (2007). Application of High Spatial Resolution Satellite Imagery for Riparian and Forest Ecosystem Classification. *Remote Sensing of Environment*. Vol. 110 (2007) pp.29-44
- Johansen, K., and Phinn, S. (2006). Linking Riparian Vegetation Spatial Structure in Australian Tropical Savannas to Ecosystem Health Indicators: Semivariogram Analysis of High Spatial Resolution Satellite Imagery. *Canadian Journal of Remote Sensing*. Vol. 32, pp. 228–243.
- John C. Russ (1995). *The Image Processing Handbook*. Second Edition, CRC Press.
- Johnson, J.V., Greenfield, P., Ellenwood, J. (2002). Using IKONOS Satellite Imagery for Forest Pest Mapping. *Proceedings of Forest Service Remote Sensing Conference*. April 9-13, 2002. San Diego, Canada.
- Jones, K. B., Ritters, K. H., Wickham, J. D., Tankersley, R. D. O'Neill, R.V., Chaloud, D. J., Smith, E. R. and Neale, A. C. (1998). *An Ecological Assessment of the United States: Mid-Atlantic Region*. Washington: EPA, 103 pp.
- JPL Spectral Library (2008). *JPL Digital Spectral Library*. Retrieved on 24 June 2008 from <http://speclib.jpl.nasa.gov/>.
- Jusoff, K. (2007). Mapping Bamboo in Berangkat Forest Reserve, Kelantan, Malaysia using Airborne Hyperspectral Imaging Sensor. *International Journal of Energy and Environment*. Vol. 1, pp. 1-6.

- Kabat, P., et al. (editors) (2004). *Vegetation, Water, Humans and the Climate: A New Perspective on an Interactive System*. Heidelberg: Springer-Verlag.
- Kalyanmoy D. (2001). *Multi-objective Optimization using Evolutionary Algorithms*. John Wiley and Son Ltd., England.
- Keshava, N. and Mustard, J.F. (2002). Spectral Unmixing. *IEEE Signal Processing*. Vol. 19(1), pp. 44 – 57
- Kelly, L. L. and Gregory, A. C. (2008). The Use of Hyperspectral Remote Sensing to Assess Vascular Plant Species Richness on Horn Island, Mississippi. *Remote Sensing of Environment*. Vol. 112, Issue 10, pp. 3908-3915.
- Key, T., Warner, T. A., McGraw, J. B. and Fajvan, M. A. (2001). A Comparison of Multispectral and Multitemporal Information in High Spatial Resolution Imagery for Classification of Individual Tree Species in a Temperate Hardwood Forest. *Remote Sensing of Environment*. Vol. 75(1), pp. 100– 112.
- Khali, A., Azman, H. H. and Chong, P.F. (1993). Monitoring of Mangrove Forest using High Resolution Satellite Imageries. In: Nyssonen., A., S. Poso and J. Rautala, (editors). *Proceedings of Iivessalo Symposium on Natural Forest Inventories*. Department of Forest Resource Management, University of Helsinki, Finland. pp 229-237.
- Khali A., Rashid, M. F., Hassan, A. and Chong, P. F., (1994). The use of Digital Remote Sensing Data for Determining Peat Swamp Forest Disturbance: A Case Study in North Selangor Peat Swamp Forest. *Proceedings of ASEAN Institute of Forest Management International Conference*. 21-24 November 1994. Kuala Lumpur, pp. 67-74.
- Kochummen, K. M., LaFrankie, J. V. and Manokaran, N. (1990). Floristic Composition of Pasoh Forest Reserve, a Lowland Rainforest in Peninsular Malaysia. *Journal of Tropical Science*. Vol. 3, pp. 1-13.



- Kondratyev, K. Y. (1998). *Multidimensional Global Change*. Chichester: Wiley/ PRAXIS Series in Remote Sensing.
- Kruse, F. A., Kierein-Young, K. S., and Boardman, J. W. (1990). Mineral Mapping at Cuprite, Nevada with a 63 Channel Imaging Spectrometer. *Photogrammetric Engineering and Remote Sensing*. Vol. 56, No. 1, pp. 83-92.
- Kruse, F. A., A. B. Lefkoff, J. B. Boardman, K. B. Heidebrecht, A. T. Shapiro, P. J. Barloon and A. F. H. Goetz, 1993. The Spectral Image Processing System (SIPS) - Interactive Visualization and Analysis of Imaging Spectrometer Data. *Remote Sensing of the Environment*. Vol. 44, pp. 145 - 163.
- Landis, J.R. and Koch, G. G. (1977). The Measurement of Observer Agreement for Categorical Data. *Biometrics*. Vol. 33, pp. 159-174
- Lau, A. M. S. (2004). *Pre-processing and Classification of Hyperspectral Data for Wetland Mapping*. Master of Science Thesis. Universiti Teknologi Malaysia, Skudai.
- Laubenfels, D.J. de. (1975). *Mapping the World's Vegetation: Regionalization of Formations and Flora*. Syracuse N.Y.: Syracuse University Press, pp. 46-74.
- Laurance, W. F. (2004). Forest-climate Interactions in Fragmented Tropical Landscapes. *Philosophical Transactions of the Royal Society, London*. Vol. B 359, pp. 345-352.
- Lewis, H.G. and Brown, M. (2001). A Generalized Confusion Matrix for Assessing Area Estimates from Remotely-sensed Data. *International Journal of Remote Sensing*. Vol. 22 (16), pp. 3223-35
- Li, X. and Strahler, A. H. (1985). Geometric-optical Modeling of a Conifer Forest Canopy. *IEEE Trans. Geoscience Remote Sensing*. GE-23(5), pp. 705-721.

- Li, X. and Strahler, A. H. (1992). Geometric-optical Bidirectional Reflectance Modeling of the Discrete-Crown Vegetation Canopy: Effect of Crown Shape and Mutual Shadowing. *IEEE Trans. Geoscience Remote Sensing*. Vol. 30, pp. 276-292.
- Liang, S. and Strahler, A. H. (1993). An Analytic BRDF model of Canopy Radiative Transfer and its Inversion. *IEEE Transaction of Geoscience Remote Sensing*. Vol. 31(5), pp. 1081-1092.
- Lillesand, T.M. and Kiefer, R.W. (2000). *Remote Sensing and Image Interpretation*. Fourth Edition. New York: John Wiley and Son.
- Lubchenco, J., Olson, A. M., Brubaker, L. B., Carpenter, S. R., Holland, M. M., Hubell, S. P., Levin, S. A., MacMahon, J. A., Matson, P. A., Melillo, J. M., Mooney, H. A., Peterson, C. H., Pullam, H. R., and Risser, P. G. (1991). The Sustainable Biosphere Initiative: an Ecological Research Agenda. *Ecology*. Vol. 72, pp. 371–412.
- Maarel, V. D. E. (2004). *Vegetation Ecology*. Oxford: Blackwell Publishers, pp. 68-135.
- McGinley, K. and Finegan, B. (2003). The Ecological Sustainability of Tropical Forest Management: Evaluation of the National Forest Management Standards of Costa Rica and Nicaragua, with Emphasis on the Need for Adaptive Management. *Forest Policy and Economics*. Vol. 5 (2003), pp. 421–431
- Madden, M., D. Jones, and L. Vilchek (1999). Photointerpretation Key for the Everglades Vegetation Classification System. *Photogrammetric Engineering and Remote Sensing*. 65(2), p. 171-177.
- Manokaran, N., and LaFrankie, J. V. (1990). Stand Structure of Pasoh Forest Reserve, a Lowland Rain Forest in Peninsular Malaysia. *Journal of Tropical Forest Science*. Vol. 3, pp. 14–24.

- Martin, M.E., Plourde, L.C., Ollinger, S.V., Smith, M.-L., McNeil, B.E. (2008). A Generalizable Method for Remote Sensing of Canopy Nitrogen across a Wide Range of Forest Ecosystems. *Remote Sensing of Environment*. Volume 112, Issue 9, pp. 3511-3519.
- Mausel, P. W., Kramberm W. J. and Lee, J. K. (1990). Optimum Band Selection for Supervised Classification of Multispectral Data. *Photogrammetric Engineering and Remote Sensing*. Vol. 56(1), pp. 55-60.
- McKeown, D. (1988). Building Knowledge-Based Systems for Detecting Manmade Structures from Remotely Sensed Imagery. *Philosophical Transactions of the Royal Society of London, Series A: Mathematical and Physical Sciences*. Vol. 324(1579), pp. 423–435.
- Miller, J.R., J. Freemantle, M.J. Belanger, C.D. Elvidge and M.G. Boyer (1991). Potential for Determination of Leaf Chlorophyll Content Using AVIRIS. *Proceedings of the Second Airborne Visible/Infrared Imaging Spectrometer (AVIRIS) Workshop*. Pasadena, Calif. USA, 4-8 June, 1990. pp. 72-77.
- Miller, J.R., and O'Neill, N.T. (1997). Multi-Altitude Airborne Observations of the Insolation Effects of Forest Fire Smoke Aerosols at BOREAS: Estimates of Aerosol Optical Parameters. *Journal of Geophysics Resources*. Vol. 102, No. D24, pp. 729–736.
- Myers, B. J. and Benson, M. L. (1981). Rainforest Species on Large-scale Colour Photos. *Photogrammetric Engineering and Remote Sensing*, Vol. 47, pp. 505–513.
- Nagendra, H. (2001). Using Remote Sensing to Assess Biodiversity. *International Journal of Remote Sensing*. Vol. 22, No. 12, pp. 2377–2400.

- Newman, R. M., Thompson, D. C., and Richman, D. B. (1998). Conservation Strategies for the Biological Control of Weeds. *Conservation Biological Control*. Academic Press, New York, NY, 371-396.
- Newstrom, L. E., Frankie, G. W., Baker, H. G. and Colwell, R. K. (1994). Diversity of Long-term Flowering Patterns. *Ecology and Natural History of A Neotropical Rain Forest*. pp. 142– 160.
- Nuraznin, A. and Hashim, M. (2007). Mapping of Urban above-ground Biomass with High Resolution Remote Sensing Data. Proceedings of 28<sup>th</sup> Asian Conference on Remote Sensing (ACRS). 12-16 November. Kuala Lumpur, 7pp.
- Okada, T. and Tomita, S., (1985). An Optimal Orthonormal System for Discriminant-Analysis. *Pattern Recognition*. Vol.18(2), pp. 139-144.
- Okuda, T., Hashim, M., Suzuki, M., Adachi, N., Yoshida, K., Niiyama, K., Nur Supardi, M. N. and Manokaran, N.(eds.) (2003). Logging History and Its Impact on Forest Structure and Species Composition in the Pasoh Forest Reserve-Implication for the Sustainable Management of Natural Resources and Landscapes. *Ecology and Natural History of a Southeast Asian Tropical Rainforest*. New York: Springer.
- Okuda, T., Yoshida, K., Numata, S., Nishimura, S. and Hashim, M (2004). Estimation of Above Ground Biomass in Logged and Primary Lowland Tropical Rainforests using 3-D Photogrammetric Analysis. *Journal of Forest Ecology and Management*. Volume 203, pp. 63-75.
- Peterson, D. L. and Running, S. W. (1989). Applications in Forest Science and Management. *Theory and Applications of Optical Remote Sensing*. Doghouses Asrar, New York: John Wiley & Sons. pp. 4210-473.
- Plaza, A., Martinez, P., Perez, R. & Plaza, J. (2002). Spatial/Spectral Endmember Extraction by Multidimensional Morphological Operations. *IEEE Transactions on Geoscience and Remote Sensing*. Vol. 40 (9), pp. 2025 – 2041.

- Prabhu, R., Colfer, C.J.P., Venkateswarlu, P., Tan, L.C., Soekmadi, R., Wollenberg, E. (1996). *Testing Criteria and Indicators for the Sustainable Management of Forest, Phase 1* (Final Report). Bogor, Indonesia: CIFOR Special Publications.
- Razak, A., M.A., Woon, W-C. & Lim, H-F (2002). Challenges in Implementing Forestry-related Policies in Malaysia. *International Workshop on Forest Science and Forest Policy in the Asia Pacific Region: Building Bridges to Sustainable Future*. 16–19 July, 2002, Chennai, India.
- Radzali, M.M., Ismail, A.B. and Razak, A. A. (1992). Application of Remote Sensing Technique for Peatland Characterization and Mapping. In: Tan, S.L., B.Aziz, J. Samy, Z. Salmah, H. Siti Petimah and S.T. Choo (eds.). *Proceedings of the International Symposium on Tropical Peatland*. Kuching, Sarawak. pp 101-106.
- Ravan, S. A., Roy, P. S., and Sharma, C. M. (1995). Space Remote Sensing for Spatial Vegetation Characterization. *Journal of Bioscience*. Vol. 20, 427–438.
- Read, J. M., Clark, D. B., Venticinque, E. M., and Moreira, M. P. (2003). Application of Merged 1-m and 4-m Resolution Satellite Data to Research and Management in Tropical Forests. *Journal of Applied Ecology*. Vol. 20, pp. 592–600.
- Research Systems Inc (2000). *ENVI Tutorials Version 3.4*. September, 2000 Edition.
- Revilla (1994). Issues on the Inventory and Management of Tropical Forests: Asia Pacific Region. *Proceedings of ASEAN Institute of Forest Management International Conference*. 21-24 November 1994. Kuala Lumpur, pp. 233-236.
- Richards, J.A. (1999). *Remote Sensing Digital Image Processing: An Introduction*. Berlin: Springer-Verlag.

- Ritter, R. and E. L. Lanzer (1997). Remote Sensing of Nearshore Vegetation in Washington State's Puget Sound. *Proceedings of the 1997 Geospatial Conference*. Seattle, WA, Vol.3 pp. 527-539.
- Roberts, D. A., Gardner, M., Church, R., Ustin, S., Scheer, G. and Green, R. O. (1998). Mapping Chaparral in the Santa Monica Mountains using Multiple Endmember Spectral Mixture Models. *Remote Sensing of Environment*. Vol. 65(3), pp. 267– 279.
- Roberts, D. A., Ustin, S. L., Ogunjemiyo, S., Greenberg, J., Dobrowski, S. Z. and Chen, J. (2004). Spectral and Structural Measures of Northwest Forest Vegetation at Leaf to Landscape Scales. *Ecosystems*. Vol. 7(5), pp. 545–562.
- Roberts, D. A., Numata, I., Holmes, K., Batista, G., Krug, T. and Moteiro, A. (2002). Large Area Mapping of Land-Cover Change in Rondonia Using Multitemporal Spectral Mixture Analysis and Decision Tree Classifiers. *Journal of Geophysical Research*. Vol. 107(20), pp. 8073.
- Roberts, D. A., Yamaguchi, Y., and Lyon, R. J. P. (1985). Calibration of Airborne Imaging Spectrometer Data to Percent of Reflectance Using Field Measurements. *Proceedings of 19<sup>th</sup> International Symposium on Remote Sensing of Environment*, Ann Arbor, MI, October 21-25, 1985.
- Salleh, M. M. and Musa, S. (1994). Issues on the Inventory and Management of Tropical Forests in Malaysia. *Proceedings of ASEAN Institute of Forest Management International Conference*. 21-24 November 1994. Kuala Lumpur, pp. 134-146.
- Sawada, H., H., Khali, A., Nakakita, O., Takao, G., Awaya, Y. and Azman, H. (1991). Development of Monitoring System for Tropical Rain Forest Environment in Peninsular Malaysia. *Technical Report of Joint Research on the Enhancement and Application of the Remote Sensing Technology with ASEAN countries*. Science and Technology Agency of Japan. pp 97-118.

- Schowengerdt R. A., (2007). *Remote Sensing Models and Methods for Image Processing*. Academic Press, United States of America, Third Edition, pp. 16-30, 77-81.
- Shafri., H. Z. M, Suhaili, A., and Mansor, S. (2007). The Performance of Maximum Likelihood, Spectral Angle Mapper, Neural Network and Decision Tree Classifiers in Hyperspectral Image Analysis. *Journal of Computer Science*. Vol. 3 (6), pp. 419-423.
- Shakoor, A. (2003). Subpixel Classification of Ground Surface Features. *GIS Development*. Vol. 7(11), pp. 20–24
- Skov F. and Svenning J. S. (2003). Predicting Plant Species Richness in a Managed Forest. *Forest Ecology and Management*. Vol. 180, pp. 583–593.
- Smith, S. J. (2003). The Evaluation of Greenhouse Gas Indices. *Climatic Change*. Vol. 58 (3), pp. 261-265
- Staenz K., Neville R., J. Lévesque, T. Szeredi, V. Singhroy, G. Borstad, P. Hauff. (1999). Evaluation of CASI and SFSI Hyperspectral Data for Environmental and Geological Applications - Two Case Studies. *Canadian Journal of Remote Sensing*, Vol. 25, No. 3, 1999, pp. 311-322.
- Steininger, M. K., Tucker, C. J., Townshend, J. R. G., Killeen, T. J., Desch, A. and Bell, V. (2001). Tropical Deforestation in the Bolivian Amazon. *Environmental Conservation*. Vol. 28(2), pp. 127–134.
- Stoms, D. M., and Estes, J. E. (1993). A Remote Sensing Research Agenda for Mapping and Monitoring Biodiversity. *International Journal of Remote Sensing*. Vol. 14, pp. 1839–1860.

- Suhaili, A., Shafri., H. Z. M, Ainuddin, N. A., Noor, A. G. A. and Hanum, I. F. (2006). Improving Species Spectral Discrimination using Derivatives Spectra for Mapping of Tropical Forest from Airborne Hyperspectral Imagery. *Proceeding of Map Malaysia 2006*.
- Theiler J. and Lavenier D. (2000). FPGA Implementation of the Pixel Purity Index Algorithm for Hyperspectral Images. *Tech Report*. LA-UR-00-2426, Los Alamos National Laboratory.
- Thenkabail, P. S., Enclona, E. A., Ashton, M. S., Legg, C. (2004). Hyperion, IKONOS, ALI, and ETM+ Sensors in the Study of African Rainforests. *Remote Sensing of Environment*. Vol. 90, pp. 23–43.
- Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., and Collingham, Y. C. (2004). Extinction Risk from Climate Change. *Nature*, Vol. 427(6970), pp. 145– 148.
- Tuomisto, H., Poulsen, A. D., Ruokolainen, K., Moran, R. C., Quintana, C. and Celi, J. (2003). Linking Floristic Patterns with Soil Heterogeneity and Satellite Imagery in Ecuadorian Amazonia. *Ecological Applications*. Vol 13(2), pp. 352–371.
- UNFCCC (2005). *Kyoto Protocol*. Retrieved on 17 August, 2008 from [http://unfccc.int/kyoto\\_protocol/items/2830.php](http://unfccc.int/kyoto_protocol/items/2830.php)
- USGS Spectral Library (2008). *USGS Digital Spectral Library*. Retrieved on 24 June, 2008, from <http://speclab.cr.usgs.gov/>.
- Vaughan, R. G., Hook, S. J., Calvin, W. M., Taranik, J. V. (2005). Surface Mineral Mapping at Steamboat Springs, Nevada, USA, with Multi-Wavelength Thermal Infrared Images. *Remote Sensing of Environment*. Volume 99, Issues 1-2, pp. 140-158.



- Verhoer, W. (1984). Light Scattering by Leaf Layers with Application to Canopy Reflectance Modeling: The SAIL Model. *Remote Sensing of Environment*. Vol. 16, pp. 125-141.
- Wang, L., Wayne, P., Sousa, P.G. and Gregory, S.B. (2004). Comparison of IKONOS and QuickBird Images for Mapping Mangrove Species on the Caribbean Coast of Panama. *Remote Sensing of Environment*. Vol. 91(3-4), pp. 432-440.
- Wang, J., L. Zhang, and Q. Tong (1998). The Derivative Matching for Wetland Vegetation Identification and Classification by Hyperspectral Data. *Proceedings of SPIE, Hyperspectral Remote Sensing Application*. Vol. 3502 p. 280-288.
- Way, J., Paris, J., Kasischke, E., Slaughter, C., Viereck, L., Christensen, N., Dobson, M. C., Simonett, D., Hoffer, R. and Weber, J. (1990). The Effect of Changing Environmental Conditions on Microwave Signatures of Forest Ecosystems: Preliminary Results of the March 1988 Alaskan Aircraft SAR Experiment. *International Journal of Remote Sensing*. Vol. 11 Issue 7 1990, pp. 1119-1144.
- Welch, R. (1996). GPS, Images Processing and GIS Techniques for Coastal Wetland Mapping Applications. *International Archives of Photogrammetry and Remote Sensing*. 31 (B4), p.931-933.
- Whitmore, T. C. (1990). *An Introduction to Tropical Rain Forests*. Oxford Iarendon Press.
- Williams, D. L. (1991). A Comparison of Spectral Reflectance Properties at the Needle, Branch, and Canopy Level for Selected Conifer Species. *Remote Sensing of Environment*. Vol. 35, pp. 79-91.
- Wilson, E. O. (1988). *Biodiversity*. Washington, D.C.: National Academy Press.

- Woodcock, C. E., Collins, J. B., Jakabhazy, V. D., Li, X., Macomber, S. A. and Wu, Y. (1997). Inversion of the Li-Strahler Canopy Reflectance Model for Mapping Forest Structure. *IEEE Transactions on Geoscience and Remote Sensing*. GE-35(2), pp. 405-414.
- Woolley, J. T. (1971). Reflectance and Transmittance of Light by Leaves. *Plant Physiology*, Vol. 47, pp. 656– 662.
- Wulder, M. A., Hall, R. J., Coops, N., & Franklin, S. (2004). High Spatial Resolution Remotely Sensed Data for Ecosystem Characterization. *Bioscience*. Vol. 54, pp.511–521.
- Wulder, M. A., LeDrew, E. L., Franklin, S. E., & Lavigne, M. B. (1998). Aerial Image Texture Information in the Estimation of Northern Deciduous and Mixed Wood Forest Leaf Area Index (LAI). *Remote Sensing of Environment*. Vol. 64, pp. 64–76.
- WWF, World Wide Fund for Nature Web (2005). *Forest Ecosystem*. Retrieved on 12 December, 2005 from [http://www.panda.org/about\\_wwf/what\\_we\\_do/forests/about\\_forests/](http://www.panda.org/about_wwf/what_we_do/forests/about_forests/).
- Yamada I. (1997). *Tropical Rain Forest of Southeast Asia: A Forest Ecologist's View*, Honolulu: University of Hawaii Press, pp 292-299.
- Yu, B., Ostland, I.M., Gong, P. and Pu, R.L., (1999). Penalized Discriminant Analysis of in Situ Hyperspectral Data for Conifer Species Recognition. *IEEE Transactions On Geoscience And Remote Sensing*. Vol. 37(5), pp. 2569-2577.
- Zhang, Y., Zhang H., Zhuo, P. B. and Yang, H. (2003). Soil Erosion and its Impacts on Environment in Yixing Tea Plantation of Jiangsu Province. *Chinese Geographical Science*, Volume 13, Number 2. Science Press, pp. 43-57.