

PROJECTIONS OF FUTURE EXTREME RAINFALL EVENTS
USING STATISTICAL DOWNSCALING IN MALAYSIA

SYAFRINA ABDUL HALIM

UNIVERSITI TEKNOLOGI MALAYSIA

PROJECTIONS OF FUTURE EXTREME RAINFALL EVENTS USING
STATISTICAL DOWNSCALING IN MALAYSIA

SYAFRINA ABDUL HALIM

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (*Generic*)

Razak School of Engineering and Advanced Technology
Universiti Teknologi Malaysia

FEBRUARY 2016

Specially dedicated to my beloved parents *Mama and Abah*, my beloved sister, *Syazana Abdul Halim* and brothers *Abdul Shahnaz, Abdul Halim and Abdul Shakin Abdul Halim*, my beloved husband, *Mohd Noreffendy Jayah* and my adorable kids *Umar, Ulya and Zara*.

ACKNOWLEDGEMENT

This thesis has been compiled with contribution from many people whom I wish to acknowledge for their dedication. I wish to express my sincere appreciation to my supervisor, Assoc. Prof. Dr. Zalina Mohd Daud for her continuous guidance, motivation and support during my study. She has been so understanding and has been a tremendous mentor for me. I would like to thank her for encouraging my research and for allowing me to grow as a research scientist. Her advice on both research as well as on my future career have been priceless. I would also like to thank my co-Supervisor, Dr Liew Juneng, who has also helped me a lot in MATLAB programming. His views and tips are useful indeed.

A special thanks also goes to my family. Words cannot express how grateful I am to my mother, Pn Robiah Hassan, father, Abdul Halim A. Karim, my mother-in law, Hjh. Sarah Ayob and father-in law, Hj. Jayah Said for all of the sacrifices that they have made on my behalf. Their prayers for me were what sustained me thus far. I would also like to thank all of my friends who supported me in writing, and inspired me to strive towards my goal. Their contribution in this thesis whether direct or indirectly has helped greatly. Last but not least, I would like to express my appreciation to my beloved husband, Mohd Noreffendy Jayah who spent sleepless nights with me and has always supported me in the moments when there was no one to answer my queries. Not to forget, my deepest love also goes to my lovely kids, Umar, Ulya and Zara. They have taught me how to be patient and how to manage my time efficiently along my PhD journey.

Yet, I am grateful to Allah for enabling me to realize these goals and for the strength He has given me to complete this thesis. Syukur Alhamdulillah.

ABSTRACT

Climate change is one of the greatest challenges for water resources management. Intensity and frequency of extreme rainfalls are increasing due to enhanced greenhouse gas effect caused by climate change. A lot of research has been done in developing innovative methods for assessing the impacts of climate change on rainfall extremes. Climate change strongly depends on General Circulation Model (GCM) outputs since they play a pivotal role in the understanding of climate change. However due to their coarse resolution, statistical downscaling is widely applied to match the scale between the GCM and the station scale. This research proposed to establish statistical downscaling model that was able to generate hourly rainfall data for future projection of hourly extreme rainfall in Peninsular Malaysia. An Advanced Weather Generator (AWE-GEN) built on stochastic downscaling principles was applied for simulating hourly rainfall data. The model construction involved 40 stations over Peninsular Malaysia with observations from 1975 to 2005. To account for uncertainties, an ensemble of multi-model namely GFDL-CM3, IS-CM5A-LR, MIROC5, MRI-CGCM3 and NorESM1-M were obtained from the dataset compiled in the WCRP's, CMIP5. The projections of extreme precipitation were based on the RCP 6.0 scenario (2081-2100). To address the problem of unavailability of rainfall data at remote areas over Peninsular Malaysia, this research also examined the spatial variability of rainfall and temperature parameters using Locally Weighted Regression. Results of the AWE-GEN showed its capability to simulate rainfall for Peninsular Malaysia. Both hourly and 24 hour extreme rainfall showed an increase for future. Extremes of dry spell was projected to decrease in future whereas extremes of wet spell was expected to remain unchanged. Simulations of present climate using interpolated parameters showed promising results for the studied regions.

ABSTRAK

Perubahan iklim adalah salah satu cabaran terbesar bagi pengurusan sumber air. Intensiti dan kekerapan hujan ekstrim semakin meningkat disebabkan peningkatan kesan gas rumah hijau yang disebabkan oleh perubahan iklim. Banyak kajian telah dilakukan dalam membangunkan kaedah inovatif untuk menilai kesan perubahan iklim ke atas hujan ekstrim. Perubahan iklim sangat bergantung kepada output Model Edaran Umum (GCM) kerana ia memainkan peranan penting dalam pemahaman perubahan iklim. Walaubagaimanapun, oleh kerana resolusi GCM yang kasar, kaedah penurunan statistik digunakan secara meluas untuk padanan skala antara GCM dan skala stesen. Kajian ini mencadangkan untuk membina model penurunan statistik yang mampu menjana data hujan seterusnya mengunjur hujan ekstrim pada selang masa satu jam di masa depan untuk Semenanjung Malaysia. Kaedah Penjana Cuaca Termaju (AWE-GEN) yang dibina atas prinsip stokastik digunakan untuk simulasi data hujan setiap jam. Pembinaan model melibatkan 40 stesen di Semenanjung Malaysia dengan data cerapan dari tahun 1975 hingga 2005. Bagi mengambil kira ketidaktentuan, pelbagai model iaitu GFDL-CM3, IS-CM5A-LR, MIROC5, MRI-CGCM3 dan NorESM1-M telah diperolehi daripada WCRP's, CMIP5. Unjuran hujan ekstrim adalah berdasarkan senario RCP 6.0 (2081-2100). Bagi menangani masalah ketiadaan data hujan di kawasan terpencil, kajian ini juga mengkaji ubahan ruang parameter hujan dan suhu dengan menggunakan kaedah Regresi Tempatan Wajaran. Keputusan hasil AWE-GEN menunjukkan keupayaan untuk menjana simulasi hujan di seluruh Semenanjung Malaysia. Kedua-dua siri hujan ekstrim bagi satu jam dan 24 jam menunjukkan peningkatan pada masa akan datang. Musim kering ekstrim dijangka berkurangan sementara musim hujan ekstrim dijangka kekal tidak berubah. Hasil bagi simulasi iklim semasa menggunakan parameter berinterpolasi menunjukkan keputusan yang agak memberangsangkan bagi rantau kajian.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	xi
	LIST OF FIGURES	xiii
	LIST OF ABBREVIATIONS	xvii
	LIST OF SYMBOLS	xix
	LIST OF APPENDICES	xxiii
1	INTRODUCTION	1
	1.1 Introduction	1
	1.2 Problem Statement	4
	1.3 Aim and Objectives of Research	5
	1.4 Scope of Research	6
	1.5 Significance of Research	7
	1.6 Summary	7
2	LITERATURE REVIEW	9
	2.1 Introduction	9
	2.2 Analysis on Precipitation Trend	9
	2.2.1 Trend Analysis of Extreme Precipitation	11
	2.2.2 Methods in Calculating the Trend	13

2.3	Statistical Downscaling Study	14
2.3.1	Statistical Downscaling Methods	15
2.3.2	Statistical Downscaling of Extreme Precipitation	18
2.3.3	Uncertainties in Statistical Downscaling	20
2.4	Weather Generator	21
2.5	Spatial Interpolation of Weather Generator Parameters	25
2.6	Summary	26
3	METHODOLOGY	27
3.1	Introduction	27
3.2	Trend Analysis	27
3.3	Advanced Weather Generator (AWE-GEN)	31
3.3.1	Historical Data	33
3.3.2	Statistical Properties of Observed Data (OBS) and AWE-GEN Parameters	33
3.3.2.1	Intra-Annual Variability of Precipitation	34
3.3.2.2	Inter-Annual Variability of Precipitation	36
3.4	Downscaling	39
3.4.1	GCMs Data	39
3.4.2	Statistical Properties of GCM	40
3.4.3	Factor of Change (FC)	41
3.4.4	Bayesian Approach	42
3.4.4.1	Likelihood Functions	43
3.4.4.2	Prior Distributions	44
3.4.4.3	Posterior Distributions	45
3.4.5	Extension to Finer Temporal Scale	48
3.4.5.1	The Scaling Properties	49
3.4.6	Return Period	51

3.5	Spatial Interpolation of Rainfall and Temperature Parameters	52
3.5.1	Locally Weighted Regression (LWR)	52
3.5.1.1	Weight of Training Data	
	Location	53
3.5.1.2	Weighted Regression	
	Coefficients	53
3.5.1.3	Local Generalized Cross-Validation (LGCV) of Weighted Regression and Weighted Average	54
3.5.2	Global Regression	56
3.5.3	Domain Average	56
3.6	Summary	56
4	HISTORICAL TREND OF EXTREME RAINFALL EVENTS IN PENINSULAR MALAYSIA	61
4.1	Introduction	61
4.2	Study Region	61
4.3	Spatial Profile of Extreme Rainfall Indices	65
4.4	Trend Analysis of Extreme Rainfall Indices	77
4.5	The Significant Contribution of Extreme Indices to Malaysia's Extreme Rainfall	87
4.6	Summary	93
5	AWE-GEN SIMULATIONS OF PRESENT AND FUTURE EXTREME CLIMATE TIME SERIES	94
5.1	Introduction	94
5.2	Study Region	94
5.3	Generation of Current Scenario using AWE-GEN	97
5.3.1	Monthly Rainfall for Present Climate	98
5.3.2	Statistical Properties	100
5.3.3	Overall Performance of AWE-GEN	104
5.4	Generation of Future Scenarios	108

5.4.1	Factor of Change	108
5.4.2	Monthly Rainfall for Future Climate	116
5.4.3	Results of AWE-GEN for CTS and FUT	117
5.5	Extreme Rainfall	122
5.6	Summary	123
6	SPATIAL INTERPOLATIONS OF RAINFALL AND TEMPERATURE PARAMETERS	125
6.1	Introduction	125
6.2	Performance of LWR, Domain Average and Global Regression	125
6.3	Spatial Interpolation using LWR	129
6.3.1	Monthly Rainfall for Present Climate	129
6.3.2	Statistical Properties	134
6.3.3	Results of Locally Weighted Regression for OBS and INT	138
6.4	Monthly Rainfall for Future Climate	143
6.5	Extreme Rainfall	144
6.6	Summary	145
7	CONCLUSION	147
7.1	Introduction	147
7.2	Conclusion	147
7.3	Recommendations for Future Work	149
	REFERENCES	150
	Appendices A - H	168 - 245

LIST OF TABLES

TABLE NO.	TITLE	PAGE
1.1	Definition of RCP scenario	4
2.1	The advantages and disadvantages of each statistical downscaling method	17
3.1	Extreme rainfall indices with definitions and units	29
3.2	AWE-GEN parameters	32
3.3	Names and measurement units of meteorological variables	33
3.4	GCMs that will be used for future projections	39
4.1	Name of stations with longitude and latitude	64
4.2	Extreme rainfall indices	65
4.3	Significant hourly extreme indices for each station during NEM	88
4.4	Significant hourly extreme indices for each station during SWM	89
4.5	Significant hourly extreme indices for each station during MA	90
4.6	Significant hourly extreme indices for each station during SO	91
5.1	List of rainfall stations with longitude, latitude and elevation	96
5.2	Observed and simulated mean monthly rainfall for station 3516022	98
5.3	Estimated rainfall parameters of the AWE-GEN model	99

5.4	Statistical properties at aggregation time of 1 hour, 24 hour and 48 hour	102
5.5	Observed and future monthly rainfall for station 3516022	117
5.6	Percentage of change in future mean rainfall (%)	121
6.1	RMSE values of LWR, domain average and global regression for all parameters	128
6.2	The selected coefficients for each rainfall parameter weighted average is represented as 'wa'. Weighted regression is represented as 'wr'	131
6.3	Observed rainfall (OBS) and estimated rainfall using interpolated parameters (INT)	133
6.4	Estimated rainfall parameters of the AWE-GEN model	134
6.5	Mean of rainfall intensity	134
6.6	Statistical properties at aggregation time of 1 hour, 24 hour and 48 hour	136
6.7	Observed rainfall (OBS), simulated rainfall using interpolated parameters (INT) and future rainfall using interpolated parameters (FUT INT)	144

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
3.1	A scheme for the NSRP model	34
3.2	Autoregressive lag-1 model	38
3.3	AWE-GEN process in generating rainfall at present climate	57
3.4	Downscaling and generating future rainfall time series process	58
3.5	Locally weighted regression (LWR)	59
3.6	Global regression	60
3.7	Domain average	60
4.1	Location of stations for trend analysis	63
4.2	Spatial distribution of Hr Max index for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	66
4.3	Spatial distribution of total amount for Tot> 95 th index, for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	68
4.4	Spatial distribution of total amount for Tot> 99 th index for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	69
4.5	Spatial distribution of 5-Hr Max index for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	71
4.6	Spatial distribution of 24-Hr Max index for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	72
4.7	Spatial distribution of Freq> 20 index for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	74

4.8	Spatial distribution of Freq> 95 th for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	75
4.9	Spatial distribution of Freq> 99 th for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	76
4.10	Spatial pattern of trends for Hr Max for (a) NEM, (b) SWM, (c) MA and (d) SO (1975-2010)	78
4.11	Spatial pattern of trends for Tot> 95 th for (a) NEM, (b) SWM, (c) MA and (d) SO	80
4.12	Spatial pattern of trends for Tot> 99 th for (a) NEM, (b) SWM, (c) MA and (d) SO	81
4.13	Spatial pattern of trends for 5-Hr Max for (a) NEM, (b) SWM, (c) MA and (d) SO	82
4.14	Spatial pattern of trends for 24-Hr Max for (a) NEM, (b) SWM, (c) MA and (d) SO	83
4.15	Spatial pattern of trends for Freq> 20 for (a) NEM, (b) SWM, (c) MA and (d) SO	84
4.16	Spatial pattern of trends for Freq> 95 th for (a) NEM, (b) SWM, (c) MA and (d) SO	85
4.17	Spatial pattern of trends for Freq> 99 th for (a) NEM, (b) SWM, (c) MA and (d) SO	86
4.18	Significant trends of extreme rainfall indices during (a) NEM, (b) SWM, (c) MA and (d) SO (Circle/triangle indicates larger number of significant extreme rainfall indices using linear regression test where ‘circle’ represents significant positive trend, ‘triangle’ represents significant negative trend)	92
5.1	Location of rainfall stations	95
5.2	A comparison between observed and simulated monthly rainfall for station 3516022. The red and green circles and vertical bars denote the mean and standard deviation of the monthly values for observed and simulated, respectively	98
5.3	A comparison between OBS and CTS monthly statistics of precipitation (mean, variance, lag-1 autocorrelation, skewness, frequency of no rainfall, transition probability wet-wet) for station 3516022, for the aggregation period of (a) 1 hour, (b) 24 hour and (c) 48 hour	101

5.4	Mean monthly rainfall of 40 stations for OBS and CTS	108
5.5	The posterior PDF of mean temperature and precipitation obtained from the multi-model ensemble for station 3516022	115
5.6	A comparison between observed and future monthly rainfall for station 3516022. The circles and vertical bars denote the mean and standard deviations of the monthly values respectively	116
5.7	Range of mean monthly rainfall of 40 stations for CTS and FUT period with percentage of change in future mean rainfall (%)	120
5.8	A comparison between the OBS, CTS and FUT values of extreme rainfall at (a) 1 and (b) 24 hours aggregation periods; (c) extremes of dry and (d) wet spell durations for station 3516022	123
6.1	RMSE values of LWR, DA and Domain-wide (global) regression for each rainfall parameters (a) α , (b) β , (c) η , (d) λ , (e) μ_c and (f) θ for station 3516022	127
6.2	A comparison between OBS and simulated monthly precipitation using interpolated parameters (INT) for station 3516022. The red and cyan circles denote the mean of the monthly values for observed and simulated, respectively. The red and cyan vertical bars denote the standard deviations of the monthly values, respectively	132
6.3	A comparison between OBS and INT monthly statistics of precipitation (mean, variance, lag-1 autocorrelation, skewness, frequency of non-precipitation, transition probability wet-wet), for the aggregation period of (a) 1, (b) 24 and (c) 48 hours for station 3516022	136
6.4	Range of mean monthly rainfall of 40 stations for OBS and INT	142
6.5	A comparison between observed and future monthly precipitation using interpolated parameters for station 3516022. The vertical bars denote the standard deviations of the monthly values	143

- 6.6 A comparison between the OBS, simulated of extreme rainfall using interpolated parameters (INT) and projected of future extreme rainfall using interpolated parameter (FUT INT) at (a) 1 and (b) 24 hours aggregation periods; (c) extremes of dry and (d) wet spell durations for station 3516022

LIST OF ABBREVIATIONS

ANN	-	Artificial Neural Network
AOGCM	-	Atmospheric Ocean General Circulation Model
AR1	-	Autoregressive Lag-1 model
AWE-GEN	-	Advanced Weather Generator
CGCM2	-	Canadian Coupled General Circulation Model
CMIP3	-	Coupled Model Intercomparison Project Phase 3
CMIP5	-	Coupled Model Intercomparison Project Phase 5
CTS	-	GCM control period
DID	-	Malaysia Drainage and Irrigation Department
ENSO	-	El Niño-Southern Oscillation
FC	-	Factors of change
FUT	-	GCM future period
GCM	-	General Circulation Model
GFDL-CM3	-	NOAA Geophysical Fluid Dynamics Laboratory, United States
GHG	-	Greenhouse gas
HadCM3	-	Hadley Centre Coupled Model, Version 3
INT	-	Interpolated
IPCC	-	Intergovernmental Panel on Climate Change
IPSL-CM5A-LR	-	Institut Pierre-Simon Laplace, Paris
LARS-WG	-	Long Ashton Research Station-Weather Generator
LGCV	-	Local Generalized Cross-Validation
LWR	-	Locally Weighted Regression
MA	-	March-April
MCMC	-	Markov Chain Monte Carlo

MIROC5	-	Atmosphere and Ocean Research Institute (The University of Tokyo), National Institute for Environmental Studies, and Japan Agency for Marine-Earth Science and Technology, Japan
MJO	-	Madden-Julian Oscillation
M-K	-	Mann-Kendall
MM5	-	Fifth-Generation Penn State/NCAR Mesoscale Model
MMD	-	Malaysia Meteorological Department
MRI-CGCM3	-	Meteorological Research Institute, Japan
NCEP	-	National Centers for Environmental Predictions
NEM	-	Northeast monsoon
NorESM1-M	-	Norwegian Climate Centre, Norway
NSRP	-	Neyman-Scott Rectangular Pulses
OBS	-	Observations
PDFs	-	Probability density functions (PDFs)
RCM	-	Regional Climate Model
RCP	-	Representation Concentration Pathways
SDSM	-	Statistical Downscaling Model
SLP	-	Sea-level pressure
SO	-	September-October
SRES	-	Special Report on Emission Scenarios
SST	-	Sea Surface Temperature
SWM	-	Southwest monsoon
WCRP	-	World Climate Research Programme

LIST OF SYMBOLS

\bar{D}_0	-	Observed mean distance between each station and their nearest neighbor
d_i	-	Distance between station i and its nearest station
\bar{D}_E	-	Expected mean distance for the stations given a random pattern
z_{ANN}	-	Score for the statistic for average nearest neighborhood
y_i	-	Observed values of the response variable for individual i
\hat{y}_i	-	Predicted values of the response variable for individual i
t	-	Test statistic for t-test
$C_h(h)$	-	Coefficient of variation
$\rho(h)$	-	The lag-1 auto-correlation
$\kappa(h)$	-	Skewness
$\Phi(h)$	-	Probability that an arbitrary interval of length is dry
λ^{-1}	-	Mean storm origin arrivals for precipitation (h)
β^{-1}	-	Mean waiting time for cell origins after the origin of the storm for precipitation (h)
η^{-1}	-	Mean duration of the cell for precipitation (h)
μ_c	-	Mean number of cell per storm for precipitation [-]
α	-	Shape parameter of the Gamma distribution of rainfall intensity for precipitation [-]

θ	-	Scale parameter of the Gamma distribution of rainfall intensity for precipitation ($\text{mm } h^{-1}$)
M_0	-	Mean fair weather cloudiness for cloud cover
σ_m	-	Standard deviation fair weather cloudiness for cloud cover
ρ_m	-	Lag-1 autocorrelation of mean fair weather cloudiness
$\gamma = \zeta$	-	Cloudiness decay rate (h^{-1})
J_1	-	Length transition period (h)
B	-	Regression coefficients of deterministic component of temperature first order differential equation for air temperature
$\overline{dT_h}$	-	Mean random temperature deviate
ρ_{dT}	-	Lag-1 autocorrelation random temperature deviate
$\sigma_{dT,h}$	-	Standard deviation random temperature deviate
dT	-	First step temperature ($^{\circ}C$)
μ_0	-	Ozone for solar radiation (cm)
μ_n	-	Nitrogen dioxide for solar radiation (cm)
α_{Λ}	-	Angstrom turbidity parameter for solar radiation
$\omega_{\Lambda 1}, \omega_{\Lambda 2}$	-	Single scattering albedos for solar radiation
ρ_g	-	Surrounding ground albedo for solar radiation
β_{Λ}	-	Angstrom turbidity parameter for solar radiation
LWP_R	-	Liquid water path with cloudiness for solar radiation (g/m^2)
$a_i, i = 0,1,2,3$	-	Regression coefficients of deterministic component of vapor pressure
$\overline{d\Delta e}$	-	Average of vapor pressure deficit deviations

$\sigma_{d\Delta e}$	-	Standard deviation of vapor pressure deficit deviations
$\rho_{d\Delta e}$	-	Lag-1 autocorrelation of the process
$c_i (i = 0, \dots, 4)$	-	Regression coefficients of deterministic component of wind speed
$\overline{dW_s}$	-	Average wind speed deviation
σ_{dW_s}	-	Standard deviation of wind speed
ρ_{dW_s}	-	Lag-1 autocorrelation of the process of wind speed
γ_{dW_s}	-	Skewness of wind speed
\overline{P}_{atm}	-	Average atmospheric pressure
$\sigma_{P_{atm}}$	-	Standard deviation of atmospheric pressure
$\rho_{P_{atm}}$	-	Lag-1 autocorrelation of the process of atmospheric pressure
\overline{P}_{yr}	-	Average annual precipitation (mm)
$\sigma_{P_{yr}}$	-	Standard deviation of annual precipitation
$\rho_{P_{yr}}$	-	Lag-1 autocorrelation of the process of annual precipitation
$\gamma_{P_{yr}}$	-	Skewness of annual precipitation
μ_h	-	Mean at a given aggregation time interval h
$\gamma_{h,l}$	-	Covariance at a given aggregation time interval and h lag l
ξ_h	-	Third moment of precipitation process
$\mu_{Pr}(h)$	-	Mean of precipitation
$\sigma^2(h)$	-	Variance of precipitation
$\rho_{Pr}(h)$	-	Lag-1 autocorrelation of precipitation

$\kappa_{Pr}(h)$	-	Skewness of precipitation
$\Phi_{Pr}(h)$	-	Probability that an arbitrary interval of length is dry
$\mu_T(h)$	-	Mean of temperature
$\nu, i = 1, \dots, 5$	-	Precision parameters
μ	-	Present temperature
ν	-	Future temperature
ν_0	-	Natural variability
k	-	Number of GCM
ϕ	-	Degrees of freedom
h	-	Aggregation time interval (hourly)
N	-	Total number of stations
w_i	-	Weight of training data
n	-	Training data
K_i	-	Biweight kernel function
D_{\max}	-	The largest of the n distances to the interpolation point
n_{\min}	-	Minimum number of training data
ε	-	Random error
b_0 and b_1	-	Coefficients of LWR equation

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Legend of rainfall stations in Appendices	168
B	A comparison between observed and simulated monthly precipitation for 39 stations. The vertical bars denote the standard deviations of the monthly values	169
C	A comparison between OBS and CTS monthly statistics of precipitation for the aggregation period of (a) 1, (b) 24 and (c) 48 hours for 39 stations	174
D	The posterior PDF of mean temperature and precipitation obtained from the multi-model ensemble for 39 stations from January to December (A-L)	188
E	A comparison between OBS and FUT monthly precipitation for 39 stations. The vertical bars denote the standard deviations of the monthly values	215
F	A comparison between the OBS, CTS and FUT values of extreme precipitation at (a) 1 and (b) 24 hours aggregation periods; (c) extremes of dry and (d) wet spell durations for 39 stations	220
G	A comparison between OBS and simulated monthly precipitation using interpolated parameters (INT) for 39 stations. The vertical bars denote the standard deviations of the monthly values	227
H	A comparison between OBS and INT monthly statistics of precipitation (mean, variance, lag-1 autocorrelation, skewness, frequency of non-precipitation, transition probability wet-wet), for the aggregation period of (a) 1-h, (b) 24-h and (c) 48-h for 39 stations	232

CHAPTER 1

INTRODUCTION

1.1 Introduction

Climate change is a crucial problem as it requires us to adapt our activities to uncertain future climate scenarios. Several sectors such as water resources, agriculture, energy and tourism face the severe impacts caused by climate change. Climate change and global warming occur when the global atmospheric concentrations of carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O) increases due to human activities. Consequently, the observed global average temperatures has also been increasing since the mid-20th century due to the rise of anthropogenic greenhouse gas (GHG) concentrations (IPCC, 2007). The warming has been shown to affect many natural systems which includes notable changes in snow, ice and frozen ground, increased runoff and changes in both terrestrial and marine ecosystems (Tangang *et al.*, 2012), changes in rainfall and risks of flooding (Willem *et al.*, 2011) and changes of occurrences of extreme precipitation (Huang *et al.*, 2011). Besides that, the notable changes in marine and freshwater ecosystems are also related to changes in temperature. Among the major impacts of global warming are the increased frequency and intensity of extreme rainfall events (Sen Roy, 2009; Cheng *et al.*, 2011). Extreme rainfall is one of the main causes of natural disasters such as flooding. Therefore, to date, considerable attention has been paid to the modeling of extreme rainfall for preventive measures of massive flooding as well as for projecting future extreme rainfall (Chu *et al.*, 2012). This has led to numerous collaborations between meteorologist and hydrologists to establish hydrological model of spatial and temporal precipitation extremes.

General Circulation Model (GCM) are widely used in providing outputs of global climate across the world which consist of hydrometeorological variables such as precipitation, air temperature, relative humidity, wind speed, geopotential height and cloud cover. GCM is a numerical model which comprises of different earth frameworks; for example, air, sea, surface area and ocean ice. Information on the important processes about global and continental scale atmosphere can be projected by GCM for future atmosphere under different emission scenarios. Despite numerous uncertainties in different GCMs (Chu *et al.*, 2010), these outputs provide hydrologists with the desired information. Unfortunately, GCMs are usually at resolution that is too coarse for many climate change impact studies (Fowler *et al.*, 2007; Hessami *et al.*, 2008; Hashmi *et al.*, 2009; Chu *et al.*, 2010; Hashmi *et al.*, 2010; Fatichi *et al.*, 2011). The internal relationships between the model's variables produced from GCM also may not always be the same as those found in the observational data. As a result, their simulations of current regional climate can often be inaccurate for sub-grid scales (Chu *et al.*, 2010; Guo *et al.*, 2011).

The discrepancy between the GCM scale and the scale that is required for most impact studies has led to the development of downscaling methodologies. In order to match the scale between the GCM outputs and hydrological process at smaller scale, downscaling must be employed. In particular, downscaling is used to model the hydrometeorological variables, at a smaller scale from a large scale. There are two approaches for downscaling: dynamical downscaling and statistical downscaling. Dynamical downscaling or known as Regional Climate Model (RCM) simulates climate at resolution of 50 km or less where the GCMs provide the boundary conditions to RCMs (Fowler and Wilby, 2010). Meanwhile, statistical downscaling is an empirical method that defines the statistical relationships between the large-scale climate features and the hydrological variables (Wilby *et al.*, 2004; Sunyer *et al.*, 2011). There are various discussions and debates on these two approaches. However, statistical downscaling requires less computational effort and is cheaper to employ (Wilby and Wigley, 2000; Huang *et al.*, 2011). Advantages of statistical downscaling also include the opportunity to use ensemble GCM results which takes into account average results from more than one model. Projections from ensemble model is better as compared to projections from individual model

where the uncertainties from different GCM models could be taken into account (Wibig *et al.*, 2015).

Previously, future projections are based on Special Report of Emission Scenarios (SRES) scenarios. All SRES scenarios are non-intervention scenarios with an increasing forcing path during the 21st century while Representative Concentration Pathways (RCP) span a large range of stabilization, mitigation and non-mitigation pathways (Rogelj *et al.*, 2012). There are some similarities and differences between temperature projections for SRES scenarios and RCPs. As stated in Rogelj *et al.* (2012), the RCP 8.5 is equivalent to SRES A1F1 scenario which represent high-emission, non-mitigation future where by 2100 the range of temperature is between 4.0 to 6.0 °C. RCP 6.0 temperature projections are equivalent to SRES B2. Likewise, RCP 4.5 temperature projections are equivalent to SRES B1. However, the lowest RCP scenario is basically different from the SRES scenario. In spite of having similarities, there are some particular differences between these two scenarios in terms of the rate of median temperature rises.

For instance, the median temperatures in RCP 8.5 rise slower than in SRES A1F1 during the period between 2035 and 2080, and faster during other periods of the 21st century. On the other hand, the median temperatures in RCP 6.0 rise faster than in SRES B2 during the three decades between 2060 and 2090 while slower during other periods of the 21st century. Similarly, the median temperatures in RCP 4.5 rise faster than in SRES B1 until mid-century, and slower afterwards. RCP scenarios are more focusing on process begins with pathways of radiative forcing, yet not detailed socioeconomic narratives or scenarios as in SRES scenarios. Table 1.1 summarizes the definition of each RCP scenarios which will be the baseline of the future climate condition. The basis of the scenario name is according to their 2100 radiative forcing level based on the forcing of greenhouse gases and other forcing agents.

Table 1.1: Definition of RCP scenario

Name	Radiative forcing ¹	Concentration ²	Pathway	SRES temperature anomaly equivalent
RCP 8.5	>8.5 W/m ² in 2100	> approx. 1370 CO ₂ -eq in 2100	Rising	SRES A1F1
RCP 6.0	Approximate 6 W/m ² at stabilization after 2100	Approximate 850 CO ₂ -eq (at stabilization after 2100)	Stabilizing without overshoot	SRES B2
RCP 4.5	Approximate 4.5 W/m ² at stabilization after 2100	Approximate 650 CO ₂ -eq (at stabilization after 2100)	Stabilizing without overshoot	SRES B1
RCP 2.6-PD ³	peak at approximate 2.6W/m ² before 2100 and then decline	peak at approximate 490 CO ₂ -eq before 2100 and then decline	Peak and decline	None

¹Approximate radiative forcing levels were defined as $\pm 5\%$ of the stated level in W/m². Radiative forcing values include the net effect of all anthropogenic GHGs and other forcing agents.

²Approximate CO₂ equivalent (CO₂-eq) concentrations. The CO₂-eq concentrations were calculated with the simple formula $\text{Conc} = 278 * \exp(\text{forcing}/5.325)$. Note that the best estimate of CO₂-eq concentration in 2005 for long-lived GHGs only is about 455 ppm, while the corresponding value including the net effect of all anthropogenic forcing agents (consistent with the table) would be 375 ppm CO₂-eq.

³PD = peak and decline.

1.2 Problem Statement

Malaysia experiences massive floods occurred during monsoon seasons and flash floods brought by convective rainfall occurred during intermonsoon seasons. Flash floods are frequently associated with convective storms which tend to be of short durations (IASH, 1974; Jamaluddin, 1985). Major concern on flash floods include structural and erosional damage, loss of life and property and disruption of socio-economic activity (Jamaluddin, 1985). This is an indication that Malaysia will face a higher probability of damages from extreme rainfall in the future. Thus,

understanding the patterns of extreme rainfall and their future behaviour is of importance to policy makers in Malaysia.

Malaysia currently ranks 52nd in the Climate Change Performance Index 2015 according to their emissions level, emissions development, renewable energy, efficiency and policy (CCPI, 2015). The change of rainfall patterns as well as temperature changes in future might be useful inputs to policy makers to initiate the mitigation and adaptation strategies in order to adapt with the future uncertain climate change. Due to the weaknesses of GCM as mentioned earlier, downscaling is used to match the scale between the GCM and the station scale since the GCM scale is much coarser compared to station scale. Two approaches of downscaling known as dynamical and statistical are extensively applied by the climatologist and hydrologist. In this study, the statistical downscaling approach will be adopted. There are three types of statistical downscaling methods: regression, weather typing scheme and weather generator. Weather generators involve stochastic process (Wilks and Wilby, 1999) that can be used to produce long time series of simulated weather variables and simulating future climate by perturbing weather parameters or by fitting to perturbed statistics (Michelle *et al.*, 2012). This study uses the hourly rainfall data as inputs to the weather generator. Research done by Fatichi *et al.* (2011) have demonstrated the capacity of an hourly climate generator in reproducing a wide set of climate statistics over a range of temporal scales including extreme variables. This method adopts stochastic models, often referred as stochastic downscaling. The absence of such methodology being used in climate projections for Malaysia is the basis of undertaking this study. Projection of future extreme rainfall events for the country is crucial, and the appropriate method for that purpose needs to be developed.

1.3 Aim and Objectives of Research

The overall aim of this research is to establish the statistical downscaling model that is able to generate hourly rainfall data at present climate and project the future hourly extreme rainfall in Peninsular Malaysia. This research also aims to

examine the potential of spatial variables in the model for simulation of ungauged sites using locally weighted regression.

The objectives of the research are:

1. To study on statistical downscaling model focusing on the Advanced Weather Generator (AWE-GEN) method.
2. To explore the ability of AWE-GEN method in projecting future extreme rainfall events in Malaysia using new model parameters.
3. To investigate and determine spatial interpolation methods using physical elements as variables for simulating extreme rainfall events for ungauged sites.

1.4 Scope of Research

This research will focus on one of the statistical downscaling model which is the weather generator. The weather generator that will be used in this research is the hourly AWE-GEN which combines the physically-based and stochastic approaches. The precipitation that will be considered in this research is hourly rainfall. Other meteorological data required in this study are hourly air temperature, hourly wind speed, hourly relative humidity, hourly atmospheric pressure, hourly cloud cover and hourly solar radiation. Hourly rainfall, temperature, wind speed and relative humidity data are from the Malaysia Meteorological Data (MMD). Meanwhile, hourly atmospheric pressure, cloud cover and solar radiation are adopted from Fatichi *et al.* (2011). Data will be limited to sites in Peninsular Malaysia. GCMs realizations will be obtained from the dataset compiled in the World Climate Research Programme's (WCRP's), Coupled Model Intercomparison Project phase 5 (CMIP5). An ensemble of multi-model (i.e. more than one GCM model will be used) in this study. They are GFDL-CM3 (United States), IS-CM5A-LR (Paris), MIROC5 (Japan), MRI-CGCM3 (Japan) and NorESM1-M (Norway). RCP 6.0 scenario will be used for future projections.

1.5 Significance of Research

Recognizing the needs and significance of having adequate rainfall data, this research attempts to propose a suitable hourly weather generator model which could generate a wide set of climate statistics over a range of temporal scales, from extremes to low-frequency inter-annual variability for the whole of Peninsular Malaysia. Such information would be beneficial especially to hydrologists and environmentalists. Moreover, realizing the importance of understanding and predicting climate change, this research also attempts to simulate the future climate scenarios, as inferred from climate models, using the proposed model. The proposed model is able to quantify uncertainties by estimating the weighted averages based on outputs of different climate models using Bayesian theories. Having insufficient climate data is a critical problem in hydrological studies (Ming Kang and Fadhilah, 2012). Thus, the output of this study which will be simulated time series of climate data will be able to alleviate this problem. This study will also extend the application of AWE-GEN by interpolating the AWE-GEN parameters to simulate weather time series at remote areas where meteorological data do not exist. This will be invaluable for hydrological studies done in such locations.

1.6 Summary

This chapter discusses issues of climate change occurring around the world, including Malaysia. Problems related to climate change which are of concern to the policy makers were also discussed. GCM outputs which provides information on climate at global scale need to be downscaled to finer scale in order to match scale required for hydrological modeling at local scale. Therefore, statistical downscaling will be applied in this study. This chapter outlined the aim and the objectives of the research, the scope and the significance of the research. Therefore, Chapter 2 will cover the literature reviews of rainfall studies as well as statistical downscaling studies. In addition to that, past researches done on spatial interpolation methods will also be discussed in Chapter 2. Next, the methodology to be applied in this research will be discussed in detail in Chapter 3 including the theories and assumptions

involved. The discussions of results will be given in Chapters 4, 5 and 6. Finally, conclusion and recommendations will be discussed in Chapter 7.

7.3 Recommendations for Future Work

Further research could identify the best fit distribution to represent rain cell intensity in AWE-GEN besides the Gamma distribution. For instance, Weibull, Generalized Pareto, Exponential and Mixed-Exponential distributions could be fitted to the rain cell intensity. Malaysia has different seasonal variation of rainfall and different geographical where local climates are affected by the presence of mountain ranges throughout Malaysia which can be divided into three groups which are the highlands, the lowlands, and coastal regions. For the spatial interpolation methods, further research could also identify the relationship between the rainfall and temperature parameters with longitude and latitude of the stations.

Furthermore, other variables such as geographical and seasonality factors could be incorporated in LWR in order to establish a more robust interpolation model for Peninsular Malaysia. Moreover, besides rainfall and temperature, other parameters such as solar radiation and wind speed could also be considered to be incorporated in the model. The range of future changes of extreme climate under certain level of radiative forcing with certain level of economic and population growth would be more beneficial for climatologists and meteorologists for designing the mitigation plan and coping with the future risks. Further research could also project future extreme precipitation under other different RCP scenarios such as RCP 2.6, RCP 4.5, RCP 6.0 and RCP 8.5. The results from the different scenarios could be assessed and a range of future values developed to give a more comprehensive projection of future extreme values.

REFERENCES

- Ababaei, B., Sohrabi, T. M., Mirzaei, F. and Karimi, B. (2010). Evaluation of a Stochastic Weather Generator in Different Climates. *Computer and Information Science*. 3, 217-229.
- Adamowski, K. and Bougadis, J. (2003). Detection of Trends in Annual Extreme Rainfall. *Hydrological Processes*. 17(18), 3547-3560.
- Aisar Ashra Muhammad Ashri, Aminuddin Mohd Baki and Ismail Atan (2011). Synthetic Simulation of Stochastic Rainfall-runoff Model. *Paper presented at UMTAS*.
- Ambun Dindang, Azlai bin Taat, Phuah Eng Beng, Atifah bt Mohd Alwi, Alliscia Alk Mandai, Siti Fauziah binti Mat Adam, Farah Safura binti Othman, Dayang Norazila binti Awang Bina and Delan Lah (2013). Statistical Trend Analysis of Rainfall Data in Kuching, Sarawak from 1968-2010.
- Arritt, R. W. and A. R. Daniel (2014). Precipitation Intensity Changes in Regional Climate Projections for North America. *EGU General Assembly Conference Abstracts vol. 16, abstract EGU 2014- 13825. Presented at European Geoscience Union General Assembly 2014. Vienna, Austria. 27 April - 2 May 2014*.
- Asefa, T. and Adams, A. (2013). Reducing Bias-Corrected Precipitation Projection Uncertainties: A Bayesian-Based Indicator-Weighting Approach. *Regional Environmental Change*. 13, S111-S120.
- Barrow, E. M. and Semenov, M. A. (1995). Climate-Change Scenarios with High Spatial and Temporal Resolution for Agricultural Applications. *Forestry*. 68(4), 349-360.
- Barrow, E. M. (2002). Obtaining Finer Resolution Scenarios of Climate Change: A Review of Downscaling Methodologies. *A Report prepared for the*

- International Joint Commission's Lake Ontario-St. Lawrence River Study*, 33.
- Beecham, S., Rashid, M. and Chowdhury, R. K. (2014). Statistical Downscaling of Multi-Site Daily Rainfall in a South Australian Catchment using A Generalized Linear Model. *International Journal Climatology Royal Meteorological Society*. <http://dx.doi.org/10.1002/joc.3933>.
- Brissette, F. P., Khalili, M. and Leconte, R. (2007). Efficient Stochastic Generation of Multi-Site Synthetic Precipitation Data. *Journal of Hydrology*. 345(3-4), 121-133.
- Burton, A., Fowler, H. J., Blenkinsop, S. and Kilsby, C. G. (2010). Downscaling Transient Climate Change using a Neyman–Scott Rectangular Pulses Stochastic Rainfall Model. *Journal of Hydrology*. 381(1-2), 18-32.
- Busuioc, A., Tomozeiu, R. and Cacciamani, C. (2008). Statistical Downscaling Model Based on Canonical Correlation Analysis for Winter Extreme Precipitation Events in the Emilia-Romagna Region. *International Journal of Climatology*. 28(4), 449-464.
- CCPI (2015). The Climate Change Performance Index: Results 2015. *Germanwatch*. Retrieved on 20th January, 2015.
- Chen, J., Brissette, F. P. and Leconte, R. (2011). Uncertainty of Downscaling Method in Quantifying the Impact of Climate Change on Hydrology. *Journal of Hydrology*. 401, 190-202.
- Cheng, C. S., Auld, H., Li, Q. and Li, G. (2011). Possible Impacts of Climate Change on Extreme Weather Events at Local Scale in South-Central Canada. *Climatic Change*. 112, 963-979.
- Chu, P. S., Xin, Z., Ying, R. and Melodie, G. (2010). Extreme Rainfall Events in the Hawaiian Islands. *Journal of Applied Meteorology and Climatology*. 48(3), 502-516.
- Chu, H. J., Pan, T. Y. and Liou, J. J. (2012). Change-Point Detection of Long-Duration Extreme Precipitation and the Effect on Hydrologic Design: A Case Study of South Taiwan. *Stochastic Environmental Research and Risk Assessment*. 26(8), 1123-1130.
- Chun, K. P., Wheeler, H. S. and Onof, C. (2013). Comparison of Drought Projections using Two UK Weather Generators. *Hydrological Sciences Journal*. 58(2), 295-309.

- Cleveland, W. S. and Devlin, S. J. (1988). Locally Weighted Regression: An Approach to Regression Analysis by Local Fitting. *Journal of the American Statistical Association*. 83(403), 596-610.
- Cowpertwait, P. S. P., O'Connell, P. E., Metcalfe, A. V. and Mawdsley, J. A. (1996a). Stochastic Point Process Modelling of Rainfall. I. Single-Site Fitting and Validation. *Journal of Hydrology*. 175(1), 17-46.
- Cowpertwait, P. S. P., O'Connell, P. E., Metcalfe, A. V. and Mawdsley, J. A. (1996b). Stochastic Point Process Modelling of Rainfall. II. Regionalisation and Disaggregation. *Journal of Hydrology*. 175(1-4), 47-65.
- Cox, D. R. and Isham, V. (1980). Point Processes. *Chapman and Hall*. New York: CRC Press.
- Craven, P. and Wahba, G. (1978). Smoothing Noisy Data with Spline Functions. *Numerische Mathematik*. 31(4), 377-403.
- DeGaetano, A. T. and Wilks, D. S. (2009). Radar-Guided Interpolation of Climatological Precipitation Data. *International Journal of Climatology*. 29(2), 185-196.
- Deni, S. M., Suhaila, J., Zin, W. Z. W. and Jemain, A. A. (2010). Spatial Trends of Dry Spells over Peninsular Malaysia during Monsoon Seasons. *Theoretical and Applied Climatology*. 99(3-4), 357-371.
- Diong Jeong Yik, Subramaniam Moten, Munirah Ariffin and Siva Shangari Govindan (2010). Trends in Intensity and Frequency of Precipitation Extremes in Malaysia from 1951 to 2009. *Technical Reports, Malaysia Meteorological Department*.
- Duan, J., McIntyre, N. and Onof, C. (2012). Resolving Non-Stationarity in Statistical Downscaling of Precipitation under Climate Change Scenarios. *British Hydrology Society, BHS Eleventh National Symposium, Hydrology for a Changing World, Dundee 2012*. ISBN: 1903741181, doi: 10.7558/bhs.2012.ns16.
- Dubrovský, M. (1997). Creating Daily Weather Series with Use of the Weather Generator. *Environmetrics*. 8(5), 409-424.
- Elison Timm, O., Takahashi, M., Giambelluca, T. W. and Diaz, H. F. (2013). On the Relation between Large-Scale Circulation Pattern and Heavy Rain Events over the Hawaiian Islands: Recent Trends and Future Changes. *Journal of Geophysical Research: Atmospheres*. 118(10), 4129-4141.

- Ewona, I. O., Osang, J. E. and Udo, S. O. (2014). Trend Analyses of Rainfall Patterns in Nigeria using Regression Parameters. *International Journal of Technology Enhancements and Emerging Engineering Research*. 2(5), 2347-4289.
- Fan, L., Lu, C. H., Yang, B. and Chen, Z. (2012). Long-Term Trends of Precipitation in the North China Plain. *Journal of Geographical Science*. 22(6), 989-1001.
- Fatichi, S., Ivanov, V. Y. and Caporali, E. (2011). Simulation of Future Climate Scenarios with a Weather Generator. *Advances in Water Resources*. 34(4), 448-467.
- Fernández-Ferrero, A., Sáenz, J., Ibarra-Berastegi, G. and Fernández, J. (2009). Evaluation of Statistical Downscaling in Short Range Precipitation Forecasting. *Atmospheric Research*. 94(3), 448-461.
- Fowler, H. J., Blenkinsop, S. and Tebaldi, C. (2007). Linking Climate Change Modelling to Impacts Studies: Recent Advances in Downscaling Techniques for Hydrological Modelling. *International Journal of Climatology*. 27(12), 1547-1578.
- Fowler, H. J. and Wilby (2010). Detecting Changes in Seasonal Precipitation Extremes using Regional Climate Model Projections: Implications for Managing Fluvial Flood Risk. *Water Resources Research*, 46, W03525, doi: 10.1029/2008WR007636.
- Friederichs, P. (2010). Statistical Downscaling of Extreme Precipitation Events using Extreme Value Theory. *Extremes*. 13(2), 109-132.
- Gädeke, A., Hölzel, H., Koch, H., Pohle, I. and Grünewald, U. (2014). Analysis of Uncertainties in the Hydrological Response of a Model-Based Climate Change Impact Assessment in a Subcatchment of the Spree River, Germany. *Hydrological Processes*. 28(12), 3978-3998.
- Giorgi F. and Mearns L. O. (2002). Calculation of Average, Uncertainty Range, and Reliability of Regional Climate Changes from AOGCM Simulations via the Reliability Ensemble Averaging (REA) Method. *Journal of Climate*. 15, 1141-1158.
- Griffiths, G. M., Salinger, M. J., and Leleu, I. (2003). Trends in Extreme Daily Rainfall across the South Pacific and Relationship to the South Pacific Convergence Zone. *International Journal of Climatology*. 23(8), 847-869.

- Guenni, L. (1997). Spatial Interpolation of Stochastic Weather Model Parameters. *Journal of Environmental Management*. 49(1), 31-42.
- Guenni, L. and Hutchinson, M. F. (1998). Spatial Interpolation of the Parameters of a Rainfall Model from Ground-Based Data. *Journal of Hydrology*. 212(1-4), 335-347.
- Gunarathna, M. H. J. P. and Kumari, M. K. N. (2013). Rainfall Trends in Anuradhapura: Rainfall Analysis for Agricultural Planning. *Rajarata Unuversity Journal*. 1, 38-44.
- Guo, J., Chen, H., Xu, C. -Y., Guo, S. and Guo, J. (2011). Prediction of Variability of Precipitation in the Yangtze River Basin under the Climate Change Conditions based on Automated Statistical Downscaling. *Stochastic Environmental Research & Risk Assessment*. 26, 157-176.
- Gutierrez-Ruacho, O. G., Brito-Castillo, L., Diaz-Castro, S. C. and Watts, C. J. (2010). Trends in Rainfall and Extreme Temperatures in Northwestern Mexico. *Climate Research*. 42(2), 133-142.
- Haktanir, T., Bajabaa, S. and Masoud, M. (2013). Stochastic Analyses of Maximum Daily Rainfall Series Recorded at Two Stations Across the Mediterranean Sea. *Arabian Journal of Geosciences*. 6(10), 3943-3958.
- Hashmi, M. Z., Shamseldin, A. Y. and Melville, B. W. (2009). Downscaling of future rainfall extreme events: a weather generator based approach. *18th World IMACS Congress and MODSIM09 International Congress on Modelling and Simulation: Interfacing Modelling and Simulation with Mathematical and Computational Sciences*. 3928-3934.
- Hashmi, M. Z., Shamseldin, A. Y. and Melville, B. W. (2010). Comparison of SDSM and LARS-WG for Simulation and Downscaling of Extreme Precipitation Events in a Watershed. *Stochastic Environmental Research and Risk Assessment*. 25(4), 475-484.
- Hashmi, M. Z., Shamseldin, A. Y. and Melville, B. W. (2013). Statistically Downscaled Probabilistic Multi-Model Ensemble Projections of Precipitation Change in A Watershed. *Hydrological Processes*. 27(7), 1021-1032.
- Hassan, Z. and Harun, S. (2012). Application of Statistical Downscaling Model for Long Lead Rainfall Prediction in Kurau River Catchment of Malaysia. *Malaysian Journal of Civil Engineering*. 24(1), 1-12.

- Hassan, Z., Shamsudin, S. and Harun, S. (2014). Application of SDSM and LARS-WG for Simulating and Downscaling of Rainfall and Temperature. *Theoretical and Applied Climatology*. 116(1-2), 243-257.
- Haylock, M. and Nicholls, N. (2000). Trends in Extreme Rainfall Indices for an Updated High Quality Data Set for Australia, 1910-1998. *International Journal of Climatology*. 20(13), 1533-1541.
- Hessami, M., Gachon, P., Ouarda, T. B. M. J., and St-Hilaire, A. (2008). Automated Regression-Based Statistical Downscaling Tool. *Environmental Modelling & Software*. 23(6), 813-834.
- Ho Ming Kang and Fadhilah Yusof (2012). Application of Self Organizing Map (SOM) in Missing Daily Rainfall Data. *International Journal of Computer Application (IJCA)*, 0975-888. 48(5).
- Huang, C. Y., Wong, C. S. and Yeh, T. C. (2011). Extreme Rainfall Mechanisms Exhibited by Typhoon Morakot (2009). *Terrestrial Atmospheric and Ocean Sciences*. 22, 613-632.
- Hubert, P., Y. Tessier, S. Lovejoy, D. Schertzer, F. Schmitt, P. Ladoy, J. P. Carbonnel, Violette, S. and Desurogne, I. (1993). Multifractals and Extreme Rainfall Events. *Geophysical Research Letters*. 20, 931-934.
- Hundecha, Y., and Bárdossy, A. (2008). Statistical Downscaling of Extremes of Daily Precipitation and Temperature and Construction of their Future Scenarios. *International Journal of Climatology*. 28(5), 589-610.
- Hussain, M. S. and Lee, S. (2013). The Regional and the Seasonal Variability of Extreme Precipitation Trends in Pakistan. *Asia-Pacific Journal of Atmospheric Sciences*. 49(4), 421-441.
- IASH (1974). Flash Floods. *Proceeding of the Paris Symposium*. September 1974. IASH/UNESCO/WMO.
- Iizumi, T., Takayabu, I., Dairaku, K., Kusaka, H., Nishimori, M., Sakurai, G., Ishizaki, N. N., Adachi, S. A. and Semenov, M. A. (2012). Future Change of Daily Precipitation Indices in Japan: A Stochastic Weather Generator-Based Bootstrap Approach to provide Probabilistic Climate Information. *Journal of Geophysical Research Atmospheres*. 117.
- IPCC (2007). Climate Change 2007: The Physical Science Basis. Summary for Policymakers, Contribution of Working Group I to the Fourth Assessment

- Report of the Intergovernmental Panel on Climate Change, IPCC Secretariat, Geneva, Switzerland.
- IPCC (2012). Summary for Policymakers. In: Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation [Field, C. B., V. Barros, T. F. Stocker, D. Qin, D. J. Dokken, K. L. Ebi, M. D. Mastrandrea, K. J. Mach, G. K. Plattner, S. K. Allen, M. Tignor and P. M. Midgley (eds.)]. *A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, USA. 1-19.
- Ivanov, V. Y, Bras, R. L. and Curtis D. C. (2007). A Weather Generator for Hydrological, Ecological and Agricultural Applications. *Water Resources Research*. 43, W10,406 doi:10.1029/2006WR005,364.
- Jamaluddin, M. J. (1985). Flash Flood Problems and Human Responses to the Flash Flood Hazard in Kuala Lumpur Area, Peninsular Malaysia. *Akademika*. 26, 45-62.
- Jeong, D. I., St-Hilaire, A., Ouarda, T. B. M. J., and Gachon, P. (2013). A Multi-Site Statistical Downscaling Model for Daily Precipitation using Global Scale GCM Precipitation Outputs. *International Journal of Climatology*. 33(10), 2431-2447.
- Joshi, S., Kumar, K., Joshi, V. and Pande, B. (2014). Rainfall Variability and Indices of Extreme Rainfall-Analysis and Perception Study for Two Stations over Central Himalaya, India. *Natural Hazards*. 72(2), 361-374.
- Juneng, L., Tangang, F. T. and Reason, C. J. C. (2007). Numerical Case Study of an Extreme Rainfall Event during 9-11 December 2004 over the East Coast of Peninsular Malaysia. *Meteorological Atmospheric Physics*. 98, 81-98.
- Juneng, L., Tangang, F. T., Kang, H., Lee, W. J. and Seng, Y. K. (2010). Statistical Downscaling Forecasts for Winter Monsoon Precipitation in Malaysia Using Multimodel Output Variables. *Journal of Climate*. 23(1), 17-27.
- Katz, R. W., Parlange, M. B. and Naveau, P. (2002). Statistics of Extremes in Hydrology. *Advances in Water Resources*. 25(8-12), 1287-1304.
- Khan, M. S., Coulibaly, P. and Dibike, Y. (2006). Uncertainty Analysis of Statistical Downscaling Methods using Canadian Global Climate Model Predictors. *Hydrological Processes*. 20(14), 3085-3104.

- Khazaei, M. R., Zahabiyou, B. and Saghafian, B. (2012). Assessment of Climate Change Impact on Floods using Weather Generator and Continuous Rainfall-Runoff Model. *International Journal of Climatology*. 32(13), 1997-2006.
- Kilsby, C. G, Jones P., Burton, A., Ford, A., Fowler H, Harpham C, James, P., Smith, A. and Wilby, R. L. (2007). A Daily Weather Generator for Use in Climate Change Studies. *Environmental Modelling and Software*. 22, 1705–1719.
- Kim, B., Kim, H., Seoh, B. and Kim, N. (2007). Impact of Climate Change on Water Resources in Yongdam Dam Basin, Korea. *Stochastic Environmental Research and Risk Assessment*. 21(4), 355-373.
- Kim, B. S., Kim, B. K. and Kwon, H. H. (2011). Assessment of the Impact of Climate Change on the Flow Regime of the Han River Basin using Indicators of Hydrologic Alteration. *Hydrological Processes*. 25(5), 691-704.
- Kuchar, L., Iwanski, S., Jelonek, L. and Szalinska, W. (2014). Application of Spatial Weather Generator for the Assessment of Climate Change Impacts on a River Runoff. *Geografie*. 119(1), 1-25.
- Kysely, J. (2009). Trends in Heavy Precipitation in the Czech Republic over 1961-2005. *International Journal of Climatology*. 29(12), 1745-1758.
- Laguardia, G. (2011). Representing the Precipitation Regime by Means of Fourier Series. *International Journal of Climatology*. 31(9), 1398-1407.
- Lakshmi Kumar, T. V., Koteswara Rao, K., Barbosa, H. and Uma, R. (2014). Trends and Extreme Value Analysis of Rainfall Pattern over Homogeneous Monsoon Regions of India. *Natural Hazards*. 73(2), 1003-1017.
- Lall, U., Moon, Y. I., Kwon, H. H. and Bosworth, K. (2006). Locally Weighted Polynomial Regression: Parameter Choice and Application to Forecasts of The Great Salt Lake. *Water Resources Research*. 42(W05422), doi:10.1029/2004WR003782.
- Larsen, A. N., Gregersen, I. B., Christensen, O. B., Linde, J. J. and Mikkelsen, P. S. (2009). Potential Future Increase in Extreme One-Hour Precipitation Events over Europe due to Climate Change. *Water Science & Technology*. 60(9), 2205-2216.
- Li, Z., Zheng, F. L., Liu, W. Z. and Flanagan, D. C. (2010). Spatial Distribution and Temporal Trends of Extreme Temperature and Precipitation Events on the Loess Plateau of China during 1961-2007. *Quaternary International*. 226(1-2), 92-100.

- Li, J., Zhang, Q., Chen, Y. D., Xu, C. Y. and Singh, V. P. (2013). Changing Spatiotemporal Patterns of Precipitation Extremes in China during 2071-2100 based on Earth System Models. *Journal of Geophysical Research: Atmospheres*. 118(22), 2013JD020300.
- Liu, Z., Xu, Z., Charles, S. P., Fu, G. and Liu, L. (2011). Evaluation of Two Statistical Downscaling Models for Daily Precipitation over an Arid Basin in China. *International Journal of Climatology*. 31(13), 2006-2020.
- Longobardi, A. and Villani, P. (2010). Trend Analysis of Annual and Seasonal Rainfall Time Series in the Mediterranean Area. *International Journal of Climatology*. 30(10), 1538-1546, doi:10.1002/joc.2001.
- Lupikasza, E. B., Hansel, S. and Matschullat, J. (2011). Regional and Seasonal Variability of Extreme Precipitation Trends in Southern Poland and Central-Eastern Germany 1951-2006. *International Journal of Climatology*. 31(15), 2249-2271.
- Madsen, H., P. S., Mikkelsen, D., Rosbjerg and P. Harremoës (2002). Regional Estimation of Rainfall Intensity-Duration-Frequency Curves using Generalized Least Squares Regression of Partial Duration Series Statistics. *Water Resources Research*. 38(11), 1239, 21-1–21-11.
- Manning, L. J., Hall, J. W., Fowler, H. J., Kilsby, C. G. and Tebaldi, C. (2009). Using Probabilistic Climate Change Information from a Multimodel Ensemble for Water Resources Assessment. *Water Resources Research*. 45(11), W11411.
- Marani M. (2003). On the Correlation Structure of Continuous and Discrete Point Rainfall. *Water Resources Research*. 39(5), doi:10.1029/2002WR001456.
- Marani M. (2005). Non-Power-Law-scale Properties of Rainfall in Space and Time. *Water Resources Research*. 41(W08413), doi:10.1029/2004WR003822.
- Mareuil, A., Leconte, R., Brissette, F. and Minville, M. (2007). Impacts of Climate Change on the Frequency and Severity of Floods in the Châteauguay River Basin, Canada. *Canadian Journal of Civil Engineering*. 34(9), 1048-1060.
- McGinley, M. (2011). Climate of Malaysia. Retrieved from <http://www.eoearth.org/view/article/151260>.
- McGree, S., Whan, K., Jones, D., Alexander, L. V., Imielska, A., Diamond, H., Ene, E., Finaulahi, S., Inape, K., Jacklick, L., Kumar, R., Laurent, V., Malala, H., Malsale, P., Moniz, T., Ngemaes, M., Peltier, A., Porteous, A., Pulehetoa-

- Mitiepo, R., Seuseu, S., Skilling, E. Tahani, L., Teimitsi, F., Toorua, U. and Vaiimene, M. (2014). An Updated Assessment of Trends and Variability in Total and Extreme Rainfall in the Western Pacific. *International Journal of Climatology*. 34(8), 2775-2791.
- Meehl, G. A., G. J. Boer, C. Covey, M. Latif and R. J. Stouffer (2000). The Coupled Model Intercomparison Project (CMIP). *Bulletin of the American Meteorological Society*. 81, 313-318.
- Menabde, M., D. Harris, A. Seed, G. Austin and D. Stow (1997). Multiscaling Properties of Rainfall and Bounded Random Cascades. *Water Resources Research*. 33, 2823-2830.
- Michaels, P. J., Knappenberger, P. C., Frauenfeld, O. W. and Davis, R. E. (2004). Trends in Precipitation on the Wettest Days of the Year across the Contiguous USA. *International Journal of Climatology*. 24(15), 1873-1882.
- Michelle, T. H. Van Vliet, Stephen Blenkinsop, Aidan Burton, Colin Harpham, Hans Peter Broers and Hayley J. Fowler (2012). A Multi-model Ensemble of Downscaled Spatial Climate Change Scenarios for the Dommel Catchment, Western Europe. *Climatic Change*. 111, 249-277.
- Min, Y. M., Kryjov, V., An, K. H., Hameed, S., Sohn, S. J., Lee, W. J., Lee, W. J. and Oh, J. H. (2011). Evaluation of the Weather Generator CLIGEN with Daily Precipitation Characteristics in Korea. *Asia-Pacific Journal of Atmospheric Sciences*. 47(3), 255-263.
- Minville, M., Brissette, F and Leconte, R. (2008). Uncertainty of the Impact of Climate Change on the Hydrology of a Nordic Watershed. *Journal of Hydrology*. 358(1-2), 70-83.
- Mohtar, Z. A., A. S. Yahaya, F. Ahmad, S. Suri and M. H. Halim (2014). Trends for Daily Rainfall in Northern and Southern Region of Peninsular Malaysia. *Journal of Civil Engineering Research*. 4, 222-227.
- Nóbrega, J. N. d., Santos, C. A. C. d., Gomes, O. M., Bezerra, B. G. and Brito, J. I. B. D. (2014). Extreme Precipitation Events in the Mesoregions of Paraíba and Its Relationship with the Tropical Oceans SST. *Revista Brasileira de Meteorologia*. 29, 197-208.
- Norton, C. W., Chu, P.-S. and Schroeder, T. A. (2011). Projecting Changes in Future Heavy Rainfall Events for Oahu, Hawaii: A Statistical Downscaling

- Approach. *Journal of Geophysical Research-Atmospheres*. 116(D17110), doi:10.1029/2011JD015641.
- Nurunnabi, A., G. West, and D. Belton (2013). Robust Locally Weighted Regression for Ground Surface Extraction in Mobile Laser Scanning 3D Data. *ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial Information Sciences, Volume II-5/W2, 2013ISPRS Workshop Laser Scanning 2013*. 11 - 13 November 2013. Antalya, Turkey, 217-222.
- Nyatuame, M., Owusu-Gyimah, V. and Ampiaaw, F. (2014). Statistical Analysis of Rainfall Trend for Volta Region in Ghana. *International Journal of Atmospheric Sciences*. 2414 Article ID 203245, 11.
- Nyenje, P. M. and Batelaan, O. (2009). Estimating the Effects of Climate Change on Groundwater Recharge and Baseflow in the Upper Ssezibwa Catchment, Uganda. *Hydrological Sciences Journal-Journal Des Sciences Hydrologiques*. 54(4), 713-726.
- Olsson, J. (1995). Limits and Characteristics of the Multifractal Behaviour of a High Resolution Rainfall Time-Series. *Nonlinear Processes in Geophysics*. 2, 23-29.
- Oscá, J., Romero, R. and Alonso, S. (2013). Precipitation Projections for Spain by Means of a Weather Typing Statistical Method. *Global and Planetary Change*. 109(0), 46-63.
- Pal, I. and Al-Tabbaa, A. (2010). Regional Changes in Extreme Monsoon Rainfall Deficit and Excess in India. *Dynamics of Atmospheres and Oceans*. 49(2-3), 206-214.
- Parajuli, K. and Kang, K. (2014). Application of Statistical Downscaling in GCMs at Constructing the Map of Precipitation in the Mekong River Basin. *Russian Meteorology and Hydrology*. 39(4), 271-282.
- Parey, S., Hoang, T.T.H. and Dacunha-Castelle, D. (2014). Validation of a Stochastic Temperature Generator Focusing on Extremes, and an Example of Use for Climate Change. *Climate Research*. 59(1), 61-75.
- Patil, S. and Stieglitz, M. (2012). Controls on Hydrologic Similarity: Role of Nearby Gauged Catchments for Prediction at an Ungauged Catchment. *Hydrology and Earth System Sciences*. 16(2), 551-562.

- Pervez, M. S. and Henebry, G. M. (2014). Projections of the Ganges–Brahmaputra Precipitation-Downscaled from GCM Predictors. *Journal of Hydrology*. 517(0), 120-134.
- Porto de Carvalho, J. R., Assad, E. D., Medeiros Evangelista, S. R. and Pinto, H. D. S. (2013). Estimation of Dry Spells in Three Brazilian Regions - Analysis of Extremes. *Atmospheric Research*. 132, 12-21.
- Pour, S, H., Harun, S. B., Shahid, S. (2014). Genetic Programming for the Downscaling of Extreme Rainfall Events on the East Coast of Peninsular Malaysia. *Atmosphere*. 5, 914-936.
- Pourtouiserkani, A. and Rakhshandehroo, G. (2014). Investigating Climate Change Impact on Extreme Rainfall Events Case Study: Chenar-Randar Basin, Fars, Iran. *Scientia Iranica*. 21(3), 525-533.
- Prudhomme, C. and Davies, H. (2009). Assessing Uncertainties in Climate Change Impact Analyses on the River Flow Regimes in the UK. Part 2: Future Climate. *Climatic Change*. 93(1-2), 197-222.
- Qian, B.D., Hayhoe, H. and Gameda, S. (2005). Evaluation of the Stochastic Weather Generators LARS-WG and AAFC-WG for Climate Change Impact Studies. *Climate Research*. 29(1), 3-21.
- Quintana-Segui, P., Habets, F. and Martin, E. (2011). Comparison of Past and Future Mediterranean High and Low Extremes of Precipitation and River Flow Projected using Different Statistical Downscaling Methods. *Natural Hazards and Earth System Sciences*. 11(5), 1411-1432.
- Re, M. and Barros, V. R. (2009). Extreme Rainfalls in SE South America. *Climatic Change*. 96(1-2), 119-136.
- Ribalaygua, J., Torres, L., Pórtoles, J., Monjo, R., Gaitán, E. and Pino, M. R. (2013). Description and Validation of a Two-Step Analogue/Regression Downscaling Method. *Theoretical and Applied Climatology*. 114(1-2), 253-269.
- Rodriguez-Iturbe, I., Eagleson, P. S. (1987). Mathematical Models of Rainstorm Events in Space and Time. *Water Resources Research*. 23(1), 181-190.
- Rogelj, J., Meinshausen, M. and Knutti, R. (2012). Global Warming under Old and New Scenarios using IPCC Climate Sensitivity Range Estimates. *Nature Climate Change*. 2(4), 248-253.

- Saadatkath, Nader, Kassim, Azman, and Lee, Lee Min (2014). Hulu Kelang, Malaysia Regional Mapping of Rainfall-Induced Landslides using TRIGRS Model. *Arabian Journal of Geosciences*. 1-12.
- Salahuddin, A., and Curtis, S. (2011). Climate Extremes in Malaysia and the Equatorial South China Sea. *Global and Planetary Change*. 78(3-4), 83-91.
- Schertzer, D. and Lovejoy, S. (1987). Physical Modelling and Analysis of Rain and Clouds by Anisotropic Scaling Multiplicative Processes. *Journal Geophysical Research*. 92, 9693-9714.
- Schmidli, J., and Frei, C. (2005). Trends of Heavy Precipitation and Wet and Dry Spells in Switzerland during the 20th century. *International Journal of Climatology*. 25(6), 753-771.
- Schmidli, J., Goodess, C. M., Frei, C., Haylock, M. R., Hurrell, J. W., Ribalaygua, J. and Schmith, T. (2007). Statistical and Dynamical Downscaling of Precipitation: An Evaluation and Comparison of Scenarios for the European Alps. *Journal of Geophysical Research*. 112, D04105.
- Schnur, R. and Lettenmaier, D. P. (1998). A Case Study of Statistical Downscaling in Australia using Weather Classification by Recursive Partitioning. *Journal of Hydrology*. 212–213(0), 362-379.
- Semenov, M. A., and Barrow, E. M. (1997). Use of Stochastic Weather Generator in the Development of Climate Change Scenarios. *Climatic Change*. 35, 397-414.
- Semenov, M. A., Brooks, R. J., Barrow, E. M. and Richardson, C. W. (1998). Comparison of the WGEN and LARS-WG Stochastic Weather Generators in Diverse Climates. *Climate Research*. 10, 95-107.
- Semenov, M. A. and Brooks, R. J. (1999). Spatial Interpolation of the LARS-WG Stochastic Weather Generator in Great Britain. *Climate Research*. 11(2), 137-148.
- Semenov, M. A. and Barrow, E. M. (2002). LARS-WG: A Stochastic Weather Generator for Use in Climate Impact Studies, Version 3.0, User's Manual.
- Semenov, M. A. (2008). Simulation of Extreme Weather Events by a Stochastic Weather Generator. *Climate Research*. 35, 203-212.
- Sen Roy, S. (2009). A Spatial Analysis of Extreme Hourly Precipitation Patterns in India. *International Journal of Climatology*. 29, 345–355.

- Shahid, S. (2010). Rainfall Variability and the Trends of Wet and Dry Periods in Bangladesh. *International Journal of Climatology*. 30, 2299-2313, doi: 10.1002/joc.2053.
- Siti Musliha M.Rasit, Zalina Mohd Daud and Norzaida Abas (2013). A Regionalized Spatial Temporal Model for Hourly Rainfall Process. *International Conference on Computing, Mathematics and Statistics ICMS 2013*. P.Pinang, 28-29 August 2013.
- Souvignet, M., Gaese, H., Ribbe, L., Kretschmer, N. and Oyarzun, R. (2010). Statistical Downscaling of Precipitation and Temperature in North-Central Chile: An Assessment of Possible Climate Change Impacts in an Arid Andean Watershed. *Hydrological Sciences Journal- Journal Des Sciences Hydrologiques*. 55(1), 41-57.
- Souvignet, M. and Heinrich, J. (2011). Statistical Downscaling in the Arid Central Andes: Uncertainty Analysis of Multi-Model Simulated Temperature and Precipitation. *Theoretical and Applied Climatology*. 106(1-2), 229-244.
- Stephenson, D. B. (2008). Definition, Diagnosis and Origin of Extreme Weather and Climate Events. *Climate Extremes and Society*, Diaz HF, Murnane RJ (eds). Cambridge University Press, New York, 348.
- Suhaila, J., Deni, S. M., Zin, W. Z. W. and Jemain, A. A. (2010). Trends in Peninsular Malaysia Rainfall Data during the Southwest Monsoon and Northeast Monsoon Seasons: 1975-2004. *Sains Malaysiana*. 39(4), 533-542.
- Suhaila, J. and Jemain, A. A. (2012). Spatial Analysis of Daily Rainfall Intensity and Concentration Index in Peninsular Malaysia. *Theoretical and Applied Climatology*. 108(1-2), 235-245.
- Sunyer, M. A., Madsen, H. and Ang, P. H. (2011). A Comparison of Different Regional Climate Models and Statistical Downscaling Methods for Extreme Rainfall Estimation under Climate Change. *Atmospheric Research*. 103, 1-128.
- Sunyer, M. A., Madsen, H. and Ang, P. H. (2012). A Comparison of Different Regional Climate Models and Statistical Downscaling Methods for Extreme Rainfall Estimation under Climate Change. *Atmospheric Research*. 103(0), 119-128.

- Suppiah, R. and Hennessy, K. J. (1998). Trends in Total Rainfall, Heavy Rain Events and Number of Dry Days in Australia, 1910-1990. *International Journal of Climatology*. 18(10), 1141-1164.
- Tangang, F. T., Juneng, L., Salimun, E., Sei, K. M., Le, L. J. and Muhamad, H. (2012). Climate Change and Variability over Malaysia. *Sains Malaysiana*. 41(11), 1355-1366.
- Tatsumi, K., Oizumi, T. and Yamashiki, Y. (2014). Assessment of Future Precipitation Indices in the Shikoku Region using a Statistical Downscaling Model. *Stochastic Environmental Research and Risk Assessment*. 28(6), 1447-1464.
- Taylor, G. I. (1935). Statistical Theory of Turbulence. *Proceedings of the Royal Society*. London. Series A, 151, 421-478.
- Tebaldi, C, Smith, R. L., Nychka, D. and Mearns L.O. (2005). Quantifying Uncertainty in Projections of Regional Climate Change: A Bayesian Approach to the Analysis of Multi-Model Ensembles. *Journal of Climate*. 18, 1524-1540.
- Teegavarapu, R. S. V., Tufail, M. and Ormsbee, L. (2009). Optimal Functional Forms for Estimation of Missing Precipitation Data. *Journal of Hydrology*. 374(1-2), 106-115.
- Teegavarapu, R. S. V. (2012). Spatial Interpolation using Nonlinear Mathematical Programming Models for Estimation of Missing Precipitation Records. *Hydrological Sciences Journal*. 57(3), 383-406.
- Timbal, B., A. Dufour, and B. McAvaney (2003). An Estimate of Future Climate Change for Western France using a Statistical Downscaling Technique. *Climate Dynamic*. 20(7-8), 807-823.
- Timbal, B. and Jones, D. A. (2008). Future Projections of Winter Rainfall in Southeast Australia using a Statistical Downscaling Technique. *Climatic Change*. 86(1-2), 165-187.
- Tryhorn, L. and DeGaetano, A. (2011). A Comparison of Techniques for Downscaling Extreme Precipitation over the Northeastern United States. *International Journal of Climatology*. 31, 1975-1989.
- Tseng, H.-W., Yang, T.-C., Kuo, C.-M. and Yu, P.-S. (2012). Application of Multi-site Weather Generators for Investigating Wet and Dry Spell Lengths under

- Climate Change: A Case Study in Southern Taiwan. *Water Resources Management*. 26(15), 4311-4326.
- Tukimat, N. N. A., Harun, S. and Shahid, S. (2012). Comparison of Different Methods in Estimating Potential Evapotranspiration at Muda Irrigation Scheme of Malaysia. *Journal of Agriculture and Rural Development in the Tropics and Subtropics*. 113(1), 77-85.
- Turner, A J., Arlene M. Fiore, Larry W. Horowitz, and M. Bauer. (2013). Summertime cyclones over the Great Lakes Storm Track from 1860–2100: Variability, Trends, and Association with Ozone Pollution. *Atmospheric Chemistry and Physics*. 13(2), doi:10.5194/acp-13-565-2013.
- Vanmarke, E. (1983). *Random Fields: Analysis and Synthesis*, MIT Press. Cambridge, Mass.
- Vanuytrecht, E., Raes, D., Willems, P. and Semenov, M.A. (2014). Comparing Climate Change Impacts on Cereals based on CMIP3 and EU-ENSEMBLES Climate Scenarios. *Agricultural and Forest Meteorology*. 195, 12-23.
- Wibig, J., Mauran, D., Benestad, R., Kjellström, E. Lorenz, P. and Christensen, O. B. (2015). Projected Change-Models and Methodology. In Second Assessment of Climate Change for the Baltic Sea Basin. The BACC II Author Team. *Regional Climate Studies*. 189-216, doi: 10.1007/978-3-319-16006-1_102015.
- Wilby, R. L., Hassan, H. and Hanaki, K. (1998). Statistical Downscaling of Hydrometeorological Variables using General Circulation Model Output. *Journal of Hydrology*. 205, 1-19.
- Wilby, R. L. and Wigley, T. M. L. (2000). Precipitation Predictors for Downscaling: Observed and General Circulation Model Relationships. *International Journal of Climatology*. 20, 641-661.
- Wilby, R. L., Dawson, C. W. and Barrow, E. M. (2002). SDSM - A Decision Support Tool for the Assessment of Regional Climate Change Impacts. *Environmental Modelling & Software*. 17, 147-159.
- Wilby, R. L., Wedgbrow, C. S. and Fox, H. R. (2004). Seasonal Predictability of the Summer Hydrometeorology of the River Thames, UK. *Journal of Hydrology*. 295, 1-16.

- Wilks, D.S. and Wilby, R.L. (1999). The Weather Generation Game: A Review of Stochastic Weather Models. *Progress in Physical Geography*. 23(3), 329-357.
- Wilks, D.S. (2008). High-Resolution Spatial Interpolation of Weather Generator Parameters Using Local Weighted Regressions. *Agricultural and Forest Meteorology*. 148, 111-120.
- Wilks, D.S. (2010). Use of Stochastic Weather Generators for Precipitation Downscaling. *Wiley Interdisciplinary Reviews-Climate Change*. 1(6), 898-907.
- Willems, P., Arnbjerg-Nielsen, K., Olsson, J. and Nguyen, V. T. V. (2011). Climate Change Impact Assessment on Urban Rainfall Extremes and Urban Drainage: Methods and Shortcomings. *Atmospheric Research*. 103, 106-118.
- Williams, C. J. R., R. P. Allan and Kniveton, D. R. (2012). Diagnosing Atmosphere-Land Feedbacks in CMIP5 Climate Models. *Environmental Research Letters*. 7, 1-9.
- Wong, L., R. Venneker, S. Uhlenbrook, A. B. M. Jamil and Y. Zhou (2009). Variability of Rainfall in Peninsular Malaysia. *Hydrology and Earth System Sciences Discussions*. 6, 5471-5503.
- Xu, H., Corte-Real, J. and Qian, B.D. (2007). Developing Daily Precipitation Scenarios for Climate Change Impact Studies in the Guadiana and the Tejo Basins. *Hydrology and Earth System Sciences*. 11(3), 1161-1173.
- Yang, W., Bardossy, A. and Caspary, H.-J. (2010). Downscaling Daily Precipitation Time Series using A Combined Circulation- and Regression-Based Approach. *Theoretical and Applied Climatology*. 102(3-4), 439-454.
- Yin, C., Li, Y., Ye, W., Bornman, J. F. and Yan, X. (2011). Statistical Downscaling of Regional Daily Precipitation over Southeast Australia Based on Self-Organizing Maps. *Theoretical and Applied Climatology*. 105, 11-26.
- Yue, S., and C. Y. Wang. (2002). The Applicability of Pre-Whitening to Eliminate the Influence of Serial Correlation on the Mann-Kendall Test. *Water Resources Research*. 38(6), 1068, doi:10.1029/2001WR000861.
- Zalina, M. D., Mohd. Nor, D. and Nguyen, V. T. V. (2002). Statistical Analysis of At-Site Extreme Rainfall Processes in Peninsular Malaysia. *FRIEND*. 61-68.

- Zarghami, M., Abdi, A., Babaeian, I., Hassanzadeh, Y. and Kanani, R. (2011). Impacts of Climate Change on Runoffs in East Azerbaijan, Iran. *Global and Planetary Change*. 78(3-4), 137-146.
- Zhao, Y. F., Zou, X. Q., Cao, L. G. and Xu, X. W. H. (2014). Changes in Precipitation Extremes over the Pearl River Basin, Southern China, during 1960-2012. *Quaternary International*. 333, 26-39.
- Zin, W. Z. W., Jamaludin, S., Deni, S. M. and Jemain, A. A. (2010). Recent Changes in Extreme Rainfall Events in Peninsular Malaysia: 1971-2005. *Theoretical and Applied Climatology*. 99(3-4), 303-314.
- Zvolensky, M., Kohnova, S., Hlavcova, K., Szolgay, J. and Parajka, J. (2008). Regionalisation of Rainfall-Runoff Model Parameters based on Geographical Location of Gauged Catchments. *Journal of Hydrology and Hydromechanics*. 56(3), 176-189.