

**THE POTENTIAL USE OF LOCAL INDEX OF BIOLOGICAL INTEGRITY
FOR TROPICAL RIVER ASSESSMENT**

NOR ZAIHA BINTI ARMAN

UNIVERSITI TEKNOLOGI MALAYSIA

THE POTENTIAL USE OF LOCAL INDEX OF BIOLOGICAL INTEGRITY FOR
TROPICAL RIVER ASSESSMENT

NOR ZAIHA BINTI ARMAN

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Doctor of Philosophy (Civil Engineering)

Faculty of Civil Engineering
Universiti Teknologi Malaysia

MAY 2017

To my beloved family and friends

In thankful appreciation for your endless love, support and encouragement

ACKNOWLEDGEMENT

In the name of Allah, the Most Gracious and the Most Merciful

Alhamdulillah, all praises to Allah for the strengths and His blessing in completing this thesis. First and foremost I offer my sincerest gratitude to my supervisor, Assoc. Prof. Dr. Mohd. Ismid Mohd. Said, for thoughtful guidance, constructive comments and motivation throughout the completion of this research. Also a million thanks to my co-supervisor, Dr. Salmiati for her patience, encouragement, support and friendship. My acknowledgement also goes to all staffs and technicians, especially to the Environmental Engineering Laboratory, FKA UTM for their assistance and co-operations.

I greatly appreciate the Research University Grant Scheme (GUP) for the funding received with grant number Q.J130000.2522.00H99 and Ministry of Higher Education (MOHE) for scholarship assistance. Sincere thanks to my fellow postgraduate students, colleagues and others who have provided assistance at various occasions. Thank you for the great moments we have shared together.

Last but not least, heartfelt thanks to my beloved family for their continuous support and encouragement. To those who indirectly contributed in this research, your kindness means a lot to me. Thank you very much.

ABSTRACT

Economic and industrial growth in this country has led to environmental problems which in turn affect the quality of human life and nature. Particularly, the management plans for river rehabilitation are less effective in addressing issues related to rivers in Malaysia. To date, most of the efforts undertaken focused on evaluating physicochemical properties of water whereas biological evaluation in determining the health of a river ecosystem is given less attention. Therefore, this study aims to integrate assessment tools of physical, chemical and biological approaches by highlighting benthic macroinvertebrates as key indicator. This study was conducted at four rivers in the state of Johor, which represents different gradients of land use, namely Sungai Ayer Hitam Besar, Sungai Berasau, Sungai Mengkibol and Sungai Melana. *In-situ* and *ex-situ* monitoring were carried out with ten (10) sampling frequencies for each river, starting from 2010 to 2013. Benthic macroinvertebrates were sampled using rectangular dip-net with frame dimension 0.5 m x 0.3 m based on the United States Environmental Protection Agency Rapid Bioassessment Protocol whilst water samples were analysed according to Standard Methods for the Examination of Water and Wastewater. River habitat assessments were also conducted concurrently with the sampling of benthic macroinvertebrates and water quality whereas, secondary data such as rainfall, land use plans and maps of catchment area were obtained from various agencies. Assessment results varied among study sites. Non-parametric statistical analysis, Spearman correlation coefficients was also conducted to measure the correlation between biological and environmental endpoints, hence the selection of most suitable and effective indices for the development of macroinvertebrate-based multimetrics of Index Biotic Integrity (IBI) and the preliminary establishment of the biocriteria. Integration of six potential metrics within three categories (taxonomic richness, taxonomic composition and tolerance and intolerance index) had produced four-level discriminatory biocriteria for river health assessment. The re-scored result showed that water quality of Sungai Ayer Hitam Besar was in non-impaired condition with excellent water quality. Both Sungai Berasau and Sungai Mengkibol showed slightly impaired water quality, whereas Sungai Melana was moderately impaired with fair to poor water quality. The validation of IBI was conducted using different data thus demonstrated that the establishment of IBI and biological criteria could provide indispensable information in managing river ecosystem more effectively.

ABSTRAK

Pertumbuhan sektor ekonomi dan perindustrian di negara ini telah membawa kepada masalah alam sekitar seterusnya menjejaskan kualiti kehidupan manusia dan alam semulajadi. Pelan pengurusan bagi pemuliharaan sungai khususnya dilihat masih kurang berkesan dalam menangani isu berkaitan sungai di Malaysia. Sehingga kini kebanyakan usaha yang dijalankan lebih tertumpu kepada menilai ciri fisikokimia air, manakala penilaian secara biologi kurang diberi perhatian dalam menentukan status kesihatan sesebuah ekosistem sungai. Oleh itu, kajian ini bertujuan untuk mengintegrasikan kaedah penilaian secara fizikal, kimia dan biologi dengan menggunakan makroinvertebrata sebagai organisma petunjuk. Kajian ini telah dijalankan di empat batang sungai di negeri Johor yang mewakili guna tanah yang berbeza, iaitu Sungai Ayer Hitam Besar, Sungai Berasau, Sungai Mengkibol dan Sungai Melana. Pemantauan dijalankan secara *in-situ* dan *ex-situ* dengan kekerapan sepuluh (10) kali persampelan di setiap sungai, bermula dari 2010 sehingga 2013. Sampel makroinvertebrata diambil menggunakan jaring sauk berbentuk segi empat tepat bersaiz 0.5 m x 0.3 m berdasarkan *United States Environmental Protection Agency Rapid Bioassessment Protocol* sementara sampel air diuji mengikut kaedah piawai pengukuran kualiti air *APHA Standard Method*. Penilaian habitat dijalankan serentak dengan persampelan makroinvertebrata dan kualiti air, sementara data sekunder seperti taburan hujan, pelan guna tanah dan peta kawasan tadahan diperolehi daripada pelbagai agensi. Hasil penilaian adalah berbeza di antara kawasan kajian. Analisis statistik bukan parametrik, pekali korelasi Spearman turut diukur untuk melihat hubungan antara parameter biologi dan alam sekitar, seterusnya memilih indeks yang paling sesuai dan berkesan untuk pembangunan multimetrik berdasarkan makroinvertebrata bagi Indeks Biotik Integriti (IBI) dan penubuhan awal kriteria biologi. Integrasi enam metrik berpotensi yang mewakili tiga kategori (kekayaan taksonomi, komposisi taksonomi dan indeks toleransi dan intoleransi) telah menghasilkan empat peringkat diskriminasi kriteria biologi bagi penilaian kesihatan sungai. Hasil pengiraan semula menunjukkan bahawa kualiti air Sungai Ayer Hitam Besar adalah tidak terjejas dengan keadaan kualiti air yang cemerlang. Kedua-dua Sungai Berasau dan Sungai Mengkibol menunjukkan kualiti air sedikit terjejas, manakala Sungai Melana adalah sederhana terjejas dengan kualiti air yang sederhana. Pengesahan IBI turut dijalankan dengan menggunakan data yang berbeza, sekali gus menunjukkan bahawa penubuhan IBI dan kriteria biologi boleh memberikan maklumat yang berguna dalam menguruskan ekosistem sungai dengan lebih berkesan.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLE	xi
	LIST OF FIGURES	xiv
	LIST OF ABBREVIATIONS	xvii
	LIST OF SYMBOLS	xix
	LIST OF APPENDICES	xx
1	INTRODUCTION	1
	1.1 Research Background	1
	1.2 Problem Statement	3
	1.3 Aims and Objectives	5
	1.4 Scope of Study	5
	1.5 Significance of Study	6
2	LITERATURE REVIEW	8
	2.1 Overview	8
	2.2 River Ecosystem Concept	9

2.3	Overview	8
2.4	River Ecosystem Concept	9
2.5	Threat to Stream Biodiversity and Conservation	
	Approaches	15
2.3.1	Deforestation	16
2.3.2	Agriculture	17
2.3.3	Urban and Industrial Development	18
2.3.4	Flow Regulation and River Fragmentation	19
2.4	Biological Assessment	20
2.5	Biological Criteria	23
2.6	Macroinvertebrate as Bioindicator	27
2.7	Development of Benthic Macroinvertebrate	30
2.8	Benthic Macroinvertebrate Functional Feeding Groups	32
2.9	Factors Affecting Benthic Macroinvertebrate	
	Distribution	33
2.9.1	Temperature	34
2.9.2	Dissolved Oxygen	35
2.9.3	pH	35
2.9.4	River Discharge	36
2.9.5	Substrate	37
2.9.6	Food	38
2.9.7	Vegetation	38
2.9.8	Droughts	39
2.9.9	Floods	40
2.10	Assessment Approaches Based on Macroinvertebrate	41
2.10.1	Saprobic Approach	41
2.10.2	Diversity Approach	43
2.10.3	Biotic Approach	44
2.10.4	Multimetric Approach	45
2.11	Water Quality	46
2.12	Conclusion	48

3	METHODOLOGY	51
3.1	Introduction	51
3.2	Material	51
3.3	Analytical Method	54
3.3.1	Water Quality Analysis	54
3.3.2	Benthic Macroinvertebrate Analysis	56
3.3.3	Particle Size Distribution	59
3.3.4	Data Analysis Technique	60
3.4	Study Area	61
3.5	Experimental Procedure	61
3.5.1	Water Quality Sampling	62
3.5.2	Benthic Macroinvertebrate Sampling	63
3.5.3	River Habitat Assessment	64
3.5.4	Sediment Sampling	65
3.5.5	River Discharge	67
4	RESULTS AND DISCUSSION	68
4.1	Overview of the Study Area	68
4.1.1	Sg Ayer Hitam Besar, Gunung Pulai Forest Reserve	68
4.1.2	Sg Berasau, Ulu Sedili Plantation Forest	70
4.1.3	Sg Mengkibol, Kluang	71
4.1.4	Sg Melana, Skudai	73
4.2	Benthic Macroinvertebrates Composition	74
4.3	Ecological Indices	86
4.4	Physicochemical Characteristics for Water Quality	93
4.5	Comparison of Water Quality Classification Based on Different Indices	101
4.6	River Habitat Survey	106
4.7	River Discharge	113
4.8	Particle Size Distribution	114

4.9	The Influence of Hydrological, Physicochemical and Habitat Characteristics on the Aquatic Macroinvertebrates	117
4.10	Establishment of Biological Criteria	123
5	CONCLUSIONS AND RECOMMENDATIONS	131
5.1	Conclusions	131
5.2	Recommendations	132
	REFERENCES	134
	Appendices A - E	162 - 202

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	Summary of the River Continuum Concept (RCC)	11
2.2	Summary of river concept characteristics	14
2.3	Biological assessment studies using different type of bioindicators	23
2.4	Advantages and disadvantages of using fish, macrophytes, macroinvertebrate, plankton and periphyton as bioindicator in biological assessment (Abbasi and Abbasi, 2012)	24
2.5	Advantages and disadvantages of narrative or numeric criteria (Bauder and Pearson, 2002)	27
2.6	Traditional Saprobic System Classes and typical biological oxygen demand (BOD) and dissolved oxygen (DO) levels (Gordon <i>et al.</i> , 2004)	42
2.7	Comparison of BMWP and FBI	45
3.1	List of equipments used in this study	52
3.2	Basic soil type groups based on BS5390:1981 (Craig, 2013)	60
3.3	Coordinate of sampling sites	61
3.4	Physical and chemical observations measured alongside the RBP habitat assessment	66
4.1	Benthic macroinvertebrates composition in Sg Ayer Hitam Besar	76
4.2	Benthic macroinvertebrates composition in Sg Berasau	79
4.3	Benthic macroinvertebrates composition in Sg Mengkibol	81

4.4	Benthic macroinvertebrates composition in Sg Melana	84
4.5	Diversity, richness and evenness indices of Sg Ayer Hitam Besar	88
4.6	Diversity, richness and evenness indices of Sg Berasau	89
4.7	Diversity, richness and evenness indices of Sg Mengkibol	91
4.8	Diversity, richness and evenness indices of Sg Melana	93
4.9	Assessment of water quality according to applied indices	102
4.10	Assessment of Spearman correlation between physicochemical parameters and applied indices	105
4.11	Mean \pm standard deviation values for instream features and habitat quality for Sg Ayer Hitam Besar	109
4.12	Mean \pm standard deviation values for instream features and habitat quality for Sg Berasau	110
4.13	Mean \pm standard deviation values for instream features and habitat quality for Sg Mengkibol	111
4.14	Mean \pm standard deviation values for instream features and habitat quality for Sg Melana	112
4.15	River discharge data (m^3/s) between 2010 and 2013 from all studied rivers	114
4.16	Value of D_{60} , D_{30} and D_{10} , C_u and C_c for Sg Ayer Hitam Besar	116
4.17	Value of D_{60} , D_{30} and D_{10} , C_u and C_c for Sg Berasau	116
4.18	Value of D_{60} , D_{30} and D_{10} , C_u and C_c for Sg Mengkibol	116
4.19	Value of D_{60} , D_{30} and D_{10} , C_u and C_c for Sg Melana	117
4.20	Summary statistic for the canonical correspondence analysis (CCA) relating aquatic macroinvertebrates-environmental variables (11 variables) of four investigated rivers in Johor, Malaysia	118
4.21	Candidate metric of IBI development	125
4.22	Metric values of both, the reference and test sites according to analysis	125
4.23	Percent similarity of the study site metrics to the reference site metrics	126
4.24	Percent ranges and corresponding point values	126
4.25	Scoring point for each sampling site	127

4.26	Results of comparison to reference stream points	127
4.27	Recommended biocriteria for river health assessment	128
4.28	Step 1: Metric values	128
4.29	Step 2: Percent similarity	129
4.30	Step 3: Scoring point	129
4.31	Step 4: Comparison to reference stream points	129

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	A generalized depiction of the relationship between stream size (order), energy inputs, and ecosystem function as predicted by the river continuum concept (Vannote <i>et al.</i> , 1980)	10
2.2	A typical stream food web (Avril and Barten, 2007)	29
2.3	Metamorphosis (a) Complete metamorphosis, midge (b) Incomplete metamorphosis, damselfly (McCafferty, 1998)	31
2.4	Major morphological features of all aquatic insects (Whitlock, 2010)	32
3.1	Flowchart of study	53
3.2	Example of physical characteristics of channel unit	66
3.3	Diagram of channel cross-section using the mechanical current-meter method (source: US Geological Survey)	67
4.1	The Sg Ayer Hitam Besar and its location in Sg Pontian Besar river basin	69
4.2	The Sg Berasau and its location in Sungai Ulu Sedili Besar river basin	71
4.3	The Sg Mengkibol and its location in Sungai Endau river basin	72
4.4	The Sg Melana and its location in Sungai Skudai river basin	74

4.5	Benthic macroinvertebrate assemblages sampled from Sg Ayer Hitam Besar	76
4.6	Distribution pattern of benthic macroinvertebrate assemblages in Sg Ayer Hitam Besar	77
4.7	Macrobenthic assemblages sampled from Sg Berasau	78
4.8	Distribution pattern of macrobenthic assemblages in Sg Berasau	80
4.9	Macrobenthic assemblages sampled from Sg Mengkibol	81
4.10	Distribution pattern of macrobenthic assemblages in Sg Mengkibol	83
4.11	Macrobenthic assemblages sampled from Sg Melana	83
4.12	Distribution pattern of macrobenthic assemblages in Sg Melana	85
4.13	Biotic indices for Sg Ayer Hitam Besar	87
4.14	Biotic indices for Sg Berasau	89
4.15	Biotic indices for Sg Mengkibol	90
4.16	Biotic indices for Sg Melana	92
4.17	pH values from all sampling rivers and events	94
4.18	DO values from all sampling rivers and events	96
4.19	BOD ₅ values from all sampling rivers and events	97
4.20	COD values from all sampling rivers and events	98
4.21	TSS values from all sampling rivers and events	99
4.22	NH ₃ N values from all sampling rivers and events	100
4.23	WQI values from all sampling rivers and events	102
4.24	Riparian vegetation from second sampling station of Sg Ayer Hitam Besar	107
4.25	Riparian vegetation from second sampling station of Sg Berasau	107
4.26	Post-harvested land area in Sg Berasau	108
4.27	Channel stabilization of man-made structures for flood prevention (a) gabion at Sg Mengkibol (b) concrete wall at Sg Melana	108

4.28	Riverbank erosion resulting in soil deposition in (a) Sg Mengkibol, and (b) Sg Melana	112
4.29	Before and after channel alteration and riverbank strengthening works in Sg Melana	113
4.30	The nMDS Euclidean distance plot of the studied rivers based on their hydrological and physicochemical variables	119
4.31	The canonical correspondence analysis (CCA) biplot of the aquatic macroinvertebrates in relation to the hydrological and physicochemical parameters in four investigated rivers in Johor, Malaysia	120
4.32	Illustrative schematic of the potential interactions between the threats, impact, and the response to the stream macroinvertebrate assemblage	122

LIST OF ABBREVIATIONS

4WD	-	Four Wheel Drive
Ag ⁺	-	Silver
ALUs	-	Aquatic Life Uses
APHA	-	American Public Health Association
ASPT	-	Average Score Per Taxon
BMWP	-	Biological Monitoring Working Party
BOD ₅	-	Biochemical Oxygen Demand
CA	-	Correlation Analysis
Cc	-	Coefficient of Curvature
CCA	-	Canonical Correspondence Analysis
COD	-	Chemical Oxygen Demand
CPOM	-	Coarse Particulate Organic Matter
Cu	-	Coefficient of Uniformity
CWA	-	Clean Water Act
DID	-	Department of Irrigation and Drainage
DO	-	Dissolved Oxygen
DOE	-	Department of Environmental Malaysia
DOM	-	Dissolved Organic Matter
EIA	-	Environmental Impact Assessment
FBI	-	Family Biotic Index
FFG	-	Functional Feeding Groups
FPOM	-	Fine Particulate Organic Matter

H'	-	Shannon's Diversity Index
H ⁺	-	Hydrogen Ions
H ₂ SO ₄	-	Sulfuric Acid
IBI	-	Index of Biotic Integrity
INWQS	-	Interim National Water Quality Standards
IRBM	-	Integrated River Basin Management
IWRM	-	Integrated Water Resources Management
J	-	Pielou Evenness Index
JMG	-	Department of Minerals and Geosciences
K ₂ Cr ₂ O ₇	-	Potassium Dichromate
LQI	-	Lincoln Quality Index
M	-	Margalef Richness Index
MetMalaysia	-	Malaysian Meteorological Department
MyGAP	-	Malaysian Good Agricultural Practice
NH ₃ N	-	Ammoniacal Nitrogen
NH ₄ ⁺	-	Ammonium
nMDS	-	Non-Multidimensional Scale
NRE	-	Natural Resources and Environment
OH ⁻	-	Hydroxyl Ions
OPP3	-	Third Outline Perspective Plan
OQR	-	Overall Quality Rating
RBP	-	Rapid Bioassessment Protocol
RCC	-	River Continuum Concept
RHS	-	River Habitat Survey
SI	-	Sub Index
SOM	-	Malaysian Organic Scheme
TAN	-	Total Ammonia Nitrogen
TSS	-	Total Suspended Solids
USA	-	United States of America
USEPA	-	United States Environmental Protection Agency
WQI	-	Water Quality Index

LIST OF SYMBOLS

>	-	more than
<	-	less than
mm	-	millimetre
μm	-	micrometer
%	-	percent
mg/l	-	milligram per litre
$^{\circ}\text{C}$	-	degree Celsius
N	-	total number of individuals
Σ	-	sum of the calculations
ln	-	natural logarithm
ppm	-	parts per million
km^2	-	square kilometre
ha	-	hectare
g	-	gram
D_{10}	-	grain diameter at 10% passing
D_{30}	-	grain diameter at 30% passing
D_{60}	-	grain diameter at 60% passing
d	-	depth
Q	-	discharge rate
A	-	area
V	-	velocity
m/s	-	metre per second
r	-	coefficient of correlation
p	-	correlation of significant

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	Biotic Indices	162
B	Malaysia Water Quality Standard	169
C	Particle Size Distribution Curve	173
D	River Habitat Assessmet	174
E	Photos of Major Taxa From Four Studied Rivers	180
F	Water Quality Analysis Result	182
G	Particle Size Distribution	187
H	List of Papers Published	200

CHAPTER 1

INTRODUCTION

1.1 Research Background

Since time immemorial, rivers play a major role in the development of human society, serving as transport routes, water supply for domestic and agricultural use and yielding an important source of protein for human consumption. Rivers have similarly played an important role in the growth of towns and cities in Malaysia, with early settlements springing up along river banks and estuaries (Chan *et al.*, 2003). The discovery of tin deposits in the flood plains and river valleys also encouraged settlements to mushroom in these areas, leading to a booming tin-mining industry in the 1800s till 1980s which made the country the largest producer of tin in the world.

Malaysia has grown rapidly over the last three decades, transforming from a rural economy based on agriculture and tin mining to an export-based, manufacturing economy. Unfortunately, rapid changes of land use, especially on forested land and food crops to plantations as well as urban development have triggered river erosion, surface runoff and sedimentation of rivers, resulting eventually in overstressed river systems. River basins are frequently facing problems arising from flooding. Many rivers are gradually losing their ability to supply fresh water and as a result, these

rivers are now mainly used for transportation (Sulong *et al.*, 2005). Although raw water supply is not yet depleted, clean water that can be safely consumed by humans is becoming hard to come by.

The major causes of water pollution in Malaysia include effluent from sewage treatment plants, discharge from agro-based industries and livestock farming, land clearing activities and domestic sewage (Jabatan Perangkaan Malaysia, 2013). Rivers in both urban and rural areas are experiencing the same problems. Although environmental issues in Malaysia raise serious concerns, the measures taken to address the problem thus far have been fragmented and inadequate. In tandem with the growing global population and improvement of living standards, the increasing demand for fresh water has been said to overshadow the concerns of the warming effect of climate change (Nienhuis, 2006). This critical situation calls for water resource management to be addressed in order to meet growing consumer demand.

In earlier studies, the evaluation of river water quality relies merely on physicochemical characteristics. However, taking into consideration the physicochemical aspects alone are not sufficient to indicate a healthy ecosystem as a whole. Healthy water bodies exhibit ecological integrity, which consist of three components, namely chemical, physical and biological integrity. When one or more of these components are degraded, health of the water body is affected and in most cases aquatic life living in it will reflect the degradation. According to Gordon *et al.*, (2013), stream health measurement takes into consideration the water quality, habitat availability and suitability, energy sources, hydrology and the biota themselves.

In order to achieve a comprehensive evaluation of healthy water bodies, biological assessment tool should be carried out simultaneously with the standard physicochemical method. Biological assessment, the primary tool to evaluate the biological condition of a water body, comprise surveys and other direct measurements through biological communities such as plankton, periphyton, microphytobenthos, macrozoobenthos, aquatic macrophytes and fish. Among all,

benthic macroinvertebrates are the most favored in freshwater monitoring and are widely used to evaluate the water body health and condition (Conti, 2008; Meng *et al.*, 2009). The advantages of using biological indicators, particularly macroinvertebrates are biological communities reflect the overall ecological quality and provides a broad measurement of fluctuating environmental conditions. In addition, the result of biological monitoring is reliable and relatively inexpensive compared to toxicity testing (Iliopoulou-Georgudaki *et al.*, 2003).

The history of biological monitoring methods for assessing water quality began more than a century ago by Kolenati (1848) and Cohn (1853) (both quoted by Liebmann, 1962). However, such studies in Malaysia is still very limited and started relatively late with the earliest documented was in the early 90's (Khan, 1990; Arsad *et al.*, 2012). After year 2000, interest on this topic is gaining attention and grows, and example of studies can be seen in (Al-Shami *et. al.*, 2010; Al-Shami *et al.*, 2011; Zaiha *et al.*, 2015; Sharifah Aisyah *et al.*, 2015; Ghani *et al.*, 2016). More recently, in 2009, DID in collaboration with Universiti Sains Malaysia, has produced a Guideline for Using Macroinvertebrates for Estimation of Streams Water Quality. The guideline provides simple, inexpensive and easy approach to estimate water quality through the identification of freshwater macroinvertebrates. This government's effort is an initial step to the development of such studies in Malaysia and proving biological methods in the study of water quality began to be accepted.

1.2 Problem Statement

In recent years, there is a growing interest in protecting and managing rivers to be more sustainable, both ecologically and aesthetically. River management and technologies in Malaysia has experienced paradigm shift from the artificial and engineering-based to be more ecological friendly approaches. Although the method using the restoration concept managed to improve the water quality and amenity of

rivers, its usefulness for the reinstatement of the river biota is rarely observed when biological aspect is given little consideration. Many river rehabilitation projects tend to focus on improving water quality and beautification of the riverbanks without taking into account the importance of habitat to aquatic life. The notion that the abiotic factors (particularly water quality) have greater influence on the structure of stream community led to many river rehabilitation activities focusing more on altering the instream habitat. Simplification of stream habitat not only affects the function of ecosystem, but also structural ecosystem features that include species composition and diversity. Furthermore, placement of physical materials such as large woody debris (LWD), boulders and other materials as an effort to improve fish and other aquatic organism for habitat enhancement still not enough to obtain suitable environment for species population. Such practices are impossible to achieve the river rehabilitation goal without initial assessment of what are the factors leading to habitat degradation, disturbance in the watershed and how to cope the issue, and disruption factors of a physical and biological production of a system. Besides, the use of single classification index such as Water Quality Index (WQI) is inappropriate, especially when only six parameters are taken into account. Though WQI is a useful tool for instantaneous water quality benchmarking, it's still not enough to represent the overall health of river ecosystem. Hence, for a better decision making tool, the National Water Quality Standards (NWQS) should be referred to achieve the desired target class. Unfortunately, these standards are still poorly understood among the authorities, making the preservation efforts in the country will continue to be hampered. Moreover, physicochemical assessment provides only a partial perspective of water degradation and regarded as one of the limitation factor of this method. Thus far, the integrated assessment of ecosystem health in the country is very limited and lack of quantity with no comprehensive assessment has been developed. On top of that, assessment practices in Malaysia is much dependent on the physical and chemical based techniques, with biological methods are often ignored.

1.3 Aims and Objectives

This study aims to evaluate the ecological health of river ecosystem by the integration of a comprehensive assessment method. Therefore, the objectives are:

- i. To assess and describe the status of the river in the state of water quality and habitat quality, based on the integration of physicochemical, biological and habitat assessment approach.
- ii. To establish the relationship between macroinvertebrates composition, water quality and habitat quality condition.
- iii. To evaluate the ecological quality and describe biological communities through the development of multi-metric Index of Biological Integrity (IBI) using benthic macroinvertebrate.
- iv. To propose a baseline data of biological criteria for river health assessment that can be applied as a measurement of water resource management.

1.4 Scope of Study

This study focused on the description of the existing ecological environment of four rivers with different environmental gradient ecosystem, *viz.* Sg Ayer Hitam Besar as reference site and Sg Berasau, Sungai Mengkibol and Sg Melana as impacted sites. Three main processes explored in this study were consisted of physical characteristic (general characteristics that are important in influencing the river's aquatic ecology such as channel forms, instream habitats, substrates, riverbank vegetation and structure. Additional habitat attributes such as anthropogenic alterations to the river were briefly described), biological characteristics (focusing on the composition and abundance of macroinvertebrates

species) and chemical characteristics (documentation of existing conditions related to commonly observed water quality parameters). The study also investigated the correlation between the physicochemical attributes and variations in the macroinvertebrates assemblages. In addition, the main objective that needs to be achieved in the end of the study is to develop a benthic macroinvertebrate-based multimetrics for the bioassessment of stream water quality in Malaysia.

1.5 Significance of Study

Generally, the Water Quality Index (WQI) is used as a measurement of water quality status and classifying rivers in the country. The basic foundation of WQI is solely based on physicochemical parameters alone, comprises of biological oxygen demand, chemical oxygen demand, organics substance, suspended solids, alkalinity, temperature and dissolved oxygen, are then calculated and evaluated. Unfortunately, the physicochemical analysis cannot yield enough information on the overall health of the river ecosystem. Therefore, this approach is not strong enough as a main tool of evaluation status of aquatic ecosystems. This is because the index could not clearly define and justify the condition of aquatic life either may survive in the water body or vice versa. Therefore, integration of biological monitoring is necessary to complete the information. The integration between the physicochemical and biological assessment gives a broader perspective to describe the state of the environment.

There is still a lack of information regarding the integration of assessment tools to ascertain the health of the river ecosystem. Most biological monitoring studies conducted were involved simple qualitative system based on the absence or presence of indicator species according to the environmental gradient regardless of

the causal factors of the environment. Thus, the development of biological assessment methods such as multimetric index is appropriate to reflect all kinds of degradation and cumulative effect at the ecosystem level. A multimetric index consisted of several metrics related to the biological properties (i.e. taxa richness, taxa composition and pollution tolerance) that changes in expected way with increasing anthropogenic interference. This method can be widely used in the valuation of the ecosystem because it involves multiple measurements and provides comprehensive information related to the pre-determined criteria derived from reference conditions. Furthermore, the methods developed will facilitate the monitoring work of river water quality status in the country, and easily understood by general users.

REFERENCES

- Abbasi, T., and Abbasi, S. A. (2012). *Water quality indices*. Elsevier.
- Acuña, V., Muñoz, I., Giorgi, A., Omella, M., Sabater, F. and Sabater, S. (2005). Drought and post drought recovery cycles in an intermittent Mediterranean stream: structural and functional aspects. *Journal of the North American Benthological Society*. 24(4): 919 - 933.
- Agostinho, A. A., Thomaz, S. M. and Gomes, L. C. (2004). Threats for biodiversity in the floodplain of the Upper Parana River: effects of hydrological regulation by dams. *Ecohydrology and Hydrobiology*. 4: 267 - 280.
- Allan, D., Erickson, D. and Fay, J. (1997). The influence of catchment land use on stream integrity across multiple spatial scales. *Freshwater Biology*. 37(1): 149 - 161.
- Allan, J. D. (1995). *Stream ecology: structure and function of running waters*. London: Chapman & Hall.
- Al-Shami, S. A., Md Rawi, C. S., Ahmad, A. H., Abdul Hamid, S. and Mohd Nor, S. A. (2011). Influence of agricultural, industrial, and anthropogenic stresses on the distribution and diversity of macroinvertebrates in Juru River Basin, Penang, Malaysia. *Ecotox Environ Safe*. 74: 1195 – 1202.
- Al-Shami, S. A., Rawi, C. S. M., Hassan Ahmad, A. and Nor, S. A. M. (2010). Distribution of Chironomidae (Insecta: Diptera) in polluted rivers of the Juru River Basin, Penang, Malaysia. *J. Environ. Sci-China*. 22: 1718 - 1727.
- Andaya, B. W. and Andaya, L. Y. (2001). *A history of Malaysia*. London: Macmillan.
- Anderson, C. W. (2005). Turbidity (version 2). US Geological Survey Techniques of Water Resources Investigations Book 9, chapter A6, section 6.7. Retrieved from http://water.usgs.gov/owq/FieldManual/Chapter6/6.7_contents.html
- Angelier, E. (2003). *Ecology of Streams and Rivers*. Enfield, USA: Science Publishers Inc.

- APHA. (2005). *Standard methods for the examination of water and wastewater (21st ed.)*. Washington, DC: American Public Health Association.
- Araki, S., Numata, M., Wada, O. and Kankyokagakuziten (1985). *A Dictionary of Environmental Sciences*. Kagakudozin, Tokyo.
- Armitage, P. D., Moss, D., Wright, J. F. and Furse, M. T. (1983). The performance of a new biological water quality score system based on macroinvertebrates over a wide range of unpolluted running waters sites. *Water Res.* 17: 333 - 347.
- Arsad, A., Abustan, I., Rawi, C. S. M., & Syafalni, S. (2012). Integrating Biological Aspects into River Water Quality Research in Malaysia: An Opinion. *OIDA International Journal of Sustainable Development*, 4(02), 107-122.
- Avril, L., & Barten, P. K. (2007). *Land use effects on streamflow and water quality in the northeastern United States*. CRC Press.
- Aziz, H. A., Yusoff, M. S., Adlan, M. N., Adnan, N. H., and Alias S. (2004). Physico-chemical removal of iron from semi-aerobic landfill leachate by limestone filter. *Waste Management*. 24: 353 - 358.
- Azrina, M. Z., Yap, C. K., Rahim Ismail, A., Ismail, A. and Tan, S. G. (2006). Anthropogenic impacts on the distribution and biodiversity of benthic macroinvertebrates and water quality of the Langat River, Peninsular Malaysia. *Ecotoxicology and environmental safety*. 64(3): 337 - 47.
- Badcock, R. (1953). Observation of oviposition under water of the aerial insect *Hydropsyche angustipennis* (Curtis) (Trichoptera). *Hydrobiologia*. 5(1): 222 - 225.
- Bae, M. J., Chon, T. S. and Park, Y. S. (2014). Characterizing differential responses of benthic macroinvertebrate communities to floods and droughts in three different stream types using a Self-Organizing Map. *Ecohydrology*. 7(1): 115 - 126.
- Barbour, M. T., Gerritsen, J., Griffith, G. E., Frydenborg, R., Mccarron, E., White, J. S. and Bastian, M. L. (1996). A framework for biological criteria for Florida streams using benthic macroinvertebrates.. *J. North Am. Benthol. Soc.* 15: 185 - 211.
- Barbour, M. T., Stribling, J. B. and Verdonschot, P. F. M. (2006). The Multihabitat Approach of USEPA's Rapid Bioassessment Protocols: Benthic Macroinvertebrates. *Limnetica*. 25(3): 839 - 850.

- Barton, D. R., Taylor, W. D. and Biette, R. M. (1985). Dimensions of riparian buffer strips required to maintain trout habitat in Southern Ontario streams. *N. Am. J. Fish. Manage.* 5: 364 - 378.
- Bauder, J. W., & Pearson, K. E. (2002). Justification for numeric standards of salinity and sodicity of water to be considered for beneficial use for irrigation. *Departement of Land Resources and Environmental Sciences, Montana State University-Bozeman, MT.*
- Behmer, D. J. and Hawkins, C. P. (1986). Effects of overhead canopy on macroinvertebrate production in a Utah stream. *Freshwater Biol.* 16: 287 - 300.
- Benke, A. C. (2001). Importance of flood regime to invertebrate habitat in an unregulated river floodplain ecosystem. *J. N. Am. Benthol. Soc.* 20: 225 - 240.
- Benstead, J. P. and Pringle, C. M. (2004). Deforestation alters the resource base and biomass of endemic stream insects in eastern Madagascar. *Freshwater Biology.* 49(4): 490 - 501.
- Benstead, J. P., Douglas, M. M. and Pringle, C. M. (2003a). Relationships of stream invertebrate communities to deforestation in eastern Madagascar. *Ecological Applications.* 13: 1473 - 1490.
- Benstead, J. P., De Rham, P. H., Gattolliat, J. L., Gibon, F. M., Loiselle, P. V., Sartori, M., Sparks, J. S., and Stiasny, M. L. J. (2003b). Conserving Madagascar's freshwater biodiversity. *BioScience* 53, 1101–1111.
- Bilby, R. E. and Ward, J. W. (1991). Characteristics and function of large woody debris in streams draining old growth, clear cut, and second-growth forests in southwestern Washington. *Can. J. Fish. Aquat. Sci.* 48: 2499 - 2508.
- Bishop, J. E. (1973). *Limnology of Small Malayan river, Sungai Gombak.* The Hague, Netherlands: Dr Junk W. Publishers.
- Boon, P. J., Davies, B. R. and Petts, G. E. (2000). *Global perspectives on river conservation: science, policy and practice.* New York, USA: John Wiley & Sons.
- Bonada, N., Prat, N., Resh, V. H., and Statzner, B. (2006). Developments in aquatic insect biomonitoring: a comparative analysis of recent approaches. *Annu. Rev. Entomol.*, 51, 495-523.
- Boulton, A. J. (2003). Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages. *Freshwater Biology.* 48(7): 1173 - 1185.

- Boulton, A. J., Peterson, C. G., Grimm, N. B. and Fisher, S. G. (1992). Stability of an aquatic macroinvertebrate community in a multiyear hydrologic disturbance regime. *Ecology*. 73: 2192 - 2207.
- Brewin, P. A., Buckton, S. T. and Ormerod, S. J. (2000). The seasonal dynamics and persistence of stream macroinvertebrates in Nepal: do monsoon floods represent disturbance? *Freshwater Biology*. 44: 581 - 594
- Brown, A. V. and Matthews, W. J. (1995). Stream ecosystem of the central United States. In Cushing, C. E., Cummins, K. W. and Minshall, G. W. (Eds). *Ecosystems of the World* (pp. 79 - 110). New York: Elsevier.
- Buffagni, A., Erba, S., Cazzola, M., Murray-Bligh, J., Soszka, H., & Genoni, P. (2006). The STAR common metrics approach to the WFD intercalibration process: Full application for small, lowland rivers in three European countries. In *The Ecological Status of European Rivers: Evaluation and Intercalibration of Assessment Methods* (pp. 379-399). Springer Netherlands.
- Bunn, S. E. and Arthington, A. H. (2002). Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environmental Management*. 30: 492 - 507.
- Cairns, J. Jr. and Pratt, J. R. (1993). A history of biological monitoring using benthic macroinvertebrates. In: Rosenberg, D. M and Resh, V. H. (Eds.). *Freshwater Biomonitoring and Benthic Macroinvertebrates* (pp. 10 - 27). New York: Chapman and Hall.
- Caldecott, J. (1996). *Designing Conservation Projects*. Cambridge, UK: Cambridge Univ. Press.
- Canadian Council of Ministers of the Environment (2006). *Developing Biocriteria as a Water Quality Assessment Tool in Canada: Scoping Assessment*. Canada: Canadian Council of Ministers of the Environment.
- Carter, J. L., and Resh, V. H. (2001). After site selection and before data analysis: sampling, sorting, and laboratory procedures used in stream benthic macroinvertebrate monitoring programs by USA state agencies. *Journal of the North American Benthological Society*, 20(4), 658-682.
- Castella, E., Adalsteinsson, H., Brittain, J. E., Gislason, G. M., Lehmann, A., Lencioni, V., Lods-Crozet, B., Maiolini, B., Milner, A. M., Olafsson, J. O., Saltveit, S. J. and Snook, D. L. (2001). Macrobenthic invertebrate richness and

- composition along a latitudinal gradient of European glacier-fed streams. *Freshwater Biology*. 46: 1811 - 1831.
- Chambers, P. A., Lacoul, P., Murphy, K. J., and Thomaz, S. M. (2008). Global diversity of aquatic macrophytes in freshwater. *Hydrobiologia*, 595(1), 9-26.
- Chan Ngai Weng, Abdullah, A. L., Ibrahim, A. L. and Ghazali, S. (2003). River Pollution and Restoration Towards Sustainable Water Resources Management in Malaysia. *National Seminar on Society, Space and Environment in a Globalised World: Prospects & Challenges*. 29 - 30 May. The City Bayview Hotel, Penang, 208 - 219.
- Chesapeake Bay Program. (2012). Water Clarity. In The Bay Ecosystem. Retrieved from <https://www.chesapeakebay.net/discover/bayecosystem/waterclarity>.
- Chiew, F. C., and Shamsudin, M. N. (2007). Education for sustainable development: An overview of Malaysia's experiences in sustainable agricultural development and environmental conservation. *Obihiro Univ. for Agriculture and Veterinary Medicine*, 53-70.
- Chow, M. F., Yusop, Z., and Shirazi, S. M. (2013). Storm runoff quality and pollutant loading from commercial, residential, and industrial catchments in the tropic. *Environmental monitoring and assessment*, 185(10), 8321-8331.
- Christensen, M. S. (1993). The artisanal fishery of the Mahakam River floodplain in East Kalimantan, Indonesia. *Journal of Applied Ichthyology*. 9(3 - 4): 185 - 192.
- Clements, W. H. (1994). Benthic invertebrate community responses to heavy metals in the Upper Arkansas River Basin, Colorado. *J. N. Am. Benthol. Soc.* 13: 30 - 44.
- Collier, M. R., Webb, R. H. and Schmidt, J. C. (1996). Dams and rivers: a primer on the downstream effects of dams. Circular 1126, U. S. Geological Survey, Menlo Park, CA.
- Collins, D. E. (1999). Turtles in peril: the China crisis. *Vivarium*. 10(4): 6 - 9.
- Conti, M. E. (2008). *Biological monitoring: theory & applications: bioindicators and biomarkers for environmental quality and human exposure assessment* (Vol. 17). WIT Press.
- Couceiro, S. R. M., Hamada, N., Luz, S. L. B., Forsberg, B. R. and Pimentel, T. P. (2007). Deforestation and sewage effects on aquatic macroinvertebrates in urban streams in Manaus, Amazonas, Brazil. *Hydrobiologia*. 575: 271 - 284.

- Covich, A. P., Palmer, M. A., and Crowl, T. A. (1999). The role of benthic invertebrate species in freshwater ecosystems: zoobenthic species influence energy flows and nutrient cycling. *BioScience*, 49(2), 119-127.
- Cummins, K. W., Wilzbach, M. A., Gates, D. M., Perry, J. B. and Taliaferro, W. B. (1989). Shredders and riparian vegetation. *Bioscience*. 39: 24 - 30.
- Dahm, C. N., Baker, M. A., Moore, D. I. and Thibault, J. R. (2003). Coupled biogeochemical and hydrological responses of streams and rivers to drought. *Freshwater Biology*. 48(7): 1219 - 1231.
- Daniel, M.H., Montebelo, A.A., Bernardes, M.C., Ometto, J.P., Camargo, P.B.D., Krusche, A.V., Ballester, M.V., Victoria, R.L. and Martinelli, L.A., (2002). Effects of urban sewage on dissolved oxygen, dissolved inorganic and organic carbon, and electrical conductivity of small streams along a gradient of urbanization in the Piracicaba river basin. *Water, Air, & Soil Pollution*, 136(1), pp.189-206.
- Davis, J. M. and Cornwell, D. A. (1991). *Introduction to Environmental Engineering*. Boston Massachusetts: P.W.S.
- de Bikuña, B.G., Arrate, J., Martínez, A., Agirre, A., Azpiroz, I. and Manzanos, A., (2017). Development of a fish-based index to assess the ecological status of oceanic-temperate streams in the Northern Iberian Peninsula. *Fundamental and Applied Limnology/Archiv für Hydrobiologie*.
- Degefe, G., Tamire, G., Hassen, S.M. and Haile, A., (2017). Evaluation of the Use of Some Benthic Macro-invertebrate Multimetric Indices Associations with Environmental Variables in Ecological Assessment of Some Ethiopian Rivers. *Hydro Nepal: Journal of Water, Energy and Environment*, 20, pp.49-54.
- De Pauw, N., Gabriels, W., and Goethals, P. L. (2006). River monitoring and assessment methods based on macroinvertebrates. *Biological monitoring of rivers*, 113-134.
- Department of Irrigation and Drainage Malaysia (2004). *Towards the Making of Rivers Rich in Nature. Foundation for riverfront improvement and restoration (Eds.)*. Kuala Lumpur: Department of Irrigation and Drainage Malaysia.
- DeShon, J. E. (1995). Development and application of the invertebrate community index (ICI). In Davis, W. S. and Simon, T. P. (Eds). *Biological assessment and criteria: Tools for water resource planning and decision making* (pp. 217 - 243). Boca Raton, Florida: Lewis Publishers.

- Diehl, S. (1992). Fish predation and benthic community structure: the role of omnivory and habitat complexity. *Ecology*. 1646 - 1661.
- Donohue, I., McGarrigle, M. L. and Mills, P. (2006). Linking catchment characteristics and water chemistry with the ecological status of Irish rivers. *Water Res.* 40: 91 - 98.
- Dudgeon, D. (1992). Endangered ecosystems: a review of the conservation status of tropical Asian rivers. *Hydrobiologia*. 248: 167 - 191.
- Dudgeon, D. (2000a). The ecology of tropical Asian rivers and streams in relation to biodiversity conservation. *Annual Review of Ecology and Systematics*. 31: 239 - 263.
- Dudgeon, D. (2000b). Riverine biodiversity in Asia: a challenge for conservation biology. *Hydrobiologia*. 418: 1 - 13.
- Dudgeon, D. (2003). The contribution of scientific information to the conservation and management of freshwater biodiversity in tropical Asia. *Hydrobiologia*. 500: 295 - 314.
- Dudgeon, D. (2006). The impacts of human disturbance on stream benthic invertebrates and their drift in North Sulawesi, Indonesia. *Freshw. Biol.* 51: 1710 - 1729.
- Dudgeon, D. (2008). *Tropical stream ecology*. California: Elsevier Inc.
- Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z., Knowler, D. J., Leveque, C., Naiman, R. J., Prieur-Richard, A. H., Soto, D., Stiassny, M. L. and Sullivan, C. A. (2006). Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews*. 81:163 - 182.
- Dugan, P. J., Barlow, C. and Agostinho, A. A. (2010). Fish migration, dams and loss of ecosystem services in the Mekong Basin. *Ambio*. 39: 344 - 348.
- Elwood, J. W., J. D. Newbold, R.V. O'Neill, and W. Van Winkle. (1983). Resource spiraling: An operational paradigm for analyzing lotic ecosystems. Pp. 3–27 in Thomas D. Fontaine III and Steven M. Bartell, eds., *Dynamics of Lotic Ecosystems*. Ann Arbor Science Publishers, Ann Arbor, Mich. 494 pp.
- Environment Agency (2003). River Habitat Survey in Britain and Ireland. Field Survey Guidance Manual: Bristol.
- EPA (2012a). 5.5 Turbidity. In Water: Monitoring & Assessment. Retrieved from <http://water.epa.gov/type/rsl/monitoring/vms55.cfm>.

- EPA (2012b). 5.8 Total Solids. In *Water: Monitoring and Assessment*. Retrieved from <http://water.epa.gov/type/rsl/monitoring/vms58.cfm>.
- Feldman, R. S. (2001). Taxonomic and size structures of phytophilous macroinvertebrate communities in Vallisneria and Trapa beds of the Hudson River, New York. *Hydrobiologia*, 452(1-3), 233-245.
- Flecker, A. S. and Feifarek, B. (1994). Disturbance and temporal variability of invertebrate assemblages in two Andean streams. *Freshwater Biology*. 31: 131-142.
- Fossati, O., Wasson, J. G., Cecile, H., Salinas, G. and Marin, R. (2001). Impact of sediment releases on water chemistry and macroinvertebrate communities in clear water Andean streams (Bolivia). *Archive für Hydrobiologie*. 151: 33 - 50.
- Frissell, C. A., Liss, W. J., Warren, C. E., and Hurley, M. D. (1986). A hierarchical framework for stream habitat classification: viewing streams in a watershed context. *Environmental management*, 10(2), 199-214.
- Fritz, K. M. and Dodds, W. K. (2004). Resistance and resilience of macroinvertebrate assemblages to drying and flood in a tallgrass prairie stream system. *Hydrobiologia*. 527: 99 - 112.
- Fulazzaky, M. A., Seong, T. W., and Masirin, M. I. M. (2010). Assessment of water quality status for the Selangor River in Malaysia. *Water, Air, and Soil Pollution*, 205(1-4), 63-77.
- Gardiner, J. L. (1990). *River projects and conservation: a manual for holistic appraisal*. John Wiley and Sons Ltd.
- Ghani, W. M. H. W. A., Rawi, C. S. M., Hamid, S. A., & Al-Shami, S. A. (2016). Efficiency of Different Sampling Tools for Aquatic Macroinvertebrate Collections in Malaysian Streams. *Tropical life sciences research*, 27(1), 115.
- Ghani, A. A. B., Zakaria, N. A., Chang, C. K. and Ainan, A. (2008). Sustainable Urban Drainage Systems (SUDS) - Malaysian experiences. *Proc. 11th International Conference on Urban Drainage*. 31 August - 5 September. Edinburgh, Scotland: UK, 1 - 10.
- Global Water Partnership (2012). *GWP in action 2012 Annual Report*. Stockholm, Sweden: GWP.
- Gopal, B. (2002). Conserving biodiversity in Asian wetlands: issues and approaches. In *The Asian Wetlands: Bringing Partnerships into Good Wetland Practices*, edited by Ali Ahyaudin, Che Salmah Md Rawi, Mashhor Mansor, Reiko

- Nakamura, Sundari Ramakrishna & Taej Mundkur. Pulau Pinang, Malaysia: Penerbit Universiti Sains Malaysia.
- Gordon, N. D., McMahon, T. A., Finlayson, B. L., Gippel, C. J. and Nathan, R. J. (2013). *Stream hydrology: an introduction for ecologists* (2nd ed). England: John Wiley & Sons.
- Gordon, N.D., McMahon, T.A. and Finlayson, B.L., (2004). *Stream hydrology: an introduction for ecologists*. John Wiley and Sons.
- Government to investigate on fishes found dead in river (2015, April 25). *The Star Online*. Retrieved April 27, 2015, from <https://translate.google.com/#en/ms/retrieve>.
- Greathouse, E. A., Pringle, C. M. and Holmquist, J. G. (2006). Conservation and management of migratory fauna: dams in tropical streams in Puerto Rico. *Aquatic Conservation: Marine and Freshwater Systems*. 16: 695 - 712.
- Grenier, P. (1949). Contribution à l'étude biologique des Simuliides de France. *Physiol. Comp. Oekol.* 1: 165 - 330.
- Griffith, M. B., Hill, B. H., McCormick, F. H., Kaufmann, P. R., Herlihy, A. T. and Selle, A. R. (2005). Comparative application of indices of biotic integrity based on periphyton, macroinvertebrates, and fish to southern Rocky Mountain streams. *Ecological Indicators*, 5(2), 117-136.
- Hall, R. J., Likens, G. E., Fiance, S. B. and Hendrey, G. R. (1980). Experimental acidification of a stream in the Hubbard Brook experimental forest, New Hampshire. *Ecology*. 61: 976 - 989.
- Halpern, M., Gasith, A., and Broza, M. (2002). Does the tube of a benthic chironomid larva play a role in protecting its dweller against chemical toxicants?. *Hydrobiologia*, 470(1-3), 49-55.
- Hamidi Abdul Aziz (1999). Rapid field assessment of organic pollution with a family level biotic index. *Journal North America Benthological Society*. 7(1): 65 - 68.
- Hamza-Chaffai, A. (2014). Usefulness of bioindicators and biomarkers in pollution biomonitoring. *International Journal of Biotechnology for Wellness Industries*, 3(1), 19.
- Hanski, I., Koivulehto, H., Cameron, A. and Rahagalala, P. (2007). Deforestation and apparent extinctions of endemic forest beetles in Madagascar. *Biology Letters*. 3(3):344 - 347.

- Harun, S., Baker, A., Bradley, C., and Pinay, G. (2016). Spatial and seasonal variations in the composition of dissolved organic matter in a tropical catchment: the Lower Kinabatangan River, Sabah, Malaysia. *Environmental Science*
- Harun, S., Al-shami, S. A., Dambul, R., Mohamed, M., and Abdullah, M. H. (2015). Water Quality and Aquatic Insects Study at the Lower Kinabatangan River Catchment, Sabah: In Response to Weak La Niña Event. *Sains Malaysiana*, 44(4): 545-558.
- Hauer, F.R., Stanford, J.A. (1982). Ecological responses of hydroptychid caddisflies to stream regulation. *Can. J. Fish. Aquat. Sci.* 39, 1235-1242.
- Healey, M. (1984). Fish predation on aquatic insects.
- Hellawell, J. M. (1986). *Biological indicators of freshwater pollution and environmental management*. New York: Elsevier.
- Hilsenhoff, W. (1988). Rapid field assessment of organic pollution with a family-level biotic index. *J. N. Am. Benthol. Soc.* 7: 65 - 68.
- Hoang, H. (2009). *Monitoring and assessment of macroinvertebrate communities in support of river management in northern Vietnam*. Ghent University.
- Hose, G., Turak, E. and Waddell, N. (2004). Reproducibility of AUSRIVAS rapid bioassessments using macroinvertebrates. *Journal of the North American Benthological Society*. 23: 126 - 139.
- Hughes, R. M., Paulsen, S. G. and Stoddard, J. L. (2000). EMAP-Surface Waters: a multi-assemblage, probability survey of ecological integrity in the USA. *Hydrobiologia*, 422, 429-443.
- Huet, M. (1954). Biologie, profils en long et en travers des eaux courantes. *Bulletin français de Pisciculture*, (175), 41-53.
- Hughes, R. M., Larsen, D. P. and Omernik, J. M. (1986). Regional reference sites: A method for assessing stream potentials. *Environmental Management*. 10: 629 - 635.
- Hussain Qazi A (2011). *An Ecological Study of Doodhganga and its Drainage Basin - A Lotic System of Kashmir*. PhD. Thesis, P. G. Department of Environmental Science, University of Kashmir, Srinagar.
- Huston, M.A. (1994). *Biological Diversity*. Cambridge, UK: Cambridge University Press.

- Hutan Lipur Gunung Pulai 1 mungkin dibuka semula (2014, Mac 10). *Sinar Harian*. Retrieved May 14, 2015, from <http://www.sinarharian.com.my/edisi/johor/hutan-lipur-gunung-pulai-1-mungkin-dibuka-semula-1.258859>.
- Hynes, H. B. N. (1970). *The ecology of running waters*. Liverpool: Liverpool University Press.
- Iliopoulou-Georgudaki, J., Kantzaris, V., Katharios, P., Kaspiris, P., Georgiadis, T. and Montesantou, B., (2003). An application of different bioindicators for assessing water quality: a case study in the rivers Alfeios and Pineios (Peloponnisos, Greece). *Ecological indicators*, 2(4), pp.345-360.
- Illies, J., & Lazare, B. (1963). Problèmes et méthodes de la classification et de la zonation écologique des eaux courantes, considérées surtout-du point de vue faunistique.
- Inisiatif pemuliharaan sungai memberangsangkan (2013, September 5). *Sinar Harian*. Retrieved May 26, 2016, from <http://www.sinarharian.com.my/rencana/inisiatif-pemuliharaan-sungai-memberangsangkan-1.198987>.
- Ismail, A. and Mohamad, A. B. (1994). *Ekologi air tawar*. Kuala Lumpur: Dewan Bahasa dan Pustaka.
- Iwata, T., Nakano, S. and Inoue, M.(2003). Impacts of past riparian deforestation on stream communities in a tropical rain forest in Borneo. *Ecological Applications*. 13: 461–473.
- Jabatan Perangkaan Malaysia (2013). *Kompendium perangkaan alam sekitar Malaysia*. Kuala Lumpur: Jabatan Perangkaan Malaysia.
- Jergentz, S., Mugni, H., Bonetto, C. and Schulz, R. (2004). Runoff-related endosulfan contamination and aquatic macroinvertebrate response in rural basins near Buenos Aires, Argentina. *Archives of Environmental Contamination and Toxicology*. 46: 345 - 352.
- Joshi, P. C., Negi, R. K. and Negi, T. (2007). Seasonal variations in benthic macroinvertebrates and their correlation with the environmental variables in a freshwater stream in Garhwal Region (India). *Life Sci. J.* 4(4): 85 - 89.
- Judy Jr, R. D., Seeley, P. N., Murray, T. M., Svirsky, S. C., Whitworth, M. R. and Ischinger, L. S. (1982). National fisheries survey. *US Fish and Wildlife Service FWS/OBS*, 84(06).

- Junk, W. J., Bayley, P. B., and Sparks, R. E. (1989). The flood pulse concept in river-floodplain systems. *Canadian special publication of fisheries and aquatic sciences*, 106(1), 110-127.
- Karr, J. R. and Dudley, D. R. (1985). Ecological perspectives on water quality goals. *Environmental Management*. 5: 55 - 68.
- Karr, J. R., Fausch, K. D., Angermeier, P. L., Yant, P. R. and Schlosser, I. J. (1986). *Assessing biological integrity in running waters: A method and its rationale*. Illinois Natural History Survey, Champaign, IL, Special publication 5.
- Kasangaki, A., Babaasa, D., Efitre, J., McNeilage, A. and Bitariho, R. (2006). Links between anthropogenic perturbation and benthic macroinvertebrate assemblages in Afromontane forest streams in Uganda. *Hydrobiologia*. 563: 231 - 245.
- Kenney, M. A., Sutton-Grier, A. E., Smith, R. F., and Gresens, S. E. (2009). Benthic macroinvertebrates as indicators of water quality: the intersection of science and policy. *Terrestrial Arthropod Reviews*. 2(2): 99 - 128.
- Khan, I. S. A. (1990). Assessment of water pollution using diatom community structure and species distribution—A case study in a tropical river basin. *Internationale Revue der gesamten Hydrobiologie und Hydrographie*, 75(3), 317-338.
- Khazanah Nasional (2006). *Comprehensive Development Plan for South Johor Economic Region 2006 - 2025*. Kuala Lumpur: Khazanah Nasional.
- Khopkar, S. M. (2007). *Environmental pollution monitoring and control*. New Delhi, India: New Age International.
- Laabs, V. W., Amelung, A. A., Pinto, M., Wantzen, K. M., da Silva, C. J. and Zech, W. (2002). Pesticides in surface water, sediments and rainfall of the north-eastern Pantanal basin, Brazil. *Journal of Environmental Quality*. 31: 1636 - 1648.
- Lake, P. S. (2000). Disturbance, patchiness, and diversity in streams. *Journal of the North American Benthological Society*. 19: 573 - 592.
- Lake, P. S. (2003). Ecological effects of perturbation by drought in flowing waters. *Freshwater Biology*. 48(7): 1161 - 1172.
- Lancaster, J., Real, M., Juggins, S., Monteith, D. T., Flower, R. J., & Beaumont, W. R. C. (1996). Monitoring temporal changes in the biology of acid waters. *Freshwater Biology*, 36(1), 179-201.

- Larned, S. T., Schmidt, J., Datry, T. K., Konrad, C. P., Dumas, J. K. and Diettrich, J. C. (2011). Longitudinal river ecohydrology: flow variation down the lengths of alluvial rivers. *Ecohydrology*. 4(4): 532 - 548.
- LeCraw, R. and Mackreth, R. (2010). Sources of small-scale variation in the invertebrate communities of headwater streams. *Freshwater Biol.* 55: 1219 - 1233.
- Lenat, D. R. (1988). Water quality assessment of streams using a qualitative collection method for benthic macroinvertebrates. *J. North Am. Benthol. Soc.* 7: 222 - 233.
- Lenntech, (2013). Acids and Alkalis in freshwater. In Water Treatment Solutions. Retrieved from <http://www.lenntech.com/aquatic/acids-alkalis.htm>
- Lessard, J. L., Hayes, D. B. (2003). Effects of elevated water temperature on fish and macroinvertebrate communities below small dams. *River Research and Applications*. 19: 721 - 732.
- Li, J., Herlihy, A. T., Gerth, W., Kaufmann, P., Gregory, S., Urquhart, S. and Larsen, D. P. (2001). Variability in stream macroinvertebrates at multiple spatial scales. *Freshwater Biol.* 46: 87 - 97.
- Li, Y., Li, Y., Xu, Z., and Li, L. (2016). Assessment of the Huntai River in China using a multimetric index based on fish and macroinvertebrate assemblages. *Journal of Freshwater Ecology*, 31(2), 169-190.
- Liebmann, H. (1962). Handbuch der Frischwasser und Abwasser biologie. In *Handbuch der frischwasser und abwasser biologie*. R. Oldenbourg.
- Lim, S. H., Abdullah, S. and Mohd. Rozali, O. (2001). Kesihatan ekosistem Sungai Labu dari aspek kualiti airnya. *Malays. J. Anal. Sci.* 7(1): 157 - 168.
- Ling, T. Y., Soo, C. L., Sivalingam, J. R., Nyanti, L., Sim, S. F., and Grinang, J. (2016). Assessment of the Water and Sediment Quality of Tropical Forest Streams in Upper Reaches of the Baleh River, Sarawak, Malaysia, Subjected to Logging Activities. *Journal of Chemistry*, 2016.
- Liow, L. H., Sodhi, N. S. and Elmquist, T. (2001). Bee diversity along a disturbance gradient in tropical lowland forests of south-east Asia. *Journal of Applied Ecology*. 38:180 - 192.
- Lorenz, C. M., Van Dijk, G. M., Van Hattum, A. G. M., and Cofino, W. P. (1997). Concepts in river ecology: implications for indicator development. *Regulated Rivers: Research & Management*. 13(6): 501 - 516.

- Macan, T. T. (1977). The influence of predation on the composition of freshwater animal communities. *Bio. Rev.* 52: 45 - 70.
- Macedo, D.R., Hughes, R.M., Ferreira, W.R., Firmiano, K.R., Silva, D.R., Ligeiro, R., Kaufmann, P.R. and Callisto, M., (2016). Development of a benthic macroinvertebrate multimetric index (MMI) for Neotropical Savanna headwater streams. *Ecological Indicators*, 64, pp.132-141.
- Macionis, J. J. and Parrillo, V. N. (2004). *Cities and Urban Life*. Upper Saddle River, NJ: Prentice-Hall.
- Mackenthun, K. M. (1969). The practice of water pollution biology. FWPCA: Washington, DC, USA
- Magurran, A. E. (2004). Measuring biological diversity. Oxford, U.K: Blackwell Publishing.
- Mandaville, S. M. (2002). *Benthic Macroinvertebrates in Freshwaters - Taxa Tolerance Values, Metrics, and Protocols*. Soil & Water Conservation Society of Metro Halifax.
- Manios, T., Stentiford, E. I. and Millner, P. A. (2002). The removal of indicator microorganisms from primary treated wastewater in subsurface reed beds using different substrates. *Environmental Technology*. 23: 767 - 774.
- March, J. G., Benstead, J. P., Pringle, C. M. and Scatena, F. N. (2003). Damming tropical island streams: problems, solutions and alternatives. *BioScience*. 53: 1069 - 1078.
- Margalef, R. (1951). Role des entomostraces dans la régénération des phosphates. *Verh. Internat. Verein. Limnol.* 11: 246 - 247.
- Marziali, L., Armanini, D. G., Cazzola, M., Ebra, S., Toppi, E., Buffagni, A. and Rossaro, B. (2010). Response of chironomid larvae (Insecta: Diptera) to ecological quality in Mediterranean river mesohabitats (South Italy). *River Res. Appl.* 26: 1036 - 1051.
- Mazlin, M., Ismail, B. and Ng, C. H. (2001). Pengelasan kualiti air 1998: Dari pantai ke Kuala Linggi. *Malays. J. Anal. Sci.* 6(1): 178 - 187.
- Mazor R.D., Reynoldson T.B. Rosenberg D.M. and Resh V.H. (2006) Effects of biotic assemblage, classification, and assessment method on bioassessment performance. *Canadian Journal of Fisheries and Aquatic Sciences*, 63, 394–411
- McCafferty, W. P. (1998). *Aquatic entomology: the fishermen's and ecologists' illustrated guide to insects and their relatives*. Jones and Bartlett.

- McIntosh, M. D., Benbow, M. E. and Burky, A. J. (2002). Effects of stream diversion on riffle macroinvertebrate communities in a Maui, Hawaii, stream. *River Research and Applications*. 18: 569 - 581.
- McKenzie, D. H., Hyatt, D. E. and McDonald, V. J., (Eds.) (1992). *Ecological Indicators*. Volume 1 & 2. New York (NY): Elsevier Applied Science.
- Md Rawi, C. S., Al-Shami, S. A., Madrus, M. R., and Ahmad, A. H. (2014). Biological and ecological diversity of aquatic macroinvertebrates in response to hydrological and physicochemical parameters in tropical forest streams of Gunung Tebu, Malaysia: implications for ecohydrological assessment. *Ecohydrology*. 7(2): 496-507.
- Md Rawi, C. S., Al-Shami, S. A., Madrus, M. R. and Ahmad, A. H. (2013). Local effects of forest fragmentation on diversity of aquatic insects in tropical forest streams: implications for biological conservation. *Aquatic Ecology*. 47(1): 75 - 85.
- Meixler, M. S., Bain, M. B. and Todd Walter, M. (2009). Predicting barrier passage and habitat suitability for migratory fish species. *Ecological Modeling*. 220 (20): 2782 - 2791.
- Meng, W., Zhang, N., Zhang, Y., and Zheng, B. (2009). Integrated assessment of river health based on water quality, aquatic life and physical habitat. *Journal of Environmental Sciences*, 21(8), 1017-1027.
- Menhinick, E. F. (1964). A Comparison of Some Species-Individuals Diversity Indices Applied to Samples of Field Insects. *Ecology*. 45: 859 - 861.
- Merriam-Webster (2013). pH. In Merriam-Webster Dictionary. Retrieved from <http://www.merriam-webster.com/dictionary/ph>.
- Merritt, R.W. and Cummins, K.W. (1996). *An Introduction to the Aquatic Insects of North America. Third Edition*. Dubuque, U.S.A.: Kendall/ Hunt Publishing Company.
- Metcalf-Smith, J. L. (1994). Biological water-quality assessment of rivers: use of macroinvertebrate communities. In Calow, P. and Petts, G. E. (Eds.). *The rivers handbook Vol.II* (pp. 144 - 169). U.K : Blackwell Scientific Publications.
- Metcalf, J. L. (1989). Biological water quality assessment of running waters based on macroinvertebrate communities: history and present status in Europe. *Environmental pollution*, 60(1), 101-139.

- Meyer, J. L. and O'Hop, J. (1983). Leaf-shredding insects as a source of dissolved organic carbon in headwater streams. *American Midland Naturalist*. 175 - 183.
- Miller, R. R., Williams, J. D. and Williams, J. E. (1989). Extinctions of North American fishes during the past century. *Fisheries*. 14(6), 22 - 38.
- Milne, T. A., Briggs, S. D., Brock, H. W., Martin, M. E., Gibbs, D., Allis, C. D. and Hess, J.L. (2002). MLL targets SET domain methyltransferase activity to Hox gene promoters. *Mol. Cell*. 10: 1107 - 1117.
- Minshall, G. W., Cummins, K. W., Petersen, R. C., Cushing, C. E., Bruns, D. A., Sedell, J. R., and Vannote, R. L. (1985). Developments in stream ecosystem theory. *Canadian Journal of Fisheries and Aquatic Sciences*, 42(5), 1045-1055.
- Minshall, G. W., Cummins, K. W., Petersen, R. C., Cushing, C. E., Bruns, D. A., Sedell, J. R. and Vannote, R. L. (1985). Developments in stream ecosystem theory. *Can. J. Fish. Aquat. Sci.* 42: 045 - 1055.
- Mol, J. H. and Ouboter, P. E. (2004). Downstream effects of erosion from small-scale gold mining on the instream habitat and fish community of a small Neotropical rainforest stream. *Conservation Biology*. 18: 201 - 214.
- Montuelle, B., Dorigo, U., Bérard, A., Volat, B., Bouchez, A., Tlili, A., Gouy, V. and Pesce, S., (2010). The periphyton as a multimetric bioindicator for assessing the impact of land use on rivers: an overview of the Ardières-Morcille experimental watershed (France). *Hydrobiologia*, 657(1), pp.123-141.
- Moretti, G., Gianotti, F. S. (1962). Der Einfluss der Strömung auf die Verteilung der Trichopteren *Agapetus gr. Fuscipes* CURT. und *Silo gr. Nigricornis* PICT. Symposium über den Einfluss der Strömungsgeschwindigkeit: 467 - 484.
- Morse, J. C., Bae, Y. J., Munkhjargal, G., Sangpradup, N., Tanida, K., Vshivkova, T. S., Wang, B., Yang, L. and Yule, C. M. (2007). Freshwater biomonitoring with macroinvertebrates in East Asia. *Front Ecol Environ*. 5: 33 - 42.
- Moulton, T. P. and Wantzen, K. M. (2006). Conservation of tropical streams - special questions or conventional paradigms? *Aquatic Conservation*. 16: 659 - 663.
- Moya, N., Tomanova, S. and Oberdorff, T. (2007). Initial development of a multimetric index based on aquatic macroinvertebrates to assess stream condition in the upper Isidoro-Sécure Basin, Cochabamba, Bolivia. *Hydrobiologia*. 589(1): 107 - 116.

- Munoz, I. (2003). Macroinvertebrate community structure in an intermittent and a permanent Mediterranean stream (NE Spain). *Limnetica*. 22(3-4): 107 - 116.
- Murdoch, W. W. and Bence, J. (1987). General predators and unstable prey populations. In Kerfoot, W. C. and Sih, A. (Eds.). *Predation: Direct and Indirect Impacts on Aquatic Communities* (pp. 17 - 30). Univ. Press of New England, Hanover and London.
- Murphy, S. (2007). General Information on Solids. In City of Boulder: USGS Water Quality Monitoring. Retrieved from <http://bcn.boulder.co.us/basin/data/NEW/info/TSS.html>.
- Naiman, R. J., Lonzarich, D. G., Beechie, T. J. and Ralph, S.C., (1992). General principles of classification and the assessment of conservation potential in rivers. In: Boon, P.J., Calow, P., Petts, G.E. (Eds), *River Conservation and Management* (pp. 93-123). John Wiley: Chichester.
- Naiman, R. and Bilby, R. E. (Eds.) (2001). *River ecology and management: lessons from the Pacific coastal ecoregion*. Springer Science & Business Media.
- Ndaruga, A. M., Ndiritu, G. G., Gichuki, N. N. and Wamicha, W.N. (2004). Impact of water quality on macroinvertebrate assemblages along a tropical stream in Kenya. *Afr. J. Ecol.* 42: 208 - 216.
- Negishi, J. N., Inoue, M. and Munokawa, M. (2002). Effect of channelization on stream habitat in relation to a spate and flow refugia for macroinvertebrates in Northern Japan. *Freshwater Biol.* 47: 1515 - 1529.
- Nemerow, N. L. (1974). *Scientific Stream Pollution Analysis*. New York: McGraw-Hill.
- Newall, P., Bate, N. and Metzeling, L. (2006). A comparison of diatom and macroinvertebrate classification of sites in the Kiewa River system, Australia. *Hydrobiologia*, 572(1), 131-149.
- Newbold, J. D., Elwood, J. W., O'Neill, R. V., and Winkle, W. V. (1981). Measuring nutrient spiralling in streams. *Canadian Journal of Fisheries and Aquatic Sciences*, 38(7), 860-863.
- Newlon, T. A. and Rabe, F. W. (1977). Comparison of macroinvertebrate samplers and the relationship of environmental factors to biomass and diversity variability in a small watershed. Research Technical Completion Report, project A-049-IDA; Idaho Water Resources Research Institute, University of Idaho, Moscow.

- Nienhuis, P. H. (2006). Water and values: ecological research as the basis for water management and nature management. In Leuven, R. S. E. W., Ragas, A. M. J., Smits, A. J. M. and van der Velde, G. (Eds.). *Living Rivers: Trends and Challenges in Science and Management* (pp. 261 - 275). Netherlands: Springer.
- Nixon, S. (2003). An overview of the biological assessment of surface water quality in Europe. *Biological Evaluation and Monitoring of the Quality of Surface Waters*, 9-15.
- Norhayati, M. T., Suhaimi, S., Mohamad, A. and Ang, K. T. (2004). Studies on nitrogen-based nutrients of Paka River system, Terengganu, Malaysia. *Prosiding Seminar Tahunan KUSTEM ke 3*: 407 - 411.
- Nor Zaiha, A., Mohd Ismid, M. S., Salmiati and Shamila, A. (2013). Anthropogenic influences on aquatic life community and water quality status in Mengkibol River, Kluang Johor. *Journal of Applied Sciences in Environmental Sanitation*. 8(3): 151 - 160.
- Nor Zaiha, A., Mohd Ismid, M. S., Salmiati, and Shahrul Azri, M. S. (2015). Effects of logging activities on ecological water quality indicators in the Berasau River, Johor, Malaysia. *Environmental Monitoring and Assessment*. 187(8), 4715.
- Ometo, J. P. H., Martinelli, L. A., Ballester, M. V., Gessner, A., Krusche, A. V., Victoria, R. L. and Williams, M. (2000). Effects of land use on water chemistry and macroinvertebrates in two streams of the Piracicaba river basin, south-east Brazil. *Freshwater Biology*. 44(2): 327 - 337.
- Ophardt, C. E. (2003). pH. In Virtual Chembook. Retrieved from <http://www.elmhurst.edu/~chm/vchembook/184ph.htm>.
- Ormerod, S. J., Rundle, S. D., Wilkinson, S. M., Daly, G. P., Dale, K. M., & Juttner, I. (1994). Altitudinal trends in the diatoms, bryophytes, macroinvertebrates and fish of a Nepalese river system. *Freshwater Biology*, 32(2), 309-322.
- Pacini, N., and Harper, D. M. (2000). River conservation in central and tropical Africa. *Global Perspectives on River Conservation: Science, Policy and Practice*. John Wiley and Sons Ltd, Chichester, UK, 155-178.
- Pandit Ashok, K., Rashid, H. U., Rather, G. H. and Lone, H. A. (2007). Limnological survey of some fresh waterbodies in Kupwara region of Kashmir. *J. Himalayan Ecol. Sustain. Dev.* 2: 92 - 97.
- Peeters, E. T. H. M., Batten, S., Beijer, J. A. J. and Gardeniers, J. J. P. (2000). The combined effects of current velocity and substratum particle size on the

- performance of *Gammarus pulex*, an experimental approach. Aquatic Ecology and Water Quality Management Group, Environmental Sciences, Wageningen University, The Netherlands.
- Peiffer, S., Beierkuhnlein, C., Sandhage-Hofmann, A., Kaupenjohann, M. and Bar, S. (1997). Impact of high aluminum loading on a small catchment area (Thuringia slate mining area) - Geochemical transformations and hydrological transport. *Water Air Soil Pollut.* 94: 401 - 416.
- Pepper, I. L., Brooks, J. P. and Gerba, C. P. (2006). Pathogens in biosolids. *Advances in Agronomy.* 90: 1 - 41.
- Petersen, R. C. and Cummins, K. W. (1974). Leaf processing in a woodland stream*. *Freshwater biology.* 4(4): 343 - 368.
- Petts, G. E. (1994). Rivers: dynamic components of catchment ecosystems. *The Rivers Handbook: Hydrological and Ecological Principles, Volume Two*, 3-22.
- Petts, G. E. and Calow, P. (1996). *River Flows and Channel Form: Selected Extracts from the Rivers Handbook*. London: Blackwell Science.
- Pielou, E. C. (1966). The Measurement of Diversity in Different Types of Biological Collections. *J. Theoret. Biol.* 13: 131 - 144.
- Poff, N. L. and Ward, J. V. (1989). Implications of streamflow variability and predictability for lotic community structure: A regional analysis of streamflow patterns. *Can. J. Fish. Aquat. Sci.* 46: 1805 - 1817.
- Pollard, J. B. and Berrill, M. (1992). The distribution of dragonfly nymphs across a pH gradient in south-central Ontario lakes. *Canadian Journal of Zoology.* 70(5): 878 - 885.
- Pringle, C. M. (1997). Exploring how disturbance is transmitted upstream: going against the flow. *Journal of the North American Benthological Society.* 16: 425 - 438.
- Pringle, C. M. (2000). Riverine conservation in tropical versus temperate regions: Ecological and socioeconomic considerations. In Boon, P. J., Davis, B. R. and Petts, G. E. (Eds). *Global Perspectives on River Conservation: Science, Policy and Practice* (pp. 367 - 378). New York, USA: John Wiley and Sons Ltd.
- Pringle, C. M., Freeman, M. and Freeman, B. (2000a). Regional effects of hydrologic alterations on riverine macrobiota in the new world: tropical-temperate comparisons. *BioScience.* 50: 807 - 823.

- Pringle, C. M., Scatena, F. N., Paaby-Hansen, P. and Nuñez-Ferrera, M. (2000b). River conservation in Latin America and the Caribbean. In Boon, P. J., Davis, B. R. and Petts, G. E. (Eds). *Global Perspectives on River Conservation: Science, Policy and Practice* (pp. 41 - 77). New York, USA: John Wiley and Sons Ltd.
- Purvis, A. and Hector, A. (2000). Review article: getting the measure of biodiversity. *Nature*. 405: 212 - 219.
- Quinn, J. M., Cooper, A. B., Davies-Colley, R. J., Rutherford, J. C. and Williamson, R. B. (1997). Land use effects on habitat, water quality, periphyton, and benthic invertebrates in Waikato, New Zealand, hill-country streams. *N Z J Mar Freshw Res*. 31(5): 579 - 597.
- Rahman, H. A. (2007). Suatu Tinjauan Terhadap Isu Pencemaran Sungai Di Malaysia. *Dibentangkan di Persidangan Geografi*. 8 - 9 September. Universiti Pendidikan Sultan Idris.
- Raja, S. S., Chong, W. L., and Rus, A. K. A. B. M. (2015). The Old and New Malaya of Colonial Days and Its Continuity Into Modern Day Malaysia. *Mediterranean Journal of Social Sciences*, 6(2 S1), 161.
- Rak, A. E., Said, I., Mohamed, M., and Abas, A. (2011). Effect of Logging Activities on Water Quality and Benthic macroinvertebrate Assemblages of Madek River Basin, Kluang, Johor, Malaysia. *Journal of Applied Sciences and Environmental Management*. 15(2): 337 - 340.
- Ramachandra, T. V. and Solanki, M. (2007). Ecological assessment of lentic water bodies of Bangalore. *The Ministry of Science and Technology*.
- Ramirez, A., Pringle, C. M., and Wantzen, K. M. (2008). Tropical stream conservation. *Tropical stream ecology*, 285-304.
- Ramsey, D. L. and Brannon, D. G. (1988). Predicted acid mine drainage impacts to the Buckhannon River, W.V., USA. *Water Air Soil Pollut*. 39: 1 - 14.
- Resh, V. H. (2008). Which group is best? Attributes of different biological assemblages used in freshwater biomonitoring programs. *Environmental Monitoring and Assessment*, 138(1-3), 131-138.
- Rios, S. L. and Bailey, R. C. (2006). Relationship between riparian vegetation and stream benthic communities at three spatial scales. *Hydrobiologia*. 553: 153 - 160.

- RM2 juta atasi masalah banjir di Mengkibol (2011, September 23). *Utusan Online*.
Retrieved May 14, 2015, from
http://www.utusan.com.my/utusan/info.asp?y=2011&dt=0923&pub=Utusan_Malaysia&sec=Johor&pg=wj_04.htm.
- Robinson, C. T., Uehlinger, U. and Monaghan, M. T. (2003a). Stream ecosystem response to multiple experimental floods from a reservoir. *River Research and Applications*. 20: 359 - 377.
- Robinson, C. T., Uehlinger, U. and Monaghan, M. T. (2003b). Effects of a multiyear experimental flood regime on macroinvertebrates downstream of a reservoir. *Aquatic Sciences*. 65: 210 - 222.
- Ross, S. T. and Baker, J. A. (1983). The response of fishes to periodic spring floods in a southeastern stream. *Am. Midl. Nat.* 109: 1 - 14.
- Roy, A. H., Rosemond, A. D., Leigh, D. S., Paul, M. J. and Wallace, J. B. (2003). Habitat-specific responses of stream insects to land disturbance: Biological consequences and monitoring implications. *Journal of the North American Benthological Society*. 22: 292 - 307.
- Sandin, L. and Solimini, A. G. (2009). Freshwater ecosystem structure-function relationships: from theory to application. *Freshwater Biology*. 54: 2017 - 2024.
- Sands, R. (2005). *Forestry in a Global Context*. CABI Publishing.
- Sar, L. and Sledge, P. (1994). *Outlook for Australian woodchips. Quarterly Forest Products Statistics. June Quarter 1994*. Canberra: ABARE.
- Sawyer, J. A., Stewart, P. M., Mullen, M. M., Simon, T. P., & Bennett, H. H. (2004). Influence of habitat, water quality, and land use on macro-invertebrate and fish assemblages of a southeastern coastal plain watershed, USA. *Aquatic Ecosystem Health & Management*, 7(1), 85-99.
- Sawyer, C. N. and McCarty, P. L. (1978). *Chemistry for environmental engineers*. New York: McGraw Hill Book Company.
- Schumm, S. A. (1977). *The Fluvial System*. New York: John Wiley and Sons.
- Shaharuddin, S., Ghani, A. A., Zakaria, N. A., and Mashhor, M. (2011). Biodiversity in stormwater constructed wetlands: Case study of USM Wetland. *3rd International Conference on Managing Rivers in the 21st Century: Sustainable Solutions for Global Crisis of Flooding, Pollution and Water Scarcity*. 6 - 9 December. Pulau Pinang.

- Sharifah Aisyah, S. O., Aweng, E. R., Razak, W., and Ahmad Abas, K. (2015). Preliminary Study on Benthic Macroinvertebrates Distribution and Assemblages at Lata Meraung Waterfall, Pahang, Malaysia. *Jurnal Teknologi*, 72(5).
- Short, R. A. and Maslin, P. E. (1977). Processing of leaf litter by a stream detritivore: effect on nutrient availability to collectors. *Ecology*. 935 - 938.
- Shrestha, S. and Kazama, F. (2007). Assessment of surface water quality using multivariate statistical techniques: A case study of the Fuji river basin, Japan. *Environmental Risk and Emergency Management*. 22: 464 - 475.
- Sidek, L., Basri, H., Lee, L. K., and Foo, K. Y. (2016). The performance of gross pollutant trap for water quality preservation: a real practical application at the Klang Valley, Malaysia. *Desalination and Water Treatment*, 1-9.
- Silveira, M. P., Buss, D. F., Nessimian, J. L. and Baptista, D. F. (2006). Spatial and temporal distribution of benthic macroinvertebrates in a southeastern Brazilian river. *Braz. J. Biol.* 66(2B): 623 - 632.
- Simon, T. P. (2000). The use of biological criteria as a tool for water resource management. *Environmental Science & Policy*. 3: 43 - 49.
- Smith, J. M. (2004). *Water Quality Trends in the Blackwater River Watershed Canaan Valley, West Virginia*. Master of Science Theses. West Virginia University.
- Smith, R. L. and Smith, T. M. (2001). *Ecology and Field Biology*. San Francisco: Benjamin Cummings.
- Smith, T. M. and Smith, R. L. (2012). *Elements of Ecology (Eighth ed.)*. Boston: Benjamin Cummings.
- Snyder, C. D. and Johnson, Z. B. (2006). Macroinvertebrate assemblage recovery following a catastrophic flood and debris flows in an Appalachian mountain stream. *Journal of the North American Benthological Society*. 25: 825 - 840.
- Sodhi, N. S., Lee, T. M., Koh, L. P. and Brook, B. W. (2009). A meta-analysis of the impact of anthropogenic forest disturbance on Southeast Asia's biotas. *Biotropica*. 41:103 - 109.
- Sonneman, J. A., Walsh, C. J., Breen, P. F., & Sharpe, A. K. (2001). Effects of urbanization on streams of the Melbourne region, Victoria, Australia. II. Benthic diatom communities. *Freshwater Biology*, 46(4), 553-565.

- Sporka, F., Vlek, H. E., Bulankova, E. and Krno, I. (2006). Influence of seasonal variation on bioassessment of streams using macroinvertebrates. *Hydrobiologia*. 566: 543 - 555.
- Standley, L. J. and Sweeney, B. W. (1995). Organochlorine pesticides in stream mayflies and terrestrial vegetation of undisturbed tropical catchments exposed to long-range atmospheric transport. *Journal of the North American Benthological Society*. 14: 38 - 49.
- Stanford, J. A., and J. V. Ward. (2001). Revisiting the serial-discontinuity concept. *Regulated Rivers: Research & Management* 17: 303-310
- Stanford, J. A., and Ward, J. V. (1993). An ecosystem perspective of alluvial rivers: connectivity and the hyporheic corridor. *Journal of the North American Benthological Society*, 48-60.
- Stanford, J.V., Hauer, F.R., Ward, J.V., (1988). Serial discontinuity in a large river system. *Verh. Int. Verein.Limnol.* 23, 1114-1118.
- Statzner, B. and Higler, B. (1986). Stream hydraulics as a major determinant of benthic invertebrate zonation patterns. *Freshwater Biology*. 16: 127 - 139.
- Steinmann, P. (1907). Die Tierwelt der Gebirgsbache. Eine faunistisch- biologische Studie. *Annls Biol. Lacustre*. 2: 30 - 150.
- Steinmann, P. (1908). Die Tierwelt der Gebirgsbache. *Arch. Hydrobiol.* 3: 266 - 273.
- Stewart, P. M., Scribaillo, R. and Simon, T. P. (1999). The use of aquatic macrophytes in monitoring and in assessment of biological integrity. In Gerhardt, A. (Ed.). *Biomonitoring of Polluted Waters - Reviews on Actual Topics* (pp. 275 - 302). Environmental Science Forum 96. Uetikon - Zuerich, Switzerland: Trans Tech Publications, Ltd.
- Strayer, D. L., & Malcom, H. M. (2007). Effects of zebra mussels (*Dreissena polymorpha*) on native bivalves: the beginning of the end or the end of the beginning?. *Journal of the North American Benthological Society*, 26(1), 111-122.
- Strayer, D. L., Lutz, C., Malcom, H. M., Munger, K., and Shaw, W. H. (2003). Invertebrate communities associated with a native (*Vallisneria americana*) and an alien (*Trapa natans*) macrophyte in a large river. *Freshwater Biology*, 48(11), 1938-1949.

- Strayer, D. L. and Dudgeon, D. (2010). Freshwater biodiversity conservation: recent progress and future challenges. *Journal of the North American Benthological Society*. 29: 344 - 358.
- Subramanian, K. A., Sivaramakrishnan, K. G. and Gadgil, M. (2005). Impact of riparian land use on stream insects of Kudremukh National Park, Karnataka state, India. *J. Insect Sci.* 5: 49 - 59.
- Suhaila, A.H., Che Salmah, M.R. & Nurul Huda, A. 2014. Seasonal abundance and diversity of aquatic insects in rivers in Gunung Jerai Forest Reserve, Malaysia. *Sains Malaysiana*. 43(5): 667-674
- Suhaila, A.H., Salmah, M.R.C. and Al-Shami, S.A. (2012). Temporal distribution of Ephemeroptera, Plecoptera and Trichoptera (EPT) adults at a tropical forest stream: Response to seasonal variations. *The Environmentalist*. 32(1): 28-34
- Sullivan, S. M. P., Watzin, M. C. and Hession, W. C. (2004). Understanding stream geomorphic state in relation to ecological integrity: evidence using habitat assessments and macroinvertebrates. *Environ. Manage.* 34(5): 669 - 683.
- Sulong Mohamad, Mohd Ekhwan Toriman, Kadaruddin Aiyub dan Mokhtar Jaafar, (2005). *Sungai dan Pembangunan Tebingan Sungai Bandar Malaysia*. Bangi: Penerbit UKM.
- Sungai tercemar di Johor dibersihkan (2012, Disember 16). *Utusan Online*. Retrived May 14, 2015, from http://ww1.utusan.com.my/utusan/Johor/20121216/wj_01/Sungai-tercemar-di-Johor-dibersihkan.
- Sweeney, B. W. and Schnack, J. A. (1977). Egg development, growth, and metabolism of *Sigara alternata* (Say) (Hemiptera: Corixidae) in fluctuating thermal environments. *Ecology*. 58: 265 - 277.
- Sweeney, B.W. (1984). Factors influencing life history patterns of aquatic insects. In Resh, V. H and Rosenberg, D. M., (Eds.). *The Ecology of Aquatic Insects*. 56-100.
- Talling, J. F. and Lemoalle, J. (1998). *Ecological Dynamics of Tropical Inland Waters*. Cambridge, UK: Cambridge Univ. Press.
- Tchobanoglous, G. and Schroeder E. D. (1985). *Water Quality*. USA: Addison - Wesley Publishing Company.

- Thomsen, A. G. and Friberg, N. (2002). Growth and emergence of the stonefly *Leuctra nigra* in coniferous forest streams with contrasting pH. *Freshwater Biol.* 47: 1159 - 1172.
- Thorp, J. H. (1986). Two distinct roles of predators in freshwater assemblages. *Oikos*. 47: 75 - 82.
- Thorp, J. H. and Covich, A. P. (1991). *Ecology and Classification of North America Freshwater Invertebrates*. San Diego, California: Academic Press Inc.
- Thorp, J. H., and DeLong, M. D. (1994). The riverine productivity model: an heuristic view of carbon sources and organic processing in large river ecosystems. *Oikos*, 305-308.
- Townsend, C. R. (1996). Concepts in river ecology: pattern and process in the catchment hierarchy. *Large Rivers*, 3-21.
- Triest, L., Kaur, P., Heylen, S., & De Pauw, N. (2001). Comparative monitoring of diatoms, macroinvertebrates and macrophytes in the Woluwe River (Brussels, Belgium). *Aquatic Ecology*, 35(2), 183-194.
- Triplehorn, C. A. and Johnson, N. F. (2005). *Study of insect*. United States of America: Thompson Brooks.
- Tronstad, L. M., Tronstad, B. P. and Benke, A. C. (2005). Invertebrate seedbanks: rehydration of soil from an unregulated river floodplain in the southeastern U.S. *Freshwater Biol.* 50: 646 - 655.
- USEPA (2002). *Summary of Biological Assessment Programs and Biocriteria Development for States, Tribes, Territories, and Interstate Commissions: Streams and Wadeable Rivers*. Environmental Protection Agency, Office of Environmental Information and Office of Water, Washington, D.C. EPA-822-R-02-048. U.S.
- Vannote, R. L. and Sweeney, B. W. (1980). Geographic analysis of thermal equilibria: A conceptual model for evaluating the effect of natural and modified thermal regimes on aquatic insect communities. *Am. Nat.* 115: 667 - 695.
- Vannote, R. L., Minshall, G. W., Cummins, K. W., Sedell, J. R. and Cushing, C. E. (1980). The river continuum concept. *Canadian journal of fisheries and aquatic sciences*. 37(1): 130 - 137.
- Vesilind, P. A. (1975). *Environmental Pollution and Control*. Ann Arbor: Ann Arbor Science Publishers.

- Viswanathan, S., Voss, K. A., Pohlman, A., Gibson, D., and Purohit, J. (2009). Evaluation of the biocriteria of streams in the San Diego Hydrologic Region. *Journal of Environmental Engineering*, 136(6), 627-637.
- Waite, I. R., Herlihy, A. T., Larsen, D. P., Urquhart, N. S. and Klemm, D. J. (2004). The effect of macroinvertebrate taxonomic resolution in large landscape bioassessments: an example from Mid-Atlantic Highlands, U.S.A. *Freshwater Biol.* 49: 474 - 489.
- Wallace, J. B., Webster, J. R., and Woodall, W. R. (1977). The role of filter feeders in flowing waters. *Arch. Hydrobiol*, 79(4), 506-S32.
- Walsh, C. J., Roy, A. H., Feminella, J. W., Cottingham, P. D., Groffman, P. M. and Morgan, R. P. (2005). The urban stream syndrome: current knowledge and the search for a cure. *Journal of the North American Benthological Society*. 24: 706 - 723.
- Wantzen, K. M. (2006). Physical pollution: effects of gully erosion in a tropical clear-water stream. *Aquatic Conservation*. 16: 733 - 749.
- Ward, J. V., and Stanford, J. A. (1983). The serial discontinuity concept of lotic ecosystems. *Dynamics of lotic ecosystems*, 10, 29-42.
- Ward, J. V. (1992). *Aquatic Insect Ecology*. New York: John Wiley.
- Watanabe, N. C., Harada, S., and Komai, Y. (2000). Long-term recovery from mine drainage disturbance of a macroinvertebrate community in the Ichi-kawa River, Japan. *Hydrobiologia*, 429(1-3), 171-180.
- Witt, V.M. and Reiff, F.M., (1991). Environmental health conditions and cholera vulnerability in Latin America and the Caribbean. *Journal of public health policy*, 12(4), pp.450-463.
- WCD (World Commission on Dams) (2000). Dams and development: a framework for decision-making. www.damsreport.org [10 December 2005].
- Weber, R. E. and Vinogradov, S. N. (2001). Non-vertebrate hemoglobins: function and molecular adaptation. *Physiol. Rev.* 81: 569 - 628.
- Whitlock, D. (2010). *Trout and Their Food: A Compact Guide for Fly Fishers*. Skyhorse Publishing Inc.
- Willoughby, L. G. and Mappin, R. G. (1988). The distribution of *Ephemerella ignita* (Ephemeroptera) in streams: The role of pH and food resources. *Freshwater Biol.* 19: 145 - 155.

- Won, D. H., Jun, Y. C., Kwon, S. J., Hwang, S. J., Ahn, K. G., and Lee, J. K. (2006). Development of Korean saprobic index using benthic macroinvertebrates and its application to biological stream environment assessment. *J. Kor. Soc. Water Quality*, 22, 768-783.
- World Health Organization (2003). pH in Drinking-water. In Guidelines for drinking-water quality. Retrieved from http://www.who.int/water_sanitation_health/dwq/chemicals/en/ph.pdf
- Wowor, D., Cai, Y., and Ng, P. K. (2004). Crustacea: Decapoda, Caridea. *Freshwater Invertebrates of the Malaysian Region. Malaysian Academy of Sciences*, 337-357.
- Wowor, D., Muthu, V., Meier, R., Balke, M., Cai, Y., and Ng, P. K. (2009). Evolution of life history traits in Asian freshwater prawns of the genus *Macrobrachium* (Crustacea: Decapoda: Palaemonidae) based on multilocus molecular phylogenetic analysis. *Molecular Phylogenetics and Evolution*, 52(2), 340-350.
- Wood, M. S. (2014). Estimating suspended sediment in rivers using acoustic Doppler meters: U.S. Geological Survey Fact Sheet 2014 - 3038, 4 p.
- Wu, H. C., Chen, P. C., and Tsay, T. T. (2010). Assessment of nematode community structure as a bioindicator in river monitoring. *Environmental pollution*, 158(5), 1741-1747.
- Younes-Baraille Y, Garcia, X. F. and Gagneur, J. (2005). Impact of longitudinal and seasonal changes of water quality on the benthic macroinvertebrate assemblages of the Andorran streams. *C. R. Biologies*. 328: 963 - 976.
- Yusop, Z., Tan, L. W., Ujang, Z., Mohamed, M., and Nasir, K. A. (2005). Runoff Quality and Pollution Loadings from A Tropical Urban Catchment. *Water science and technology*. 52 (9): 125-132.
- Yousuf, A. R., Bhat, F. A. and Mahdi, M. D. (2006). Limnological features of River Jhelum and its important tributaries in the Kashmir Himalaya with a note on fish fauna. *J. Himalayan Ecol. Sustain. Dev.* 1: 37 - 50.
- Yuan, L. L. (2004). Assigning macroinvertebrate tolerance classifications using generalized additive models. *Freshwater Biol.* 49: 662 - 677.
- Yule, C. M. and Yong, H.S. (2004). *Freshwater Invertebrates of the Malaysian Region*. Kuala Lumpur: Academy of Sciences Malaysia.

- Yule, C. M. (2004). Insecta: Diptera. In Yule, C.M. and Yong, H.S. (eds.) *Freshwater Invertebrates of the Malaysian Region* (pp 610 - 612) Malaysia: Academy of Sciences Malaysia.
- Zaiha, A. N., Ismid, M. M., and Azri, M. S. (2015). Effects of logging activities on ecological water quality indicators in the Berasau River, Johor, Malaysia. *Environmental monitoring and assessment*, 187(8), 1-9.
- Zahar, A. R. (1951). The ecology and distribution of black flies (Simuliidae) in South East Scotland. *The Journal of Animal Ecology*. 33 - 62.
- Zakaria, N. A. (2009). *Sistem saliran bandar Mapan: penyelesaian banjir kilat, pencemaran sungai dan kekurangan air untuk kesejahteraan sejagat*. Penerbit Universiti Sains Malaysia.
- Zhe-ren, D. O. N. G. (2003). Diversity of river morphology and diversity of bio-communities [J]. *Journal of Hydraulic Engineering*, 11, 1-6.
- Zhou, Q., Zhang, J., Fu, J., Shi, J., and Jiang, G. (2008). Biomonitoring: an appealing tool for assessment of metal pollution in the aquatic ecosystem. *Analytica chimica acta*, 606(2), 135-150.