


MAZ (MYC Associated Zinc Finger Protein)



The human MAZ consists of six exons, the first five of which are encoding exons; total exon length is 4.57 kb (Song et al., 1998).

Description

The human MAZ contains six exons; the encoding sequence consists of 1431 bases (Song et al., 1998).

Transcription


The human gene encoding for MAZ is located on chromosome 16p11.2 and is transcribed as an mRNA of 2.7 kilobases (kb). The primary transcript encodes a 477 amino-acid long MAZ-1 protein with a 60-kDa molecular mass that contains six C2H2-type zinc-finger domains responsible for DNA binding. MAZ protein has two additional isoforms: MAZ-2 and MAZ-3. The MAZ-2 transcript is generated by an alternative splicing that results in the insertion of a new exon originating from the non-coding sequences of the intron 4. This transcript gives rise to the MAZ-2 isoform, which is a 493 amino-acids long protein with distinct carboxyl-

terminus which contains two additional zinc-finger domains (Ray et al., 2002). The MAZ-2 isoform is reported to have a higher DNA-binding activity and to act as a negative regulator of MAZ-1 function (Ray et al., 2002). The MAZ-3 transcript is expressed at very low levels under normal physiological condition in various tissues, but is highly expressed during inflammation. The MAZ-3 transcript is transcribed from a distinct upstream promoter and is processed with alternative splicing. The MAZ-3 transcript is translated from a different starting codon that gives rise to the MAZ-3 isoform of 455 amino-acids (Ray et al., 2009).

Pseudogene

No reported pseudogenes are found.

Protein




Domains of MAZ are depicted with vertical colored lines; Blacks are Poly-Alanine repeats; Green is Poly-Proline tract; Red column is Poly-Glycine repeat. C2H2-type zinc finger domains of MAZ-1 are represented in dark blue-green vertical lines

Description

The human MAZ protein contains three Poly alanine, one poly-proline and one poly-glycine domains (Song et al., 1998). Poly-alanine repeats considered to have role in cellular localization of the protein; the alteration in the intracellular distribution may contribute to diseases, including muscular

dystrophy (OPMD) (Oma et al., 2004). Similarly, poly-glycine repeats are responsible in protein targeting (Uthayakumar et al., 2012). Poly-proline tracks, on the other hand, generates structures that are predicted to have important roles in protein-protein interactions (Williamson, 1994). The human MAZ-1 contains six C2H2-type zinc finger domains (Song et al., 1998), which are frequently occurring in proteins involved in transcriptional regulation.

MAZ (MYC Associated Zinc Finger Protein)




Expression and synthesis of MAZ in various cancerous tissues. Retrieved from: <http://www.proteinatlas.org/ENSG00000103495-MAZ/cancer> on November 2, 2017.

Expression

MAZ is expressed in the human heart, brain, lungs, liver, skeletal muscle, pancreas, and prostate (Jiao et

al., 2013; Dudas et al., 2008). The synthesis of MAZ protein is observed to occur at high levels in breast, thyroid and urothelial cancers as well as in melanoma (Ugai et al., 2001)



Immunofluorescent staining of human cell line MCF7. Retrieved from: <http://www.proteinatlas.org/ENSG00000103495-MAZ/cell> on November 2, 2017. Immunofluorescent staining of MCF7 cells derived from breast adenocarcinoma shows that MAZ localizes to the nucleus.

Localisation

MAZ is located in the nucleus (Jordan-Sciutto et al, 2000).

Function

MAZ as a transcription factor interacts with a permutation of the GGGAGGG sequence motif

present in GC-rich promoter regions of target genes by unfolding of G-quadruplex structures of DNA (Cogoi et al., 2014) to activate or repress transcription. MAZ is also suggested to participate in transcription termination and polyadenylation. Several oncogenes, including MYC, HRAS, PPARG, TSG101, VEGFA, CAV1, PTHR1, NOS3, MYB, and hTER, are transcriptionally regulated by MAZ (Jun Song et al., 2001; Lee et al., 2016; Ray et al., 2002). Deregulated expression of MAZ appears to participate in the development and/or progress various tissue malignancies including the breast, thyroid, hepatocellular and urothelial cancers (Jiao et al., 2013; Dudas et al., 2008; Yu et al., 2017; Ray, 2011; Zhu et al., 2016)

Homology

The human MAZ protein is conserved 100% in chimpanzee (*P.troglodytes*), 98.4% in mouse (*M.musculus*), and 98.4% in rat (*R.norvegicus*); with conserved DNA of 99.8%, 93.2%, and 92.7%, respectively (Retrieved from: <https://blast.ncbi.nlm.nih.gov/Blast.cgi>. November 2, 2017).

Mutations

Note

Genetic mutations are not described for MAZ.

Implicated in

Prostate cancer

It was reported that the MAZ expression is higher in clinical prostate cancer (PCa) specimens than in benign prostatic hyperplasia (BPH) and adjacent normal tissues (Jiao et al., 2013). Moreover, the MAZ expression appears to be positively correlated with the expression of androgen receptor (AR), which is critical for the initiation and development of androgen-dependent PCa (Jiao et al., 2013). Extending these findings, experimental studies in cell models derived from PCa indicated that MAZ is involved in the phenotypic manifestation of PCa cell models as siRNA knockdown of MAZ levels reduces cell proliferation, migration, and invasion through mechanisms involve the expression of AR (Jiao et al., 2013).

Hepatocellular carcinoma

The expression of MAZ was reported to be upregulated in the majority (78.94%) of hepatocellular carcinoma (HCC) samples compared to normal liver samples (Dudas et al., 2008). Experimental studies using cell lines derived from HCC further suggest that MAZ-mediated regulation of PROX1, which is a transcription factor critical for the expression of a number of genes involved in hepatic metabolic functions, contributes to the progression of HCC (Dudas et al., 2008).

Breast cancer

Based on data sets in Gene expression-based Outcome for Breast Cancer Online (GOBO, <http://co.bmc.lu.se/gobo/>), the expression of MAZ is found to be correlated with distant metastasis-free survival (DMFS) in basal-like breast cancer (BLBC) patients and that the under-expression of MAZ is involved in the metastatic spread of BLBC (Yu et al., 2017). Based on these finding, it was suggested that MAZ plays dual roles in basal-like breast cancer (BLBC): it suppresses cancer progression but promotes cellular proliferation (Yu et al., 2017). Experimental studies using model cell lines derived from breast cancer indeed suggest that MAZ promotes cell proliferation yet it suppresses the aggressiveness of BLBC by controlling the transition toward a more mesenchymal phenotype (Yu et al., 2017; Ray, 2011).

Pancreatic carcinoma

Based on samples from pancreatic carcinoma patients, it was reported that the expression of MAZ is significantly higher in PC tissue compared to the adjacent non-tumor tissues (Zhu et al., 2016). Moreover, it appears that the over-expression of MAZ is associated with poor prognosis of PC patients (Zhu et al., 2016).

Hodgkin's Disease and Paraneoplastic Cerebellar Dysfunction

In neuronal cells, MAZ interacts with the Deleted in Colorectal Cancer product (DCC), the receptor for NTN1 netrin-1 which plays a central role in axonal guidance and neuronal migration as well as survival during development. Analyses of sera from patients with HD and PCD Hodgkin's disease and paraneoplastic cerebellar degeneration indicated that patient sera contain auto-antigens directed against the MAZ-DCC complex. Based on these observations, it was speculated that auto-antigens could interfere with neuronal function resulting in neuronal degeneration (Bataller and Wade, 2002).

To be noted

Expression of the MAZ gene is found to be regulated by MIR-125B, which is suggested to affect VEGF-induced angiogenesis in glioblastoma (Vandertop et al., 2017).

MIR449A targets MAZ transcripts, the down-regulation of which is reported to contribute to glioblastoma (Chen et al., 2015; Zhao et al., 2014; Yao et al., 2015).

MIR34C is also reported to target MAZ. Decrease levels of MAZ by miR-34c are suggested to impair the integrity and increased the permeability of blood-tumor barrier (Zhao et al., 2014).

References

- Bataller L, Wade DF, Graus F, Rosenfeld MR, Dalmau J. The MAZ protein is an autoantigen of Hodgkin's disease and paraneoplastic cerebellar dysfunction. *Ann Neurol*. 2003 Jan;53(1):123-7
- Bossone SA, Asselin C, Patel AJ, Marcu KB. MAZ, a zinc finger protein, binds to c-MYC and C2 gene sequences regulating transcriptional initiation and termination. *Proc Natl Acad Sci U S A*. 1992 Aug 15;89(16):7452-6
- Cogoi S, Shchekotikhin AE, Xodo LE. HRAS is silenced by two neighboring G-quadruplexes and activated by MAZ, a zinc-finger transcription factor with DNA unfolding property. *Nucleic Acids Res*. 2014 Jul;42(13):8379-88
- Chen SP, Liu BX, Xu J, Pei XF, Liao YJ, Yuan F, Zheng F. MiR-449a suppresses the epithelial-mesenchymal transition and metastasis of hepatocellular carcinoma by multiple targets. *BMC Cancer*. 2015 Oct 15;15:706
- Dudas J, Mansuroglu T, Moriconi F, Haller F, Wilting J, Lorf T, Füzesi L, Ramadori G. Altered regulation of Prox1-gene-expression in liver tumors. *BMC Cancer*. 2008 Apr 9;8:92
- Ashfield R, Patel AJ, Bossone SA, Brown H, Campbell RD, Marcu KB, Proudfoot NJ. MAZ-dependent termination between closely spaced human complement genes. *EMBO J*. 1994 Dec 1;13(23):5656-67
- Jiao L, Li Y, Shen D, Xu C, Wang L, Huang G, Chen L, Yang Y, Yang C, Yu Y, Sun Y. The prostate cancer-up-regulated Myc-associated zinc-finger protein (MAZ) modulates proliferation and metastasis through reciprocal regulation of androgen receptor. *Med Oncol*. 2013;30(2):570
- Jordan-Sciutto KL, Dragich JM, Caltagarone J, Hall DJ, Bowser R. Fetal Alz-50 clone 1 (FAC1) protein interacts with the Myc-associated zinc finger protein (ZF87/MAZ) and alters its transcriptional activity. *Biochemistry*. 2000 Mar 28;39(12):3206-15
- Song KH, Li T, Chiang JY. A Prospero-related homeodomain protein is a novel co-regulator of hepatocyte nuclear factor 4alpha that regulates the cholesterol 7alpha-hydroxylase gene. *J Biol Chem*. 2006 Apr 14;281(15):10081-8
- Lee WP, Lan KH, Li CP, Chao Y, Lin HC, Lee SD. Akt phosphorylates myc-associated zinc finger protein (MAZ), releases P-MAZ from the p53 promoter, and activates p53 transcription. *Cancer Lett*. 2016 May 28;375(1):9-19
- Ray A, Dhar S, Ray BK. Control of VEGF expression in triple-negative breast carcinoma cells by suppression of SAF-1 transcription factor activity. *Mol Cancer Res*. 2011 Aug;9(8):1030-41
- Ray A, Schatten H, Ray BK. Activation of Sp1 and its functional co-operation with serum amyloid A-activating sequence binding factor in synoviocyte cells trigger synergistic action of interleukin-1 and interleukin-6 in serum amyloid A gene expression. *J Biol Chem*. 1999 Feb 12;274(7):4300-8
- Ray BK, Murphy R, Ray P, Ray A. SAF-2, a splice variant of SAF-1, acts as a negative regulator of transcription. *J Biol Chem*. 2002 Nov 29;277(48):46822-30
- Smits M, Wurdinger T, van het Hof B, Drexhage JA, Geerts D, Wesseling P, Noske DP, Vandertop WP, de Vries HE, Reijerkerk A. Myc-associated zinc finger protein (MAZ) is regulated by miR-125b and mediates VEGF-induced angiogenesis in glioblastoma. *FASEB J*. 2012 Jun;26(6):2639-47
- Song J, Ugai H, Kanazawa I, Sun K, Yokoyama KK. Independent repression of a GC-rich housekeeping gene by Sp1 and MAZ involves the same cis-elements. *J Biol Chem*. 2001 Jun 8;276(23):19897-904
- Song J, Ugai H, Ogawa K, Wang Y, Sarai A, Obata Y, Kanazawa I, Sun K, Itakura K, Yokoyama KK. Two consecutive zinc fingers in Sp1 and in MAZ are essential for interactions with cis-elements. *J Biol Chem*. 2001 Aug 10;276(32):30429-34
- Ugai H, Li HO, Komatsu M, Tsutsui H, Song J, Shiga T, Fearon E, Murata T, Yokoyama KK. Interaction of Myc-associated zinc finger protein with DCC, the product of a tumor-suppressor gene, during the neural differentiation of P19 EC cells. *Biochem Biophys Res Commun*. 2001 Sep 7;286(5):1087-97
- Smits M, Wurdinger T, van het Hof B, Drexhage JA, Geerts D, Fearon E, Murata T, Yokoyama KK. Interaction of Myc-associated zinc finger protein with DCC, the product of a tumor-suppressor gene, during the neural differentiation of P19 EC cells. *Biochem Biophys Res Commun*. 2001 Sep 7;286(5):1087-97
- Smits M, Wurdinger T, van het Hof B, Drexhage JA, Geerts D, Fearon E, Murata T, Yokoyama KK. Interaction of Myc-associated zinc finger protein with DCC, the product of a tumor-suppressor gene, during the neural differentiation of P19 EC cells. *Biochem Biophys Res Commun*. 2001 Sep 7;286(5):1087-97
- Zhu X, Luo W, Liang W, Tang F, Bei C, Ren Y, Qin L, Tan C, Zhang Y, Tan S. Overexpression and clinical significance of MYC-associated zinc finger protein in pancreatic carcinoma. *Onco Targets Ther*. 2016;9:7493-7501
- Yao Y, Ma J, Xue Y, Wang P, Li Z, Li Z, Hu Y, Shang X, Liu Y. MiR-449a exerts tumor-suppressive functions in human glioblastoma by targeting Myc-associated zinc-finger protein. *Mol Oncol*. 2015 Mar;9(3):640-56
- Yonaha M, Proudfoot NJ. Specific transcriptional pausing activates polyadenylation in a coupled in vitro system. *Mol Cell*. 1999 May;3(5):593-600
- Yu ZH, Lun SM, He R, Tian HP, Huang HJ, Wang QS, Li XQ, Feng YM. Dual function of MAZ mediated by FOXF2 in basal-like breast cancer: Promotion of proliferation and suppression of progression. *Cancer Lett*. 2017 Aug 28;402:142-152
- Zhao L, Wang P, Liu Y, Ma J, Xue Y. miR-34c regulates the permeability of blood-tumor barrier via MAZ-mediated expression changes of ZO-1, occludin, and claudin-5. *J Cell Physiol*. 2015 Mar;230(3):716-31

This article should be referenced as such:

Karakaya, B.; Muyan, M. MAZ (MYC Associated Zinc Finger Protein). *Atlas Genet Cytogenet Oncol Haematol*. 2018; 22(8):323-327.
