



Ripley, Wendy and Wilby, Helen (2014) Family-carers' perspectives of involvement in stroke rehabilitation of a family member. In: College of Occupational Therapists 38th Annual Conference and Exhibition, 3-5 June 2014, Brighton, UK.

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/4756/>

*Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.*

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

**provided that**

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
  - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

**You may not**

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing [insight@cumbria.ac.uk](mailto:insight@cumbria.ac.uk).

**Author CVs**

Sue Parkinson is the lead author of the MOHO Screening Tool and the MOHO-ExpLOR. She works as a freelance MOHO trainer.

John Cooper is an experienced occupational therapist working in learning disability services at 2gether NHS Foundation Trust.

**Contact**

spark39@hotmail.co.uk

**Keywords**

Research, Service evaluation, Participation, Long term conditions (including neurological)

**P132****The perceived impact of type 2 diabetes upon human occupation: a role for occupational therapy**

**Purnell V, Moore B, Price L, University of Brighton, UK**

Introduction: Type 2 diabetes currently affects 90% of the 2.9 million people diagnosed with diabetes mellitus in the United Kingdom, and costs the National Health Service an estimated £13bn per year (Diabetes UK 2012, Hex et al 2012). Evidence suggests that existing Type 2 diabetes intervention programmes do not facilitate sustainable lifestyle changes that support effective diabetes self-management (Khunti et al 2012).

Aim: The aim of this research was to enhance understanding of the impact of diabetes on human occupation, and if occupational therapy could support effective diabetes self-management in the United Kingdom. It also strives to contribute to the diabetes and occupational therapy research base, which, to date, is extremely limited internationally.

Method: Ethical approval was obtained from SREGP on 07/11/2012. This research utilised a qualitative and inductive approach to undertake a Basic Interpretive Qualitative Study (Merriam 2009). Data was gathered through the use of semi-structured interviews with eight participants recruited from diabetes support groups. Interviews were recorded and transcribed verbatim.

Findings: Through thematic analysis, three main themes emerged: "Balancing diabetes and everyday life", "The effect of diabetes upon meaningful occupations", and "Coming to terms with diabetes and the importance of support". The findings revealed a myriad of difficulties diabetes imposes upon the person, daily life, and occupational participation, and highlighted the value of peer and professional support.

Conclusion: The findings identify that Type 2 diabetes can impact on engagement in meaningful occupation, occupational roles and established routines. This suggests an important role for occupational therapists practicing in the United Kingdom. They could empower individuals to self-manage their condition and support them to maintain participation in valued occupations.

**References**

Diabetes UK (2012) Diabetes in the UK: key statistics on diabetes. Available at: <http://www.diabetes.org.uk/Documents/Reports/Diabetes-in-the-UK-2012.pdf> Date accessed: 26.10.12.

Hex N, Bartlett C, Wright D, Taylor M, Varley D (2012) Health economics: estimating the current and future costs of Type 1 and Type 2 diabetes in the UK, including direct health costs and indirect societal and productivity costs, *Diabetic Medicine*, 29(7), 855–862.

Khunti K, Gray LJ, Skinner T, Carey ME, Realf K, Dallosso H, Fisher H, Campbell M, Heller S, Davies MJ (2012) Effectiveness of a diabetes education and self-management programme

(DESMOND) for people with newly diagnosed type 2 diabetes mellitus: three year follow-up of a cluster randomised controlled trial in primary care. *British Medical Journal*, 344(7860), 1–12.

Merriam SB (2009) *Qualitative research: a guide to design and implementation*. San Francisco: Jossey-Bass.

**Author CVs**

Briony Moore is an MSc Health through Occupation student at the University of Brighton. She completed her Post-Graduate Diploma in Occupational Therapy in July 2013.

Victoria Purnell is an MSc Health through Occupation student at the University of Brighton. She completed her Post-Graduate Diploma in Occupational Therapy in July 2013.

**Contact**

vsp11@uni.brighton.ac.uk

**Keywords**

Long term conditions (including neurological), Health and wellbeing, Research, Practice development

**P133****Family-carers' perspectives of involvement in stroke rehabilitation of a family member**

**Ripley W, Wilby H, University of Cumbria, UK**

Stroke has the potential to have a major impact on health and quality of life (Stroke Association, 2012). Occupational therapy supports people to participate in meaningful occupations and return home post stroke (Edmans, 2010). Since the target of 45 minutes of rehabilitation on five days out of seven was recommended (Kwakkel et al, 1997), creative ways to help meet this target have been explored and promoted. Family involvement in rehabilitation has been identified as one option for extending rehabilitation opportunities (Stroke Association, 2012). On return to home following inpatient rehabilitation, many people receive informal care from family members (Kniepmann, 2012).

The purpose of this small-scale, qualitative study was to explore the nature of carers' participation in their family members' rehabilitation, carers level of involvement, and the support and preparation families felt they needed. Family carers were recruited to the study via a voluntary organisation for people post stroke. Semi-structured interviews were used to explore participants' perspectives. Data analysis was conducted using thematic analysis. Ethical approval was gained from the University of Cumbria and permission granted by the charity. The study was self-funded by the researcher.

Four family carers participated in the study. Six key themes were identified: Carer involvement, Satisfaction with involvement, Desired level of involvement, Working hours, Carer role, and Information and support. These findings highlight the challenges raised and additional support required by carers. These findings could assist decision-making regarding the involvement of carers and the development of strategies to aid their participation. Such investment could enable and support family carers in making a valuable contribution to their family members' rehabilitation and to the achievement of the NICE rehabilitation intensity targets, both in in-patient units and on return to home.

**References**

Edmans, J. (2010). *Occupational therapy and Stroke*. Hoboken: Wiley-Blackwell.

Kniepmann, K. (2012). Female family carers for survivors of stroke: occupational loss and quality of life. *British Journal of Occupational Therapy*, Vol 75(5), p.p. 208–216.

Kwakkel, G., Wagenaar, R. C., Koelman, T. W., Lankhorst, G. J., & Koetsier, J. C. (1997). Effects of Intensity of Rehabilitation After Stroke. *Stroke*, Vol(28), pp. 1550–1556.

Stroke Association. (2012, Spring). Struggling to Recover. from Stroke Association: Available at: <http://www.stroke.org.uk/sites/default/files/files/StrugglingRptFIN%20lowres.pdf> (Accessed: September 10, 2013).

#### Author CVs

Wendy Ripley qualified as an occupational therapist in 2013 at the University of Cumbria. Her interests include neurological conditions and carer involvement.

Helen Wilby is a Senior Lecturer in Occupational Therapy, University of Cumbria. Her interests include development of research capacity, assessment and clinical reasoning.

#### Contact

wendy.ripley@tiscali.co.uk

#### Keywords

Recovery and rehabilitation, Long term conditions (including neurological), Practice development

### P134

#### Assessing research impact: myths, models and multiple dimensions

Sainty M, College of Occupational Therapists, UK

Introduction: Assessing research impact is a challenging endeavour (Beacham et al 2005, p1). A mythical view is that publications equate with impact; however, the definition of research impact used by the higher education funding bodies confirms its irrefutably multidimensional nature: an effect on, change or benefit to the economy, society, culture, public policy or services, health, the environment or quality of life beyond academia (Higher Education Funding Council for England et al 2011, p26). Models, such as the Payback Framework (Buxton and Hanney 1996) and the Becker Medical Library Model (Sarli et al 2010), identify various domains of impact, but are they suitable for assessing the impact of occupational therapy research?

Method: The Becker Model was selected, following a literature review, to gather intelligence on projects completed with support from the United Kingdom Occupational Therapy Research Foundation. Eight researchers participated, completing a form covering research outputs and knowledge advancement; clinical implementation; community and public benefit; and economic benefit. Ethics approval was obtained from the College of Occupational Therapists.

Results: The model supported reporting of more easily quantifiable indicators, such as publications, as well as profiling a range of others; for example, educational activities, engagement of practitioners or service users, and opportunities arising for follow-on funding/projects. Determining any impact on practice and service user outcomes remained, however, inherently more complicated to attribute.

Conclusion: Demonstrating research impact is complex, but vital. A multidimensional framework provides a practical means by which research impact can potentially be captured throughout, and beyond, the life of a project. Occupational therapists should consider the use of an impact assessment model to report, and share, the diversity of their research impact activities with funding bodies and the profession.

#### References

Beacham B, Kalucy L, McIntyre E (2005) Understanding and measuring research impact. (Focus On, Issue 2). Adelaide: Primary Health Care Research and Information Service.

Buxton M, Hanney S (1996) How can payback from health services research be assessed? *Journal of Health Services Research Policy* 1(1), 35–43.

Higher Education Funding Council for England, Scottish Funding Council, Higher Education Funding Council for Wales, Department for Employment and Learning (2011) Research Excellence Framework 2014: assessment framework and guidance on submissions. (REF 02.2011). Bristol: HEFCE, Scottish Funding Council, HEFCW, Department for Employment and Learning.

Sarli CC, Dubinsky EK, Holmes KL (2010) Beyond citation analysis: a model for assessment of research impact. *Journal of the Medical Library Association*, 98(1), 17–23.

#### Author CVs

Mandy Sainty, Research and Development Manager at the College of Occupational Therapists, leads on research governance and supports the development of practice guidelines.

#### Contact

mandy.sainty@cot.co.uk

#### Keywords

Research, Knowledge transfer

### P135

#### Perspectives of the VdT model of creative ability

Sherwood J, London South Bank University, UK

Originating in South Africa, the Vona du Toit Model of Creative Ability (VdT MoCA) is widely applied in OT practice and taught on the majority of undergraduate programmes in South Africa (Casteleijn & Graham 2012). The VdT MoCA is becoming an established model and the model of choice in a number of NHS trusts and independent healthcare providers in the UK.

In 2013 an on-line survey was conducted of 72 occupational therapists and support workers to:

- Gain insight into perspectives on the VdT MoCA
- Identify potential areas for development in the model
- Identify potential professional development needs of participants
- Identify potential areas for research.

The results support Chinembiri's (2011) findings that the model is predominantly applied in mental health and learning disabilities services. However, the model is also used across a broad range of fields of practice and settings, including paediatrics. This survey provides new insights into the use of the model in the UK and its perceived strengths and limitations. The data predominantly indicates that respondents perceive the model as highly valuable for increasing their clinical reasoning, confidence and effectiveness as OTs and OT support workers, thus increasing job satisfaction and professional identity. The model is perceived to have good utility for practice; however, there are also indications of the need to improve resources on the model and a need for continuing professional development opportunities.

Following this survey, occupational therapists were also surveyed regarding their perspectives of the research priorities for the