

RESEARCH

Open Access



Does antibiotic resistance impair plasma susceptibility of multi-drug resistant clinical isolates of enterococci in vitro?

Matthias Napp¹, Sebastian von Podewils^{2*}, Ingo Klare³, Hermann Haase², Richard Kasch⁴, Denis Gumbel^{1,5}, Axel Ekkernkamp^{1,5}, Michael Jünger² and Georg Daeschlein²

Abstract

Background: Cold atmospheric plasma could constitute an alternative against multi-drug resistant pathogens. Susceptibility of enterococci to cold atmospheric plasma was investigated in vitro.

Methods: 39 clinical isolates of enterococci were grouped dependent on the most important resistance patterns and treated on agar using dielectric barrier discharge plasma. These included enterococci with combined vancomycin- and high-level gentamicin resistance, high-level resistance to gentamicin (HLGR) only, vancomycin resistance alone (VRE), and enterococci susceptible to both. Susceptibility to cold atmospheric plasma was evaluated based on the zones of inhibition and examined in terms of the enterococcal group and the “degree” of drug resistance.

Results: Cold atmospheric plasma treatment killed all groups. Comparison of VRE and HLGR strains with non-VRE and non-HLGR isolates concerning zones of inhibition revealed that enterococci with special resistance patterns (VRE and HLGR) showed significantly smaller zones of inhibition than the sensitive ones. The mean of all isolates, irrespective of belonging to groups, showed smaller zones of inhibition with increasing “degree” of drug resistance.

Conclusions: Cold atmospheric plasma treatment killed all isolates of enterococci, but its efficacy depended on the “degree” of drug resistance and on membership in special resistance groups with particular clinical-outbreak importance. However, a possible role of the different genetic lineages, which might be prone to acquiring more or less resistance phenotypes, may also play a role in this context.

Keywords: Dielectric barrier discharge plasma (DBD), Cold atmospheric plasma (CAP), Antimicrobial efficacy, Plasma susceptibility, Plasma medicine, Vancomycin-resistant enterococci, Gentamicin-resistant enterococci

Background

The microorganisms most often involved in multi-drug (MDR) resistance are the so-called ESKAPE pathogens (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and enterobacteriaceae except *Klebsiella*), suggesting their proclivity in escaping antibacterial treatments [1, 2]. Beyond the search for new antibiotic drugs, novel antimicrobial strategies as alternatives

to conventional agents with antimicrobial efficacy are strongly needed [3]. One new concept is cold atmospheric plasma (CAP), which combines potent physical properties such as ultraviolet and infrared radiation (within the near infrared region), reactive oxidative species, and charged particles [4–7]. In previous studies, the present authors showed significant efficacy of CAP in vitro against bacterial and fungal species [5, 8]. Successful treatments of different infectious diseases based on the considerable antimicrobial efficacy of CAP have been documented; hence, it can be deduced that plasma could be effective in hospital hygiene and wound medicine, as chronic wounds represent an important risk factor for MDR colonization [9–16].

*Correspondence: sebastian.podewils@uni-greifswald.de

² Department of Dermatology, Ernst Moritz Arndt University, Sauerbruchstrasse, 17475 Greifswald, Germany

Full list of author information is available at the end of the article

However, to fulfil aforementioned high expectations in CAP as innovative alternative in the treatment of MDR pathogens, CAP efficacy should not be affected by resistance to antibiotics.

Therefore, two factors possibly influencing plasma susceptibility of enterococci were investigated: first, the “degree” of multi-drug resistance of enterococci, and second, the role of pathogens known as outbreak strains—the vancomycin-resistant enterococci (VRE) and/or enterococci resistant to high doses of gentamicin (HLGR).

Methods

The aim of this in vitro-study was to determine the susceptibility of clinical isolates of enterococci showing different patterns of antimicrobial resistance to CAP treatment.

Plasma source

The plasma source used was a dielectric barrier discharge (DBD) plasma device with a high voltage electrode of 20 mm in diameter developed by the CINOXY GmbH, Duderstadt, Germany. The electrode is being covered with a dielectric barrier made of an industrial ceramic. The pulse repetition rate of the device was adjusted on 250 Hz which leads to electric power dissipated in the gas discharge within the range of 167–237 mW [17]. The area covered by the visible plasma beam averaged 314 mm².

Susceptibility test model in vitro

CAP susceptibility was calculated as described detailed in a previous publication [5]. In few words, before starting plasma treatment, the strains were thawed and cultured aerobically at 36 °C overnight on sheep blood agar (Columbia agar, Biomérieux, Nürtingen, Germany). Suspensions were made out of these overnight cultures, diluted in sterile saline (1 colony in 1 ml). This suspension was diluted 1/100 in sterile saline (optical density was proved by measurement of McFarland) and 100 ml were plated on Columbia blood agar under sterile conditions. All isolates of enterococci were treated for 3 s by CAP on agar. Treatments were done under controlled

room climate with temperatures not surpassing 21 °C and a humidity of ca. 45 %.

To compute the CAP susceptibility, the diameters (mean of two measurements at a 90° angle) of the obtained circular zones of inhibition (ZOI) were measured and the area calculated.

Microbiology

A total of 39 clinical isolates were included, 16 *E. faecium* and 23 *Enterococcus faecalis*. Isolates were recovered consecutively during routine wound microbiology at the University Hospital of Greifswald, Germany. VRE identified as outbreak strains and isolated elsewhere were provided by the National Reference Center for Staphylococci and Enterococci, Robert Koch Institute, Wernigerode, Germany [18].

The different isolates were grouped according to their resistance pattern, including sensitive, vancomycin resistance (VRE) and high level gentamycin resistance (HLGR) as seen in Table 1 (membership of one isolate in several groups was possible).

Irrespective of above-mentioned resistance patterns were all 39 isolates differentiated into subgroups depending on their “degree” of antimicrobial resistance (resistance group, RG), due to the automated break-point determination test (breakpoints based on DIN and CASFM as interpretation criteria) in the VITEK[®] 2 Compact System (bioMérieux, Nürtingen, Germany). The 12 classes of antibiotic agents tested are given in Fig. 1 with the corresponding percentage of isolates showing resistance. The maximum number of resistances of one isolate in this study was 10, meaning that this isolate was placed in RG10.

Statistical analysis


A paired t test with a significance level of 5 % was applied to compare the species-specific (groups 1–6) ZOI obtained after plasma treatment. Furthermore, an analysis of variance was performed to test for significant differences related to the “degree” of resistance (RG 2–10). For all calculations, the free software package R

Table 1 Total number (n) of isolates and number of specific species within the different groups

	Group 1 other enterococci (oE) ^b	Group 2 HLGR only	Group 3 ^a group 2 + 4	Group 4 VRE and HLGR	Group 5 ^a group 6 + 4	Group 6 VRE only
n of isolates	12	11	19	8	16	8
<i>E. faecium</i>	1	5	9	4	10	6
<i>E. faecalis</i>	11	6	10	4	6	2

^a Group 3: containing all HLGR strains, group 5: containing all VRE strains

^b Enterococci with susceptibility to vancomycin and high concentrations of gentamicin



(R Development Core Team, R Foundation for Statistical Computing, Vienna, Austria) was used.

Results

In all enterococci isolates plasma treatment resulted in circular bacteria-free areas (ZOI) in the homogeneous growth zone of the agar surface within 48 h.

The ZOI of the enterococcal groups 1–6 are presented in Fig. 2. The group of “other enterococci” (group 1) showed the biggest ZOI (486 mm²). The VRE (group 6) exhibited the smallest ZOI (290 mm²). The difference between the maximum (oE, group 1) and minimum ZOI (only VRE, group 6) measured 178.4 mm², equal to 38.1 %. Additionally, *E. faecium* exhibited higher CAP resistance than *E. faecalis*.

The comparison of groups and pooled groups of enterococcal isolates with different antimicrobial resistance patterns (i.e. VRE, HLGR, etc.) revealed significant differences in sizes of ZOI (Table 2). Significantly smaller ZOI were found in VRE strains (group 5) as well as in HLGR isolates (group 3) compared with non-VRE and non-HLGR isolates, respectively. As shown by the significantly larger ZOI in group 1 compared with all other groups, the highest CAP efficacy was obtained in the

group of enterococci without the clinically important resistance patterns (group 1).

The statistical analysis of single isolates regarding the “degree” of drug resistance revealed that isolates with superior RG numbers showed smaller ZOI (p = 0.045) than isolates with lower RG numbers (Fig. 3). These findings did not occur within the statistical analysis of the enterococcal (resistance pattern) groups, probably due to the very small numbers within the different groups (Fig. 4a–f).

Discussion

Clinically relevant enterococci primarily include two species, *E. faecalis* and *E. faecium*. Both species exhibit high levels of resistance against the aminoglycosides, and *E. faecium* is inherently resistant to β-lactam antibiotics. Currently, most isolates of *E. faecium* are highly resistant to ampicillin, while a non-negligible proportion exhibit high-level resistance to aminoglycosides and are resistant to glycopeptides, causing outbreaks in clinical settings [19].

Additionally, an increasing resistance to the newer antimicrobials linezolid, daptomycin and tigecycline could be observed in many countries [20].




Table 2 Significant differences (p value) in ZOI between the different enterococcal groups

Compared groups	Mean (SD) ZOI (mm ²) of respective group	p value
Group 1 vs. 2	468.1 (71.6) vs. 334.4 (69.1)	0.000
Group 1 vs. 4	468.1 (71.6) vs. 301.0 (44.8)	0.000
Group 4 vs. 6	301.0 (44.8) vs. 289.7 (42)	0.000
Group 1 vs. all other isolates	468.1 (71.6) vs. 311.3 (56.0)	0.000
Group 3 vs. all other isolates	320.3 (61.0) vs. 396.8 (105.3)	0.017
Group 4 vs. all other isolates	301.0 (44.8) vs. 374.6 (97.7)	0.014
Group 5 vs. all other isolates	295.4 (42.4) vs. 404.2 (94.8)	0.000
Group 6 vs. all other isolates	289.7 (42.0) vs. 377.6 (95.7)	0.002



Few studies have been published on plasma susceptibility of multidrug-resistant strains (e.g. VRE), such as those referring to decontamination of surfaces in hospital environments [21, 22]. However, no data on enterococci in terms of antibiotic resistance are available. From previous studies focussing on basic plasma suitability for inactivating wound and skin bacteria [8, 20, 23], no statistically supported conclusions on enterococcal susceptibility to plasma treatment can be drawn. Therefore, the

antimicrobial efficacy of plasma from a well-researched plasma source (DBD) against a set of clinical isolates of enterococci, including VRE and HLGR, was investigated in vitro.

It was clearly demonstrated that the DBD easily eliminated all isolates of enterococci examined here, regardless of special resistance patterns such as VRE and HLGR. Interestingly, significant differences were found when the obtained ZOI diameters of distinct groups were compared.

The most obvious finding was that *E. faecium* was more resistant to plasma treatment than *E. faecalis*, which resembles the well-known greater drug susceptibility of *E. faecalis* compared to that of *E. faecium*. The reasons for the difference in plasma susceptibility of these two species could not get cleared in this study as the number of isolates did not allow deeper statistical analysis. Nevertheless, this result could be the next hint to related resistance mechanisms to CAP and antibiotics in enterococci.

The next important observation is that the authors found lower CAP susceptibility of VRE and HLGR treated with DBD in comparison with enterococci lacking these clinically important resistance patterns.



In addition, “higher” resistance to chemical antibiotics was correlated with less susceptibility to DBD (decrease in killing-zone diameter). This correlation was found only for all isolates irrespective of belonging to specific resistance groups probably due to the very small numbers within the different groups.

Based on the overall results, an influence of vancomycin resistance and HLGR on CAP susceptibility can

be assumed for DBD, since the properties “vancomycin resistance” and “HLGR alone” confer significantly smaller ZOI on agar. However, a possible influence of the different genetic lineages, which might be prone of acquiring more or less resistant phenotypes, may also play a role in this context.

Although the data presented here strongly indicate plasma susceptibility of all enterococci, the lower

susceptibility of *E. faecium* (compared to *E. faecalis*), VRE, HLGR, and generally MDR enterococci against DBD suggest the need for further research to clarify the as-yet unknown mechanisms by which bacteria are killed by CAP. This phenomenon could also help recognize and manage potential bacterial resistance mechanisms, which have not yet been documented.

Conclusion

Plasma treatment with DBD was followed by a significant reduction of all tested enterococci, irrespective of species and antibiotic resistance pattern. However, the obtained data indicate significantly less susceptibility of VRE and HLGR enterococci as well as less susceptibility of *E. faecium* to CAP derived from a DBD source and a correlation between decreasing susceptibility to CAP and increasing resistance level of enterococci (Additional file 1).

Additional file

Additional file 1. Each strain is given with their corresponding zone of inhibition and their detailed profile of antibiotic resistance.

Abbreviations

CAP: cold atmospheric plasma; DBD: dielectric barrier discharge; HLGR: high-level resistance to gentamicin; MDR: multi-drug resistant; oE: other enterococci; RG: resistance group; SD: standard deviation; VRE: vancomycin resistant enterococci; XDR: extreme drug resistant; ZOI: zones of inhibition.

Authors' contributions

MN, SP and GD made substantial contributions to conception and design as well to acquisition, analysis and interpretation of data. HH made substantial contributions to analysis of data. IK, RK, DG, AE and MJ have been involved in drafting the manuscript and revising it critically for important intellectual content. All authors read and approved the final manuscript.

Author details

¹ Department of Surgery, Ernst Moritz Arndt University, Sauerbruchstrasse, 17475 Greifswald, Germany. ² Department of Dermatology, Ernst Moritz Arndt University, Sauerbruchstrasse, 17475 Greifswald, Germany. ³ National Reference Center for Staphylococci and Enterococci, Robert-Koch-Institute, Wernigerode branch, Burgstraße 37, 38855 Wernigerode, Germany. ⁴ Department of Orthopaedics, Ernst Moritz Arndt University, Sauerbruchstrasse, 17475 Greifswald, Germany. ⁵ Unfallkrankenhaus Berlin (ukb), Warener Straße 7, 12683 Berlin, Germany.

Acknowledgements

Not applicable.

Competing interests

All authors disclose any financial and/or non-financial competing interest (e.g. employment, consulting fees, research contracts, stock ownership, patent licenses, honoraria, advisory affiliations, etc.).

Funding

This study was performed within the joint research project "Campus PlasmaMed" supported by the *German Federal Ministry of Education and Research* (Grant Nos. 13N9773 and 13N9779).

The role of the funding body did not include any influence in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Received: 10 June 2016 Accepted: 22 August 2016

Published online: 01 September 2016

References

- Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens. *Expert Rev Anti Infect Ther*. 2013;11:297–308.
- Boucher HW, Talbot GH, Bradley JS, Edwards JE, Gilbert D, Rice LB, Scheld M, Spellberg B, Bartlett J. Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America. *Clin Infect Dis*. 2009;48:1–12.
- Appelbaum PC. 2012 and beyond: potential for the start of a second pre-antibiotic era? *J Antimicrob Chemother*. 2012;67:2062–8.
- Daeschlein G, Scholz S, Arnold A, von Podewils S, Haase H, Emmert S, von Woedtke T, Weltmann K-D, Jünger M. In vitro susceptibility of important skin and wound pathogens against low temperature atmospheric pressure plasma jet (APPJ) and dielectric barrier discharge plasma (DBD). *Plasma Process Polym*. 2012;9:380–9.
- Daeschlein G, von Woedtke T, Kindel E, Brandenburg R, Weltmann K-D, Jünger M. Antibacterial activity of an atmospheric pressure plasma jet against relevant wound pathogens in vitro on a simulated wound environment. *Plasma Process Polym*. 2010;7:224–30.
- Ehlbeck J, Brandenburg R, von Woedtke T, Krohmann U, Stieber M, Weltmann KD. PLASMOSE—antimicrobial effects of modular atmospheric plasma sources. *GMS Krankenhhyg Interdisziplin*. 2008;3:14.
- Fridman G, Brooks AD, Balasubramanian M, Fridman A, Gutsol A, Vasilets VN, Ayan H, Friedman G. Comparison of direct and indirect effects of non-thermal atmospheric-pressure plasma on bacteria. *Plasma Process Polym*. 2007;4:370–5.
- Daeschlein G, Scholz S, Woedtke TV, Niggemeier M, Kindel E, Brandenburg R, Weltmann KD, Junger M. In vitro killing of clinical fungal strains by low-temperature atmospheric-pressure plasma jet. *IEEE Trans Plasma Sci*. 2011;39:815–21.
- Bowler PG, Davies BJ. The microbiology of infected and noninfected leg ulcers. *Int J Dermatol*. 1999;38:573–8.
- Gostev VA, Ignakhin VS, Popova EK, Ostashkov OA. Cold plasma—a powerful agent for biological applications. In: Güçeri S, Fridman A, Gibson K, Haas C, editors. *Plasma assisted decontamination of biological and chemical agents*. Dordrecht: Springer; 2008. p. 65–77.
- Heinlin J, Morfill G, Landthaler M, Stolz W, Isbary G, Zimmermann JL, Shimizu T, Karrer S. Plasma medicine: possible applications in dermatology. *J Dtsch Dermatol Ges*. 2010;8:968–76.
- Lademann J, Richter H, Alborova A, Humme D, Patzelt A, Kramer A, Weltmann KD, Hartmann B, Ottomann C, Fluhr JW, et al. Risk assessment of the application of a plasma jet in dermatology. *J Biomed Opt*. 2009;14:054025.
- Lademann O, Richter H, Patzelt A, Alborova A, Humme D, Weltmann KD, Hartmann B, Hinz P, Kramer A, Koch S. Application of a plasma-jet for skin antiseptics: analysis of the thermal action of the plasma by laser scanning microscopy. *Laser Phys Lett*. 2010;7:458–62.
- Melendez JH, Frankel YM, An AT, Williams L, Price LB, Wang NY, Lazarus GS, Zenilman JM. Real-time PCR assays compared to culture-based approaches for identification of aerobic bacteria in chronic wounds. *Clin Microbiol Infect*. 2010;16:1762–9.
- Stoffels E. Cold plasma treatment in wound care: efficacy and risk assessment. In: 60th gaseous electronics conference, vol 52. Arlington: Virginia Bulletin of the American Physical Society; 2007.
- Stoffels E, Kieft IE, Sladek REJ, van den Bedem LJM, van der Laan EP, Steinbuch M. Plasma needle for in vivo medical treatment: recent developments and perspectives. *Plasma Sources Sci Technol*. 2006;15:S169.
- Helmke A, Hoffmeister D, Mertens N, Emmert S, Schuette J, Vioel W. The acidification of lipid film surfaces by non-thermal DBD at atmospheric pressure in air. *New J Phys*. 2009;11:115025.
- Knoll M, Daeschlein G, Okpara-Hofmann J, Klare I, Wilhelms D, Wolf HH, Borneff-Lipp M. Outbreak of vancomycin-resistant enterococci (VRE) in a

- hematological oncology ward and hygienic preventive measures. A long-term study. *Onkologie*. 2005;28:187–92.
19. Klare I, Witte W, Wendt C, Werner G. Vancomycin-resistant enterococci (VRE). Recent results and trends in development of antibiotic resistance. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2012;55:1387–400.
 20. Marshall SH, Donskey CJ, Hutton-Thomas R, Salata RA, Rice LB. Gene dosage and linezolid resistance in *Enterococcus faecium* and *Enterococcus faecalis*. *Antimicrob Agents Chemother*. 2002;46:3334–6.
 21. Cahill OJ, Claro T, O'Connor N, Cafolla AA, Stevens NT, Daniels S, Humphreys H. Cold air plasma to decontaminate inanimate surfaces of the hospital environment. *Appl Environ Microbiol*. 2014;80:2004–10.
 22. Klampfl TG, Isbary G, Shimizu T, Li YF, Zimmermann JL, Stolz W, Schlegel J, Morfill GE, Schmidt HU. Cold atmospheric air plasma sterilization against spores and other microorganisms of clinical interest. *Appl Environ Microbiol*. 2012;78:5077–82.
 23. Daeschlein G, Scholz S, Ahmed R, Majumdar A, von Woedtke T, Haase H, Niggemeier M, Kindel E, Brandenburg R, Weltmann KD, Junger M. Cold plasma is well-tolerated and does not disturb skin barrier or reduce skin moisture. *J Dtsch Dermatol Ges*. 2012;10:509–15.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

