



Draft Genome Sequence of *Candida pseudohaemulonii* Isolated from the Blood of a Neutropenic Patient

Ratna Mohd Tap,^a Nur Amalina Kamarudin,^a Stephanie Jane Ginsapu,^a Ahmed Rafezzan Ahmed Bakri,^b Norazah Ahmad,^a Fairuz Amran,^a Matthias Sipiczki^c

^aMycology Laboratory, Bacteriology Unit, Infectious Diseases Research Centre, Institute for Medical Research, Jalan Pahang, Kuala Lumpur, Malaysia

^bMicrobiology Unit, Pathology Department, Queen Elizabeth Hospital, Kota Kinabalu, Sabah, Malaysia

^cDepartment of Genetics and Applied Microbiology, University of Debrecen, Debrecen, Hungary

ABSTRACT *Candida pseudohaemulonii* is phylogenetically close to the *C. haemulonii* complex and exhibits resistance to amphotericin B and azole agents. We report here the draft genome sequence of *C. pseudohaemulonii* UZ153_17 isolated from the blood culture of a neutropenic patient. The draft genome is 3,532,003,666 bp in length, with 579,838 reads, 130 contigs, and a G+C content of 47.15%.

Candida pseudohaemulonii was first isolated in 2006 from the blood of a patient in Thailand (1). It is phylogenetically similar to *C. haemulonii*, which is also resistant to amphotericin B. *C. pseudohaemulonii* cannot be well differentiated by current commercial identification methods from its other related species, *C. haemulonii*, *C. haemulonii* var. *vulnera*, *C. duobushaemulonii*, and *C. auris*. Thus, a molecular barcode was proposed for fungal identification, namely, the internal transcribed spacer (ITS) region, the D1/D2 domain of the large subunit (LSU) region in the rRNA gene, and the matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI-TOF MS) (2). In this article, we report the draft genome sequence of *C. pseudohaemulonii* UZ153_17 isolated from the blood culture of a neutropenic patient with short bowel syndrome.

DNA was extracted using the cetyltrimethylammonium bromide (CTAB) method (3), and a DNA library was prepared following the Illumina protocol using a TruSeq DNA LT sample prep kit (Illumina, San Diego, CA, USA). The genomic DNA was sequenced to a 451-fold depth of coverage using the Illumina HiSeq 2500 platform. From our analysis, the estimated genome size of *C. pseudohaemulonii* UZ153_17 is 14.74 Mb. The resulting draft genome's total size is 3,532,003,666 bp, with an overall G+C content of 47.15%. The genome was assembled into 130 contigs using the SOAPdenovo version 2.04 assembler (4). All of the generated contigs were formed into 90 scaffolds, of which 47 are more than 1,000 bp in length, with a scaffold N_{50} size of 799,808 bp. The total size of all the scaffolds is 12,700,538 bp. Subsequent gene prediction analysis using MAKER2 (5) yielded a total of 5,540 predicted protein-coding genes with an average length of 1,492 bp. Genome annotation on the predicted genes was carried out by BLAST similarity searches against Swiss-Prot, a protein sequence database, with a total of 60.99% genes successfully annotated.

The genome is found to contain fluconazole and multidrug resistance genes. This finding is parallel with other studies that have shown that the isolate was resistant to amphotericin B and fluconazole (1, 2). The draft genome sequence of *C. pseudohaemulonii* will contribute to further investigations into the molecular mechanisms that lead to antifungal drug resistance.

Accession number(s). The nucleotide sequence of the *C. pseudohaemulonii* UZ153_17 genome has been deposited in DDBJ/EMBL/GenBank under accession no. [PEJZ00000000](https://www.ncbi.nlm.nih.gov/nuclseq/PEJZ00000000).

Received 12 February 2018 **Accepted** 27 February 2018 **Published** 5 April 2018

Citation Mohd Tap R, Kamarudin NA, Ginsapu SJ, Ahmed Bakri AR, Ahmad N, Amran F, Sipiczki M. 2018. Draft genome sequence of *Candida pseudohaemulonii* isolated from the blood of a neutropenic patient. *Genome Announc* 6:e00166-18. <https://doi.org/10.1128/genomeA.00166-18>.

Copyright © 2018 Mohd Tap et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ratna Mohd Tap, ratna@imr.gov.my.

ACKNOWLEDGMENTS

This study was supported by a research grant from the Ministry of Health of Malaysia.

We would like to thank the Director General of Health, Malaysia and the Director of Institute for Medical Research for permission to publish this article.

REFERENCES

1. Sugita T, Takashima M, Poonwan N, Mekha N. 2006. *Candida pseudohaemulonii* sp. nov., an amphotericin B- and azole-resistant yeast species, isolated from the blood of a patient from Thailand. *Microbiol Immunol* 50:469–473. <https://doi.org/10.1111/j.1348-0421.2006.tb03816.x>.
2. Cendejas-Bueno E, Kolecka A, Alastruey-Izquierdo A, Theelen B, Groenewald M, Kostrzewa M, Cuenca-Estrella M, Gómez-López A, Boekhout T. 2012. Reclassification of the *Candida haemulonii* complex as *Candida haemulonii* (*C. haemulonii* group I), *C. duobushaemulonii* sp. nov. (*C. haemulonii* group II), and *C. haemulonii* var. *vulnera* var. nov.: three multiresistant human pathogenic yeasts. *J Clin Microbiol* 50:3641–3651. <https://doi.org/10.1128/JCM.02248-12>.
3. Lööke M, Kristjuhan K, Kristjuhan A. 2011. Extraction of genomic DNA from yeasts for PCR-based applications. *Biotechniques* 50:325–328.
4. Li R, Li Y, Kristiansen K, Wang J. 2008. SOAP: short oligonucleotide alignment program. *Bioinformatics* 24:713–714. <https://doi.org/10.1093/bioinformatics/btn025>.
5. Holt C, Yandell M. 2011. MAKER2: an annotation pipeline and genome-database management tool for second-generation genome projects. *BMC Bioinformatics* 12:491. <https://doi.org/10.1186/1471-2105-12-491>.