

4-15-2019

Probing the Nanomechanical Behavior of Cells and Cell Nuclei

Jesse Schimpf
Boise State University

Josh Newberg
Boise State University

Gunes Uzer
Boise State University

Paul H. Davis
Boise State University



Probing the Nanomechanical Behavior of Cells and Cell Nuclei

Jesse Schimpf,¹ Josh Newberg,² Gunes Uzer,² and Paul H. Davis^{1,*}

BOISE STATE UNIVERSITY

¹Micron School of Materials Science & Engineering, ²Department of Mechanical & Biomedical Engineering, *Mentor

Atomic Force Microscopy (AFM)


AFM is a scanning probe microscopy (SPM) technique used to generate high resolution topographical maps of a surface. Images are obtained by mechanically oscillating a cantilever with a sharp tip while scanning over a small (< 100 μm wide) area.



Bruker Dimension Icon FastScan AFM¹


Nanomechanical Measurements

The cantilever acts as a spring. If its spring constant (applied force required per unit distance of deflection, expressed in N/m or nN/nm) and deflection sensitivity (relative photodiode signal as a function of deflection, in V/nm) are known, the cantilever's measured deflection can be converted into a force measurement at each point along the scan. Piezoelectric actuators can precisely control the displacement of the probe, which allows the acquisition of a force-displacement curve and subsequent mechanical properties analysis. The interaction between a spherical probe and the sample can be modeled with either Hertzian or JKR (Johnson, Kendell, and Roberts) dynamics. JKR is an extension of the Hertzian model that takes into account the effect of adhesive forces between the tip and sample during the retract curve.




Nanomechanical Properties of MSC Nuclei


Mesenchymal stem cells (MSCs) are multipotent, capable of differentiating into bone, cartilage, muscle, or fat cells. A combination of mechanical and chemical signals guide this transformation; the goal of this study is to understand how mechanical signals affect MSCs.



One measurable response seen in MSCs is a change in the elastic or Young's modulus (i.e., the "stiffness") of the nucleus, which can be calculated from the slope of a force-displacement curve.




Rather than a sharp AFM tip, a 10 μm diameter glass bead is used to indent on nuclei because it has an ~5,000x larger radius of curvature. This disperses the applied force, preventing the probe from puncturing the cell membrane. It also allows a more holistic measurement of the nucleus's mechanical properties.



AFM cantilever with 10 μm diameter glass sphere attached.

Cells vs. Isolated Nuclei

Nanoindentation was performed on MSCs and isolated MSC nuclei. Noticeable structural differences were observed between whole cells and isolated nuclei, including a decrease in nuclear area.




(a) Whole MSC vs (b) isolated nucleus at same magnification.

Low Intensity Vibration and Structural Knockout

• Low intensity vibration (LIV) was applied to unaltered MSCs before isolating and testing the nuclei. LIV was performed for 20 minute intervals with 1 hour rests in between. LIV 1X represents two intervals and LIV 2X represents four intervals.

• In a separate group, structural component knockouts were performed on MSCs before isolating and testing the nuclei. The effects of both Lamin A/C and Sun-1&2 knockouts were measured (separately).



Conclusions

- Removing the cytoskeleton causes the nucleus to shrink and significantly reduces nuclear stiffness.
- Extended LIV treatment introduces a stressful mechanical environment for the nucleus, triggering an increase in nuclear stiffness.
- Lamin A/C and Sun-1&2 provide structural support for the cell. When removed, nuclear stiffness decreases significantly. This suggests that these components also impart structure to the nucleus.

Author:

Jesse Schimpf
Undergraduate Research Assistant
Surface Science Laboratory
Micron School of Materials Science & Engineering
jesseschimpf@u.boisestate.edu



Co-Authors:

Josh Newberg, joshnewberg@u.boisestate.edu
Gunes Uzer, gunesuzer@boisestate.edu
Paul H. Davis (mentor), pauldavis2@boisestate.edu

References:

1. Bruker.
2. Chan, Coucouvanis, Tousey, Andersen, and Jessie. "Improved Expansion of MSC Without Loss of Differentiation Potential" R&D Systems, Inc.

Funding:

COBRE Matrix Biology, NIH NIGMS P20GM109095
AR3T Technology Development Grant, NIH 5P2CHD086843-03 (Sub-Award 126873-13)