



Insufficient Reporting of Factors Associated with Exercise Referral Scheme Uptake, Attendance and Adherence: A Systematic Review of Reviews.

Journal:	<i>Journal of Physical Activity & Health</i>
Manuscript ID	JPAH.2018-0341.R2
Manuscript Type:	Review
Keywords:	physical activity, exercise prescription, program evaluation

SCHOLARONE™
Manuscripts

Accepted author manuscript version reprinted, by permission, from *Journal of Physical Activity and Health*, 2019, 16 (8), pp. 667-676. <https://doi.org/10.1123/jpah.2018-0341> © Human Kinetics, Inc.

1
2
3
4 **1 Abstract**

5
6 **2 Background:** Exercise Referral Schemes (ERS) are prescribed programmes aimed at tackling physical
7
8
9 **3** inactivity and associated non-communicable disease. Inconsistencies in reporting, recording and
10
11 **4** delivering ERS make it challenging to identify what works, why, and for whom.

12
13
14 **5 Methods:** PRISMA guided this narrative review of reviews. Fifteen electronic databases were
15
16 **6** searched for systematic reviews of ERS. Reviewers applied inclusion criteria and quality assessed via
17
18 **7** the AMSTAR tool. Data on uptake, attendance and adherence were extracted.

19
20
21 **8 Results:** Eleven reviews met the inclusion criteria. AMSTAR quality was medium. Definitions of
22
23 **9** uptake varied within reviews. Uptake ranged from 35%-81%. Groups reported as more likely to take
24
25 **10** up ERS included, (i) females and (ii) older adults. Attendance was defined variably but ranged from
26
27 **11** 12%-49%. Men were more likely to attend ERS. Effect of medical diagnosis upon uptake and
28
29 **12** attendance was inconsistent. Exercises prescribed were unreported and therefore, adherence to
30
31 **13** exercise prescriptions was unreported. The influence of theoretically-informed approaches on
32
33 **14** uptake, attendance and adherence was generally lacking, however, self-determination, peer support
34
35 **15** and supervision support were reported as influencing attendance.

36
37
38
39 **16 Conclusions:** There was insufficient reporting across studies about uptake, attendance and
40
41 **17** adherence. Complex interventions like ERS require consistent definitions, recording and reporting of
42
43 **18** these key facets, but this is not evident from the existing literature.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A Review of Reviews of Exercise Referral Schemes.

19 **Introduction**

20 The benefits of regular physical activity (PA) are well established ^(1,2,3,4), notably for adults who
21 engage in at least 150 minutes of moderate intensity activity or 75 minutes of vigorous intensity
22 activity per week, alongside activities that improve muscular strength on at least two days of the
23 week ⁽¹⁾. Despite the clear and causal relationship between achieving PA guidelines and health
24 outcomes ⁽¹⁾, physical inactivity is rising ⁽⁵⁾. Inactivity is now the fourth leading risk factor for global
25 mortality (6% of deaths globally), behind high blood pressure (13%), tobacco use (9%) and high
26 blood glucose (6%) ⁽⁵⁾. Contextually it must be acknowledged that while physical inactivity is a
27 standalone risk factor ⁽⁵⁾ PA status is a key determinant in moderating the additional risk factors of
28 high blood pressure and high blood glucose levels ^(5,4).

29 In recognition of this physical inactivity burden, the UK government has implemented population-
30 wide programmes ⁽⁶⁾. Additionally, at the individual level, tailored attempts to manage physical
31 inactivity has led to the expansion of exercise referral schemes (ERS). Individual's showing evidence
32 of non-communicable disease coupled with an inactive lifestyle are directly referred by allied health
33 practitioners to independent third party exercise facilities to undertake a structured and prescriptive
34 exercise programme ⁽⁷⁾. The popularity of the ERS model drove a rapid rise in the number of centres
35 offering referral programmes despite limitations presented within the literature evaluating the
36 implementation of schemes ⁽⁸⁾ and thereby limited evidence of what works, why and for whom.

37 Since 1998, a combination of systematic reviews, meta-analyses or narrative reviews of ERS have
38 been published focusing on different constructs surrounding the effectiveness of ERS ^{(2, 9, 10, 11, 12, 13, 14,}
39 ^{15, 16, 17, 18, 19)}. It is evident from this body of work that ERS is considered an important element in the
40 armoury to increase PA and tackle chronic disease ⁽²⁰⁾. However, the United Kingdom's National
41 Institute for Health and Care Excellence (NICE) have indicated that the varying nature of ERS
42 programmes makes it impossible to differentiate between the effectiveness and acceptability of
43 different schemes ⁽⁴⁾ in addressing physical inactivity and non-communicable disease. This conclusion

A Review of Reviews of Exercise Referral Schemes.

1
2
3 44 is largely due to little evidence of any effectiveness of specific schemes ⁽⁴⁾. It is crucial therefore, that
4
5 45 clear reporting of key ERS components is available to allow for components associated with
6
7 46 effectiveness to be replicated in future schemes. Factors such as reporting of referral uptake,
8
9 47 attendance and adherence, and the behaviour change techniques (BCT's) underpinning ERS uptake
10
11
12 48 and adherence are key components to understand for the following reasons:

15 49 *Referral uptake*

16
17 50 Knowledge of who does or does not take-up an ERS referral is imperative for improving the degree
18
19
20 51 of take-up ⁽¹¹⁾. Whether participant characteristics influence the proportion of ERS uptake is
21
22 52 uncertain. It is important to know what referral demographics are consistently reported, or
23
24 53 unreported, and how they relate to uptake; to gain an understanding of what and how these are
25
26 54 influencing scale of ERS uptake. In particular, it is not clear whether those with certain medical
27
28 55 conditions have a greater uptake, and whether socio-economic or demographic characteristics
29
30 56 influence an individual's uptake ⁽²¹⁾. Unless uptake and characteristics of people using ERS are
31
32
33 57 known then it is difficult to interpret the extent of generalizability of effect to groups of the
34
35 58 population i.e. understand what works, why and for whom.

38 59 *Attendance and Adherence*

39
40
41 60 Slade et al. ⁽²²⁾ described adherence as the extent to which a person's behaviour corresponds to the
42
43 61 agreed referral. It is important within the context of the present review to distinguish the difference
44
45 62 between adherence and attendance. Presently, adherence to ERS refers to the percentage of
46
47 63 sessions attended ^(11, 15), which may be more appropriately classified as attendance. Adherence then
48
49 64 would refer to the participant's behaviour in completing the prescribed PA. Attendance alone does
50
51 65 not signify a suitable level of exercise intensity undertaken to benefit from the prescribed PA. For
52
53 66 those who complete ERS, promising results were reported for reduced skinfolds, systolic blood
54
55 67 pressure and BMI ⁽²³⁾, greater self-efficacy to overcoming PA barriers ⁽²⁴⁾ and higher self-reported PA
56
57 68 levels ⁽²⁵⁾. However, failure to attend and adhere to the prescribed programme reduces the
58
59
60

A Review of Reviews of Exercise Referral Schemes.

69 opportunity a participant has for achieving these benefits. As such, it is important to understand
70 what characteristics influence participants' attendance and adherence levels. A valid and reliable
71 assessment of both attendance and adherence is essential for drawing valid conclusions about ERS
72 ⁽²²⁾.

Behaviour Change Techniques

74 In evaluating a complex intervention such as ERS, there is a requirement to understand the
75 theoretical underpinning at every stage of the intervention, what it is based upon, and the
76 mechanisms through which behaviour change is achieved ^(26, 27). In the same way that the PA
77 content of ERS schemes varies, and is determined between service provider and participant ⁽²⁸⁾, no
78 single explicit behaviour change theory or technique is embedded within ERS ⁽²⁶⁾. The explicit
79 reporting of the components within a complex intervention such as ERS, including its contextual
80 factors such as BCT's utilised, is imperative in understanding; what facilitates uptake, attendance
81 and adherence to ERS and further implementation of ERS ⁽²²⁾.

82 The current review of reviews aims to systematically collate and evaluate the evidence base of
83 review level findings around ERS. It will address what is reported and the consistent and
84 contradictory observations at both a theoretical and practical level. The review aims to: 1) describe
85 definitions rates and participant characteristics associated with ERS uptake; 2) to describe definitions
86 rates and participant characteristics associated with attendance at and adherence to ERS schemes;
87 and 3) to describe theoretically-informed approaches associated with ERS uptake and attendance.
88 Furthermore, this review will establish what key features of ERS reporting go relatively under-
89 reported to inform future studies of ERS. This review may lead to a clear consensus positively
90 affecting both research and practice in a critical area that could improve the health and wellbeing of
91 individual's with non-communicable disease referred to ERS.

A Review of Reviews of Exercise Referral Schemes.

92 **Methods**

93 **Data Sources and Search Strategy**

94 The PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) statement
95 guided the conduct of this narrative review of reviews ⁽²⁹⁾. Studies were identified by structured
96 electronic database searches. One author (RP), who is a librarian and information specialist,
97 searched 15 electronic databases (CINHAL, Medline, Embase, Cochrane Library, SportsDiscus,
98 PsychInfo, SCOPUS, HMIC, AMED, Public Health Database, ASSIA, CRD databases, Prospero, Web of
99 Knowledge, Campbell Collaboration Library and PubMed), in November 2016, for systematic reviews
100 and meta-analysis of exercise referral schemes in adults published in English language with no date
101 restriction. Additional reviews were searched until June 2017, via reference lists of included
102 literature, alongside searches of UpToDate, BMJ Best Practice and Dynamed Plus. Search strategies
103 were constructed and then amended and agreed by group consultation of all authors. As an
104 example, search terms used for Medline are available within the supplementary material
105 (Supplementary material 1). The protocol and the purpose of the current study were designed
106 beforehand; however, these were not published or uploaded on the web.

107 **Review Selection.**

108 Review inclusion and exclusion criteria were constructed by CS and then amended and agreed by
109 group consultation and described in Table 1. One author (CS) independently examined titles and
110 abstracts against the inclusion and exclusion criteria. Three authors (GH, TG & SG) independently
111 assessed a randomised sub sample of papers, amounting to 20% of the total title and abstract
112 sample. A group (CS, GH, TG & SG) consensus conferred on which papers were progressed further to
113 full text review and which excluded. Full text review articles were then obtained and assessed by
114 two authors (CS & SG) against the inclusion and exclusion criteria described in Table 1. Figure 1
115 describes the justification of excluding papers at multiple stages of the PRISMA protocol. Any
116 discrepancies were discussed until a consensus was reached.

A Review of Reviews of Exercise Referral Schemes.

117 **Data Extraction**

118 Data was extracted by one author (CS) against a data extraction template and verified by one co-
119 author (SG), with any discrepancies discussed until resolution. Eleven predefined constructs relating
120 to the effectiveness of interventions were used (supplementary material 2), taking guidance from
121 work conducted by Greaves et al. ⁽³⁰⁾.

122 **Grading of Evidence**

123 Each review was graded independently and empirically in duplicate (by CS & SG) with any
124 discrepancies discussed. The AMSTAR (A Measurement Tool to Assess Systematic Reviews) criteria
125 was used to grade reviews. AMSTAR grades scores as low quality (0-3), medium (4-7) and high (8-11)
126 ⁽³¹⁾. While the AMSTAR tool was developed to assess reviews of randomized controlled trials (RCTs)
127 only, it has been shown to be applicable to reviews of non-randomized studies, demonstrating good
128 psychometric properties ⁽³²⁾. Cohen's kappa was calculated as a measure of reliability for each
129 AMSTAR item. Kappa values less than 0 were rated as less than chance agreement; 0.01–0.20, slight
130 agreement; 0.21–0.40, fair agreement; 0.41–0.60, moderate agreement; 0.61–0.80, substantial
131 agreement; and 0.81–0.99, almost perfect agreement ⁽³²⁾. Additionally, reviews were graded on the
132 type of evidence they were reviewing, such as randomised controlled trials (see Table 2). The
133 classification of AMSTAR (e.g. 9) was then combined with the type of evidence (e.g. A). For example,
134 high-quality systematic reviews of randomised controlled trials was coded as 9A.

135 **Analysis**

136 A narrative synthesis of the results indicating the quality of the evidence was more appropriate for a
137 review of reviews and was considered for the following reasons: a lack of consistency within
138 reporting of results to undertake a meta-meta-analysis, and the variety of study interventions
139 examined within the reviews. Furthermore, the present review aimed to understand the 'how and
140 why' of ERS in order to gain an insight into the manner that ERS is reported. In accordance with
141 reporting guidelines for systematic reviews, a PRISMA checklist is available for this review.

A Review of Reviews of Exercise Referral Schemes.

142 **Results**

143 A systematic search identified 3211 potentially relevant articles (Figure 1). Following exclusion of
144 duplicates and non-relevant articles, 39 articles were retrieved in full and assessed. One additional
145 article was identified in March 2017 via a database alert that was initially set up from the original
146 database search. Eleven reviews met the pre-established inclusion criteria (Table 1).

147 **Review Characteristics**

148 All eleven reviews reported upon an aspect of effectiveness within ERS: seven reviews examined
149 referral to, uptake and attendance at ERS (2, 9, 10, 11, 12, 14, 15) with eight reporting for whom ERS is
150 successful (9, 11, 12, 13, 14, 15, 16, 18). Eleven reviews make comment on the theoretical underpinnings or
151 BCT's within ERS (2, 9, 10, 12, 13, 14, 15, 16, 18, 19). Eight reviews reported upon changes of physical activity
152 levels (10, 12, 13, 14, 15, 16, 18, 19). Reviews included data from a range of referral populations (e.g.
153 hypertension, diabetes, raised cholesterol, mental health and obesity) and delivery settings (e.g.
154 leisure centre, primary care, green settings, such as community outdoor walks or community
155 gardening). Dates of published studies included within the reviews ranged from 1966 to 2015 and
156 incorporated 221 cited papers of which 98 were duplicated across reviews. We further draw
157 attention of the reader to three linked reviews resulting from our search by Pavey *et al.* (11, 12, 13),
158 who report different aspects of the same systematic search, and one by Campbell *et al.* (14) who
159 based their initial search upon that of the Pavey papers (11, 12, 13) and built upon it. Where
160 appropriate, unique items are reported separately.

161 **Review quality**

162 The methodological quality of included reviews was generally of a medium standard (median
163 AMSTAR score = 7.50, mean AMSTAR = 7.00). Only three systematic reviews examined evidence
164 adjudged to be of high strength (Table 2). Table 3 illustrates the combined duplicate reviewer
165 AMSTAR scoring and strength of evidence within the reviews. The mean inter-rater reliability
166 (Kappa) for applying agreement on review quality was 0.67 (95% CI 0.87 to 0.48). The most common

A Review of Reviews of Exercise Referral Schemes.

167 methodological weaknesses adjudged via AMSTAR were, the lack of lists indicating the included and
168 excluded studies and the assessment of potential bias within the selection of articles (e.g. in meta-
169 analysis assessment of publication bias should include a combination of graphical aids such as funnel
170 plot and/or statistical tests such as Egger regression test or Hedges-Olken). The greatest
171 methodological strengths were the comprehensive nature of the literature searches performed and
172 the assessment and documentation of the scientific quality of the included studies. Table 4
173 highlights the reported uptake and attendance figures across reviews. Some reviews were able to
174 pool results in order to conduct meta-analysis. Other reviews were not able to achieve this due to
175 the lack of consistent reporting within the original papers. Recording and reporting methods of
176 uptake, attendance, adherence and theory varied within reviews and are detailed below.

177 **Narrative of Results**

178 **Referral uptake**

179 Uptake was defined within three linked reviews as attending the initial consultation ^(11, 12, 14). Two
180 reviews did not provide a definition for uptake ^(10, 15), while one review indicated that it struggled to
181 define uptake due to differences reported within evaluations ⁽⁹⁾. Table 4 highlights the reported
182 uptake across reviews; demonstrating a range of uptake values between 35% and 85%. Three of the
183 five reviews ^(9, 10, 12) which report on uptake highlight similar values around 65%. Importantly, no
184 review reported characteristics for participant who failed to take up ERS representing around 30-
185 40% of participants referred. Where reviews have reported characteristics relating to take-up, these
186 figures are derived from participants present within the scheme.

187 One high quality review reported that two studies demonstrated women were more likely to take-
188 up ERS than men ⁽¹²⁾. However, two studies within that review showed no association between
189 gender and uptake of ERS. Pavey et al. ⁽¹¹⁾ concluded that being female and of increasing age was a
190 positive predictor of uptake. A low quality review summarised that uptake was greater within
191 females (60 – 40% for females vs. males) ⁽⁹⁾. Their review importantly highlighted that reporting of

A Review of Reviews of Exercise Referral Schemes.

192 characteristics was poor and only reported within five primary studies ⁽⁹⁾. Pavey et al. ⁽¹¹⁾ reported
193 that participants who were more deprived and suffered from a respiratory diagnosis were more
194 likely to take-up ERS than those with the same diagnoses who were least deprived (OR 1.45, 95% CI
195 1.06 to 1.99, $p < 0.05$). A high quality review reported that pre-existing medical diagnoses were not
196 separately reported which prevented conclusions being drawn within these subgroups ⁽¹³⁾. One
197 medium quality review reported that participants referred with mental health problems were more
198 likely to participate in ERS than those with no medical referral (OR 1.79, 95% CI 1.24 to 2.39, $p < 0.01$)
199 ⁽¹¹⁾. However, participants with cardiovascular disease were more likely to engage than participants
200 with mental health problems (OR 0.33, 95% CI 0.27 to 0.57, $p < 0.01$), musculoskeletal problems (OR
201 0.75, 95% CI 0.58 to 0.99, $p < 0.05$), or overweight/obesity problems (OR 0.63, 95% CI 0.50 to 0.81,
202 $p < 0.01$). Overweight or obese participants were more likely to take-up ERS than smokers ⁽¹¹⁾. Referral
203 uptake to ERS was greater for participants referred from a cardiac nurse over those referred from
204 either a general practitioner or practice nurse ⁽¹¹⁾.

205 **Attendance and Adherence**

206 The most important finding from the reviews, was a lack of reporting on exercise prescription
207 adherence. No review detailed the type of exercise prescribed or the extent to which the participant
208 adhered to the prescription. Reviews did report the term adherence, however, it must be
209 understood that this term was a reference to attendance at the programme. As such, the term
210 adherence from the original reviews is replaced by the more appropriate term of attendance in the
211 current manuscript. Large inconsistencies were observed surrounding attendance figures. Four
212 studies failed to define attendance by determining a threshold, instead acknowledging the term as,
213 'duration of participation within ERS' ^(9, 10, 14, 15). Pavey and colleagues used a threshold of $\geq 75\%$
214 attendance of available sessions within an ERS programmes as its inclusion criteria for studies ^(11, 12).
215 Objective measures, such as the use of leisure centre records, were underutilised ⁽⁹⁾.

A Review of Reviews of Exercise Referral Schemes.

216 One high quality and one medium quality review, based upon the same initial systematic search,
217 reported that males were more likely to attend from two studies, while three further studies within
218 the reviews found no such association with attendance ^(11, 12). A further low quality review reported
219 attendance was higher in males ⁽⁹⁾. Their review did report one primary study citing higher
220 attendance in women and two RCT's finding no relationship between sexes ⁽⁹⁾. Increasing age was
221 positively associated with attendance to ERS from five studies in one review, however, it additionally
222 reported two studies suggesting no such relationship ⁽¹¹⁾. Gidlow et al. ⁽⁹⁾ reported two evaluations
223 indicating increasing age and being retired were associated with better attendance. However,
224 increasing age was found to reduce participation in PA from one RCT and one evaluation, while one
225 RCT reported no relationship ⁽⁹⁾. One medium quality review adds to these mixed views surrounding
226 increasing age by reporting that attendance appeared to be higher in older participants ⁽¹⁶⁾.
227 Increasing attendance was more prevalent in participants who had higher baseline activity levels and
228 were overweight ⁽¹⁶⁾. Gidlow et al. ⁽⁹⁾ reported one study, which indicated that socioeconomic
229 characteristics were unrelated to attendance. One high and one medium quality review indicated
230 that socioeconomic characteristics were poorly reported, preventing any clear conclusions on the
231 impact of different socioeconomic characteristics ^(16, 18).
232 One review acknowledged that a medical diagnosis was a factor that could affect attendance,
233 however they reported it was not consistently demonstrated throughout their included studies ⁽⁹⁾.
234 Participants with cardiovascular disease were more likely to attend than participants with pulmonary
235 disease ⁽¹¹⁾. Physical health problems were a greater predictor of attendance than mental health
236 issues (22% vs 34%, $p < 0.001$), however it was not reported what was defined as a physical health
237 problem ⁽¹¹⁾. Diabetic participants were less likely to attend ERS (OR 0.76, 95% CI 0.63 to 0.93,
238 $p < 0.01$) than those with cardiovascular disease ⁽¹¹⁾. Conversely, a medium quality review reported
239 that participants referred with sedentary lifestyles or diabetes demonstrated a higher adherence
240 than those with cardiovascular disease or obesity ⁽¹⁵⁾. A high quality review reported no consistent

A Review of Reviews of Exercise Referral Schemes.

241 difference in attendance rates between participants within one of three at-risk groups: smoking,
242 obesity and hypertension ⁽¹⁴⁾.

243 One high quality review identified seventeen studies which highlighted peer support and supervision
244 support from staff as a facilitator for attendance to the referral programme ⁽²⁾. Barriers to
245 attendance were noted as: the location (distance to travel); difficulties reaching the activities by
246 public transport; perceived safety of the location; timings of sessions; and cost ⁽²⁾. A medium quality
247 review indicated that European programmes had greater attendance over American, Australian or
248 Canadian programmes ⁽¹⁵⁾. A meta-regression shows the duration of an ERS programme is not
249 correlated to the attendance rate ⁽¹⁵⁾. However, a follow up period greater than six months, after the
250 cessation of the programme, did have a positive effect on attendance when the scheme was running
251 ⁽¹⁵⁾.

252 ***Behaviour Change Techniques***

253 High quality evidence from two reviews of randomised controlled trials ^(13, 18) showed that the
254 transtheoretical model of behaviour change was the most frequently utilised underlying theory of
255 intervention design, while social cognitive theory, theory of planned behaviour, and health belief
256 model were used less. One low quality review ⁽¹⁹⁾ that did not directly report on any theoretical
257 underpinnings, suggested that schemes should consider implementing theory driven approaches to
258 behaviour change. One medium ⁽¹⁰⁾ and one high ⁽¹¹⁾ quality review commented that the greatest
259 challenge for ERS was in increasing uptake and improving attendance ^(10, 11). One review reported
260 directly on how attendance were affected by theoretical techniques ⁽¹⁶⁾. Four reviews reported on
261 theoretical techniques to increase PA time or clinical markers ^(12, 13, 14, 18). With the exception of a few
262 primary studies, reviews reported little reference to the delivery, fidelity, or BCT's of health
263 behaviour change ^(10, 12, 13, 14, 18). Motivational interviewing was the most frequently utilised BCT, cited
264 within eight reviews ^(9, 10, 12, 13, 14, 16, 18, 19), with only one review indicating the fidelity of the technique
265 ⁽¹⁴⁾. One medium quality review reporting upon on a primary study, report that participants who

A Review of Reviews of Exercise Referral Schemes.

266 received motivational interviewing combined with free vouchers had a greater self-reported physical
267 activity score at 12 weeks, than a control group given advice alone (all groups combined, $P < 0.001$)
268 ⁽¹⁶⁾. The greatest increase in PA was observed in the intensive motivational interviewing and free
269 vouchers group which offered six motivational interviewing sessions (55%, $P < 0.001$). No difference
270 was evident at one year post-intervention between groups ⁽¹⁶⁾. Morgan ⁽¹⁶⁾ further reported that
271 exercise-plus-motivation vs exercise-only elicited a greater attendance over a 20 week programme
272 (62% vs 38%, $P < 0.05$). A further primary study reported that there was no significant difference
273 between three groups attending sessions when comparing, exercise alone, instructions based on the
274 'relapse prevention' model, or sessions that included re-enforcement methods (prizes for high
275 attendance). Attendance attrition was reported at 30% and 72% for all groups at 9 and 18 weeks,
276 respectively ⁽¹⁶⁾. One high quality review indicated that motivational interviewing was utilised
277 predominantly by different practitioners but provides no indication of its impact ⁽¹⁸⁾. A high quality
278 review reported two contrasting studies; one indicated that attendance was positively influenced by
279 participant levels of self-determination, and one study found no association between these
280 parameters ⁽¹²⁾. A medium quality review reported participant dissatisfaction when lacking social
281 support, or with an exercise leader lacking motivational skills ⁽¹⁰⁾.

282 Discussion

283 This review is the first to systematically identify, collate, and grade the evidence from review level
284 findings on ERS. The creation of this review highlights and summarises the consistent and
285 contradictory findings surrounding the reporting of: 1) uptake to ERS; 2) attendance and adherence
286 to ERS; and 3) BCT's relating to attendance and adherence. We discuss the characteristics of studies
287 and definitions used within them and how they affect uptake, attendance and adherence. We
288 highlight how a lack of reporting and knowledge of what is delivered within ERS limits our insight
289 into both attendance rates and adherence to the prescribed programme. Where reported,
290 considerations are made as to why participants with certain characteristics (e.g. medical referral) are
291 more likely to take up and attend schemes. Lastly, we discuss the reporting of theoretical constructs

A Review of Reviews of Exercise Referral Schemes.

292 and BCT's, proposing reporting tools that could be considered to help improve uptake and
293 attendance.

294 **Referral uptake**

295 Historically, through the reviews, it is reported that uptake of ERS sits around 60-70% of those
296 referred. This suggests that no improvements have been made in increasing the number of
297 participants starting a scheme over the years examined. Uptake to ERS was greater in randomised
298 controlled trials than observational evaluations, as reported in Table 4. Explanation for this could
299 centre on participants offering full consent, coupled with more stringent recruitment processes
300 often excluding participants, which might have elicited higher commitment to the programme ^(33, 34).
301 Importantly, no review reported detailed characteristics of participants who are referred, but fail to
302 make contact with an ERS. Instead, reviews have commented upon uptake figures from those who
303 start a scheme. While this review cannot comment on the research priorities of previous reviews or
304 individual studies, we suggest that this reflects a crucial gap in understanding within the primary
305 data. It is important to this review to question why this gap exists, because if large sub groups of the
306 population are not taking up a referral they cannot benefit from the programme.

307 Reviews have attempted to unpick participant referral characteristics, for example, there have been
308 suggestions that participants referred for a medical reason may show greater referral uptake
309 compared to those with no specific referral ⁽¹¹⁾, however, most reviews are less clear on such a
310 hierarchy of take up. Referral to ERS from a cardiac nurse appears more effective than referral from
311 any other health professional ⁽¹¹⁾, from which it could be inferred that the hospitalisation
312 surrounding the condition has served as a teachable moment. 'Teachable moments' have been
313 described within health behaviour as a noticeable feature of a patient's circumstance which prompts
314 a change of behaviour ⁽³⁵⁾. These opportunities are utilised within the patient-clinician interaction or
315 by patients alone to foster positive health behaviours, such as increasing physical activity levels ^{(35,}
316 ³⁶⁾. It is important to acknowledge that teachable moments do not occur within isolated situations.

A Review of Reviews of Exercise Referral Schemes.

317 Systemic features, such as the expectations of those involved, communication skills of the health
318 practitioner, and medical practice setting will influence the situation ⁽³⁶⁾.

319 Reviews have indicated participant profiling is varied and this lack of information on subgroups
320 prevents any inference being drawn about who ERS is best suited for. Pampel et al. ⁽³⁷⁾ comment that
321 low socioeconomic status groups have fewer opportunities to engage in services that promote good
322 health, yet the picture is not clear in ERS. For example, Pavey et al. ⁽¹¹⁾ have shown participants who
323 had a medical referral coupled with a low socioeconomic status were more likely to take up ERS over
324 participants with a higher economic status, but this was only within participants with a respiratory
325 diagnosis. Further reviews ^(16, 18) suggest that insufficient reporting of socioeconomic status within
326 ERS prevents any conclusions being reached.. The complexity of uptake of ERS, insufficient reporting
327 within ERS ^(16, 18) coupled with a lack of clear guidance within the socioeconomic and health
328 behaviour literature ⁽³⁷⁾ means it is very difficult to produce generalizable statements surrounding
329 the socioeconomic status and take up of ERS.

330 **Adherence and Attendance**

331 In contrast to uptake, attendance figures across reviews appear greater within observational studies.
332 Interpretation of attendance and completion rates should be treated cautiously due to a lack of a
333 standardised protocol (i.e. objective, standardised, universally consistent measure of attendance) for
334 reporting these figures across the literature. The lack of a standardised measure-surrounding
335 attendance could see participants being classified as completing a scheme ⁽¹²⁾, while the reality may
336 be attendance at a couple of sessions and attendance at the final exit session of the scheme. This
337 invites questions as to why there is no standardized use of an objective count for attendance, since
338 all outcomes of ERS are dependent upon attendance at, and adherence to the programme ⁽⁹⁾. The
339 review by Pavey et al. ⁽¹¹⁾ concluded that the number of exercise sessions made available within a
340 scheme might elicit higher rates of attendance. Meanwhile Arsenijevic and Groot ⁽¹⁵⁾ reported that
341 attendance rates were not correlated to the duration of the programme. Additionally, they report

A Review of Reviews of Exercise Referral Schemes.

342 that a prolonged follow up with increased contact time with researchers and exercise practitioners
343 post ERS could offer explanations to why attendance was greater during the programme ⁽¹⁵⁾. This in
344 itself is suggestive that observational studies may reflect the true nature of ERS ⁽¹¹⁾.

345 While uptake to ERS was reported to be slightly higher in females, overall, males were more likely to
346 participate within a programme ^(9, 11, 12). However, various primary studies within these reviews
347 found no association between gender and attendance. This review questions why there is such a
348 poor base level of reporting with only three reviews, of which two reviews are based upon the same
349 study search criteria ^(11, 12), reporting on gender characteristics and referral uptake ^(9, 11, 12). The
350 present review does highlight a greater number of reviews reporting upon medical conditions and
351 the impact they have upon attendance. Despite this greater breadth of reporting, no clear consensus
352 can be reached, with reviews reporting no difference in attendance between at risk groups ⁽¹⁴⁾;
353 either participants with cardiovascular disease were more likely to attend ⁽¹¹⁾; or conversely
354 participants with cardiovascular disease were less likely to attend ⁽¹⁵⁾.

355 Inference on reporting of attendance figures could be suggestive of at least two aspects of ERS. First,
356 it could suggest a referral process where the wrong participants are being referred into the system.
357 Resolution of this could be gained through a clearer and more robust understanding of who is
358 referred to ERS ⁽⁹⁾. Second, it could reflect a lack of knowledge of what participants adhere to within
359 the programme. With no understanding of what is delivered within a programme, from an exercise
360 prescription standpoint, and no reporting on the extent to which individuals adhere to the
361 prescription, few conclusions can be drawn about the effectiveness of the scheme. It is important to
362 question why this gap exists, because if sub groups of the population are not completing the
363 prescribed programme then they clearly cannot benefit from the programme. It is important to
364 acknowledge that participants will stand to benefit from any episode of PA completed, regardless of
365 meeting the prescribed dose. However, it is important to know and understand the frequency,
366 intensity, type, and time (FITT) of prescribed exercise in these programmes, but this is not commonly

A Review of Reviews of Exercise Referral Schemes.

367 recorded or reported. Knowledge of the prescribed dose could help to understand if it is too
368 demanding, thus leading to participant drop out, or insufficiently demanding to engage participants
369 or provide a clinical benefit. Development of the Consensus on Exercise Reporting Template allows
370 detailed and explicit reporting of the delivery of ERS for policy makers and practitioners alike ⁽²²⁾.
371 Implementation of the Consensus on Exercise Reporting Template will further enhance the
372 knowledge base and understanding for whom prescribed exercise is beneficial. Ultimately, it must be
373 understood that both attendance and adherence are multidimensional constructs affected by the
374 relationship between participant and practitioner intertwined with participant centric factors (e.g.
375 mood state, self-efficacy, time and forgetfulness) ⁽²²⁾ and potentially exercise prescription centric
376 factors (e.g. frequency, intensity, time and type of prescription).

377 ***Behaviour Change Techniques***

378 Development of ERS was based upon policy recommendations rather than theoretical guidelines
379 developed for health promotion interventions ⁽¹⁵⁾. Riddoch, Puig-Riberia and Cooper ⁽¹⁷⁾ reported that
380 early ERS's were not based upon any behaviour change model. Failure to acknowledge, deliver or
381 evaluate behaviour change models during a programme prevents clear evaluation of the ERS. More
382 often, the end point (e.g. PA levels, blood glucose concentration, attendance and adherence levels)
383 is the result of behaviour change. Evaluating and reporting interventions on outcomes alone is
384 problematic with an array of influences that could determine the path between behaviour and
385 outcome ⁽³⁸⁾. This is evident within the literature where success of ERS is judged as the outcome
386 evaluation without taking into consideration the methods underpinning it, such as BCT utilised to
387 motivate uptake or increase attendance levels to ERS or the quality of those delivering these BCT's.
388 Theories of behaviour change provide a validated framework to understand not only how behaviour
389 changed, but importantly why behaviour changed ⁽³⁹⁾. A review by Dombrowski et al. ⁽³⁹⁾ found that
390 the delivery and features of behaviour change techniques was heterogeneous in nature.
391 Additionally, a later review reported 44% of interventions did not report use of a theoretical

A Review of Reviews of Exercise Referral Schemes.

392 framework ⁽⁴⁰⁾. This lack of reporting within both reviews of ERS and the wider field of health
393 behaviour prevents any clear guidance upon which theory or BCT, or how it is operationalized, is
394 most effective to understand health behaviour, and health behaviour change ⁽⁴⁰⁾.

395 In the present review we have highlighted how uptake of ERS and attendance at ERS could be
396 influenced by participant characteristics or external factors. Whilst factors such as that of
397 socioeconomic status and teachable moments, have been discussed, a further consideration would
398 be the theoretical perceived risk. The health belief model and the common-sense model hypothesise
399 that perceived severity combined with perceived susceptibility and external cues to actions are
400 strong contributing factors within an individual's perception of threat ^(41, 42). While our review cannot
401 comment upon direct causality to why individuals with certain medical conditions have a greater
402 uptake or attendance rate, we can stress the importance of trying to understand these factors. This
403 understanding comes from the ability to record, report and evaluate these perceptions. Tools such
404 as the Illness Perception Questionnaire - allow for a greater understanding of what an individual
405 perceives of their condition ⁽⁴³⁾. The brief Illness Perception Questionnaire has been shown as a valid
406 and reliable measure of illness perception distinguishing between various illness groups and
407 importantly for ERS, is a predictor of attendance at cardiac rehabilitation classes ⁽⁴³⁾. Incorporation of
408 tools such as the Illness Perception Questionnaire seem essential to trying to understand theoretical
409 factors or BCT's impacting upon ERS uptake or attendance.

410 In line with Prestwich et al. ⁽⁴⁰⁾, we report that the transtheoretical model of behaviour change and
411 social cognitive theory, alongside BCT's such as motivational interviewing, are used, but ultimately
412 are underutilised tools. Fundamentally, the lack of theoretical reporting within ERS generates more
413 questions than it solves. Is the lack of reporting due to there being a lack of delivery or, more an
414 inability to efficiently test and report its delivery? Without assessing any possible association
415 between the BCT's in conjunction with the theory they are based within, we are unable to further
416 our understanding of possible effects ⁽³⁹⁾. The reporting of an intervention's BCT and delivery (e.g.

A Review of Reviews of Exercise Referral Schemes.

417 duration, intensity, setting, group vs individual, verbal vs written and the skill level of the
418 practitioner deliverers) ⁽³⁹⁾ is imperative if we are to understand which techniques are appropriate
419 within ERS for improving uptake and attendance.

420 Implications for practice and policy

421 This review highlights the need for consistent reporting methods to be implemented within ERS. The
422 ability to robustly detail participant characteristics from the point of referral to exiting the scheme
423 (at any stage) will only enhance the understanding of the ERS process. The ability to gain an
424 understanding of what is being delivered at a theoretical level (e.g. BCT's, such as goal setting or
425 relapse prevention and the delivery process of these BCT's) and physical activity level (e.g.
426 frequency, intensity, type and time) would allow strong associations to be attributed, or not, to
427 these features, notably for understanding levels of attendance and adherence. Failure to have these
428 fundamentals in order places increased pressure upon ERS, notably when the National Institute for
429 Health and Care Excellence ⁽⁴⁾ are highlighting the scarce evidence of effectiveness for ERS schemes.
430 The use of tools such as the Consensus on Exercise Reporting Template, Illness Perception
431 Questionnaire or behaviour change taxonomies will further help to understand how and why
432 behaviour changed within programmes. Greater understanding at any level will benefit
433 stakeholders, health practitioners and participants alike, by providing individualised care pathways,
434 in achieving the most effective results from stretched resources and improving programme
435 implementation and viability.

436 Strengths and limitations.

437 This is the first systematic review of reviews focusing on ERS. By taking an overarching view of all
438 the reviews we have been able to highlight key areas that require exploration to inform future
439 evaluation of ERS. However, due to the nature of reviewing systematic reviews, we are unable to
440 explore and provide detailed in-depth mechanistic knowledge surrounding ERS effectiveness.
441 Additional problems arise for evidence reporting namely around AMSTAR grading. While AMSTAR

A Review of Reviews of Exercise Referral Schemes.

1
2
3 442 has been shown to demonstrate good psychometric properties, it was developed after four of the
4
5 443 included reviews had been published. Subsequently two of these reviews were classified as low
6
7 444 quality yet they provide vital insights into ERS. Whilst this review of reviews examined 11 systematic
8
9 445 reviews, which in turn sourced 221 citations, 44% of citations were duplicated across reviews
10
11 446 reducing the breadth of data available. While this review draws attention to tools such as the Illness
12
13 447 Perception Questionnaire to assess a participant's perception towards their illness, this review is
14
15 448 unable to address participant's perspectives towards ERS. This is impart due to the scope of this
16
17 449 review. We acknowledge primary studies that have provided qualitative insight towards participant
18
19 450 perspectives of ERS^(3, 44); however, there exists a knowledge gap in utilising this for implementation
20
21 451 refinement of ERS.
22
23
24
25

452 **Directions for future research**

26
27
28 453 Research should work hand-in-hand with practice and policy makers, in the first instance, to
29
30 454 facilitate robust participant profiling: 1) to capture data on individuals referred but not taking up the
31
32 455 referral to help reduce health inequalities; 2) to generate a consensus on monitoring attendance and
33
34 456 adherence to ERS with objective measures for reporting it; and 3) to incorporate explicit reporting of
35
36 457 BCT's and what they are being used for. With a consistent and coherent basis for reporting, future
37
38 458 evaluations and RCT interventions will be able to explore a multitude of potential interventions, safe
39
40 459 in the knowledge they are grounded within consistent reporting, allowing for direct comparison
41
42 460 between schemes. Greater research is needed surrounding ERS programme content. The very
43
44 461 nature of ERS referral is for participants with lifestyle disease, yet only small to moderate clinical
45
46 462 improvements have been shown within the literature. The use of tools such as the Consensus on
47
48 463 Exercise Reporting Template will help drive research on frequency, intensity, and type of PA
49
50 464 delivered within programmes. Tools such as the Illness Perception Questionnaire have the potential
51
52 465 to build greater understanding of whether individuals are potential candidates for referral to ERS or
53
54 466 not and perhaps leading to a more efficient uptake process. These tools may help in understanding
55
56 467 who and what specifically, contributes to each part of the complexity surrounding ERS. Should
57
58
59
60

A Review of Reviews of Exercise Referral Schemes.

1
2
3 468 interventions be focused upon deliverers, the content and its delivery methods within the scheme;
4
5 469 or would research be better utilised in understanding behaviour change towards ERS. Further
6
7 470 research is required to establish methods that address the challenging nature of measuring BCT
8
9 471 within ecological valid environments, such as ERS. Other research may be able to track participants
10
11 472 over the long-term, reporting on future primary care or hospital admissions with the ability to link it
12
13 473 back to ERS data. While the potential for future research is hypothesised here, it is fundamentally
14
15 474 reliant on robust, clear, standardised and explicit reporting.
16
17
18
19

475 **Conclusions**

20
21
22 476 Exercise referral schemes benefit from not being constrained within a rigid framework, allowing a
23
24 477 varied spectrum of delivery. However, it would appear this currently may also be detrimental in
25
26 478 understanding what works, why and for whom. Complex interventions such as ERS, which involves
27
28 479 multiple input from various health practitioners, requires robust and consistent recording and
29
30 480 reporting of all its facets at every stage of the process. ERS outcomes which are often judged in
31
32 481 terms of increased physical activity, are subject to multiple, separate and complex constructs along
33
34 482 the way. We have reported that uptake levels to schemes have not changed and are potentially
35
36 483 influenced by a variety of participant characteristics. Importantly, there is a requirement to know
37
38 484 who the people are that are not taking up referral and thus missing the opportunity to potentially
39
40 485 benefit from ERS. We have identified that attendance within schemes is potentially influenced by
41
42 486 both participant characteristics and scheme definition. Critically there is no knowledge or reporting
43
44 487 to-date on what participants adhere to within ERS. We have also highlighted that BCT's are poorly
45
46 488 reported, preventing any knowledge of how and why change may have occurred. Failure to robustly
47
48 489 produce effective reporting methods or have a clear scheme wide consensus prevents any firm
49
50 490 conclusions on causal effect. Ultimately, a lack of reporting prevents any research from accurately
51
52 491 validating its hypothesis ⁽⁴⁵⁾ and within the construct of this review, assigning a true reflection on the
53
54 492 effectiveness of ERS.
55
56
57
58
59
60

A Review of Reviews of Exercise Referral Schemes.

493 **References**

- 494 1. Department of Health, Physical Activity and Health Improvement and Protection. *Start*
495 *Active, Stay Active: A report on physical activity from the four home countries' Chief*
496 *Medical Officers*. London: Department of Health. 2011.
- 497 2. Morgan F, Battersby A, Weightmam AL, Searchfield L, Turley R, Morgan H, Jagroo J, Ellis
498 S. Adherence to exercise referral schemes by participants – what do providers and
499 commissioners need to know? A systematic review of barriers and facilitators. *BMC*
500 *Public Health*. 2016;16(227):1-11.
- 501 3. Din NU, Moore GF, Murphy S, Wilkinson C, Williams NH. Health professionals'
502 perspectives on exercise referral and physical activity promotion in primary care:
503 Findings from a process evaluation of the National Exercise Referral Scheme in Wales.
504 *Health Education Journal*. 2015;74(6):743-757.
- 505 4. National Institute for Health and Care Excellence. *Physical Activity: exercise referral*
506 *schemes. Public Health Guideline*. Manchester: NICE. 2014.
- 507 5. World Health Organisation. *Global recommendations on physical activity for health*.
508 Switzerland: WHO Library Cataloguing-in-Publication. 2010.
- 509 6. National Institute for Health and Care Excellence. *Physical Activity: walking and cycling*.
510 *Public Health Guideline*. Manchester: NICE. 2012.
- 511 7. Moore GF, Moore L, Murphy S. Facilitating adherence to physical activity: exercise
512 professionals' experiences of the National Exercise Referral Scheme in Wales. A
513 qualitative study. *BMC Public Health*. 2011;11(935):1-12.
- 514 8. Ward M. Experiences in physical activity promotion in health care settings for primary
515 prevention in the UK. *Swiss Sports & Exercise Medicine*. 2014;62(2):26-28.
- 516 9. Gidlow C, Johnston LH, Crone D, James D. Attendance of exercise referral schemes in the
517 UK: A systematic review. *Health Education Journal*. 2005;64(2):168-186.

A Review of Reviews of Exercise Referral Schemes.

- 1
2
3 518 10. Williams NH, Hendry M, France B, Lewis R, Wilkinson C. Effectiveness of exercise-referral
4
5 519 schemes to promote physical activity in adults: systematic review. *British Journal of*
6
7 520 *General Practice*. 2007;57(545):979-986.
- 8
9
10 521 11. Pavey T, Taylor A, Hillsdon M, Fox K, Campbell J, Foster C, Moxham T, Mutrie N, Searle J,
11
12 522 Taylor R. Levels and predictors of exercise referral scheme uptake and adherence: A
13
14 523 systematic review. *Journal of epidemiology and community health*. 2012;66(8):737-744.
- 15
16 524 12. Pavey T, Anokye N, Taylor A, Trueman P, Moxham T, Fox K, Hillsdon M, Green C,
17
18 525 Campbell J, Foster C, Mutrie N, Searle J, Taylor R. The clinical effectiveness and cost-
19
20 526 effectiveness of exercise referral schemes: a systematic review and economic
21
22 527 evaluation. *Health Technology Assessment*. 2011;15(24):1-270.
- 23
24 528 13. Pavey TG, Taylor AH, Fox KR, Hillsdon M, Anokye N, Campbell JL, Foster C, Green C,
25
26 529 Moxham T, Mutrie N, Searle J, Trueman P, Taylor RS. Effect of exercise referral schemes
27
28 530 in primary care on physical activity and improving health outcomes: Systematic review
29
30 531 and meta-analysis. *BMJ open*. 2011;343:1-14.
- 31
32 532 14. Campbell F, Holmes M, Everson-Hock E, Davis S, Buckley Woods H, Anokye N,
33
34 533 Tappenden P, Kaltenthaler, E. A systematic review and economic evaluation of exercise
35
36 534 referral schemes in primary care: a short report. *Health Technology Assessment*.
37
38 535 2015;19(60):1-105.
- 39
40 536 15. Arsenijevic J, Groot W. Physical activity on prescription schemes (PARS): do programme
41
42 537 characteristics influence effectiveness? Results of a systematic review and meta-
43
44 538 analyses. *BMJ Open*. 2017:1-13.
- 45
46 539 16. Morgan O. Approaches to increase physical activity: reviewing the evidence for exercise-
47
48 540 referral schemes. *Public Health*. 2005;119:361-370.
- 49
50 541 17. Riddoch C, Puig-Riberia A, Cooper A. *Effectiveness of physical activity promotion schemes*
51
52 542 *in primary care: a review*. London: Health Education Authority. 1998.
- 53
54
55
56
57
58
59
60

A Review of Reviews of Exercise Referral Schemes.

- 1
2
3 543 18. Orrow G, Kinmonth AL, Sanderson S. Sutton S. Effectiveness of physical activity
4
5 544 promotion based in primary care: systematic review and meta-analysis of randomised
6
7 545 controlled trials. *BMJ Open*. 2012;344:1-17.
8
9
10 546 19. Sorensen JB, Skovgaard T, Puggaard L. Exercise on prescription in general practice: A
11
12 547 systematic review. *Scandinavian journal of primary health care*. 2006;24(2):69-74.
13
14 548 20. Jones F, Harris P, Waller H, Coggins A. Adherence to an exercise prescription scheme:
15
16 549 The role of expectations, self-efficacy, stage of change and psychological well-being.
17
18 550 *British Journal of Health Psychology*. 2005;10:359-378.
19
20
21 551 21. Dinan S, Lenihan P, Tenn T, Iliffe S. Is the promotion of physical activity in vulnerable
22
23 552 older people feasible and effective in general practice? *British Journal of General*
24
25 553 *Practice*. 2006:791-793.
26
27
28 554 22. Slade SC, Dionne CE, Underwood M Buchbinder R. Consensus on exercise reporting
29
30 555 template (CERT): Explanation and elaboration statement. *British Journal of Sports*
31
32 556 *Medicine*. 2016;50:1428-1437.
33
34
35 557 23. Taylor AH, Doust J, Webborn. Randomised controlled trial to examine the effects of a GP
36
37 558 exercise referral programme in Hailsham, East Sussex, on modifiable coronary heart
38
39 559 disease risk factors. *Journal of Epidemiology and Community Health*. 1998;52:595-601.
40
41 560 24. Edmunds J, Ntoumanis N, Duda JL. Adherence and well-being in overweight and obese
42
43 561 patients referred to an exercise on prescription scheme: A self-determination theory
44
45 562 perspective. *Psychology of Sport and Exercise*. 2007;8:722-740.
46
47
48 563 25. Leijon ME, Bendtsen P, Nilsen P, Festin K, Stahle A. Does a physical activity referral
49
50 564 scheme improve the physical activity among routine, primary health care patients?
51
52 565 *Scandinavian Journal of Medicine & Science in Sports*. 2009;19(5):627-636.
53
54
55 566 26. Littlecott H, Moore GF, Moore L, Murphy S. Psychosocial mediators of change in physical
56
57 567 activity in the Welsh national exercise referral scheme: secondary analysis of a
58
59
60

A Review of Reviews of Exercise Referral Schemes.

- 1
2
3 568 randomised controlled trial. *International Journal of Behavioural Nutrition and Physical*
4
5 569 *Activity*. 2014;11(109):1-11.
6
7
8 570 27. Moore GF, Audrey S, Barker M, Bond L, Bonell C, Cooper C, Hardeman W, Moore L,
9
10 571 O’Cathain A, Tinati T, Wright D, Baird J. Process evaluation in complex public health
11
12 572 intervention studies: the need for guidance. *Journal of Epidemiology and Community*
13
14 573 *Health*. 2014;68(2): p.1-2.
15
16 574 28. Woods C, McCaffrey N, Furlong B, Fitzsimons-D’Arcy L, Murphy M, Harrison M, Glynn L,
17
18 O’Riordan J, O’Neill B, Jennings S, Peppard C. *The National Exercise Referral Framework.*
19 575
20 576 *Health and Wellbeing Division, Health Service Executive*. Dublin: Health Service
21
22 577 Executive. 2016.
23
24 578 29. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred reporting items
25
26 579 for systematic reviews and meta-analyses: The PRISMA Statement. *PLoS Medicine*.
27
28 580 2009;6(7):1-6.
29
30 581 30. Greaves CJ, Sheppard KE, Abraham C, Hardeman W, Roden M, Evans PH, Schwarz P.
31
32 582 Systematic review of reviews of intervention components associated with increased
33
34 583 effectiveness in dietary and physical activity interventions. *BMC Public Health*.
35
36 584 2011;11(119):1-12.
37
38 585 31. Shea BJ, Grimshaw JM, Wells GA, Boers M, Andersson N, Hamel, C, Porter AC, Tugwell P,
39
40 586 Moher D, Bouter LM. Development of AMSTAR: a measurement tool to assess the
41
42 587 methodological quality of systematic reviews. *BMC Medical Research Methodology*,
43
44 588 2007;7(10):1-7.
45
46 589 32. Pieper D, Mathes T, Eikermann M. Can AMSTAR also be applied to systematic reviews of
47
48 590 non-randomized studies? *BMC Research Notes*. 2014;7(609):1-6.
49
50 591 33. Martin KA, Sinden AR. Who will stay and who will go? A review of older adults’
51
52 592 adherence to a randomized controlled trials of exercise. *Journal of Ageing and Physical*
53
54 593 *Activity*. 2001;9:91-114.
55
56
57
58
59
60

A Review of Reviews of Exercise Referral Schemes.

- 1
2
3 594 34. Benson K, Hartz AJ. A comparison of observational studies and randomized controlled
4
5 595 trials. *The New England Journal of Medicine*. 2000;345(25):1878–1886.
6
7 596 35. Flocke S, Clark E, Antognoli E, Mason MJ, Lawson PJ, Smith S, Cohen DJ. Teachable
8
9 moments for health behaviour change and intermediate patient outcomes. *Patient*
10 597
11 *Education and Counselling*. 2014;96(1):1-16.
12 598
13 599 36. Lawson PJ, Flocke S, Teachable moments for health behaviour change: a concept
14
15 600 analysis. *Patient Education and Counselling*. 2009;76(1):25-30.
16
17 601 37. Pampel FC, Krueger PM, Denney JT. Socioeconomic Disparities in Health Behaviours.
18
19 602 *Annual Review of Sociology*. 2010;36:349-370.
20
21 603 38. Michie S, Johnston M. Theories and techniques of behaviour change: Developing a
22
23 604 cumulative science of behaviour change, *Health Psychology Review*. 2012;6(1):1-6.
24
25 605 39. Dombrowski SU, Sniehotta FF, Avenell A, Johnston M, MacLennan G, Araújo-Soares V.
26
27 606 Identifying active ingredients in complex behavioural interventions for obese adults with
28
29 607 obesity-related co-morbidities or additional risk factors for co-morbidities: a systematic
30
31 608 review. *Health Psychology Review*. 2012;6(1):7-32.
32
33 609 40. Prestwich A, Sniehotta FF, Whittington C, Dombrowski SU, Rogers L, Michie S. Does
34
35 610 theory influence the effectiveness of health behaviour interventions? Meta-analysis.
36
37 611 *Health Psychology*, 2014;33(5):465-474.
38
39 612 41. Jones CJ, Smith H, Llewellyn C. Evaluating the effectiveness of health belief model
40
41 613 interventions in improving adherence: A systematic review. *Health Psychology Review*.
42
43 614 2014;8(3):253-269.
44
45 615 42. Leventhal H, Meyer D, Nerenz D. The common sense representation of illness danger. *In:*
46
47 616 *Rachman, S., Ed., Contributions to Medical Psychology*. 1980;2:17-30.
48
49 617 43. Broadbent E, Petrie, KJ, Main J, Weinman J. The brief illness perception questionnaire.
50
51 618 *Journal of Psychosomatic Research*. 2006;60:631-637.
52
53
54
55
56
57
58
59
60

A Review of Reviews of Exercise Referral Schemes.

- 1
2
3 619 44. Eynon MJ, O'Donnell C, Williams L. Gaining qualitative insight into the subjective
4
5 620 experiences of adherers to an exercise referral scheme: A thematic analysis. *Journal of*
6
7 621 *Health Psychology*. 2016;23(11):1476-1487.
8
9
10 622 45. Resnick B, Inguito P, Orwig D, Yu Yahiro J, Hawkes W, Werner M, Zimmerman S,
11
12 623 Magaziner J. Treatment fidelity in behaviour change research: A case example. *Nursing*
13
14 624 *Research*. 2005;54(2):139-143.
15
16

625 **Acknowledgments**

626 N/A

627 **Funding source**

628 This work forms part of a PhD thesis with the lead author receiving funding from the University of
629 Stirling as part of the Impact Studentship Scheme.

A Review of Reviews of Exercise Referral Schemes.

630 **Tables**

Table 1 Inclusion and Exclusion criteria used in collating Systematic reviews inclusion.

1.	Not in English language	Due to lack of translation facilities, all non-English papers were excluded.
2.	Type of study	Systematic reviews (which included RCTs, observational studies, case-controlled or other quasi-experimental studies within them) and meta-analysis were included for review.
3.	Type of Intervention	Interventions promoting changes in physical activity behaviour via a direct referral from a health practitioner within primary care to a recognised exercise programme (e.g. local leisure centre, local walking scheme). Interventions initiated from secondary care and beyond were excluded alongside any health screening programmes.
4.	Type of Participants	Adults (16 years and over) of any gender who had been diagnosed or placed at risk of non-communicable disease (e.g. type 2 diabetes, hypertension, cardiovascular disease) from their primary care practitioner. Interventions where participants were sedentary but had no further risk factors but were used as a control groups where included if reported separately.
5.	Outcomes	Reviews were selected where the primary outcome measures reported were increase in physical activity (e.g. frequency, intensity) uptake and adherence to schemes, physiological and psychological (e.g. changes in Body Mass Index or Short Form (36) health survey). Outcomes could be measured objectively (e.g. with accelerometers) or by self-reported means (e.g. questionnaires). Where reported the cost effectiveness of

A Review of Reviews of Exercise Referral Schemes.

schemes was included.

631

Table 2 Classification of strength based upon evidence included within reviews.

A	Systematic Reviews of RCTs
B	Systematic reviews of individual, non-RCTs, case-control studies, cohort studies, controlled before-and-after (CBA), correlation studies or similar.
C	Systematic reviews of both RCTs and non-RCTs, case-control studies, cohort studies, controlled before-and-after (CBA), correlation studies or similar.

632

633

A Review of Reviews of Exercise Referral Schemes.

Table 3 Total and mean AMSTAR scoring, classification of strength based upon evidence included within reviews and Cohen's Kappa for inter-rate reliability.

Review lead author and year	Reviewer	Total AMSTAR score	Mean AMSTAR score	Evidence classification	Inter-rate reliability
Arsenijevic et al., 2017	CS	6	7	C	0.62
	SG	8			
Campbell et al., 2015	CS	9	9	C	0.39
	SG	9			
Gidlow et al., 2005	CS	3	2.5	C	0.74
	SG	2			
Morgan et al., 2016	CS	9	9	C	1
	SG	9			
Morgan, 2005	CS	6	6	A	1
	SG	6			
Orrrow et al., 2012	CS	9	9	A	1
	SG	9			
Pavey et al., 2011a	CS	11	11	C	0
	SG	11			
Pavey et al., 2012	CS	4	5	C	0.49

A Review of Reviews of Exercise Referral Schemes.

	SG	6			
	CS	9			
Pavey et al., 2011b			8.5	A	0.74
	SG	8			
	CS	2			
Sorensen et al., 2006			3.5	C	0.42
	SG	5			
	CS	7			
Williams et al., 2007			7	C	1
	SG	7			

634

Table 4 Reported uptake and attendance figures for Observational studies and RCT's where reported within reviews


Review lead author and year	Strength of study	Reported uptake	Reported attendance
Campbell et al., 2015	9C	35-85%	25 - 86%
			for final assessment only
Gidlow et al., 2005	2.5C	< 67%	12 - 25%
Pavey et al., 2011a	11C	Observational studies 66% (95% CI = 57-75%) RCTs 80% (95% CI = 61-98%)	Observational studies 49% (95% CI = 40-59%) RCTs 37% (95% CI = 20-54%)

A Review of Reviews of Exercise Referral Schemes.

Pavey et al., 2012	5C	Observational studies	Observational studies
		66% (95% CI = 57-75%)	43% (95% CI = 32-54%)
		RCTs	RCTs
		81% (95% CI = 68-94%)	80% (95% CI = 61-98%)
Williams et al., 2007	7C	66%	12 – 42%

For Peer Review

A Review of Reviews of Exercise Referral Schemes.



636

Figure 1. PRISMA flow diagram depicting study selection, screening, eligibility for inclusion and analysis.

Supplementary data.

Supplementary data 1: Search strategy terms and protocol steps used within Medline database

Step	Search term
1	exp Exercise/
2	exp Exercise Movement Techniques/ or exp Exercise Therapy/
3	exp Sports/
4	exp Motor Activity/
5	exp Physical Exertion/
6	exercis* refer*.af.
7	refer* exer*.af.
8	exerc* prescr*.af.
9	prescr* exerci*.af.
10	(activ* refer* or refer* activ*).af.
11	(presc* activ* or activ* precri*).af.
12	6 or 7 or 8 or 9 or 10 or 11
13	exp Prescriptions/
14	(prescribing or prescribe or prescriptions).af.
15	13 or 14
16	or/1-5
17	15 and 16
18	12 or 17
19	limit 18 to english language
20	(metaanal: or meta-anal:).af.
21	exp Meta-Analysis/
22	20 or 21
23	limit 19 to meta analysis
24	19 and 22
25	23 or 24
26	(systematic* review* or review* systematic*).af.
27	limit 19 to systematic reviews
28	19 and 26
29	27 or 28
30	systematic review*.af.
31	19 and 30
32	27 and 31
33	29 not 25
34	25 or 29 or 33

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Supplementary data 2: Eleven predefined constructs used as an extraction tool and relating to the reporting effectiveness of reviews

1	Aim of Review
2	Theoretical basis (any stated theory)
3	Behaviour change technique (goal setting, problem solving)
4	location(s) or setting where the intervention occurred (local leisure centre, green spaces)
5	Intervention provider (should always be 3 rd party exercise provider as part of ERS)
6	Frequency of intervention (period of time including the number of sessions, their schedule, and their duration, intensity (if mentioned in initial studies or current review))
7	Mode of delivery (group, individual, supervised)
8	Characteristics of the participants (sex, disease or risk state)
9	Amount of studies within the review
10	Number of participants within the review / Sample sizes
11	Cost effectiveness if mentioned

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Supplementary data 3: Key characteristics of included systematic reviews

Review lead author and year											
Characteristics of reviews	Arsenijevic et al., 2017	Campbell et al., 2015	Gidlow et al., 2005	Morgan et al., 2016	Morgan, 2005	Orrow et al., 2012	Pavey et al., 2011a	Pavey et al., 2012	Pavey et al., 2011b	Sorensen et al., 2006	Williams et al., 2007
Type of review performed and its search period											
Descriptive	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Meta-analysis	✓	✓				✓	✓	✓	✓		✓
Economic Evaluation		✓					✓				
Date period searched	2000 onwards	October 2009 – June 2013	Origin - 2003	1995 – June 2015	1966 - 2002	Origin - May 2009	1990 – October 2009	Origin - October 2009	Origin - October 2009	1980 – June 2005	Origin – March 2007
Objectives and outcomes markers of review											
Overall effectiveness (increase of PA levels)	✓			✓	✓	✓			✓	✓	✓
Clinical effectiveness		✓					✓		✓		
Cost effectiveness		✓					✓			✓	
Uptake to ERS							✓	✓			
Attendance or completion of ERS			✓	✓			✓	✓			✓
Implementation / design ERS	✓									✓	
Reported inclusion criteria of reviews											

1												
2												
3												
4	Any study design considered	✓				✓			✓	✓	✓	✓
5												
6	RCT only									✓		
7	Males and females ≥ 18	✓	✓		19 ≥					16 ≥		
8	Peer reviewed and published in English	✓				✓						✓
9	Referral from primary care to improve clinical outcomes or physical activity levels					✓	✓	✓		✓	✓	✓
10	Programme designs more intensive than advice alone		✓							✓		✓
11	UK based studies					✓						
12												
13												
14												
15												
16												
17												
18												
19												
20												
21												
22												
23												
24												
25												
26												
27												
28												
29												
30												
31												
32												
33												
34												
35												
36												
37												
38												
39												
40												
41												
42												
43												
44												
45												
46												

Supplementary data 4: Protocol to undertake systematic review of reviews.

Insufficient Reporting of Factors Associated with Exercise Referral Scheme Uptake, Attendance and Adherence: A Systematic Review of Reviews (Protocol)

Review Question

1) Describe definitions, rates, and participant characteristics associated with ERS uptake; 2) to describe definitions, rates, and participant characteristics associated with attendance at and adherence to ERS schemes; and 3) to describe theoretically informed approaches associated with ERS uptake and attendance.

Searches

We will systematically search the Databases:

- CINHAL, Medline, Embase, Cochrane Library, SportsDiscus, PsychInfo, SCOPUS, HMIC, AMED, Public Health Database, ASSIA, CRD databases, Prospero, Web of Knowledge, Campbell Collaboration Library and PubMed.

We will perform additional non-systematic searches of UpToDate, BMJ Best Practice and Dynamed.

We will hand search reference lists of included articles for additional material. The searches will be restricted to English language, reviews published within peer viewed journals, and with no date restrictions.

Type of Study to be included

We will include systematic reviews that themselves have reviewed RCTs, observational studies, case-controlled or other experimental studies and meta-analysis.

Condition or domain being studied

Exercise referral schemes. Commonly used terms within the literature are 'Exercise on Prescription' and 'GP Referral', which we will also be search for to ensure no data is missed.

Participants/population

Inclusion: We will search for adults (18 years and over) of any gender who had been diagnosed or considered at risk of non-communicable disease (e.g. type 2 diabetes, hypertension, cardiovascular disease) by their primary care practitioner. Interventions that use participants that were sedentary with no further risk factors of non-communicable disease, but are used as a control groups will be included if reported separately.

Exclusion: We will exclude youths and children.

Intervention(s), exposure(s)

Inclusion: We will include systematic reviews that have reviewed interventions known as 'exercise referral', 'GP referral', or 'exercise on prescription'. Interventions promoting changes in physical activity behaviour via a direct referral from a health practitioner within primary care to a recognised exercise programme.

Exclusion: We will exclude reviews that report on interventions initiated from secondary care or are health-screening programmes.

Primary outcomes

- Define rates and participant characteristics associated with ERS uptake.
- Define rates and participant characteristics associated with attendance at, and, adherence to, ERS schemes.
- Report theoretically informed approaches associated with ERS uptake and attendance.

Secondary outcomes

- Establish what key features of ERS reporting go relatively under-reported.

Data extraction (selection and coding)

The PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) statement will guide the conduct of the narrative of the review of reviews. Results of the search strategy will be

1
2
3 screened for duplicates by one researcher, with duplicates being removed. One author will
4 independently examine titles and abstracts against the inclusion and exclusion criteria. Three authors
5 will independently assess a randomised sub sample of papers, amounting to 20% of the total title and
6 abstract sample. A group consensus will decide which papers will be progressed further to full text
7 review and which to exclude. Full text review articles will be obtained and assessed by two authors
8 against the inclusion and exclusion criteria. We will provide a detailed recording of the selection
9 process, which we will report via a PRISMA flow diagram and 'Characteristics of excluded studies'
10 table.
11
12
13
14
15

16 17 **Data to be extracted:**

- 18 - Aim of Review / study objectives
- 19 - Theoretical basis (any stated theory)
- 20 - Behaviour change technique (goal setting, problem solving)
- 21 - location(s) or setting where the intervention occurred (local leisure centre, green spaces)
- 22 - Intervention provider (should always be 3rd party exercise provider as part of ERS)
- 23 - Frequency of intervention (period of time including the number of sessions, their schedule
24 and their duration, intensity (if mentioned in initial studies or current review))
- 25 - Mode of delivery (group, individual, supervised)
- 26 - Characteristics of the participants (sex, disease or risk state)
- 27 - Amount of studies within the review
- 28 - Number of participants within the review / sample sizes
- 29 - Cost effectiveness if mentioned
- 30 - Author name and date
- 31 - Reporting of attendance/ adherence rates
- 32 - Reporting on non-attenders
- 33 - Attendance figures, completion rates
- 34 - Key findings from reviews

35 36 **Risk of bias (quality) assessment**

37
38 Two authors will grade each review independently and empirically and discuss any discrepancies.

39
40 We will use The AMSTAR (A Measurement Tool to Assess Systematic Reviews) criteria to grade the
41 reviews. Amstar is based on 11 questions and scores as reviews as follows: low quality (0-3), medium
42 (4-7) and high (8-11). Question within the AMSTAR are as follows:
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. Was an 'a priori' design provided?
2. Was there duplicate study selection and data extraction?
3. Was a comprehensive literature search performed?
4. Was the status of publication (i.e. grey literature) used as an inclusion criterion?
5. Was a list of studies (included and excluded) provided?
6. Were the characteristics of the included studies provided?
7. Was the scientific quality of the included studies assessed and documented?
8. Was the scientific quality of the included studies used appropriately in formulating conclusions?
9. Were the methods used to combine the findings of studies appropriate?
10. Was the likelihood of publication bias assessed?
11. Was the conflict of interest included?

Strategy for data synthesis

A table of results will display the extracted information by study. We will present the findings via a descriptive narrative of the key variables associated with uptake attendance and adherence to exercise referral schemes.