



Quantifying the potential of 'on-farm' seed priming to increase crop performance in developing countries. A meta-analysis

Javier Carrillo-Reche¹  · Mario Vallejo-Marín¹ · Richard S. Quilliam¹Accepted: 13 September 2018
© The Author(s) 2018

Abstract

Low-input agriculture in marginal areas of developing countries faces considerable challenges during crop development. A key stage in crop growth is seed germination, which is often constrained by abiotic factors such as low water potential, high temperatures and soil crusting, which can result in poor establishment. This is exacerbated by low soil fertility, salinity, drought, pests and diseases, which ultimately leads to reduced yields. Over the last 20 years, the potential of 'on-farm' seed priming, a traditional, low-cost technique, consisting of soaking seeds in water prior to sowing, has been applied to different crops and conditions with varying degrees of success. To understand the significance of this potentially transformative agronomic strategy, we have conducted a global meta-analysis of on-farm seed priming by quantifying (i) the rate of emergence, (ii) final emergence and (iii) total yield from 44 published papers on 17 crops across 10 countries. Our results show that on-farm seed priming has a significantly positive effect on crop performance: seeds emerge 22% faster, with an increased final emergence of 11%, with total yields 21% higher than conventionally sown seeds. Furthermore, sub-group analyses demonstrated that on-farm seed priming is more advantageous under stressful abiotic conditions with case studies categorized as being either 'nutrient deficient', 'salinity-stressed' or 'dry climates' gaining the highest yield improvements (22–28%). On-farm seed priming can be particularly beneficial to resource-poor farmers working in low-input agricultural systems where yield potential is limited by intrinsically stressed agronomic environments. Here, we demonstrate for the first time that on-farm seed priming is perfectly adapted to local situations in developing countries. Our results provide the evidence that on-farm seed priming could be effectively adopted by resource-poor farmers as a strategy to increase food security in some of the most marginal agricultural areas.

Keywords Crop yield · Germination · Low-input agriculture · Seed technology

Abbreviations

CI	Confidence interval
<i>df</i>	Degrees of freedom
Na ⁺	Sodium
Cl ⁻	Chloride
ZnSO ₄	Zinc sulphate
N	Nitrogen
N ₂	Atmospheric di-nitrogen

1 Introduction

Low-input agriculture in marginal and semi-arid areas of developing countries encounters many challenges that limit yield potential and thus restricts food security (Tittonnell and Giller 2013; Aune et al. 2017). This is further intensified by predicted climate change scenarios such as increasingly unpredictable rainfalls and extreme temperatures (Knox et al. 2012). For example, in semi-arid agricultural systems, important physical constraints in the seedbed, such as low water potential and soil crusting, have frequently been identified as the most significant issues for successful crop establishment (Townend et al. 1996; Tisdall 1996; Nabi et al. 2001; Passioura and Angus 2010). Tillage, fertilizers and amendments of the seedbed, together with timely irrigation, may ameliorate some of these constraints, although are often

✉ Javier Carrillo-Reche
javier.carrilloreche1@stir.ac.uk

¹ Biological and Environmental Sciences, Faculty of Natural Sciences, University of Stirling, Stirling FK9 4LA, UK

unaffordable or not accessible to smallholder farmers (Chianu et al. 2012; Titttonell and Giller 2013; Tonitto and Ricker-Gilbert 2016). Therefore, inexpensive and sustainable strategies with the potential to alleviate unfavourable conditions and reduce input (e.g., cover crops, water harvesting or organic fertilizer) are becoming more relevant for ensuring food security in semi-arid agro-ecosystems (Branca et al. 2013).

Over the past three decades, there has been a renewed interest in a traditional agronomic technique known as ‘on-farm’ seed priming, in part because of its simplicity and low-cost (Murungu et al. 2004a; Rashid et al., 2006). On-farm seed priming is a form of hydro-priming, which consists of soaking seeds in water for a number of hours, usually overnight, surface drying them (to allow limited storage) and sowing soon after (Fig. 1) (Harris, 1996). Prior soaking of seeds in water decreases the time needed for seed imbibition in the soil after sowing; thus, on-farm seed priming shortens the exposure of the seed to adverse soil conditions such as limited soil moisture (Harris et al. 2001a). On-farm seed priming technology has been tested in a wide variety of crops and environmental conditions, and has been extensively developed through participatory trials with local farmers (Harris et al., 1999, 2001a; Rashid et al., 2006a, b). Reports from participatory workshops and research-managed trials have largely agreed that crops grown from on-farm primed seeds emerge faster, obtain higher plant density and vigour, reach flowering and harvest more rapidly, and ultimately, result in higher yields compared with non-primed crops (Harris, 1996a, b; Harris et al., 1999a, b, 2001a, b; Murungu et al. 2003, 2004b; Farooq et al., 2008). However, the extent of these benefits varies widely even under similar contexts; for example, yield improvement of chickpea ranged from 25 to 67% and 4 to 35% in two different villages in India (Harris et al., 1999a, b). There have also been cases where

soaking seeds has turned out to be counterproductive, e.g., for cotton (Murungu et al. 2004b), barley (Rashid et al., 2006a, b), pearl millet (Aune and Ousman, 2011) and sesame (Ousman and Aune, 2011). Consequently, it is unclear to what extent on-farm seed priming can improve crop performance and what factors influence the outcome, and a greater understanding is needed to identify the environments where it may be detrimental and where it can best be applied.

The most important aspect of on-farm seed priming is the duration of the soaking, which must be calculated for each crop species, and even for each variety or cultivar of crop (Harris 2006). Exceeding a ‘safe limit’ of soaking will trigger premature germination, which could lead to damage of the radicle during sowing, or if seeds are left in the priming water for too long, or not surface dried properly, they will begin to rot (Harris 2006). Although farmers have some knowledge of the advantages of soaking seeds prior to sowing (Harris et al., 1999a, b, 2001a), it is often only carried out on seeds for re-sowing in order to ‘catch up’ with the rest of crop and is rarely used as a routine practice. In general, farmers who have used on-farm seed priming are unaware of safe limits and therefore have had varying degrees of success or failure with this method (Harris et al., 1999a, b, 2001a).

To date, only narrative reviews about on-farm seed priming have been published (Ashraf and Foolad 2005; Harris 2006); therefore, a more systematic approach, such as meta-analysis, is needed to quantitatively review this simple technology in terms of increased crop establishment and production. Meta-analysis is a powerful synthesis tool that is being increasingly adopted in agro-ecological disciplines (e.g. Tonitto and Ricker-Gilbert 2016), and using this approach will allow a large number of independent on-farm seed priming case studies to be objectively analysed across different crop types and

Fig. 1 ‘On-farm’ seed priming steps carried out with maize seeds in Kenya. **a** Pouring seeds into buckets. **b** Soaking seeds. **c** Surface drying after priming. **d** Effect on emergence of wheat in Pakistan: non-primed (left) vs. primed (right). (Photos courtesy of H. Wainwright and A. Rashid)



environments. A better understanding of the potential of on-farm seed priming, and in which environments it could be most usefully promoted, could provide governmental institutions and policymakers in developing countries with the evidence to promote its adoption as recommended practice. Therefore, the overarching aim of this paper was to quantify the effect of on-farm seed priming compared to conventional sowing and identify the context where it can best be applied. Specifically, our objectives were to quantify the effect of on-farm seed priming on crop performance (speed of germination, final emergence and yield) and evaluate the impact of climate, crop type and common yield-limiting factors on the final outcome of crops grown from on-farm primed seed.

2 Material and methods

2.1 Sources of data

A literature search was carried out in ‘Web of Science Core Collection’ on 15 November 2017 using the key-words: ‘on farm seed priming’, ‘on station seed priming’, ‘pre-sowing seed soak*’ or ‘hydro*priming’, which resulted in a total of 293 articles. Titles and abstracts were screened and unrelated papers, or studies focussed on tree seeds were discarded. The full text of the remaining papers was examined and had to meet the following criteria: (1) The study had to contain a dry seed sample (control) and primed seed samples (treatment) consisting of seeds submerged in water with no additional oxygenation, and (2) seeds had to be surface dried or partially dried after priming (maximum air-drying duration of less than 24 h). Artificial drying methods such as ovens or air-conditioned cabinets and seeds re-dried to their original moisture, regardless of the methods used, were not included. Other priming strategies, e.g. seeds placed between filter paper and saturated jute mat, were also excluded due to the confounding effects of matric potential. We excluded 141 articles that did not match these requirements. In addition, studies performing other types of seed treatments (19), not containing or missing data (15), lacking or giving ambiguous description of priming (8) and reviews (5) were excluded, and a further 23 papers without full-text, and five more because the same data had been used in several publications, were also excluded. Six additional papers were identified in the reference list of one of the selected papers, which gave a total of 44 valid papers available for meta-analysis (Table 1).

For each publication, three variables were recorded for both control and primed treatments: (i) final emergence, (ii) time to 50% emergence and (iii) yield (i.e. the most common unit of yield for each crop, e.g. grain for cereals, pods for legumes) giving three datasets. The mean and the number of paired observations (n) contributing to that mean value were recorded, e.g. the experimental design of Harris et al.

(2005a, b) consisted of two cultivars of chickpea with three replications during two seasons, i.e. $n = 12$. When available, standard deviation (SD), standard error (SE) or standard error of the difference in mean (SED) were also collected as a measure of the variance. Mean and variation values from published graphs were extracted taking a snapshot of the figure and scaling the axes with WebPlotDigitizer Version 3.10 (Rohatgi 2010) to obtain numerical values. In addition to the statistical data, any characteristics that may have influenced the outcome of the priming treatment and thus could potentially explain heterogeneity in effect size (moderator variables) were also recorded.

If a single publication presented several case studies, the mean effect was calculated (in order to minimize within-study dependence); however, if the moderators differed then, they were considered as independent case studies in the meta-analysis (Koricheva et al. 2013). In cases where several priming outcomes had a common control, the total number of replications of the control group was divided by the number of treatments to avoid overweighting. If papers presented results that had been carried out by distinct groups, e.g. the design of some of the on-station trials, which included both researcher-led and farmer-led experimental and participatory trials, these data were considered as independent. Although these observations cannot be considered fully independent, this approach is commonly used in both plant biology and ecology meta-analyses and allows greater statistical power (Castagneyrol and Jactel 2012; Mayerhofer et al. 2013; Shrestha et al. 2016). The resulting dataset contained 129 case studies derived from 44 papers, which covered 17 crops across ten countries.

2.2 Effect size and meta-analysis

The natural log response ratio ($\ln R$) of the experimental mean divided by the control mean was used as metric of treatment effect (Hedges et al. 1999):

$$\ln R = \ln \left(\frac{X_e}{X_c} \right)$$

where X_e and X_c are the experimental and control mean. Given that more than 50% of the case studies did not provide a measure of variance, case studies were weighted using non-parametric variance ($V_{\ln R}$) (Adams et al. 1997):

$$V_{\ln R} = \frac{n_e + n_c}{n_e * n_c}$$

where n_e and n_c are the experimental and control number of paired observations, respectively.

Bias-corrected bootstrapped 95% confidence intervals based on 10,000 iterations were calculated for overall effect sizes (Adams et al. 1997) and represented as a percentage change relative to controls (%), transforming them back by

Table 1 Data sources reviewed in the meta-analysis

Author	Journal	Country	Crop	Response variables	Study type	Yield-limiting factor ^a	Climate zone ^a
Abdalla et al. (2015)	Agronomy-Basel 5 (4):476–490	Sudan	Sorghum, groundnut, sesame, and cowpea	FE and yield	Field	Nutrient-stressed	Semi-arid
Abro et al. (2009)	Pak J Bot 41 (5):2209–2216	Pakistan	Wheat	E50 and yield	Field	Salinity	Arid
Ahmad et al. (2013)	Int J Agric Biol 15 (4):791–794	Pakistan	Rice	E50, FE, and yield	Field	Non-stressed	Arid
Ali et al. (2013)	Turk J Agric For 37 (5):534–544	Pakistan	Wheat	FE and yield	Field	Non-stressed	Arid
Ali et al. (2008a, b)	Aust J Crop Sci 2 (3):150–157	Pakistan	Wheat and maize	Yield	Field	Non-stressed and nutrient-stressed	Semi-arid
Anwar et al. (2013)	Pak J Bot 45 (1):157–162	Pakistan	Rice	FE	Lab and field		
Ashraf et al. (2003)	Agronomie 23 (3):227–234	Pakistan	Pearl millet	E50 and FE	Lab		
Aune and Ousman (2011a, b)	Exp Agr 47 (3):419–430	Sudan	Sorghum and pearl millet	FE and yield	Field	Nutrient-stressed	Arid
Aune et al. (2012)	Outlook Agr 41:103–108	Mali	Sorghum and pearl millet	Yield	Field	Nutrient-stressed	Semi-arid
Basra et al. (2011)	Int J Agric Biol 13 (6):1006–1010	Pakistan	Maize	E50 and FE	Pots		
Basu et al. (2014)	Indian J Agr Sci 74 (6):311–315	Bangladesh	Maize	FE	Field		
Chivasa et al. (2000)		Zimbabwe	Maize and sorghum	E50	Pots		
Eyob (2009)	J Med Plants Res 3 (9):652–659	Etiopia	Korarima	FE	Pots		
Farooq et al. (2008a, b)	J Agron Crop Sci 194 (1):55–60	Pakistan	Wheat	E50 and yield	Field	Non-stressed	Arid
Farooq et al. (2017a, b)	Plant Physiol Bioch 111:274–283	Pakistan	Chickpea	FE	Pots		
Fattahi et al. (2011)	Hortic Environ Biote 52 (6):559–566	Iran	<i>Dracocephalum kotschy</i> Boiss	E50 and FE	Pots		
Finch-Savage et al. (2004a, b)	Field Crop Res 90 (2–3):361–374	UK ^b	Maize	E50 and FE	Lab and pots		
Ghassemi-Golezani et al. (2008)	Research Journal of Seed Science 1 (1):34–40	Iran	Chickpea	FE and yield	Field	Non-stressed	Semi-arid
Harris (1996a, b)	Soil and Tillage Research 40 (1–2):73–88	Botswana	Sorghum	E50 and FE	Lab and field		
Harris et al. (2005a, b)	Aust J Agr Res 56 (11):1211–1218	India	Chickpea	Yield	Pots		
Harris et al. (1999a, b)	Exp Agr 35 (1):15–29	India	Chickpea	Yield	Field	Nutrient-stressed	Semi-arid
Harris et al. (2001a)	Agr Syst 69 (1–2):151–164	India	Maize	Yield	Field	Nutrient-stressed	Semi-arid
Harris et al. (2001b)	Exp Agr 37 (3):403–415	India, Nepal and Pakistan	Wheat	E50 and yield	Lab and field	Non-stressed, nutrient-stressed, and salinity	Temperate, tropical, and arid
Harris et al. (2007a, b)	Field Crop Res 102 (2):119–127	Pakistan	Maize	Yield	Field	Nutrient-stressed	Semi-arid
Harris et al. (2008a, b)	Plant Soil 306 (1–2):3–10	Pakistan	Wheat and chickpea	Yield	Field	Nutrient-stressed	Semi-arid
Iqbal and Ashraf (2005)	J Integr Plant Biol 47 (11):1315–1325	Pakistan	Wheat	Yield	Field	Non-stressed and salinity	Arid
Iqbal and Ashraf (2010)	J Agron Crop Sci 196 (6):440–454	Pakistan	Wheat	FE and yield	Lab and field	Non-stressed and salinity	Arid
Islam et al. (2015a, b)	Acta Physiol Plant 37 (8)	Pakistan	Wheat	E50, FE, and yield	Pots	Non-stressed and salinity	Arid
Khanal et al. (2004)	Proc Micronutr South and South East Asia, Kathmandu, Nepal, pp 121–132	Nepal	Chickpea	FE and yield	Field	Nutrient-stressed	Temperate

Table 1 (continued)

Author	Journal	Country	Crop	Response variables	Study type	Yield-limiting factor ^a	Climate zone ^a
Kumar et al. (2002)	Int Sorg Mill Newsl 43(1):90–92	India	Finger millet	Yield	Field	Non-stressed	Temperate
Mani et al. (2013)	J Agrometeorol 15 (2):138–141	India	Wheat	Yield	Field	Non-stressed	Semi-arid
Marwat et al. (2007)	Pak J Bot 39 (5):1583–1591	Pakistan	Maize	Yield	Field	Nutrient-stressed	Semi-arid
Murungu et al. (2004b)	Exp Agr 40 (1):23–36	Zimbabwe	Maize and cotton	E50, FE, and yield	Field	Non-stressed	Semi-arid
Murungu et al. (2004a)	Field Crop Res 89 (1):49–57	Zimbabwe	Maize	E50 and FE	Field		
Murungu and Madanzi (2010)	Afr J Agr Res 5 (17):8	Zimbabwe	Wheat	E50 and FE	Field		
Musa et al. (2001)	Exp Agr 37 (4):509–521	Bangladesh	Chickpea	FE and yield	Field	Non-stressed	Tropical
Neamatollahi et al. (2009)	Not Bot Horti Agrobo 37 (2):190–194	Iran	Fennel	FE	Lab		
Ousman and Aune (2011)	Exp Agr 47 (3):431–443	Sudan	Groundnut, sesame, cowpea	FE and yield	Field	Nutrient-stressed	Arid
Rashid et al. (2004a)	Crop Prot 23 (11):1119–1124	Pakistan	Mungbean	FE and yield	Field	Nutrient-stressed	Semi-arid
Rashid et al. (2004b)	Exp Agr 40 (2):233–244	Pakistan	Mungbean	E50, FE, and yield	Field	Nutrient-stressed	Semi-arid
Rashid et al. (2006)	Eur J Agron 24 (3):276–281	Pakistan	Barley	Yield	Field	Non-stressed, nutrient-stressed, and salinity	Semi-arid
Rehman et al. (2011a)	Int J Agric Biol 13 (5):786–790	Pakistan	Rice	E50 and yield	Field	Non-stressed	Arid
Rehman et al. (2011b)	Turk J Agric For 35 (4):357–365	Pakistan	Rice	E50, FE, and yield	Field	Non-stressed	Arid
Virk et al. (2006)	Exp Agr 42 (4):411–425	India	Horsegram	E50, FE, and yield	Lab and field	Nutrient-stressed	Temperate

E50 time to 50% emergence, FE final emergence

^a Data corresponding to response variable 'yield'

^b Simulating semi-arid climate conditions in cabinets

($\exp(LRR) - 1 \times 100$) for easier interpretation, where LRR is the weighted summary effect size across case studies. Overall effect sizes were considered significant when their confidence intervals did not overlap (Gurevitch and Hedges 1999).

A random effects model was chosen because of the high variation expected between studies due to the diversity of crops and environmental factors. In addition, the aim of this study was to obtain mean effects that can be generalized to different scenarios, which is best done with random effects models (Borenstein et al. 2009).

To investigate the relationship between emergence and yield, pairs of effect sizes of ‘time to 50% emergence’ and ‘final emergence’, and pairs of final emergence and ‘yield’ from the same case studies were analysed using time to 50% emergence and final emergence as moderators, respectively. The influence of each moderator was assessed with F_M (test of moderator) by meta-regression using restricted maximum likelihood with Knapp-Hartung adjustment (Viechtbauer 2010; Inthout et al. 2014), assuming a fixed effect across levels and a random effect within levels (Borenstein et al. 2009). Given the importance of soil interactions, papers reporting laboratory-based experiments were omitted from these specific analyses. To further quantify the extent of yield benefits that can be ascribed to emergence, a hypothesized regression line where changes in final emergence are equal to increments in yield was compared against the weighted linear regression obtained from the meta-regression using linear hypothesis testing.

All calculations were conducted with ‘metafor’ (Viechtbauer 2010), ‘car’ (Fox et al. 2016) and ‘boot’ packages (Canty and Ripley 2012) in R version 3.3.0 (R Development Core Team 2016).

2.3 Moderator variables

Sub-group analysis allowed further exploration of variables in terms of explaining variability and identification of possible trends (Borenstein et al. 2009). We considered levels within moderators to be significantly different from one another when their confidence intervals did not overlap (Gurevitch and Hedges 1999).

The effect of climate on total yield was accounted for by categorizing papers as either ‘temperate’, ‘tropical’ or ‘dry’ according to the Köppen-Geiger climate classification (Kottek et al. 2006). Dry climates were further subdivided into ‘semi-arid’ or ‘arid’ to account for potential evapotranspiration as a function of temperature and cycle of precipitation (Kottek et al. 2006). For this purpose, the high-resolution Köppen-Geiger climate world map (<http://koeppen-geiger.vu-wien.ac.at/present.htm>) was loaded into Google Earth Pro (Wuthrich 2006) and the location of the case studies in each paper used to determine the climate group. When geographical coordinates were not reported, the location of the experimental station or the nearest city at which the study took place was used.

Based on yield-limiting factors, three agronomic scenarios were commonly identified across the case studies and used for evaluation of on-farm seed priming on yield. The first scenario included case studies where crops were grown without major nutrient and water limitations. The second scenario contained case studies where crops were grown under rain-fed conditions and low soil fertility was identified as a major constraint (by authors stating that there were low levels of the main macronutrients or other known nutrient deficiencies in the area). The third scenario contained case studies where salinity was identified as the main constraint or when trials were designed to test the effect of salinity. These scenarios were named as ‘non-stressed’, ‘nutrient deficient’ and ‘salinity stressed’, respectively. Case studies not mentioning or giving ambiguous descriptions about any of these factors were omitted for categorical analyses.

2.4 Dataset overview

Overall, our analysis comprised work conducted in 10 countries across the Middle East, South Asia and sub-Saharan Africa. The three most globally cultivated cereals, wheat (*Triticum aestivum*), maize (*Zea mays*) and rice (*Oryza sativa*), comprised 46% of case studies, whilst 19% of case studies included essential cereals common in semi-arid areas: sorghum (*Sorghum bicolor*), millet (*Pennisetum glaucum* and *Eleusine coracana*) and barley (*Hordeum vulgare*). Legumes, including chickpea (*Cicer arietinum*), mungbean (*Vigna radiata*), cowpea (*Vigna unguiculata*) and horsegram (*Macrotyloma uniflorum*), represented 21% of the case studies. Cash crops, such as sesame (*Sesamum indicum*), cotton (*Gossypium hirsutum*) and groundnut (*Arachis hypogaea*), represented 11% of case studies, and minor crops, fennel (*Foeniculum vulgare*), korarima (*Aframomum corrorima*) and *Dracocephalum kotschyi* Boiss (the last two grown for their spice and medicinal properties) accounted for 3% of case studies analysed.

The dataset of time to 50% emergence was mainly characterized by case studies using small-scale trials (three to four replications) testing the response of varieties or cultivars to on-farm seed priming and, to a lesser extent, different soaking durations. The growing conditions included field trials (46%), pots trials (25%) and lab experiments (29%); most case studies were carried out with monocots (83%). The final emergence dataset encompassed small-scale trials and medium size trials repeated over two to three seasons. More than half of the case studies in this dataset were conducted in the field (61%), with fewer laboratory (22%) and pot trials (17%). Monocot (56%) and dicot species (44%) were almost equally represented in this dataset. For the yield dataset (65 case studies), most of the experiments were conducted under field conditions (with only three pot trials), in both irrigated and in rain-fed plots. Over half of the case studies were carried out at research farms, commonly testing priming treatments on different cultivars or varieties over several seasons, averaging 15

experimental replications per study. The remaining 43% of the case studies were mainly participatory trials carried out by local farmers following local practices and constraints. The average experimental replications for these case studies was 38, and the biggest study accounted for 108 trials of wheat across the state of Gujarat in India (Harris et al. 2001b).

2.5 Publication bias and sensitivity analysis

Studies showing negative results are less likely to be published; therefore, effect sizes in meta-analyses could be overestimated (Gurevitch and Hedges 1999). Consequently, indirect methods such as rank correlation tests and funnel plots of effect size vs. variance are commonly used to detect bias (Gurevitch and Hedges 1999; Koricheva and Gurevitch 2014). We conducted Kendall's tau test, where significant correlation between effect size and corresponding sample size would indicate asymmetry in the funnel plot and therefore, potential publication bias (Begg and Mazumdar 1994; Viechtbauer 2010). However, no significant relationship between effect size and increasing number of replicates for any of the three datasets in our analysis was seen. We also performed 'trim and fill' funnel plots to detect potential missing studies. The trim and fill method is a funnel-based test that imputes values that would compensate for the most extreme values in one side of the funnel (Duval et al. 2000). In our meta-analysis, trim and fill imputed 12 and 15 potential missing case studies in time to 50% emergence and yield datasets, respectively. In both cases, adjusted summary effects would further deviate from zero suggesting that our results may be conservative (summary tables and funnel plots from these analyses are available at <http://hdl.handle.net/11667/123>).

Sensitivity analyses are also important to determine the robustness of the results (Koricheva and Gurevitch 2014). Leave-one-out meta-analysis, i.e. recalculating summary effect size omitting the study with highest effect size for each variable and observing the deviation introduced by this modification, was performed to test robustness of the summary effects. The removal of the study with largest influence in the yield dataset (Harris et al. 2001b) increased the summary effect by 1.42%. The study with the largest influence on final emergence was Finch-Savage et al. (2004), whose removal changed the summary effect by 1.89%. Lastly, the study with biggest impact on the time to 50% emergence dataset was Harris et al. (2001b), and its removal decreased the overall effect size by 6.04%. In conclusion, we did not find evidence of publication bias in our datasets, and although the time to 50% emergence dataset presented some sensitivity, all three datasets were suitable for meta-analysis.

Data availability The datasets generated during and analysed during the current study are available in the Stirling Online Repository for Research Data repository, <http://hdl.handle.net/11667/123>.

3 Results and discussion

3.1 'On-farm' seed priming: an inexpensive technology for increased food security

Our meta-analysis showed that on-farm seed priming has a significantly positive effect on crop performance, from nascence until harvest, relative to conventional ('control') seed sowing (Fig. 2). Although there is substantial variation (ranging from -36 to -7%), on-farm seed priming significantly decreases the time to emergence by 22% compared with non-primed seeds. On average, the number of plants emerged increased by 11%. Ultimately, yields increased by 21% compared with non-primed seeds, and only six out the 65 case studies reported negative effects on yield (data not shown).

Improved crop performance following on-farm seed priming can have important implications for smallholders' food production. Higher yield is often accompanied by higher straw biomass, which is especially remunerative in mixed crop-livestock systems. Enhanced plant density reduces costs and the labour needed for re-sowing, and can also increase the willingness of farmers to invest in fertilizers, as the risk of plant stand failure is lower. Faster emergence typically results in plants reaching flowering and harvest stages earlier (e.g. by several weeks), giving farmers more labour flexibility, for example, by facilitating more optimal sowing for the subsequent crop or including an extra crop in rotation systems, or even by allowing migration for off-season work (Harris et al., 1999a, b, 2001a; Virk et al., 2006). Furthermore, the benefits are not restricted to the traits accounted for in our data, as faster development combined with the improved vigour and more uniform emergence in crops from on-farm primed seeds may save labour allocated to weeding. Although it is tempting to suggest that these benefits may increase net incomes, additional costs such as extra fertilizer or extra costs associated with harvesting, processing and storing greater yields, together with access to markets, will determine the final return from adopting on-farm seed priming.

3.2 Relationships between early growth and yield on crops grown from 'on-farm' primed seeds

To further investigate the relationships between rate of emergence, crop establishment and yield, we conducted separate analyses of the effect of time to 50% emergence on final emergence and the effect of final emergence on yield of crops from on-farm primed seeds relative to non-primed. Final emergence versus time to 50% emergence showed that in general, quicker emergence conferred by on-farm seed priming relative to non-primed seeds produced a higher number of successfully emerged seedlings (Fig. 3a). Although this relationship was significant ($P < 0.01$), it must be interpreted with caution due to the relatively small number of case studies.




Fig. 2 Summary analyses of the response of crops to ‘on-farm’ seed priming. Numbers in parentheses indicate number of case studies. Error bars represent back-transformed 95% bootstrap CIs




Fig. 3 **a** Relationship between final emergence and time to 50% emergence relative to crops from non-primed seeds ($n = 12$). **b** Relationship between field and final emergence relative to crops grown from non-primed seeds. Solid line represents the weighted model regression line, and dotted line represents the hypothesized regression line where changes in final emergence cause equal changes in yield ($n = 22$). Bubble size represents the weight of each study in the meta-regression

Meta-regression of yield versus final emergence (relative to crops from non-primed seeds) showed a positive relationship (Fig. 3b), although this relationship was not found significant. We found no difference between the hypothesized line and the meta-regression line ($P > 0.05$), which demonstrates that higher yields are proportional to improvements in emergence. However, in over two-thirds of the case studies, improvements in yield were proportionally higher than the expected gain due solely to improvement in final emergence. This suggests that increments in yield due to on-farm seed priming are not only a consequence of rapid and more prolific emergence, but that additional benefits may persist long after emergence.

Rapid emergence is crucial for the vulnerable seedling to avoid abiotic and biotic stresses and ensure high crop establishment (Gardarin et al. 2016). On-farm seed priming facilitates rapid emergence by accelerating germination through two complementary mechanisms. Firstly, it ensures water availability and the successful completion of phase I (the imbibition phase) prior to sowing, rather than relying on the seed imbibing soil moisture in the field where the water supply can be restricted or discontinuous (Wojtyla et al. 2016). Throughout the imbibition phase, both mechanical and biochemical changes, e.g. embryo enlargement, respiration, protein synthesis and DNA repair, are initiated (Gallardo et al. 2001; Weitbrecht et al. 2011; Steinbrecher and Leubner-Metzger 2017). All these processes prepare the seed for cell elongation (phase II, the lag phase); therefore, on-farm primed seeds are developmentally more advanced than dried seeds, resulting in a ‘head start of germination’ (Chen and Arora 2013). Secondly, on-farm primed seeds are only externally dried so that, once in the field, seeds need to absorb less water from the soil to complete phase III (the post-germination phase) when the radicle emerges from the seed coat. Furthermore, it has been reported that seed soaking enhances the production of the enzyme α -amylase (Ashraf and Foolad 2005; Farooq et al., 2017), which plays a crucial role in starch mobilization and provides the embryo with carbohydrates for

respiration during germination and seedling growth (Ashraf and Foolad 2005; Farooq et al., 2017a, b). As a result, seedlings from on-farm primed seeds have more developed roots before the common limiting factors such as declining soil moisture, crust formation and/or high salinity prevent successful emergence.

Our results suggest that the gains in yield due to on-farm seed priming can be mainly attributed to enhanced emergence, i.e. rapid emergence leads to better crop establishment, which is conducive to higher yields. However, advanced establishment may also be coupled with higher vigour of individual plants, which is translated into significantly more tillers, more fruits (cobs/panicles/pods) per plant, greater number of grain and 1000-grain weight, or straw yield (Harris 2006; Rashid et al., 2006; Harris et al., 2007; Farooq et al., 2008). In addition to these physiological benefits, other circumstantial benefits are frequently observed, for example, earlier maturation decreases crop exposure to end of season drought, disease and pest attacks (Harris et al. 2001a; Rashid et al., 2006a, b). It is also likely that seed priming exerts important metabolic changes during early plant growth that are able to persist until later stages of development (Ashraf and Foolad 2005; Chen and Arora 2013); for example, there is evidence for enhanced

disease resistance (Musa et al., 2001; Rashid et al. 2004; Harris et al., 2005) or drought tolerance (Wojtyła et al. 2016).

3.3 What modulates the 'priming' response?

It is important not only to identify the context where on-farm seed priming can best be applied, but also understand the potential situations where it can be counterproductive. Therefore, a subgroup analysis of moderators was conducted to examine potential factors that influence the effect of seed priming.

3.3.1 Climate

It is clear that yield benefits are more evident under low and unpredictable rain conditions. The largest response to on-farm seed priming was seen in areas with dry climates (Fig. 4a) with significantly higher yields for both arid (22%) and semi-arid (28%) climates compared to temperate climates (11%). Variation in yield between seasons due to on-farm seed priming has been frequently attributed to rainfall profiles, with greater yield increments commonly reported during rainy seasons with limited precipitation (Rashid et al., 2006a, b; Virk et al., 2006; Ousman and Aune, 2011). Low soil moisture and high evapotranspiration can slow and interrupt imbibition, which is conducive to emergence failure (Harris 1996); however, on-farm seed priming can offset a lack of soil moisture, as seeds have already imbibed water prior to sowing.

Importantly, in crust-prone soils, if rainfall occurs before emergence, shoots from on-farm primed seeds could be mechanically impeded, whilst the later emerging non-primed seedling may find more favourable soil strength (Murungu et al. 2004a). Equally, if rainfall is considerably delayed after sowing, seedlings from on-farm primed seeds may be damaged as germination has already been initiated and a lack of water could kill the developing seedling, whereas non-primed seeds will not initiate germinate until the rain comes (Murungu et al. 2003; Rashid et al., 2006a, b). However, the occurrence of these events seems to be very rare (Murungu et al. 2003, 2004a), and our data at emergence stage is consistent with the yield subgrouping, i.e. showing the higher benefits under dry climates (data not shown).

The interaction between soil temperature and on-farm seed priming, however, is less clear. Primed maize seed is more sensitive to elevated temperature under both dry and wet soil conditions (Finch-Savage et al., 2004a, b; Murungu et al. 2004b). For the former, internal seed moisture may induce heat stress by acting as a thermal conductor in soils of higher temperatures, while wet soils may exacerbate prolonged hypoxia (Finch-Savage et al., 2004a, b). Conversely, late sown wheat and chickpea plants from on-farm primed seeds have shown increased tolerance to chilling temperatures (Farooq et al., 2008a, b, Farooq et al., 2017a, b), possibly due to enhanced carbohydrate supply to the germinating embryo, which together




Fig. 4 Sub-grouped summary effect sizes and 95% CIs for priming effect on crop yield. Comparisons among **a** levels of climate, **b** levels of yield-limiting factors, and **c** levels of plant type. Numbers in parentheses indicate number of case studies. Error bars represent back-transformed 95% bootstrap confidence intervals

with an accumulation of trehalose, can protect proteins and membranes from oxidative damage under abiotic stress.

3.3.2 Yield-limiting factors

Figure 4b shows that crops from on-farm seed grown under 'salinity stress' or in nutrient deficient soils had significantly higher yields compared to crops from on-farm seed grown under non-stressed environments (approximately 16% difference). In saline environments, germination is delayed or inhibited through reduced water availability and/or accumulation of toxic Na^+ and Cl^- . However, primed seeds are already hydrated and therefore less subjected to these constraints (Ibrahim 2016; Savvides et al. 2016). Importantly, case studies growing crops in conditions defined as non-stressed were mostly from research-managed trials using fertilizers and pesticides, whilst case studies grouped as nutrient deficient were mainly from farmer-managed trials with limited access to fertilizers and pesticides, and therefore more accurately reflect resource-poor farming conditions in marginal areas. These data indicate that on-farm seed priming can compensate, to some extent, for low-yielding environments and the lack of inputs that would further limit yields. Under low fertility environments, the quicker development of seedlings from on-farm primed seeds allows greater uptake from fertilizers, before nutrients are leached from the soil surface or become volatilized (Harris et al. 2001b; Rashid et al., 2006a, b).

Declining soil fertility together with limited access to affordable mineral fertilizers is a major constraint for achieving optimal yields in marginal areas of developing countries (Chianu et al. 2012). However, low-cost strategies that combine on-farm seed priming with low amounts of inorganic fertilizers have been carried out to alleviate nutrient deficiencies with promising results (Aune and Ousman, 2011a, b; Ousman and Aune, 2011a, b). On-farm seed priming in combination with micro-dosing, i.e. application of small amounts of fertilizer in the planting pocket, demonstrated greater fertilizer use efficiency than micro-dosing alone for all the crops tested (Aune and Ousman, 2011a, b; Ousman and Aune, 2011a, b). Small amounts of micronutrients added to the water used for on-farm seed priming, e.g. ZnSO_4 , can also be highly cost-effective (Harris et al., 2007a, b, Farooq et al., 2008a, b).

3.3.3 Plant type

On-farm seed priming of all the major tropical crops produces similar or greater yields than traditionally sown crops in almost all cases (data not shown). However, decreased performance following on-farm seed priming has also been occasionally reported for barley (Rashid et al., 2006a, b), pearl millet (Aune and Ousman, 2011a, b), rice (Rehman et al. 2011), sesame (Ousman and Aune, 2011a, b), maize (Ali et al., 2008), wheat (Islam et al., 2015), and cotton

(Murungu et al. 2004b), although for each of these crops, there are also studies showing an increased performance (e.g. Harris et al., 2007a, b; Farooq et al., 2008a, b; Rashid et al., 2006a, b). Importantly, negative results are rarely attributed to the incompatibility of priming with the crop, but rather to untimely adverse environmental conditions. The largest yield loss due to on-farm seed priming was 8% for pearl millet in a series of on-station trials; however, in this study, the farmer-managed replicates registered a 30% increase in yield (Aune and Ousman, 2011a, b). Therefore, we have found no consistent evidence of negative interactions between specific crops and on-farm seed priming, which suggests that this is therefore a safe practice for all crop species trialled so far.

The effect of categorizing case studies by plant type on total yield is shown in Fig. 4c. On average, the yield increase of cereals (monocots) was 13% less than dicots. Dicot plants, broadly represented by legumes with 18 out of 23 case studies, responded better to on-farm seed priming averaging a 28% yield increase. This is in line with our final emergence data where greater effect sizes generally belonged to dicotyledonous crops (data not shown). Cereals were commonly grown with irrigation or during the rainy season, whilst legumes were sown as a component of the rotation after cereals in the post-rainy season or in fallow lands that were unsuitable for the main crop. In these marginal contexts, the benefit of seed being hydrated prior to sowing leads to more rapid emergence and establishment.

We cannot conclude from our data whether specific crops are more responsive to on-farm seed priming than others; however, on-farm seed priming may facilitate the use of legumes into rotational and intercropping systems. Currently, in both rotational and intercropping systems, the adoption of legumes is largely discouraged due to poor establishments of the legume component. In rotation, legumes are commonly grown utilizing residual soil moisture remaining during the dry season, and with no additional fertilization, whilst in intercropping systems, the planting of a legume companion is delayed in order to avoid shading and competition (Masvaya et al. 2017). Therefore, on-farm seed priming may ameliorate these unfavourable planting conditions and boost the benefits of cereal-legume cropping systems, e.g. by improving soil fertility and providing an additional income.

4 Conclusion

In developing regions of the world, tackling yield reductions due to both natural and socio-economic constraints, e.g. increasingly unpredictable rainfalls, declining soil fertility and limited access to inputs and resources, requires inexpensive and sustainable strategies to ensure food production and self-sufficiency. This is the first study quantifying the potential of on-farm seed priming for sustainably increasing food production at a global scale, and our results have shown that it is a

valid approach to closing yield gaps. The literature considered in our meta-analysis encompassed a representative number of agro-environments where on-farm seed priming can be practiced and gives us the basis to draw the following conclusions.

On-farm seed priming attenuates the negative effects of adverse planting conditions, and low inputs, by facilitating rapid and enhanced crop establishment that may also result in improved individual plant performance. These effects are more evident in semi-arid and arid regions and, given that millions of hectares in dry climates are experiencing yield reductions, these findings could have important implications. Our results have also highlighted that crops grown in marginal lands can especially benefit from this intervention. This is particularly important for farmers with limited access to mineral fertilizers where to a large extent an input of N is dependent on N₂ fixed by legumes.

On-farm seed priming can be seen as a starting point towards sustainable intensification in marginal areas of the developing world. This technology requires very few resources and technical knowledge, and its benefits would be compatible with a range of other sustainable strategies such as smart-use of farmyard manure, micro-dosing and water harvesting practices. Therefore, our results provide the evidence needed to encourage governmental institutions and policymakers in developing countries to promote the adoption of on-farm seed priming as recommended practice.

Acknowledgments The authors would like to thank the Ekhaga Foundation (Sweden) for their financial support.

Compliance with ethical standards

This study was funded by Ekhaga Foundation (2015-60).

Conflict of interest The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Adams DC, Gurevitch J, Rosenberg MS (1997) Resampling tests for meta analysis of ecological data. *Ecology* 78:1277–1283. <https://doi.org/10.2307/2265879>
- Ali S, Khan AR, Mairaj G et al (2008a) Assessment of different crop nutrient management practices for yield improvement. *Aust J Crop Sci* 2:150–157
- Ashraf M, Foolad MR (2005) Pre-sowing seed treatment—a shotgun approach to improve germination, plant growth, and crop yield under saline and non-saline conditions. *Adv Agron* 88:223–271. [https://doi.org/10.1016/S0065-2113\(05\)88006-X](https://doi.org/10.1016/S0065-2113(05)88006-X)
- Aune JB, Coulibaly A, Giller KE (2017) Precision farming for increased land and labour productivity in semi-arid West Africa. A review. *Agron Sustain Dev* 37(3):16. <https://doi.org/10.1007/s13593-017-0424-z>
- Aune JB, Ousman A (2011a) Effect of seed priming and micro-dosing of fertilizer on sorghum and pearl millet in Western Sudan. *Exp Agric* 47:419–430. <https://doi.org/10.1017/S0014479711000056>
- Begg CB, Mazumdar M (1994) Operating characteristics of a rank correlation test for publication bias. *Biometrics* 50:1088–1101. <https://doi.org/10.2307/2533446>
- Borenstein M, Hedges LV, Higgins JPT, Rothstein HR (2009) Introduction to meta-analysis. *Psychother Res J Soc Psychother Res* 19:421. <https://doi.org/10.1002/9780470743386>
- Branca G, Lipper L, McCarthy N, Jolejole MC (2013) Food security, climate change, and sustainable land management. A review. *Agron Sustain Dev* 33:635–650. <https://doi.org/10.1007/s13593-013-0133-1>
- Canty A, Ripley B (2013). boot: Bootstrap R (S-Plus) Functions. R package version 1.3-9, <http://CRAN.R-project.org/package=boot>
- Castagneyrol B, Jactel H (2012) Unraveling plant–animal diversity relationships: a meta-regression analysis. *Ecology* 93:2115–2124. <https://doi.org/10.2307/41739269>
- Chen K, Arora R (2013) Priming memory invokes seed stress-tolerance. *Environ Exp Bot* 94:33–45. <https://doi.org/10.1016/j.envexpbot.2012.03.005>
- Chianu JN, Chianu JN, Mairura F (2012) Mineral fertilizers in the farming systems of sub-Saharan Africa. A review. *Agron Sustain Dev* 32:545–566. <https://doi.org/10.1007/s13593-011-0050-0>
- Duval S, Tweedie R, Taylor S (2000) Trim and fill: a simple funnel plot based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 56:455–463. <https://doi.org/10.1111/j.0006-341x.2000.00455.x>
- Farooq M, Basra SMA, Rehman H, Saleem BA (2008a) Seed priming enhances the performance of late sown wheat (*Triticum aestivum* L.) by improving chilling tolerance. *J Agron Crop Sci* 194:55–60. <https://doi.org/10.1111/j.1439-037X.2007.00287.x>
- Farooq M, Hussain M, Nawaz A, Lee DJ, Alghamdi SS, Siddique KHM (2017a) Seed priming improves chilling tolerance in chickpea by modulating germination metabolism, trehalose accumulation and carbon assimilation. *Plant Physiol Biochem* 111:274–283. <https://doi.org/10.1016/j.plaphy.2016.12.012>
- Finch-Savage WE, Dent KC, Clark LJ (2004a) Soak conditions and temperature following sowing influence the response of maize (*Zea mays* L.) seeds to on-farm priming (pre-sowing seed soak). *F Crop Res* 90:361–374. <https://doi.org/10.1016/j.fcr.2004.04.006>
- Fox J, Weisberg S, Adler D, et al (2016) Package “car.” In: CRAN Repos. <https://cran.r-project.org/web/packages/car/car.pdf>
- Gallardo K, Job C, Groot SPC, Puype M, Demol H, Vandekerckhove J, Job D (2001) Proteomic analysis of arabidopsis seed germination and priming. *Plant Physiol* 126:835–848. <https://doi.org/10.1104/pp.126.2.835>
- Gardarin A, Coste F, Wagner MH, Dürr C (2016) How do seed and seedling traits influence germination and emergence parameters in crop species? A comparative analysis. *Seed Sci Res* 26:317–331. <https://doi.org/10.1017/S0960258516000210>
- Gurevitch J, Hedges L (1999) Statistical issues in ecological meta-analyses. *Ecology* 80:1142–1149. [https://doi.org/10.1890/0012-9658\(1999\)080\[1142:SIEMA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1142:SIEMA]2.0.CO;2)
- Harris D (1996a) The effects of manure, genotype, seed priming, depth and date of sowing on the emergence and early growth Sorghum bicolor (L.) Moench in semi-arid Botswana. *Soil Tillage Res* 40:73–88. [https://doi.org/10.1016/S0167-1987\(96\)01047-1](https://doi.org/10.1016/S0167-1987(96)01047-1)

- Harris D (2006) Development and testing of “on-farm” seed priming. *Adv Agron* 90:129–178. [https://doi.org/10.1016/S0065-2113\(06\)90004-2](https://doi.org/10.1016/S0065-2113(06)90004-2)
- Harris D, Breese WA, Rao JVDKK (2005a) The improvement of crop yield in marginal environments using “on-farm” seed priming: nodulation, nitrogen fixation and disease resistance. *Aust J Agric Res* 56:1211–1218. <https://doi.org/10.1071/AR05079>
- Harris D, Joshi A, Khan PA, Gothkar P, Sodhi PS (1999a) On-farm seed priming in semi-arid agriculture: development and evaluation in maize, rice and chickpea in India using participatory methods. *Exp Agric* 35:15–29. <https://doi.org/10.1017/S0014479799001027>
- Harris D, Pathan AK, Gothkar P, Joshi A, Chivasa W, Nyamudeza P (2001a) On-farm seed priming: using participatory methods to revive and refine a key technology. *Agric Syst* 69:151–164. [https://doi.org/10.1016/S0308-521X\(01\)00023-3](https://doi.org/10.1016/S0308-521X(01)00023-3)
- Harris D, Raghuvanshi BS, Gangwar JS, Singh SC, Joshi KD, Rashid A, Hollington PA (2001b) Participatory evaluation by farmers of “on-farm” seed priming in wheat in India. *Exp Agric* 37:403–415. <https://doi.org/10.1017/S0014479701003106>
- Harris D, Rashid A, Miraj G, Arif M, Shah H (2007a) “On-farm” seed priming with zinc sulphate solution—a cost-effective way to increase the maize yields of resource-poor farmers. *F Crop Res* 102:119–127. <https://doi.org/10.1016/j.fcr.2007.03.005>
- Harris D, Rashid A, Miraj G, Arif M, Yunas M (2008a) “On-farm” seed priming with zinc in chickpea and wheat in Pakistan. *Plant Soil* 306:3–10. <https://doi.org/10.1007/s11104-007-9465-4>
- Hedges LV, Gurevitch J, Curtis PS (1999) The meta-analysis of response ratios in experimental ecology. *Ecology* 80:1150–1156. [https://doi.org/10.1890/0012-9658\(1999\)080\[1150:TMAORR\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1150:TMAORR]2.0.CO;2)
- Ibrahim EA (2016) Seed priming to alleviate salinity stress in germinating seeds. *J Plant Physiol* 192:38–46. <https://doi.org/10.1016/j.jplph.2015.12.011>
- Inthout J, Ioannidis JP, Borm GF (2014) The Hartung-Knapp-Sidik-Jonkman method for random effects meta-analysis is straightforward and considerably outperforms the standard DerSimonian-Laird method. *BMC Med Res Methodol* 14:1–12. <https://doi.org/10.1186/1471-2288-14-25>
- Islam F, Yasmeen T, Ali S, Ali B, Farooq MA, Gill RA (2015a) Priming-induced antioxidative responses in two wheat cultivars under saline stress. *Acta Physiol Plant* 37:1–12. <https://doi.org/10.1007/s11738-015-1897-5>
- Knox J, Hess T, Daccache A, Wheeler T (2012) Climate change impacts on crop productivity in Africa and South Asia. *Environ Res Lett* 7:034032-. <https://doi.org/10.1088/1748-9326/7/3/034032>
- Koricheva J, Gurevitch J (2014) Uses and misuses of meta-analysis in plant ecology. *J Ecol* 102:828–844. <https://doi.org/10.1111/1365-2745.12224>
- Koricheva J, Gurevitch J, Mengersen K (2013) Handbook of meta-analysis in ecology and evolution. Princeton University Press, New Jersey, Princeton
- Kottek M, Grieser J, Beck C, Rudolf B, Rubel F (2006) World map of the Köppen-Geiger climate classification updated. *Meteorol Z* 15:259–263. <https://doi.org/10.1127/0941-2948/2006/0130>
- Masvaya EN, Nyamangara J, Descheemaeker K, Giller KE (2017) Is maize-cowpea intercropping a viable option for smallholder farms in the risky environments of semi-arid southern Africa? *F Crop Res* 209:73–87. <https://doi.org/10.1016/j.fcr.2017.04.016>
- Mayerhofer MS, Kernaghan G, Harper KA (2013) The effects of fungal root endophytes on plant growth: a meta-analysis. *Mycorrhiza* 23:119–128. <https://doi.org/10.1007/s00572-012-0456-9>
- Murungu FS, Chiduzza C, Nyamugafata P, Clark LJ, Whalley WR, Finch-Savage WE (2004a) Effects of “on-farm seed priming” on consecutive daily sowing occasions on the emergence and growth of maize in semi-arid Zimbabwe. *F Crop Res* 89:49–57. <https://doi.org/10.1016/j.fcr.2004.01.020>
- Murungu FS, Chiduzza C, Nyamugafata P et al (2004b) Effect of on-farm seed priming on emergence, growth and yield of cotton and maize in a semi-arid area of Zimbabwe. *Exp Agric* 40. S0014479703001509, <https://doi.org/10.1017/S0014479703001509>
- Murungu FS, Nyamugafata P, Chiduzza C et al (2003) Effects of seed priming, aggregate size and soil matric potential on emergence of cotton (*Gossypium hirsutum* L.) and maize (*Zea mays* L.). *Soil Tillage Res* 74:161–168. <https://doi.org/10.1016/j.still.2003.06.003>
- Musa AM, Harris D, Johansen C, Kumar JVDK (2001a) Short duration chickpea to replace fallow after aman rice: the role of on-farm seed priming in the high Barind tract of Bangladesh. *Exp Agric* 37:509–521. <https://doi.org/10.1017/S0014479701000448>
- Nabi G, Mullins CE, Montemayor MB, Akhtar MS (2001) Germination and emergence of irrigated cotton in Pakistan in relation to sowing depth and physical properties of the seedbed. *Soil Tillage Res* 59:33–44. [https://doi.org/10.1016/S0167-1987\(00\)00182-3](https://doi.org/10.1016/S0167-1987(00)00182-3)
- Ousman A, Aune JB (2011a) Effect of seed priming and micro-dosing of fertilizer on groundnut, sesame and cowpea in Western Sudan. *Exp Agric* 47:431–443. <https://doi.org/10.1017/S0014479711000056>
- Passioura JB, Angus JF (2010) Improving productivity of crops in water limited environments. *Adv Agron* 106:37–75. [https://doi.org/10.1016/S0065-2113\(10\)06002-5](https://doi.org/10.1016/S0065-2113(10)06002-5)
- R Development Core Team (2016) R: A Language and Environment for Statistical Computing. R Found Stat Comput, Vienna 0: {ISBN} 3–900051–07-0. <https://doi.org/10.1038/sj.hdy.6800737>
- Rashid A, Harris D, Hollington P, Ali S (2004) On-farm seed priming reduces yield losses of mungbean (*Vigna radiata*) associated with mungbean yellow mosaic virus in the north west Frontier Province of Pakistan. *Crop Prot* 23:1119–1124. <https://doi.org/10.1016/j.cropro.2004.04.002>
- Rashid A, Hollington PA, Harris D, Khan P (2006a) On-farm seed priming for barley on normal, saline and saline-sodic soils in north west Frontier Province, Pakistan. *Eur J Agron* 24:276–281. <https://doi.org/10.1016/j.eja.2005.10.006>
- Rehman H, Basra SMA, Farooq M et al (2011) Seed priming with CaCl₂ improves the stand establishment, yield and quality attributes in direct seeded rice (*Oryza sativa*). *Int J Agric Biol* 13:786–790
- Rohatgi A (2015) WebPlotDigitizer 3.8. <http://arohatgi.info/WebPlotDigitizer>
- Savvides A, Ali S, Tester M, Fotopoulos V (2016) Chemical priming of plants against multiple abiotic stresses: mission possible? *Trends Plant Sci* 21:329–340. <https://doi.org/10.1016/j.tplants.2015.11.003>
- Shrestha U, Augé RM, Butler DM (2016) A meta-analysis of the impact of anaerobic soil disinfestation on pest suppression and yield of horticultural crops. *Front Plant Sci* 7:1–20. <https://doi.org/10.3389/fpls.2016.01254>
- Steinbrecher T, Leubner-Metzger G (2017) The biomechanics of seed germination. *J Exp Bot* 68:765–783. <https://doi.org/10.1093/jxb/erw428>
- Tisdall JM (1996) Crop establishment—a serious limitation to high productivity. *Soil Tillage Res* 40:1–2. [https://doi.org/10.1016/S0167-1987\(96\)80002-X](https://doi.org/10.1016/S0167-1987(96)80002-X)
- Tittonell P, Giller KE (2013) When yield gaps are poverty traps: the paradigm of ecological intensification in African smallholder agriculture. *F Crop Res* 143:76–90. <https://doi.org/10.1016/j.fcr.2012.10.007>
- Tonitto C, Ricker-Gilbert JE (2016) Nutrient management in African sorghum cropping systems: applying meta-analysis to assess yield and profitability. *Agron Sustain Dev* 36:1–19. <https://doi.org/10.1007/s13593-015-0336-8>
- Townend J, Mtakwa PW, Mullins CE, Simmonds LP (1996) Soil physical factors limiting establishment of sorghum and cowpea in two contrasting soil types in the semi-arid tropics. *Soil Tillage Res* 40:89–106. [https://doi.org/10.1016/S0167-1987\(96\)01048-3](https://doi.org/10.1016/S0167-1987(96)01048-3)
- Viechtbauer W (2010) Conducting meta-analyses in R with the metafor package. *J Stat Softw* 36:1–48
- Virk DS, Chakraborty M, Ghosh J, Harris D (2006a) Participatory evaluation of horsegram (*Macrotyloma uniflorum*) varieties and their

- on-station responses to on-farm seed priming in eastern India. *Exp Agric* 42:411. <https://doi.org/10.1017/S0014479706003838>
- Weitbrecht K, Müller K, Leubner-Metzger G (2011) First off the mark: early seed germination. *J Exp Bot* 62:3289–3309. <https://doi.org/10.1093/jxb/err030>
- Wojtyła Ł, Lechowska K, Kubala S, Garmczarska M (2016) Molecular processes induced in primed seeds—increasing the potential to stabilize crop yields under drought conditions. *J Plant Physiol* 203: 116–126. <https://doi.org/10.1016/j.jplph.2016.04.008>
- Wutrich D (2006) Google earth pro. *Geospatial Solut* 16:30–32
- ## References of the meta-analysis
- Abdalla EA, Osman AK, Maki MA, Nur FM, Ali SB, Aune JB (2015) The response of sorghum, groundnut, sesame, and cowpea to seed priming and fertilizer micro-dosing in South Kordofan state, Sudan. *Agronomy-Basel* 5(4):476–490. <https://doi.org/10.3390/agronomy5040476>
- Abro SA, Mahar AR, Mirbahar AA (2009) Improving yield performance of landrace wheat under salinity stress using on-farm seed priming. *Pak J Bot* 41(5):2209–2216
- Ahmad R, Hussain S, Farooq M, Atique-Ur-Rehman JA (2013) Improving the performance of direct seeded system of rice intensification by seed priming. *Int J Agric Biol* 15(4):791–794
- Ali H, Iqbal N, Shahzad AN, Sarwar N, Ahmad S, Mehmood A (2013) Seed priming improves irrigation water use efficiency, yield, and yield components of late-sown wheat under limited water conditions. *Turk J Agric For* 37(5):534–544. <https://doi.org/10.3906/tar-1207-70>
- Ali S, Khan R, Mairaj G, Arif M, Fida M, Bibi S (2008b) Assessment of different crop nutrient management practices for yield improvement. *Aust J Crop Sci* 2(3):150–157
- Anwar S, Iqbal M, Raza SH, Iqbal N (2013) Efficacy of seed preconditioning with salicylic and ascorbic acid in increasing vigor of rice (*Oryza sativa* L.) seedling. *Pak J Bot* 45(1):157–162
- Ashraf M, Kausar A, Ashraf MY (2003) Alleviation of salt stress in pearl millet (*Pennisetum glaucum* (L.) R. Br.) through seed treatments. *Agronomie* 23(3):227–234. <https://doi.org/10.1051/agro:2002086>
- Aune JB, Ousman A (2011b) Effect of seed priming and micro-dosing of fertilizer on sorghum and pearl millet in Western Sudan. *Exp Agric* 47(3):419–430. <https://doi.org/10.1017/S0014479711000056>
- Aune JB, Traoré CO, Mamadou S (2012) Low-cost technologies for improved productivity of dryland farming in Mali. *Outlook Agric* 41(2):103–108. <https://doi.org/10.5367/oa.2012.0084>
- Basra SMA, Iftikhar MN, Afzal I (2011) Potential of Moringa (*Moringa oleifera*) leaf extract as priming agent for hybrid maize seeds. *Int J Agric Biol* 13(6):1006–1010
- Basu S, Sharma SP, Dadlani M (2014) Effect of seed-invigorating treatments on field emergence and crop performance of maize (*Zea mays*) parental lines in monsoon, winter and spring-summer season. *Indian J Agric Sci* 74(6):311–315
- Chivasa W, Harris D, Chiduzo C, Mashingaidze AB, Nyamudeza P (2000) Determination of optimum on-farm seed priming time for maize (*Zea mays* L.) and -Sorghum (*Sorghum bicolor* [L.] Moench) for use to improve stand establishment in semi-arid agriculture. *Tanzan J Agric Sci* 2:103–112
- Eyob S (2009) Promotion of seed germination, subsequent seedling growth and in vitro propagation of korarima *Aframomum corrorima* (Braun) P. C. M. Jansen. *J Med Plant Res* 3(9):652–659
- Farooq M, Basra SMA, Rehman H, Saleem BA (2008b) Seed priming enhances the performance of late sown wheat (*Triticum aestivum* L.) by improving chilling tolerance. *J Agron Crop Sci* 194(1):55–60. <https://doi.org/10.1111/j.1439-037X.2007.00287.x>
- Farooq M, Hussain M, Nawaz A, Lee D-J, Alghamdi SS, Siddique KHM (2017b) Seed priming improves chilling tolerance in chickpea by modulating germination metabolism, trehalose accumulation and carbon assimilation. *Plant Physiol Biochem* 111:274–283. <https://doi.org/10.1016/j.plaphy.2016.12.012>
- Fattahi M, Nazeri V, Sefidkon F, Zamani Z, Palazon J (2011) The effect of pre-sowing treatments and light on seed germination of *Dracocephalum kotschyi* Boiss: an endangered medicinal plant in Iran. *Hortic Environ Biotechnol* 52(6):559–566. <https://doi.org/10.1007/s13580-011-0057-0>
- Finch-Savage WE, Dent KC, Clark L (2004b) Soak conditions and temperature following sowing influence the response of maize (*Zea mays* L.) seeds to on-farm priming (pre-sowing seed soak). *Field Crop Res* 90(2–3):361–374. <https://doi.org/10.1016/j.fcr.2004.04.006>
- Ghassemi-Golezani K, Sheikhzadeh-Mosaddegh P, Valizadeh M (2008) Effects of Hydropriming duration and limited irrigation on field performance of chickpea. *Res J Seed Sci* 1(1):34–40. <https://doi.org/10.3923/rjss.2008.34.40>
- Harris D (1996b) The effects of manure, genotype, seed priming, depth and date of sowing on the emergence and early growth of Sorghum bicolor (L.) Moench in semi-arid Botswana. *Soil Tillage Res* 40(1–2):73–88. [https://doi.org/10.1016/S0167-1987\(96\)01047-1](https://doi.org/10.1016/S0167-1987(96)01047-1)
- Harris D, Breese WA, Rao JVDKK (2005b) The improvement of crop yield in marginal environments using ‘on-farm’ seed priming: nodulation, nitrogen fixation, and disease resistance. *Aust J Agric Res* 56(11):1211–1218. <https://doi.org/10.1071/Ar05079>
- Harris D, Joshi A, Khan PA, Gothkar P, Sodhi PS (1999b) On-farm seed priming in semi-arid agriculture: development and evaluation in maize, rice and chickpea in India using participatory methods. *Exp Agric* 35(1):15–29. <https://doi.org/10.1017/S0014479799001027>
- Harris D, Pathan AK, Gothkar P, Joshi A, Chivasa W, Nyamudeza P (2001a) On-farm seed priming: using participatory methods to revive and refine a key technology. *Agric Syst* 69(1–2):151–164. [https://doi.org/10.1016/S0308-521x\(01\)00023-3](https://doi.org/10.1016/S0308-521x(01)00023-3)
- Harris D, Raghuvanshi BS, Gangwar JS, Singh SC, Joshi KD, Rashid A, Hollington PA (2001b) Participatory evaluation by farmers of on-farm seed priming in wheat in India, Nepal and Pakistan. *Exp Agric* 37(3):403–415. <https://doi.org/10.1017/S0014479701003106>
- Harris D, Rashid A, Miraj G, Arif M, Shah H (2007b) ‘On-farm’ seed priming with zinc sulphate solution—a cost-effective way to increase the maize yields of resource-poor farmers. *Field Crop Res* 102(2):119–127. <https://doi.org/10.1016/j.fcr.2007.03.005>
- Harris D, Rashid A, Miraj G, Arif M, Yunas M (2008b) ‘On-farm’ seed priming with zinc in chickpea and wheat in Pakistan. *Plant Soil* 306(1–2):3–10. <https://doi.org/10.1007/s11104-007-9465-4>
- Iqbal M, Ashraf M (2005) Presowing seed treatment with cytokinins and its effect on growth, photosynthetic rate, ionic levels and yield of two wheat cultivars differing in salt tolerance. *J Integr Plant Biol* 47(11): 1315–1325. <https://doi.org/10.1111/j.1744-7909.2005.00163.x>
- Iqbal M, Ashraf M (2010) Changes in hormonal balance: a possible mechanism of pre-sowing chilling-induced salt tolerance in spring wheat. *J Agron Crop Sci* 196(6):440–454. <https://doi.org/10.1111/j.1439-037X.2010.00434.x>
- Islam F, Yasmeen T, Ali S, Ali B, Farooq MA, Gill RA (2015b) Priming-induced antioxidative responses in two wheat cultivars under saline stress. *Acta Physiol Plant* 37(8). <https://doi.org/10.1007/s11738-015-1897-5>
- Khanal N, Joshi KD, Harris D, Chand SP (2004) Effect of micronutrient loading, soil application, and foliar sprays of organic extracts on grain legumes and vegetable crops under marginal farmers’ conditions in Nepal. In: *Proceedings of Micronutrients in South and South East Asia*, Kathmandu, Nepal, pp 121–132
- Kumar A, Gangwar JS, Prasad SC, Harris D (2002) On-farm seed priming increases yield of direct-sown finger millet in India. *Int Sorg Mill*

- News1 43(1), 90–92 <https://doi.org/10.1017/CBO9781107415324.004>
- Mani JK, Singh R, Singh D, Rao VUM (2013) Variation in radiation use efficiency of wheat (*Triticum aestivum* L.) as influenced by thermal stress management strategies under late sown conditions. *J Agrometeorol* 15(2):138–141
- Marwat KB, Arif M, Khan MA (2007) Effect of tillage and zinc application methods on weeds and yield of maize. *Pak J Bot* 39(5):1583–1591
- Murungu FS, Chiduzza C, Nyamugafata P, Clark LJ, Whalley WR (2004b) Effect of on-farm seed priming on emergence, growth and yield of cotton and maize in a semi-arid area of Zimbabwe. *Exp Agric* 40. S0014479703001509, <https://doi.org/10.1017/S0014479703001509>
- Murungu FS, Chiduzza C, Nyamugafata P, Clark LJ, Whalley WR, Finch-Savage WE (2004d) Effects of ‘on-farm seed priming’ on consecutive daily sowing occasions on the emergence and growth of maize in semi-arid Zimbabwe. *Field Crop Res* 89(1):49–57. <https://doi.org/10.1016/j.fcr.2004.01.020>
- Murungu FS, Madanzi T (2010) Seed priming, genotype and sowing date effects on emergence, growth and yield of wheat in a tropical low altitude area of Zimbabwe. *Afr J Agric Res* 5(17):8
- Musa AM, Harris D, Johansen C, Kumar J (2001b) Short duration chickpea to replace fallow after aman rice: the role of on-farm seed priming in the high Barind tract of Bangladesh. *Exp Agric* 37(4):509–521
- Neamatollahi E, Bannayan M, Ghanbari A, Haydari M, Ahmadian A (2009) Does hydro and Osmo-priming improve fennel (*Foeniculum vulgare*) seeds germination and seedlings growth? *Not Bot Horti Agrobo* 37(2):190–194
- Ousman A, Aune JB (2011b) Effect of seed priming and micro-dosing of fertilizer on groundnut, sesame and cowpea in Western Sudan. *Exp Agric* 47(3):431–443. <https://doi.org/10.1017/S0014479711000068>
- Rashid A, Harris D, Hollington P, Ali S (2004a) On-farm seed priming reduces yield losses of mungbean (*Vigna radiata*) associated with mungbean yellow mosaic virus in the north west Frontier Province of Pakistan. *Crop Prot* 23(11):1119–1124. <https://doi.org/10.1016/j.cropro.2004.04.002>
- Rashid A, Harris D, Hollington PA, Rafiq M (2004b) Improving the yield of mungbean (*Vigna radiata*) in the north west Frontier Province of Pakistan using on-farm seed priming. *Exp Agric* 40(2):233–244. <https://doi.org/10.1017/S0014479703001546>
- Rashid A, Hollington PA, Harris D, Khan P (2006b) On-farm seed priming for barley on normal, saline and saline-sodic soils in north west Frontier Province, Pakistan. *Eur J Agron* 24(3):276–281. <https://doi.org/10.1016/j.eja.2005.10.006>
- Rehman H, Basra SMA, Farooq M, Ahmed N, Afzal I (2011a) Seed priming with CaCl₂ improves the stand establishment, yield and quality attributes in direct seeded rice (*Oryza sativa*). *Int J Agric Biol* 13(5):786–790
- Rehman HU, Basra SMA, Farooq M (2011b) Field appraisal of seed priming to improve the growth, yield, and quality of direct seeded rice. *Turk J Agric For* 35(4):357–365. <https://doi.org/10.3906/tar-1004-954>
- Virk DS, Chakraborty M, Ghosh J, Harris D (2006b) Participatory evaluation of horsegram (*Macrotyloma uniflorum*) varieties and their on-station responses to on-farm seed priming in eastern India. *Exp Agric* 42(4):411–425. <https://doi.org/10.1017/S0014479706003838>