

Geophysical Research Abstracts  
Vol. 15, EGU2013-1819-3, 2013  
EGU General Assembly 2013  
© Author(s) 2013. CC Attribution 3.0 License.



## Vesta in the Light of Dawn

Christopher Russell (1), Carol Raymond (2), Harry McSween (3), Ralf Jaumann (4), Andreas Nathues (5), Maria DeSanctis (6), Tom Prettyman (7), Simone Marchi (8), Nico Schmedemann (9), Diego Turrini (6), Jennifer Scully (1), Martin Hoffman (10), and Debra Buczowski (11)

(1) University of California, Los Angeles, Institute of geophysics, Earth and Space Science, Los Angeles, United States (ctrussell@igpp.ucla.edu, 1-310-206-3051), (2) Jet Propulsion Laboratory, California Institute of Technology, Pasadena CA 91109-8099 USA, (3) University of Tennessee, Department of Earth and Planetary Sciences, Knoxville, TN 37996-1410, USA, (4) Deutsches Zentrum für Luft- und Raumfahrt, Institute of Planetary Research, Berlin, Germany, (5) Max Planck Institut für Sonnensystemforschung, Katlenburg-Lindau, Germany, (6) Istituto di Astrofisica e Planetologia Spaziali, Istituto Nazionale di Astrofisica, Rome, Italy, (7) Planetary Science Institute, Tucson, AZ 85719, USA, (8) NASA Lunar Science Institute, Center for Lunar Origin and Evolution, Boulder, CO 80302 USA, (9) Freie Universität, Berlin, Department of Earth Sciences, Institute of Geosciences, Planetary Sciences and Remote Sensing, Berlin, Germany, (10) NASA Dryden Flight Research Center, Edwards, CA 93523-0273 USA, (11) Johns Hopkins University Applied Physics Laboratory, Laurel, MD 20723-6099 USA

Dawn completed its observation of Vesta in September 2012, after nearly 14 months in orbit. What Dawn discovered ranged from the anticipated to the totally unexpected. Vesta is an intact, almost complete, basaltic protoplanet, with the properties inferred from the study of the HED meteorites. It has a pyroxene crust consisting of an outer eucritic component and an inner diogenitic layer overlying an olivine mantle. The fact that the eucritic crust is still quite extensive places important limits on the formation of Jupiter and its role in the evolution of the main belt. The crust in the southern hemisphere has been strongly excavated by the two events that formed the Veneneia and Rheasilvia basins. Most of the northern crust is older than the southern basins and the cratering record is still in production down to about 2 km crater size on ~3.8 Ga old surfaces. The cratering record is consistent with some of the models of the evolution of the asteroid belt and planetary migration theory and puts important constraints on the history evolution of the solar system. Vesta has a 100 km radius iron core and this may have aided Vesta's survival through its heavy bombardment early in its history. The biggest surprises have come from a detailed high-resolution study of the surface: the discovery of pits as opposed to craters; the different styles of gully formation; and discovery of odd crater morphologies.