

Quantum many-body systems out of equilibrium

J. Eisert, M. Friesdorf, and C. Gogolin

Dahlem Center for Complex Quantum Systems, Freie Universität Berlin, 14195 Berlin, Germany

Closed quantum many-body systems out of equilibrium pose several long-standing problems in physics. Recent years have seen a tremendous progress in approaching these questions, not least due to experiments with cold atoms and trapped ions in instances of quantum simulations. This article provides an overview on the progress in understanding dynamical equilibration and thermalisation of closed quantum many-body systems out of equilibrium due to quenches, ramps and periodic driving. It also addresses topics such as the eigenstate thermalisation hypothesis, typicality, transport, many-body localisation, universality near phase transitions, and prospects for quantum simulations.

How do closed quantum many-body systems out of equilibrium eventually equilibrate? How and in precisely what way can such quantum systems with many degrees of freedom thermalise? It was established early on how to capture equilibrium states corresponding to some temperature in the framework of quantum statistical mechanics, but it seems much less clear how such states are eventually reached via the local dynamics following microscopic laws. This dichotomy gives rise to some tension between a microscopic description and one in terms of the familiar ensembles of statistical mechanics. It also leads to the question of how the actual dynamics of quantum phase transitions can be grasped. Asking questions of this type has a long tradition, but many problems have remained largely unresolved. It was only relatively recently that they moved back into the focus of attention. This development is driven by the availability of experiments that allow to probe such questions under controlled conditions, but also by new developments in theoretical physics. The present review aims at providing a first impression of important progress in this rapidly developing field of research, it emphasises promising perspectives, and collects a number of key references. Given its brevity, it can by no means be faithful to all developments or cover the whole literature of the field.

Local quantum many-body systems

The physical systems considered here are *local quantum many-body systems* with finite-range interactions, ubiquitous in condensed-matter physics. They can be described by some interaction graph, with physical constituents at the vertices. The system is governed by a Hamiltonian that can be written as

$$H = \sum_j h_j. \quad (1)$$

Most commonly considered are nearest-neighbour interactions, where each h_j acts non-trivially on finitely many sites. Paradigmatic examples of this type are spin lattice systems,

such as the Ising model, as well as bosonic or fermionic lattice systems, prominently Fermi- and Bose-Hubbard models. In the latter case, the Hamiltonian takes the form

$$H = J \sum_{\langle j,k \rangle} (b_j^\dagger b_k + b_k^\dagger b_j) + U \sum_j n_j (n_j - 1), \quad (2)$$

with $n_j = b_j^\dagger b_j$, b_j denoting the bosonic annihilation operator at site j , the parameters U, J being real and the sum $\langle j, k \rangle$ running over nearest neighbours only. Common variants of this are models with other short range interactions, or long range interactions with a strength decaying polynomially in the distance.

Equilibration after quenches

Investigating the fundamental connection between thermodynamics and the dynamics of closed quantum systems has a long tradition¹ but has also been a key topic in recent years. A very clean setting for non-equilibrium dynamics is the one emerging from a sudden *global quench*^{2-9,11-16}: Initially, the system is in a state ρ_0 , which could be the ground state of a local Hamiltonian. Then one quickly alters the system's parameters globally and considers the many-body unitary time evolution under some local Hamiltonian H . Of specific interest are expectation values of observables A at later times

$$\langle A(t) \rangle = \text{tr}(e^{-itH} \rho_0 e^{itH} A). \quad (3)$$

A main question is to what extent and for what times the situation can be described by a suitable equilibrium ensemble. Notably, the dynamics is time reversal invariant and for finite systems recurrent, so a priori it seems far from clear how and in what sense equilibrium can be reached dynamically.

A first important insight is that such systems generically indeed relax and equilibrate in the following sense: Even though the dynamics is entirely unitary, following transient non-equilibrium dynamics, it is expectation values $\langle A(t) \rangle$ of *many observables* that equilibrate. This is specifically true for

local observables which are supported only on a small number of sites. The apparent long time equilibrium state has to be equal to the time average

$$\omega = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt e^{-itH} \rho_0 e^{itH}. \quad (4)$$

This state is the *maximum entropy state*, holding all constants of motion fixed¹⁷. In case of non-degenerate eigenvalues of the Hamiltonian, this ensemble is also called the *diagonal ensemble*¹⁸. This feature is reminiscent of a dynamical Jaynes' principle: A many-body system is pushed out of equilibrium and follows unitary dynamics. Yet, for most times, the systems appears as if it had equilibrated to a maximum entropy state.

The general expectation that many-body systems equilibrate can be made rigorous in a number of senses: For some free models, specifically for the case of the integrable non-interacting limit of the Bose-Hubbard model, equilibration is proven to be true in the strong sense of equilibration during intervals: Here, one can show for local observables A that $|\langle A(t) \rangle - \text{tr}(\omega A)|$ is arbitrarily small after a known initial relaxation time, and remains so for a time that grows linearly with the system size^{6,19}. One merely has to assume that the otherwise arbitrary initial state has suitable polynomially decaying correlations. At the heart of such a rigorous argument are Lieb-Robinson bounds as well as non-commutative central limit theorems.

If no locality structure is available or is made use of, one can still show equilibration on average for all Hamiltonians that have non-degenerate energy gaps: One can bound the time average $\mathbb{E}_{t \in [0, \infty)} |\langle A(t) \rangle - \text{tr}(\omega A)|$ by a quantity that is usually exponentially small in the system size^{20,21}. This means that out of equilibrium quantum systems appear relaxed for overwhelmingly most times, even if by that statement alone no information on time scales can be deduced. More refined bounds have been derived, also providing some information about the relevant time scales of equilibration^{22,23}, but they are far off from those observed in numerical simulations^{4,7,12,16} and experiments^{14,24,25}. These findings are also related to further efforts of describing the size of the time fluctuations after relaxation^{26,27}.

By no means are sudden global quenches the only relevant setting of quantum many-body systems out of equilibrium. Similarly important are scenarios of *local quenches*^{28,29}, where not the entire system is uniformly modified, but rather the system is locally suddenly driven out of equilibrium by a change limited to a bounded number of lattice sites. *Ramps*^{12,30-33}, during which the Hamiltonian is changed according to some schedule, are of interest in studies of the dynamics of quantum phase transitions. The response of a system to a sudden alteration of its geometry can be studied by *geometric quenches*^{5,34,35}. *Time-dependent periodic driving* is discussed in a separate section. In this brief review, closed many-body systems are in the focus of attention. It should be clear that specifically *pump-probe* type settings³⁶ generate an out of equilibrium situation that can be used as a diagnostic tool in the condensed-matter context, giving rise to an inter-

esting research topic in its own right.

Thermalisation

The success of thermodynamics in describing large scale systems indicates that often the equilibrium values can be described by few parameters such as the global temperature and particle number. We use the term *thermalisation* to refer to the equilibration towards a state ω that is in a suitable sense close to being indistinguishable from a thermal equilibrium state proportional to $e^{-\beta H}$ for some inverse temperature $\beta > 0$. There are several mechanisms that can lead to thermalisation in this sense: The first is based on the *eigenstate thermalisation hypothesis* (ETH)^{9,37-40}. It conjectures that sufficiently complex quantum systems have eigenstates that — for physically relevant observables such as local ones — are practically indistinguishable from thermal states with the same average energy. The ETH and its breakdown in integrable systems have recently been investigated extensively, mostly with numerical methods. Supporting evidence for it has been collected in a plethora of models^{9,12,41-43}.

Without invoking the ETH one can still rigorously show thermalisation in weakly interacting systems under stronger conditions on the initial state⁴⁴. In addition, dynamical thermalisation of local expectation values to those in the global thermal state⁴⁵ has been shown under stronger conditions, most notably translation invariance and the existence of a unique Gibbs state. The notion of *relative thermalisation*⁴⁶ focuses on the decoupling of a subsystem from a reference system. So far, there is no compelling proof that globally quenched non-integrable systems dynamically thermalise.

These dynamical approaches are complemented by works based on *typicality* arguments^{47,48}. Here, rather than following the dynamical evolution of a system, one tries to justify the applicability of ensembles by showing that most states drawn randomly according to some measure have the same physical properties as some appropriate ensemble such as the canonical one.

If a quantum system is integrable⁴⁹ in the sense that it has local conserved quantities, one should not expect the system to thermalise: These constants of motion prohibit full thermalisation to the canonical ensemble. One can still expect the system to equilibrate to the maximum entropy state given these locally conserved quantities^{15,17}, a so called *generalized Gibbs ensemble*^{6,18}. Effective thermalisation has been put into context with unitary quantum dynamics under conditions of classical chaos⁵⁰.

Several quantum many-body systems exhibit some instance of *pre-thermalisation*: This term refers to any apparent equilibration to a meta-stable state on a short time scale, before — on a long time scale — the system relaxes to a state indistinguishable from a genuine thermal state. Examples are almost integrable systems^{51,52} and continuous systems of coupled Bose-Einstein condensates²⁵.

“Light cones” and entanglement dynamics

An important stepping stone for a better understanding of the non-equilibrium dynamics of local Hamiltonian models is the insight that *Lieb-Robinson bounds*⁵³⁻⁵⁵ limit the speed of information propagation in such systems. There are several ways of stating such bounds: One is to say that for any two observables A and B

$$\| [A(t), B] \| \leq c \|A\| \|B\| \min\{|A|, |B|\} e^{-\mu(d(A,B)-v|t|)}, \quad (5)$$

where $d(A, B)$ is the distance between the support of the observables, $|A|$ and $|B|$ the size of their supports, $v \geq 0$ takes the role of a group velocity, and $c, \mu > 0$ are constants. That is to say, information propagation outside a “light-cone” is exponentially suppressed. It immediately follows that correlation functions can significantly grow in time only inside the light cone⁵⁵. While these bounds are not necessarily tight, the picture that the local dynamics leads to excitations traveling through the system with velocity v offers a lot of explanatory potential for non-equilibrium dynamics^{3,6,7,19,24,56,57}. Such light-cone dynamics has been experimentally observed in optical lattice systems using novel tools that allow for single-site addressing⁵⁸ and in continuous systems of cold atoms²⁵.

The light-cone-like propagation of information has implications on *entanglement dynamics*. Many-body states are said to satisfy an *area law* for the entanglement entropy, if for a subset of sites the latter scales only like the boundary “area” of the subset rather than its “volume”^{59,60}. It has been shown that if initially such a law holds, this will remain to be true for longer times^{55,61}. Yet, at the same time, the pre-factor of this area law is expected to grow exponentially⁶². This is the ultimate reason why numerical *tensor network methods*^{63,64}, such as the *density-matrix renormalisation group method*⁶⁵, can simulate out of equilibrium dynamics efficiently for short times on a classical computer, while long times are not accessible. This is a feature shared by other powerful numerical methods as well, such as *Monte-Carlo*⁶⁶ and *dynamical mean field theory (DMFT)*⁶⁷. This is in line with complexity-theoretic arguments that provide evidence that generic quantum long-time dynamics cannot be efficiently classically simulated⁶⁸. If interactions are not strictly local, modified Lieb-Robinson bounds still hold⁵³, giving rise to a rich phenomenology which has been experimentally explored in systems of trapped ions exhibiting long-range order^{57,69}.

Transport

Understanding electronic transport is one of the main motivations that started the field of condensed matter physics and gave rise to a large body of work trying to accurately capture the conductivity properties of these systems. Most relevant for the purpose of this review are the transport properties of simple paradigmatic models such as the Heisenberg spin chain or Hubbard-type models. In these models, transport is expected to lead to equilibration, as it allows the system to locally forget its precise initial configuration^{6,70}. In spin chains, transport typically refers to a spreading of magnetised domains⁷¹. In

optical lattices one commonly studies the transport of particles starting from an initially trapped situation^{72,73}, or of quasi-particles^{57,58}. Two of the key aims are to distinguish between diffusive and ballistic transport and to understand the limitations of *linear response theory*⁷¹. While simple, integrable models can be solved exactly and thus provide an excellent benchmark, our understanding of transport in realistic setups heavily relies on numerical tools. Tools explicitly storing the full wave function in the memory, such as exact diagonalisation, are limited to comparably small systems. Tensor network tools such t-DMRG, TEBD, and variants thereof^{63,65} allow to parametrize the state more efficiently and are accurate up to machine precision as long as entanglement entropies are sufficiently small⁷⁴. Thus, for short and intermediate times these tools allow for an investigation of transport and equilibration in systems with several hundred sites^{14,16,71,73}.

Absence of thermalisation and many-body localisation

Crucial for transport properties of materials is the influence of irregularities such as defects. The impact of random potentials on particle mobility has thus been the subject of intensive study over the last century, culminating in the famous result of Anderson, showing full localisation both of the eigenfunctions as well as of the dynamics⁷⁵ for single particle *disordered models*. In the case of non-interacting fermions in one dimension or equivalently, via the Jordan-Wigner transformation, for free spin chains, Anderson’s result directly gives strong localisation of all fermionic modes. This leads to the entanglement entropy remaining upper bounded from above⁷⁶ and a vanishing Lieb-Robinson velocity⁷⁷. These systems thus fail to serve as their own heat bath and transport of any kind is completely suppressed. For interacting many-body systems, the effects of disorder are still much less clear, despite great efforts⁷⁸. Interestingly, both the support of time evolved local observables as well as the entanglement entropy are found to grow logarithmically in the interacting case and are thus unbounded⁷⁶, in stark contrast to the non-interacting case. Dynamical aspects of many-body localisation are expected to go along with localisation of the eigenstates. In several recent studies localisation effects due to disorder have been associated with a lack of entanglement in many energy eigenstates^{17,79,80} or alternatively only in those below a so called *mobility edge*^{77,78}. This could lead to a phase transition between a disordered insulator with exactly zero conductivity and a conducting phase⁷⁸. The localisation of eigenstates leads to a clear violation of the ETH and connects to an absence of thermalisation effect for localising many-body systems and the existence of local constants of motion in these models^{17,79}. While these signatures of many-body localisation have been extensively explored, a comprehensive definition of many-body localisation is still lacking.

Dynamics of quantum phase transitions

Exploring the dynamical signatures of phase transitions is an interesting problem in its own right. It connects to the important question of how slow an experimental ramp needs to

be to avoid the creation of defects when preparing a quantum phase and whether this is at all possible in the thermodynamic limit⁸¹. A key theoretical model for these transitions, based on the adiabatic theorem, has been provided by Kibble and Zurek³⁰ for the important case of moving from a gapped phase into criticality. Based solely on the critical exponents of the model, this strikingly simple formalism predicts the number of introduced defects as a function of the velocity of the ramp. It has originally been developed for thermal transitions, where it has also been experimentally tested^{82–84}. For quantum phase transitions such a scaling can be derived in the limit of infinitely slow ramps, based on universality arguments together with adiabatic perturbation theory⁸¹ and seems to be capable of accurately describing the dynamics of quantum phase transitions when crossing a single critical point in some models³⁰. For strongly correlated systems in realistic experimental settings, the dynamics of quantum phase transitions is more elusive and there are indications that the complexity of the dynamics cannot fully be captured by employing simple scaling arguments based only on the critical exponents of the model^{31,85}. The transition out of criticality has also been investigated, both numerically⁸⁶ and experimentally³². Despite these first promising results, the dynamics of quantum phase transitions and how to capture them in terms of simple scaling laws is still a largely open problem.

Quantum simulations

When investigating the non-equilibrium behaviour of large-scale many-body systems, analytical and numerical tools quickly reach their limits, especially for lattice models in more than one dimension. A particularly exciting perspective arises from the idea that controlled quantum many-body systems constitute instances of *analogue quantum simulators*^{14,31,87,88} overcoming these limitations. These devices mimic natural interacting quantum many-body systems by reconstructing their Hamiltonian, but now under precisely controlled conditions. This is in contrast to *digital quantum simulators*, anticipated devices that approximate quantum dynamics by suitable quantum circuits. Such analogue systems enable one to investigate some of the questions central to the research summarised in this review.

Ultra-cold atoms in optical lattices provide a very promising platform for quantum simulators in this sense⁸⁹. With present technology they can be used to investigate dynamics under Bose- and Fermi-Hubbard Hamiltonians^{2,14,31,32,58,72,73,85,90–92} as well as for emulating spin chains⁹³. Due to the flexibility of optical lattice systems, e.g., by making use of Feshbach resonances, periodic driving, and superlattices, even sophisticated quantum simulations are conceivable, e.g., the simulation of *lattice gauge theories*⁹⁴. Continuous optical setups allow for the investigation of out-of-equilibrium behaviour of quantum fields⁹⁵ and have successfully been employed to investigate (pre-)thermalisation²⁵ and out of equilibrium dynamics not leading to equilibration on relevant time scales⁹⁶.

Another promising architecture is provided by *ion chains*⁹⁷, allowing for the precise control of the individual

constituents^{83,98}. Due to the involved Coulomb interaction, these systems are, among many other applications, well suited to explore the validity of Lieb-Robinson bounds and their breakdown for long-range interactions^{57,69}. *Photonic architectures* as well as arrays of *superconducting qubits* offer further promising platforms. While the precise *computational complexity* of analogue simulators has not been identified yet, and while it is unclear to what extent genuinely reliable analogue simulation is possible in the absence of fully-fledged quantum error correction, there is evidence that such analogue simulators have the potential of outperforming classical computers. Quantum many-body dynamics can already be probed in experimental settings under precisely controlled conditions for longer times than can be kept track of using state-of-the-art tensor network methods on modern classical supercomputers¹⁴.

Periodically driven systems

Periodically driving a well controlled quantum system opens up new vistas for quantum simulations. Consider a system with a Hamiltonian that satisfies $H(t) = H(t + \tau)$ for all times t for some period $\tau > 0$. The eigenstates of such systems are usually constantly changing in time, but the long time dynamics can be inferred from their *Floquet operator*^{99,100} $U(\tau)$, which is the time evolution operator over one period of driving. The Floquet operator gives rise to an effective Hamiltonian H_{eff} via

$$U(\tau) = e^{-i\tau H_{\text{eff}}}, \quad (6)$$

and the time evolution of the driven system can be seen as a stroboscopic simulation^{101,102} of the time evolution under H_{eff} . The effective Hamiltonian can be tuned by choosing an appropriate driving schedule with great flexibility concerning the model parameters. This allows the study of topological effects¹⁰² in driven systems and can be used to engineer gauge fields¹⁰¹. This has led to an experimental realisation of the *topological Haldane model* in a periodically modulated optical honeycomb lattice¹⁰³ as well as to experimental setups featuring *Hofstadter bands*¹⁰⁴ and *spin-orbit coupling*¹⁰⁵ in optical lattice architectures¹⁰⁶. The synchronisation of the motion of periodically driven systems with the drive allows the description of the long time behavior by a *periodic ensemble*. For free systems it turns out to be given by a maximum entropy ensemble given the constants of motion¹⁰⁷ — reminiscent of Jaynes' principle for systems in equilibrium and the generalised Gibbs ensemble emerging during equilibration after a quench.

Perspectives for open and closed systems

The research field of quantum many-body systems out of equilibrium is rapidly developing and is taking new and unexpected directions at a remarkably high rate. This is partially driven by the impetus the field is subjected to through novel experimental developments. In optical lattice systems, *single-site addressing*^{32,58,108} opens up new possibilities for

probing the out of equilibrium dynamics; settings with periodic driving now allow for the study of topologically non-trivial situations¹⁰² and to develop a deeper understanding of particle physics and condensed matter systems¹⁰¹. *Continuous systems* can be controlled with ever increasing precision^{25,95}, suggesting that such systems can serve as a platform for probing even effects of *gravity*, such as the Unruh effect^{109,110}. Hybrid architectures, such as cold atoms in optical lattices inside an optical cavity, constitute a particularly promising avenue for experimental research¹¹¹. Crucial for further progress will be a deeper understanding of the computational capability of analogue quantum simulators and how their correct functioning can be certified in regimes that are classically not addressable^{14,31,88,90}.

At the same time, new theoretical developments keep the subject exciting, with *quantum information theory* providing a fresh point of view and new methods. *Entanglement theory* can deliver important new insights specifically for the study of information propagation^{56,59–61}. *Quantum thermodynamics*¹¹² — exploring the ultimate limits of thermodynamic processes in the quantum regime — is now seeking contact with this field¹¹³, with interesting perspectives opening up. The *AdS-cft-correspondence* also contributes a novel tool to study strongly correlated quantum systems out of equilibrium¹¹⁴.

In spite of the great progress, many questions are still wide open, even very basic ones. While it is understood under which conditions many-body systems *equilibrate*, it is far from clear on what *time scales* they do so. The main reason for this is that the role of *locality* for equilibration and thermalisation is, at least from a rigorous perspective, far from being understood. Steps in this direction have been taken only

very recently^{45,53–55,61,115}. The conditions for *thermalisation* are less clear and it is questionable whether (non-)integrability is able to capture this phenomenon in all of its facets^{17,49}. The problem of *many-body localisation* has recently moved in the focus of attention^{79,80}, but it seems fair to say that in the light of the various pictures of the phenomenon in the literature, a unifying framework is still missing.

A research area that deserves special attention is based on the idea that noise and dissipation are not necessarily detrimental to attempts to prepare and preserve interesting states of matter. In fact such preparations can be facilitated by suitably *engineered local Markovian noise*¹¹⁶ that can be rigorously captured in terms of a Liouvillian. In this mindset, an open system quantum simulator with trapped ions has been realised¹¹⁷, as well as one based of cold atoms controlled with localized dissipation¹¹⁸. In principle, even symmetry-protected topologically ordered states can be prepared by purely dissipative local Liouvillian dynamics¹¹⁹.

It goes without saying that this perspectives article merely touches the surface of this exciting and rapidly developing field. Quantum many-body systems out of equilibrium allow us to probe long-standing questions in the foundations of statistical mechanics, relate to open questions in condensed-matter theory, and provide a promising platform for analogue quantum simulators outperforming classical computers.

We would like to thank many colleagues for numerous discussions over the years, and E. Bergholtz, E. T. Campbell, A. del Campo, J.-S. Caux, F. Essler, T. Farrelly, M. Rigol, and R. Moessner for comments on the manuscript, and the EU (RAQUEL, SIQS), the ERC, the BMBF, and the Studienstiftung des Deutschen Volkes for support.

-
- ¹ v. Neumann, J. Beweis des Ergodensatzes und des H-Theorems in der neuen Mechanik. *Z. Phys.* **57**, 30 (1929).
 - ² Greiner, M., Mandel, O., Hänsch, T. W. & Bloch, I. Collapse and revival of the matter wave field of a Bose-Einstein condensate. *Nature* **419**, 51 (2002).
 - ³ Calabrese, P. & Cardy, J. Time dependence of correlation functions following a quantum quench. *Phys. Rev. Lett.* **96**, 136801 (2006).
 - ⁴ Kollath, C., Läuchli, A. & Altman, E. Quench dynamics and non equilibrium phase diagram of the Bose-Hubbard model. *Phys. Rev. Lett.* **98**, 180601 (2007).
 - ⁵ Rigol, M., Dunjko, V., Yurovsky, V. & Olshanii, M. Relaxation in a completely integrable many-body quantum system: An ab initio study of the dynamics of the highly excited states of 1D lattice hard-core bosons. *Phys. Rev. Lett.* **98**, 050405 (2007).
 - ⁶ Cramer, M., Dawson, C. M., Eisert, J. & Osborne, T. J. Quenching, relaxation, and a central limit theorem for quantum lattice systems. *Phys. Rev. Lett.* **100**, 030602 (2008).
 - ⁷ Flesch, A., Cramer, M., McCulloch, I. P., Schollwöck, U. & Eisert, J. Probing local relaxation of cold atoms in optical superlattices. *Phys. Rev. A* **78**, 033608 (2008).
 - ⁸ Moeckel, M. & Kehrein, S. Interaction quench in the Hubbard model. *Phys. Rev. Lett.* **100**, 175702 (2008).
 - ⁹ Rigol, M., Dunjko, V. & Olshanii, M. Thermalization and its mechanism for generic isolated quantum systems. *Nature* **452**, 854 (2008).
 - ¹⁰ Barthel, T., & Schollwöck, U. Dephasing and the steady state in quantum many-particle systems. *Phys. Rev. Lett.* **100**, 100601 (2008).
 - ¹¹ Manmana, S. R., Wessel, S., Noack, R. M. & Muramatsu, A. Time evolution of correlations in strongly interacting fermions after a quantum quench. *Phys. Rev. B* **79**, 155104 (2009).
 - ¹² Polkovnikov, A., Sengupta, K., Silva, A. & Vengalattore, M. Non-equilibrium dynamics of closed interacting quantum systems. *Rev. Mod. Phys.* **83**, 863 (2011).
 - ¹³ Calabrese, P., Essler, F. H. L. & Fagotti, M. Quantum quench in the transverse-field Ising chain. *Phys. Rev. Lett.* **106**, 227203 (2011).
 - ¹⁴ Trotzky, S. *et al.* Probing the relaxation towards equilibrium in an isolated strongly correlated one-dimensional Bose gas. *Nature Phys.* **8**, 325 (2012).
 - ¹⁵ Caux, J.-S. & Essler, F. H. L. Time evolution of local observables after quenching to an integrable model. *Phys. Rev. Lett.* **110**, 257203 (2013).
 - ¹⁶ Barmettler, P., Punk, M., Gritsev, V., Demler, E. & Altman, E. Relaxation of antiferromagnetic order in spin- $\frac{1}{2}$ chains following a quantum quench. *Phys. Rev. Lett.* **102**, 130603 (2009).
 - ¹⁷ Gogolin, C., Mueller, M. P. & Eisert, J. Absence of thermalization in non-integrable systems. *Phys. Rev. Lett.* **106**, 040401 (2011).
 - ¹⁸ Cassidy, A. C., Clark, C. W. & Rigol, M. Generalized thermalization in an integrable lattice system. *Phys. Rev. Lett.* **106**, 140405 (2011).

- (2011).
- 19 Cramer, M. & Eisert, J. A quantum central limit theorem for non-equilibrium systems: Exact local relaxation of correlated states. *New J. Phys.* **12**, 055020 (2010).
 - 20 Reimann, P. Foundation of statistical mechanics under experimentally realistic conditions. *Phys. Rev. Lett.* **101**, 190403 (2008).
 - 21 Linden, N., Popescu, S., Short, A. J. & Winter, A. Quantum mechanical evolution towards thermal equilibrium. *Phys. Rev. E* **79**, 061103 (2009).
 - 22 Reimann, P. & Kastner, M. Equilibration of isolated macroscopic quantum systems. *New J. Phys.* **14**, 043020 (2012).
 - 23 Short, A. J. & Farrelly, T. C. Quantum equilibration in finite time. *New J. Phys.* **14**, 013063 (2012).
 - 24 Langen, T., Geiger, R., Kuhnert, M., Rauer, B. & Schmiedmayer, J. Local emergence of thermal correlations in an isolated quantum many-body system. *Nature Phys.* **9**, 640 (2013).
 - 25 Geiger, R., Langen, T., Mazets, I. & Schmiedmayer, J. Local relaxation and light-cone-like propagation of correlations in a trapped one-dimensional Bose gas. *New J. Phys.* **16**, 053034 (2014).
 - 26 Torres-Herrera, E. J., Kollmar, D. & Santos, L. F. Relaxation and thermalization of isolated many-body quantum systems. ArXiv:1403.6481.
 - 27 Venuti, L. C. & Zanardi, P. Universal time-fluctuations in near-critical out-of-equilibrium quantum dynamics. *Phys. Rev. E* **89**, 022101 (2014).
 - 28 Torres-Herrera, E. J. & Santos, L. F. Local quenches with global effects in interacting quantum systems. *Phys. Rev. E* **89**, 062110 (2014).
 - 29 Stephan, J.-M. & Dubail, J. Local quantum quenches in critical one-dimensional systems: Entanglement, the Loschmidt echo, and light-cone effects. *J. Stat. Mech.* P08019 (2011).
 - 30 del Campo, A. & Zurek, W. H. Universality of phase transition dynamics: Topological defects from symmetry breaking. *Int. J. Mod. Phys. A* **29**, 1430018 (2014).
 - 31 Braun, S. *et al.* Emergence of coherence and the dynamics of quantum phase transitions. ArXiv:1403.7199.
 - 32 Bakr, W. S. *et al.* Probing the superfluid-to-Mott insulator transition at the single-atom level. *Science* **329**, 547 (2010).
 - 33 Dóra, B., Haque, M. & Zaránd, G. Crossover from adiabatic to sudden interaction quench in a Luttinger liquid. *Phys. Rev. Lett.* **106**, 156406 (2011).
 - 34 Mossel, J., Palacios, G. & Caux, J.-S. Geometric quenches in quantum integrable systems. *J. Stat. Mech.* L09001 (2010).
 - 35 Alba, V. & Heidrich-Meisner, F. Entanglement spreading after a geometric quench in quantum spin chains. ArXiv:1402.2299.
 - 36 Bovensiepen, U., Petek, H. & Wolf, M. (eds.) *Dynamics at solid state surfaces and interfaces* (Wiley-VCH, Weinheim, Germany, 2010).
 - 37 Deutsch, J. M. Quantum statistical mechanics in a closed system. *Phys. Rev. A* **43**, 2046 (1991).
 - 38 Srednicki, M. Chaos and quantum thermalization. *Phys. Rev. E* **50**, 888 (1994).
 - 39 Tasaki, H. From quantum dynamics to the canonical distribution: General picture and a rigorous example. *Phys. Rev. Lett.* **80**, 1373 (1998).
 - 40 Rigol, M. & Srednicki, M. Alternatives to eigenstate thermalization. *Phys. Rev. Lett.* **108**, 110601 (2012).
 - 41 Rigol, M. Breakdown of thermalization in finite one-dimensional systems. *Phys. Rev. Lett.* **103**, 100403 (2009).
 - 42 Steingeweg, R., Khodja, A., Niemeyer, H., Gogolin, C. & Gemmer, J. Pushing the limits of the eigenstate thermalization hypothesis towards mesoscopic quantum systems. *Phys. Rev. Lett.* **112**, 130403 (2014).
 - 43 Beugeling, W., Moessner, R. & Haque, M. Finite-size scaling of eigenstate thermalization. *Phys. Rev. E* **89**, 042112 (2014).
 - 44 Riera, A., Gogolin, C. & Eisert, J. Thermalization in nature and on a quantum computer. *Phys. Rev. Lett.* **108**, 080402 (2012).
 - 45 Mueller, M. P., Adlam, E., Masanes, L. & Wiebe, N. Thermalization and canonical typicality in translation-invariant quantum lattice systems. ArXiv:1312.7420.
 - 46 del Rio, L., Hutter, A., Renner, R. & Wehner, S. Relative thermalization (2014). ArXiv:1401.7997.
 - 47 Goldstein, S. Canonical typicality. *Phys. Rev. Lett.* **96**, 050403 (2006).
 - 48 Popescu, S., Short, A. J. & Winter, A. The foundations of statistical mechanics from entanglement: Individual states vs. averages. *Nature Phys.* **2**, 754 (2006).
 - 49 Caux, J.-S. & Mossel, J. Remarks on the notion of quantum integrability. *J. Stat. Mech.* P02023 (2011).
 - 50 Altland, A. & Haake, F. Quantum chaos and effective thermalization. *Phys. Rev. Lett.* **108**, 073601 (2012).
 - 51 Marcuzzi, M., Marino, J., Gambassi, A. & Silva, A. Prethermalization in a non-integrable quantum spin chain after a quench. *Phys. Rev. Lett.* **111**, 197203 (2013).
 - 52 Essler, F. H. L., Kehrein, S., Manmana, S. R., Robinson, N. J., Quench dynamics in a model with tuneable integrability breaking. *Phys. Rev. B* **89**, 165104 (2014).
 - 53 Hastings, M. B. & Koma, T. Spectral gap and exponential decay of correlations. *Commun. Math. Phys.* **265**, 781 (2006).
 - 54 Sidoravicius, V. (ed.). *Locality estimates for quantum spin systems* (Springer Verlag, 2009).
 - 55 Bravyi, S., Hastings, M. B. & Verstraete, F. Lieb-Robinson bounds and the generation of correlations and topological quantum order. *Phys. Rev. Lett.* **97**, 050401 (2006).
 - 56 Calabrese, P. & Cardy, J. Evolution of entanglement entropy in one-dimensional systems. *J. Stat. Mech.* P04010 (2005).
 - 57 Jurcevic, P. *et al.* Quasiparticle engineering and entanglement propagation in a quantum many-body system. *Nature* **511**, 202 (2014).
 - 58 Cheneau, M. *et al.* Light-cone-like spreading of correlations in a quantum many-body system. *Nature* **481**, 484 (2012).
 - 59 Hastings, M. B. An area law for one-dimensional quantum systems. *J. Stat. Mech.* P08024 (2007).
 - 60 Eisert, J., Cramer, M. & Plenio, M. B. Area laws for the entanglement entropy. *Rev. Mod. Phys.* **82**, 277 (2010).
 - 61 Eisert, J. & Osborne, T. J. General entanglement scaling laws from time evolution. *Phys. Rev. Lett.* **97**, 150404 (2006).
 - 62 Schuch, N., Wolf, M. M., Vollbrecht, K. G. H. & Cirac, J. I. On entropy growth and the hardness of simulating time evolution. *New J. Phys.* **10**, 033032 (2008).
 - 63 Orus, R. A practical introduction to tensor networks: Matrix product states and projected entangled pair states. *Ann. Phys.* **349**, 117 (2014).
 - 64 Banuls, M. C., Hastings, M. B., Verstraete, F. & Cirac, J. I. Matrix product states for dynamical simulation of infinite chains. *Phys. Rev. Lett.* **102**, 240603 (2009).
 - 65 Schollwöck, U. The density-matrix renormalization group in the age of matrix product states. *Ann. Phys.* **326**, 96 (2011).
 - 66 Troyer, M., Alet, F., Trebst, S. & Wessel, S. Non-local updates for quantum Monte Carlo simulations. *AIP Conf. Proc.* **690**, 156 (2003).
 - 67 Aoki, H. *et al.* Non-equilibrium dynamical mean-field theory and its applications. *Rev. Mod. Phys.* **86**, 779 (2014).
 - 68 Nagaj, D. Fast universal quantum computation with railroad-switch local Hamiltonians. *J. Math. Phys.* **51**, 62201 (2010).
 - 69 Richerme, P. *et al.* Non-local propagation of correlations in long-

- range interacting quantum systems (2014). ArXiv:1401.5088.
- ⁷⁰ Gogolin, C. & Eisert, J. Pure state statistical mechanics. *In preparation* (2014).
- ⁷¹ Langer, S., Heidrich-Meisner, F., Gemmer, J., McCulloch, I. P. & Schollwöck, U. Real-time study of diffusive and ballistic transport in spin- $\frac{1}{2}$ chains using the adaptive time-dependent density matrix renormalization group method. *Phys. Rev. B* **79**, 214409 (2009).
- ⁷² Schneider, U. *et al.* Fermionic transport and out-of-equilibrium dynamics in a homogeneous Hubbard model with ultra-cold atoms. *Nat. Phys.* **8**, 213 (2012).
- ⁷³ Ronzheimer, J. P. *et al.* Expansion dynamics of interacting bosons in homogeneous lattices in one and two dimensions. *Phys. Rev. Lett.* **110**, 205301 (2013).
- ⁷⁴ Schuch, N., Wolf, M. M., Verstraete, F. & Cirac, J. I. Entropy scaling and simulability by matrix product states. *Phys. Rev. Lett.* **100**, 030504 (2008).
- ⁷⁵ Lagendijk, A., van Tiggelen, B. & Wiersma, D. S. Fifty years of Anderson localization. *Physics Today* **62**, 24 (2009).
- ⁷⁶ Bardarson, J. H., Pollmann, F. & Moore, J. E. Unbounded growth of entanglement in models of many-body localization. *Phys. Rev. Lett.* **109**, 017202 (2012).
- ⁷⁷ Hamza, E., Sims, R. & Stolz, G. Dynamical localization in disordered quantum spin systems. *Commun. Math. Phys.* **315**, 215 (2012).
- ⁷⁸ Aleiner, I. L., Altshuler, B. L. & Shlyapnikov, G. V. A finite-temperature phase transition for disordered weakly interacting bosons in one dimension. *Ann. Phys.* **321**, 1126 (2006).
- ⁷⁹ Nandkishore, R. & Huse, D. A. Many body localization and thermalization in quantum statistical mechanics. ArXiv:1404.0686.
- ⁸⁰ Bauer, B. & Nayak, C. Area laws in a many-body localized state and its implications for topological order. *J. Stat. Mech.* P09005 (2013).
- ⁸¹ Polkovnikov, A. & Gritsev, V. Breakdown of the adiabatic limit in low-dimensional gapless systems. *Nature Phys.* **4**, 477 (2008).
- ⁸² Ruutu, V. M. H. *et al.* Vortex formation in neutron-irradiated superfluid ^3He as an analogue of cosmological defect formation. *Nature* **382**, 334 (1996).
- ⁸³ Ulm, S. *et al.* Observation of the Kibble-Zurek scaling law for defect formation in ion crystals. *Nat. Comm.* **4**, 2290 (2013).
- ⁸⁴ Pyka, K. *et al.* Topological defect formation and spontaneous symmetry breaking in ion coulomb crystals. *Nat. Comm.* **4** (2013).
- ⁸⁵ Chen, D., White, M., Borries, C. & DeMarco, B. Quantum quench of an atomic Mott insulator. *Phys. Rev. Lett.* **106**, 235304 (2011).
- ⁸⁶ Bernier, J.-S., Poletti, D., Barmettler, P., Roux, G. & Kollath, C. Slow quench dynamics of Mott-insulating regions in a trapped Bose gas. *Phys. Rev. A* **85**, 033641 (2012).
- ⁸⁷ Cirac, J. I. & Zoller, P. Goals and opportunities in quantum simulation. *Nature Phys.* **8**, 264 (2012).
- ⁸⁸ Hauke, P., Cucchietti, F. M., Tagliacozzo, L., Deutsch, I. & Lewenstein, M. Can one trust quantum simulators? *Rep. Prog. Phys.* **75**, 082401 (2012).
- ⁸⁹ Bloch, I., Dalibard, J. & Nascimbene, S. Quantum simulations with ultracold quantum gases. *Nature Phys.* **8**, 267 (2012).
- ⁹⁰ Trotzky, S. *et al.* Suppression of the critical temperature for superfluidity near the Mott transition: validating a quantum simulator. *Nature Phys.* **6**, 998 (2010).
- ⁹¹ Esslinger, T. Fermi-Hubbard physics with atoms in an optical lattice. *Ann. Rev. Cond. Mat. Phys.* **1**, 129 (2010).
- ⁹² Pertot, D. *et al.* Relaxation dynamics of a Fermi gas in an optical superlattice. ArXiv:1407.6037.
- ⁹³ Meinert, F. *et al.* Quantum quench in an atomic one-dimensional ising chain. *Phys. Rev. Lett.* **111**, 053003 (2013).
- ⁹⁴ Zohar, E., Cirac, J. I. & Reznik, B. A cold-atom quantum simulator for $\text{su}(2)$ Yang-Mills lattice gauge theory. *Phys. Rev. Lett.* **110**, 125304 (2013).
- ⁹⁵ Hofferberth, S., Lesanovsky, I., Fischer, B., Schumm, T. & Schmiedmayer, J. Non-equilibrium coherence dynamics in one-dimensional Bose gases. *Nature* **449**, 324 (2007).
- ⁹⁶ Kinoshita, T., Wenger, T. & Weiss, D. S. A quantum Newton's cradle. *Nature* **440**, 900 (2006).
- ⁹⁷ Blatt, R. & Roos, C. F. Quantum simulations with trapped ions. *Nature Phys.* **8**, 277 (2012).
- ⁹⁸ Islam, R. *et al.* Onset of a quantum phase transition with a trapped ion quantum simulator. *Nat. Commun.* **2**, 377 (2011).
- ⁹⁹ Maricq, M. M. Application of average Hamiltonian theory to the NMR of solids. *Phys. Rev. B* **25**, 6622 (1982).
- ¹⁰⁰ Arimondo, E., Ciampini, D., Eckardt, A., Holthaus, M. & Morsch, O. Kilohertz-driven Bose-Einstein condensates in optical lattices. *Adv. At. Mol. Opt. Phys.* **61**, 515 (2012).
- ¹⁰¹ Goldman, N. & Dalibard, J. Periodically-driven quantum systems: Effective Hamiltonians and engineered gauge fields. ArXiv:1404.4373.
- ¹⁰² Kitagawa, T., Berg, E., Rudner, M. & Demler, E. Topological characterization of periodically driven quantum systems. *Phys. Rev. B* **82**, 235114 (2010).
- ¹⁰³ Jotzu, G. *et al.* Experimental realization of the topological Haldane model. ArXiv:1406.7874.
- ¹⁰⁴ Aidelsburger, M. *et al.* Revealing the topology of Hofstadter bands with ultra-cold atoms. ArXiv:1407.4205.
- ¹⁰⁵ Struck, J., Simonet, J. & Sengstock, K. Spin orbit coupling in periodically driven optical lattices. ArXiv:1407.1953.
- ¹⁰⁶ Goldman, N. *et al.* Direct imaging of topological edge states in cold-atom systems. *PNAS* **110**, 6736 (2013).
- ¹⁰⁷ Lazarides, A., Das, A. & Moessner, R. Periodic thermodynamics of isolated systems. *Phys. Rev. Lett.* **112**, 150401 (2014).
- ¹⁰⁸ Endres, M. *et al.* Single-site- and single-atom-resolved measurement of correlation functions. ArXiv:1303.5652.
- ¹⁰⁹ Martin-Martinez, E., Fuentes, I. & Mann, R. B. Using Berry's phase to detect the Unruh effect at lower accelerations. *Phys. Rev. Lett.* **107**, 131301 (2011).
- ¹¹⁰ Agarwal, K. *et al.* Chiral prethermalization in supersonically split condensates. ArXiv: 1402.6716.
- ¹¹¹ Ritsch, H., Domokos, P., Brennecke, F. & Esslinger, T. Cold atoms in cavity-generated dynamical optical potentials. *Rev. Mod. Phys.* **85**, 553 (2013).
- ¹¹² Linden, N., Popescu, S. & Skrzypczyk, P. How small can thermal machines be? The smallest possible refrigerator, *Phys. Rev. Lett.* **105**, 130401 (2010).
- ¹¹³ Gallego, R., Riera, A. & Eisert, J. Correlated thermal machines in the micro-world (2013). Arxiv.1310.8349.
- ¹¹⁴ Hubeny, V. E. & Rangamani, M. A holographic view on physics out of equilibrium. *Adv. High En. Phys.* **2010**, 297916 (2010).
- ¹¹⁵ Kliensch, M., Gogolin, C., Kastoryano, M. J., Riera, A. & Eisert, J. Locality of temperature. *Phys. Rev. X* **4**, 031019 (2014).
- ¹¹⁶ Diehl, S. *et al.* Quantum states and phases in driven open quantum systems with cold atoms. *Nature Phys.* **4**, 878 (2008).
- ¹¹⁷ Barreiro, J. T. *et al.* An open-system quantum simulator with trapped ions. *Nature* **470**, 486 (2011).
- ¹¹⁸ Barontini, G. *et al.* Controlling the dynamics of an open many-body quantum system with localized dissipation. *Phys. Rev. Lett.* **110**, 035302 (2013).
- ¹¹⁹ Diehl, S., Rico, E., Baranov, M. A. & Zoller, P. Topology by dissipation in atomic quantum wires. *Nature Phys.* **7**, 971 (2011).