



Microsatellites within the feline androgen receptor are suitable for X chromosome-linked clonality testing in archival material

Nadine M Farwick¹, Robert Klopffleisch¹,
Achim D Gruber¹ and Alexander Th A Weiss²

Journal of Feline Medicine and Surgery
1–7

© The Author(s) 2016
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/1098612X16634386
jfms.com



Abstract

Objectives A hallmark of neoplasms is their origin from a single cell; that is, clonality. Many techniques have been developed in human medicine to utilise this feature of tumours for diagnostic purposes. One approach is X chromosome-linked clonality testing using polymorphisms of genes encoded by genes on the X chromosome. The aim of this study was to determine if the feline androgen receptor gene was suitable for X chromosome-linked clonality testing.

Methods The feline androgen receptor gene, was characterised and used to test clonality of feline lymphomas by PCR and polyacrylamide gel electrophoresis, using archival formalin-fixed, paraffin-embedded material.

Results Clonality of the feline lymphomas under study was confirmed and the gene locus was shown to represent a suitable target in clonality testing.

Conclusions and relevance Because there are some pitfalls using X chromosome-linked clonality testing, further studies are necessary to establish this technique in the cat.

Accepted: 26 January 2016

Introduction

Most tumours arise from a single cell of origin and are therefore considered clonal proliferations.^{1–3} Clonality testing has developed into a significant accessory technique in human cancer diagnostics.⁴ Clonality testing is especially important for the diagnosis of haematological malignancies such as lymphoma and leukaemia because lymphatic neoplasia and hyperplasia are not always easily differentiated histologically.^{5,6}

To improve lymphoma diagnostics in veterinary medicine several techniques have been applied to analyse clonal antigen receptor rearrangement, including Southern blotting^{7–10} and PCR.^{7,9–16} Further techniques have been used for clonality testing in veterinary medicine and targeted clonally integrated feline leukaemia virus (FeLV) provirus within the host's genome,¹⁷ mutations within *c-kit*,¹⁸ clonal polymorphisms within mitochondrial DNA,¹⁹ and polymorphisms of microsatellites and mitochondrial DNA in the transmissible venereal tumour.²⁰ Clonality testing involving X-linked genes has also been used in dogs and cats,^{21,22} both employing the androgen receptor gene.

To date, there is no routinely implementable clonality test for non-lymphoid neoplasms in cats.²³ The analysis of clonal inactivation of genes located on the X chromosome in females (X-linked clonality testing) has nevertheless been applied to many human tumours.^{24,25} This approach is based on the fact that females carry two copies of the genes located on the X chromosome, of which one is inactivated to avoid an 'overdose' of gene expression. The inactivated allele of X-chromosomal genes is randomly selected during early embryogenesis and inactivation is based on methylation of cytosine residues

¹Department of Veterinary Pathology, Free University of Berlin, Berlin, Germany

²Chemical and Veterinary Analytical Institute Muensterland-Emscher-Lippe, Muenster, Germany

Corresponding author:

Alexander Weiss DVM, PhD, Chemical and Veterinary Analytical Institute Muensterland-Emscher-Lippe, Joseph-Koenig-Str 40, Muenster, Germany
Email: alexander.weiss@cvua-mel.de

within CpG-rich islands.²⁶ To test for clonality, genomic DNA is digested by methylation-sensitive endonucleases, such as *HpaII* or *HhaI*. In the following PCR only the inactivated allele of the corresponding gene can be amplified.²⁷ One gene with such a polymorphism located next to a CpG island that is heterozygous in a large fraction of individuals is the human androgen receptor. This gene features a polymorphic short tandem repeat of CAG,²⁷ which has been used to demonstrate clonality of tumours in women.²⁸ This test has been termed the HUMARA assay.²⁸

The androgen receptor gene of cats has already been sequenced.²⁹ Similar to its canine counterpart it includes two stretches of CAG repeats. However, possible polymorphic microsatellites within exon 1 have not yet been reported. Therefore, we investigated exon 1 of the feline androgen receptor for potentially useful microsatellites and applied them in clonality tests. We then designed the assay such that archival formalin-fixed and paraffin embedded material could be tested.

Materials and methods

Case selection

To assess the presence of microsatellites within the feline androgen receptor we used the DNA of 42 cats from a previous study.³⁰ The newly developed assay was then applied to 50 feline lymphomas and lymph nodes with reactive lymphocytic hyperplasia that had already been thoroughly investigated.^{15,31}

Nucleic acid extraction, PCR amplification of the feline androgen receptor exon 1 and polyacrylamide gel electrophoresis

Genomic DNA was extracted from macro-dissected, formalin-fixed and paraffin-embedded tissue samples with

the Genra Puregene Tissue Kit (Qiagen), including RNase treatment as described.³⁰

Primers spanning both CAG repeats and primers for the first CAG region within exon 1 of the feline androgen receptor were newly designed according to a publicly available sequence (Table 1; GenBank Accession: AJ893545) using GeneFisher.³³ Primers for the second CAG stretch of the feline androgen receptor were adapted from primers designed for the canine androgen receptor.³²

PCR was performed as described elsewhere.²¹ In short, the master mix consisted of 2.0 mM MgCl₂, 0.2 mM of each dNTP, 0.2 μM of each primer and 0.024 U/μl Taq DNA polymerase (GoTaq; Promega). Cycling conditions encompassed an initial denaturing at 94°C for 5 mins followed by 40 cycles of DNA melting at 92°C for 30 s, annealing for 15 s, amplification at 72°C for 80 s in the case of the long amplicons (>600 base pairs) and 20 s in the case of the short ones. This was followed by a final amplification at 72°C for 5 mins. Annealing temperatures for specific primer combinations are summarised in Table 1.

Using a semi-nested PCR protocol, products of the first round of PCR were diluted 1:500 with 5 mM Tris buffer. Cycling conditions were modified in the second round of PCR by using only 30 cycles, 59°C as the primer annealing temperature and 15 s elongation time. After amplification, PCR products were screened by agarose gel electrophoresis using 2% gels impregnated with ethidium bromide.

To detect deactivated alleles of the canine androgen receptor genomic DNA was digested prior to PCR amplification with the methylation-sensitive endonuclease *HpaII* (restriction site: CCGG) (Fermentas). Genomic DNA (1 μg) was digested with 10 U endonuclease overnight at 37°C in a final volume of 20 μl.

Table 1 Primer sequences, melting temperatures, amplicon sizes and positions

No	Identification	Primer sequence	Temperature (°C)	Amplicon size (bp) in AJ893545
Outer primers				
1	FeARof1*	5'-CAAGACCTATCGAGGAGCTTTC-3'	60–55 [†]	645
2	FeARor1*	5'-GTCGAACTGCCACCTAGGTAAC-3'		
First polymorphic CAG repeat – FELARA1				
3	FELARAf1	5'- TATTCCAGAGCGTGCGCGAAG -3'	57	245
4	FELARAr1	5'- GCTGTTGTGAAGGCTGCTGTTTC -3'		
Second polymorphic CAG repeat – FELARA2 ³²				
5	FELARAf2*	5'-GACTCAGCTGCCCCATCCAC-3'	57	250
6	FELARAr2*	5'-GGTAACTGTCTTGGAGGAGG-3'		
Additional combinations used				
3/6	FELARAf1 and FELARAr2		59	606
1/4	FeARof1 and FELARAr1		57	269
5/2	FELARAf2 and FeARor1		57	265

*Designed for the canine androgen receptor but also works with the feline counterpart

[†]A touch-down protocol was applied starting at 60°C, reducing the temperature by 0.5°C per cycle for 10 cycles followed by 30 cycles at 55°C
bp = base pairs




Figure 4 Clonality analysis of feline lymphomas (1–3) and lymphatic hyperplasias. Each sample comprises two lanes from two different PCRs. The left includes native DNA the right *Hpa*II-digested DNA. Samples 1 and 2 display non-random deactivation of one androgen receptor gene; that is, they are clonal (one band after *Hpa*II digestion). Sample 3 displays a visibly weakened second band, probably due to a strong non-neoplastic lymphatic infiltration of the clonal, neoplastic cell population. Samples 4 and 5 display random inactivation of the androgen receptor gene (two bands in both lanes), resulting in amplification of both alleles, indicating a polyclonal cell population. bp = base pairs

recent findings in the analysis of the heterozygosity of dogs, where we identified 19% of animals as heterozygous,²¹ while older reports identified 43% of animals as heterozygous.³⁴

The lack of interspersed 'CAA' triplets within the second microsatellite in cats when compared with dogs (see Figure 1b), results in a higher rate of heterozygous cats, leading to better applicability of this test in cats than in dogs.

Thorough digestion of the non-deactivated X chromosome prior to PCR by a methylation-sensitive endonuclease has been identified as a crucial aspect for the assay presented here. In this study, as in a previous one,²¹ digestion was most effective using the long fragments for PCR amplification. When using shorter fragments, digestion had to be extended or higher concentrations of endonuclease had to be used. In this case, digestion controls from male animals have to be used. This phenomenon is a result of the different number of restriction sites covered by the PCR fragments. The long fragment includes three restriction sites, whereas the short

fragments FELARA1 and FELARA2 each contain only one restriction site. As we used archival material we had to cope with limited amounts of starting material and therefore had to use nested PCR to produce enough material for the PAGE. However, digestion should be no problem using fresh or freshly frozen material and the long PCR fragment. Another alternative could be direct amplification of the short fragments.

All lymphomas from heterozygous animals that were tested revealed a non-random deactivation of one allele of the feline androgen receptor and, therefore, are of clonal origin. In contrast, the two lymphatic hyperplasias analysed revealed a random deactivation and therefore, as was expected, are of polyclonal origin.

One case of lymphoma in this study revealed only a weakened band upon digestion; this could be due to a reactive infiltration by lymphocytes. Incomplete digestion of the active allele could be a further reason for this result.

It is absolutely vital with this technique to first check whether informative pattern, that is, heterozygous alleles, are present in the animal to be tested. Both samples should be directly compared upon PAGE to visualise clearly any changes after digestion by a methylation-sensitive endonuclease.

This study demonstrates the usefulness of the polymorphic tandem repeat within exon 1 of the feline androgen receptor for clonality analysis, and it can be expected that the assay can also be applied to other tumour types. Because of contrary results regarding the rate of heterozygosity, further studies are needed to establish the true extent to which this test can be utilised in routine diagnostics.

However, there are further caveats that need to be addressed. In humans, a skewing of the X chromosome inactivation pattern in haematopoietic cells, especially granulocytes, of some ageing women has been identified.³⁵ It needs to be determined if such a phenomenon exists in cats. This should be visible in a PAGE of PCR products obtained from non-digested DNA. To be certain only to identify truly clonal populations, total loss of a second band could be regarded as hallmark of a clonal cell population.³⁵

Furthermore, the two polymorphic tandem repeats within the androgen receptor gene are approximately 1 kb downstream from the promoter region of the gene. This region is vital in gene regulation; that is, activation and inactivation. In some humans methylation patterns of the promoter region of the androgen receptor gene and within exon 1 are not completely concordant.³⁶ In the future it should be determined whether this rather remote location exhibits the same methylation pattern as the direct vicinity of the promoter region in the cat. Additionally, more than one polymorphic locus should be studied to determine truly monoclonal cell populations.

Conclusions

Until now there have been no routine assays to evaluate clonality in feline neoplasms with the exception of lymphomas;²² therefore, it is necessary to further establish X chromosome-linked clonality testing for basic research of carcinogenesis in this species.

Conflict of interest The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding The authors received no financial support for the research, authorship, and/or publication of this article.

References

- Garcia SB, Novelli M and Wright NA. **The clonal origin and clonal evolution of epithelial tumours.** *Int J Exp Pathol* 2000; 81: 89–116.
- Marusyk A and Polyak K. **Tumor heterogeneity: causes and consequences.** *Biochim Biophys Acta* 2010; 1805: 105–117.
- Nowell PC. **The clonal evolution of tumor cell populations.** *Science* 1976; 194: 23–28.
- Leedham SJ and Wright NA. **Human tumour clonality assessment – flawed but necessary.** *J Pathol* 2008; 215: 351–354.
- Diaz-Cano SJ, Blanes A and Wolfe HJ. **PCR techniques for clonality assays.** *Diagn Mol Pathol* 2001; 10: 24–33.
- Jevremovic D and Viswanatha DS. **Molecular diagnosis of hematopoietic and lymphoid neoplasms.** *Hematol Oncol Clin North Am* 2009; 23: 903–933.
- Ishiguro N, Matsui T and Shinagawa M. **Differentiation analysis of bovine T-lymphosarcoma.** *Vet Immunol Immunopathol* 1994; 41: 1–17.
- Ishiguro N, Shinagawa T, Matsui T, et al. **Putative bovine B cell lineage tumor in sporadic bovine leukosis.** *Vet Immunol Immunopathol* 1994; 42: 185–197.
- Levesque KS, Bonham L and Levy LS. **Flvi-1, a common integration domain of feline leukemia virus in naturally occurring lymphomas of a particular type.** *J Virol* 1990; 64: 3455–3462.
- Momoi Y, Nagase M, Okamoto Y, et al. **Rearrangements of immunoglobulin and T-cell receptor genes in canine lymphoma/leukemia cells.** *J Vet Med Sci* 1993; 55: 775–780.
- Burnett RC, Vernau W, Modiano JF, et al. **Diagnosis of canine lymphoid neoplasia using clonal rearrangements of antigen receptor genes.** *Vet Pathol* 2003; 40: 32–41.
- Henrich M, Hecht W, Weiss AT, et al. **A new subgroup of immunoglobulin heavy chain variable region genes for the assessment of clonality in feline B-cell lymphomas.** *Vet Immunol Immunopathol* 2009; 130: 59–69.
- Moore PF, Woo JC, Vernau W, et al. **Characterization of feline T cell receptor gamma (TCRG) variable region genes for the molecular diagnosis of feline intestinal T cell lymphoma.** *Vet Immunol Immunopathol* 2005; 106: 167–178.
- Vernau W and Moore PF. **An immunophenotypic study of canine leukemias and preliminary assessment of clonality by polymerase chain reaction.** *Vet Immunol Immunopathol* 1999; 69: 145–164.
- Weiss AT, Klopffleisch R and Gruber AD. **T-cell receptor gamma chain variable and joining region genes of subgroup 1 are clonally rearranged in feline B- and T-cell lymphoma.** *J Comp Pathol* 2011; 144: 123–134.
- Werner JA, Woo JC, Vernau W, et al. **Characterization of feline immunoglobulin heavy chain variable region genes for the molecular diagnosis of B-cell neoplasia.** *Vet Pathol* 2005; 42: 596–607.
- Casey JW, Roach A, Mullins JI, et al. **The U3 portion of feline leukemia virus DNA identifies horizontally acquired proviruses in leukemic cats.** *Proc Natl Acad Sci U S A* 1981; 78: 7778–7782.
- Zavodovskaya R, Chien MB and London CA. **Use of kit internal tandem duplications to establish mast cell tumor clonality in 2 dogs.** *J Vet Intern Med* 2004; 18: 915–917.
- Bertagnolli AC, Soares P, van Asch B, et al. **An assessment of the clonality of the components of canine mixed mammary tumours by mitochondrial DNA analysis.** *Vet J* 2009; 182: 269–274.
- Murgia C, Pritchard JK, Kim SY, et al. **Clonal origin and evolution of a transmissible cancer.** *Cell* 2006; 126: 477–487.
- Delcour NM, Klopffleisch R, Gruber AD, et al. **Canine cutaneous histiocytomas are clonal lesions as defined by X-linked clonality testing.** *J Comp Pathol* 2013; 149: 192–198.
- Mochizuki H, Goto-Koshino Y, Takahashi M, et al. **X-chromosome inactivation pattern analysis for the assessment of cell clonality in cats.** *Vet Pathol* 2012; 49: 963–970.
- Moore PF. **Immunological concepts applied to pathologic diagnosis of proliferative diseases of the immune system.** *Cancer Therapy* 2008; 6: 263–270.
- Mannhalter C and Mitterbauer G. **Analysis of clonal cell growth using X-chromosome inactivation patterns.** *Onkologie* 1998; 21: 380–386.
- Wainscoat JS and Fey MF. **Assessment of clonality in human tumors: a review.** *Cancer Res* 1990; 50: 1355–1360.
- Leeb M and Wutz A. **Mechanistic concepts in X inactivation underlying dosage compensation in mammals.** *Heredity* 2011; 105: 64–70.
- Allen RC, Zoghbi HY, Moseley AB, et al. **Methylation of HpaII and HhaI sites near the polymorphic CAG repeat in the human androgen-receptor gene correlates with X chromosome inactivation.** *Am J Hum Genet* 1992; 51: 1229–1239.
- Peng H, Du M, Diss TC, et al. **Clonality analysis in tumours of women by PCR amplification of X-linked genes.** *J Pathol* 1997; 181: 223–227.
- Poux C, Madsen O, Marquard E, et al. **Asynchronous colonization of Madagascar by the four endemic clades of Primates, Tenrecs, Carnivores, and Rodents as inferred from nuclear genes.** *Syst Biol* 2005; 54: 719–730.
- Weiss AT, Delcour NM, Meyer A, et al. **Efficient and cost-effective extraction of genomic DNA from formalin-fixed and paraffin-embedded tissues.** *Vet Pathol* 2011; 48: 839–843.
- Weiss AT, Klopffleisch R and Gruber AD. **Prevalence of feline leukaemia provirus DNA in feline lymphomas.** *J Feline Med Surg* 2010; 12: 929–935.
- Wagner LA, Kiem HP, McSweeney P, et al. **Development of a clonality assay for canine hematopoietic stem cells.** *Blood* 1995; 86: 3265–3267.

-
- 33 Giegerich R, Meyer F and Schleiermacher C. **GeneFisher – software support for the detection of postulated genes.** *Proc Int Conf Intell Syst Mol Biol* 1996; 4: 68–77.
- 34 Shibuya H, Nonneman DJ, Huang TH, et al. **Two polymorphic microsatellites in a coding segment of the canine androgen receptor gene.** *Anim Genet* 1993; 24: 345–348.
- 35 Chen GL and Prchal JT. **X-linked clonality testing: interpretation and limitations.** *Blood* 2007; 110: 1411–1419.
- 36 Swierczek SI, Piterkova L, Jelinek J, et al. **Methylation of AR locus does not always reflect X chromosome inactivation state.** *Blood* 2012; 119: e100–109.