

Aus dem Institut für Medizinische Psychologie
der Medizinischen Fakultät Charité – Universitätsmedizin Berlin

DISSERTATION

Characterization in humans of *in vitro* leukocyte maximal telomerase activity
capacity (mTAC) and association with stress

zur Erlangung des akademischen Grades
Doctor rerum medicinalium (Dr. rer. medic.)

vorgelegt der Medizinischen Fakultät
Charité – Universitätsmedizin Berlin

von

Karin de Punder
aus Purmerend

Datum der Promotion: 14.09.2018

Table of Contents

| | |
|---|----|
| Abstract (German) | 3 |
| Abstract (English) | 5 |
| Affidavit | 7 |
| Excerpt from the Journal Summary List (ISI Web of KnowledgeSM) | 9 |
| Decision Letter | 13 |
| Characterization in humans of <i>in vitro</i> leukocyte maximal telomerase activity capacity (mTAC) and association with stress | 14 |
| Supplemental material | 43 |
| Curriculum Vitae | 46 |
| List of Publications | 47 |
| Acknowledgements | 52 |

Abstract (German)

Hintergrund: Telomerase, ein zelluläres Transkriptionenzym, das der Telomerverkürzung entgegenwirkt und wichtig ist für gesunde Zellfunktion, ist ein vielversprechender Biomarker hinsichtlich pathologischer Folgen von Stress auf zelluläre Alterungsprozesse. Bisherige Studien zur Messung von Telomeraseaktivität oder -expression haben dies meistens unter basalen Bedingungen durchgeführt. Basale Telomerase-Werte sind jedoch aufgrund der niedrigen Expressionsrate und der dynamischen Regulation der Telomerase in Leukozyten sehr schwierig zu messen und zu interpretieren. Im Gegensatz zur basalen Expressionsrate kann Telomerase-Aktivität im Rahmen eines ökologisch-validen *Challenge Tests* wie der *in vitro* Mitogen-Stimulation quantifiziert werden. Ein potentieller Vorteil dieses Verfahrens ist, dass dadurch die o.a. Einschränkungen der Interpretierbarkeit der basalen Telomerase-Aktivität umgangen werden können, und man somit die Möglichkeit hat, individuelle Unterschiede zu erfassen in der Kapazität des Telomersystems auf einem immunologischen Stimulus zu reagieren.

Ziel: Das Ziel der Studie bestand in der Entwicklung und Validierung eines Maßes der maximalen Telomerase-Aktivitätskapazität (mTAC) zur Nutzung in Humanstudien im Bereich der Telomer-Biologie. Außerdem wurde der Zusammenhang dieses Maßes mit Stress bzw. Stressreaktivität untersucht.

Methoden: Zuerst wurde der optimale post-stimulative Zeitverlauf zur mTAC-Charakterisierung mit einem *in vitro* Mitogen-Challenge-Protokoll (PHA ergänzt durch Interleukin(IL)-2) bestimmt. Danach wurden sowohl mTAC in Leukozyten und Cortisolkonzentrationen im Speichel von 28 jungen gesunden Proband/innen zu verschiedenen Messzeitpunkten im Verlauf des Tages und im Rahmen eines standardisierten Paradigmas zur Induktion psychosozialen Stresses im Labor gemessen. Darüber hinaus wurde in einem Teil der Stichprobe Durchflusszytometrie verwendet, um die Immunzell-Verteilung vor Stimulation zu bestimmen. Die wahrgenommene (chronische) Stressbelastung wurde durch die *Perceived Stress Scale* erfasst.

Resultate: Der optimale Zeitpunkt zur Quantifizierung von mTAC in humanen Leukozyten liegt bei 72 Stunden nach Mitogen-Stimulation. mTAC weist eine erhebliche Stabilität innerhalb von Personen auf und wird nicht durch situationsbedingte Faktoren wie Tageszeit, Cortisol, akute Stressexposition und Immunzell-Verteilung beeinflusst. Ein signifikanter Anteil der Varianz zwischen Personen in mTAC wird durch chronische Stressbelastung und biologische

Stressreaktivität erklärt. Insbesondere waren mTAC-Werte bei Personen mit hoher wahrgenommener Stressbelastung um 25% niedriger im Vergleich zu Probanden mit mittlerem oder niedriger wahrgenommener Stressbelastung. Außerdem erklärten individuelle Unterschiede in der Cortisolreaktion auf Laborstressexposition 32% der Varianz von mTAC.

Fazit: Basierend auf diesen Ergebnissen kann man schlussfolgern, dass mTAC einen nützlichen individuellen Marker für stressbezogene Humanstudien im Bereich der Telomerbiologie darstellt.

Abstract (English)

Background: Telomerase, a cellular reverse transcriptase enzyme that can counteract telomere shortening and, in addition, preserves healthy cell function, constitutes a promising target for research into the pathological effects of stress on cellular aging. Previous studies that have included measures of telomerase have typically measured telomerase expression or activity under basal (resting) conditions. It is, however, challenging to reliably quantify or interpret these data because leukocyte telomerase is typically expressed at very low levels and is dynamic in nature. In contrast to basal measures, telomerase activity can be quantified in response to an ecologically-valid challenge such as mitogen stimulation *in vitro*. The potential advantage of this approach is that it may bypass the above-mentioned limitations to provide an indicator of individual differences in the *capacity* of the telomere biology system to respond to an immunological challenge.

Objective: The aim of this study was to validate an *in vitro* measure of leukocyte maximal telomerase activity capacity (mTAC) for use in human studies of telomere biology, and to determine its association with measures of stress and stress responsivity.

Methods: First, the optimal post-stimulation time course to characterize mTAC was established using an *in vitro* mitogen challenge (phytohemagglutinin (PHA) supplemented with interleukin(IL)-2). Next, mTAC was measured in leukocytes and cortisol concentrations were assessed in saliva obtained from 28 healthy young women and men at different times of the day and before and after a standardized laboratory stressor. In addition, immune cell distributions prior to mitogen stimulation were determined by flow cytometry in a subset of the participants. Perceived (chronic) stress also was assessed using the *Perceived Stress Scale*.

Results: The optimal time point to quantify human leukocyte mTAC was 72 hours after mitogen stimulation. mTAC exhibited substantial within-subject stability across time and was not influenced by situational factors including time of day, cortisol concentration, acute stress exposure, and immune cell distribution. A significant proportion of the between-subject variability in mTAC was associated with measures of stress and stress responsivity. Particularly, there was a 25% difference in mTAC between subjects reporting high compared to medium or low levels of perceived (chronic) stress. Also, individual differences in the cortisol response to stress-exposure accounted for as much as 32% of the variation in mTAC.

Conclusion: Based collectively on these findings, it appears that mTAC may represent a potentially useful individual difference measure in stress-related studies of the human telomere biology system.

Affidavit

I, Karin de Punder certify under penalty of perjury by my own signature that I have submitted the thesis on the topic: „Characterization in humans of *in vitro* leukocyte maximal telomerase activity capacity (mTAC) and association with stress“. I wrote this thesis independently and without assistance from third parties, I used no other aids than the listed sources and resources.

All points based literally or in spirit on publications or presentations of other authors are, as such, in proper citations (see "uniform requirements for manuscripts (URM)" the ICMJE www.icmje.org) indicated. The section on methodology (in particular practical work, laboratory requirements, statistical processing) and results (in particular images, graphics and tables) corresponds to the URM (s.o) and are answered by me. My contribution in the selected publication for this dissertation corresponds to those that are specified in the following joint declaration with the responsible person and supervisor.

The importance of this affidavit and the criminal consequences of a false affidavit (section 156,161 of the Criminal Code) are known to me and I understand the rights and responsibilities stated therein.

Date

Signature

Detailed Declaration of Contribution

Karin de Punder had the following share in the following publication:

Publikation: Karin de Punder, Christine Heim, Ingo Przesdzing, Pathik D. Wadhwa, Sonja Entringer (2018). Characterization in humans of *in vitro* leukocyte maximal telomerase activity capacity (mTAC) and association with stress. *Philos Trans R Soc Lond B Biol Sci.* 5;373(1741) pii: 20160441.

<http://dx.doi.org/10.1098/rstb.2016.0441>

Contribution in detail:

Karin de Punder helped writing the ethical application for this study. She participated in the design of the study and recruited all the study participants. She planned and coordinated the study and guided and prepared the participants during both test days. In addition, she carried out the laboratory assays (leukocyte isolation and stimulation, telomerase measurements, salivary cortisol measurements, antibody stainings for the flow cytometry experiments), performed the statistical data analyses and drafted the manuscript. She also made the tables and figures presented in the manuscript.

Signature, date and stamp of the supervising University teacher

Signature of the doctoral candidate

Excerpt from the Journal Summary List (ISI Web of KnowledgeSM)

Journal Data Filtered By: **Selected JCR Year: 2016** Selected Editions: SCIE,SSCI
 Selected Categories: **“BIOLOGY”** Selected Category Scheme: WoS **Gesamtanzahl: 72**
Journalle

| Rank | Full Journal Title | Total Cites | Journal Impact Factor | Eigenfactor Score |
|----------|--|---------------|-----------------------|-------------------|
| 1 | Physics of Life Reviews | 1,327 | 13.840 | 0.003180 |
| 2 | BIOLOGICAL REVIEWS | 9,669 | 11.615 | 0.017300 |
| 3 | PLOS BIOLOGY | 26,893 | 9.797 | 0.067500 |
| 4 | eLife | 15,730 | 7.725 | 0.130380 |
| 5 | BMC BIOLOGY | 4,588 | 6.779 | 0.016120 |
| 6 | PHILOSOPHICAL TRANSACTIONS OF THE ROYAL SOCIETY B-BIOLOGICAL SCIENCES | 36,908 | 5.846 | 0.069600 |
| 7 | FASEB JOURNAL | 42,242 | 5.498 | 0.057660 |
| 8 | BIOSCIENCE | 15,467 | 5.378 | 0.013660 |
| 9 | PROCEEDINGS OF THE ROYAL SOCIETY B-BIOLOGICAL SCIENCES | 44,463 | 4.940 | 0.088440 |
| 10 | BIOESSAYS | 9,199 | 4.441 | 0.016940 |
| 11 | QUARTERLY REVIEW OF BIOLOGY | 3,683 | 4.250 | 0.001500 |
| 12 | Current Opinion in Insect Science | 665 | 3.660 | 0.002870 |
| 13 | JOURNAL OF BIOLOGICAL RHYTHMS | 2,617 | 3.500 | 0.004220 |
| 14 | Geobiology | 1,798 | 3.462 | 0.004150 |
| 15 | BIOELECTROCHEMISTRY | 3,916 | 3.346 | 0.003820 |
| 16 | JOURNAL OF EXPERIMENTAL BIOLOGY | 30,619 | 3.320 | 0.036880 |
| 17 | Biology Letters | 8,242 | 3.089 | 0.021760 |
| 18 | Biology Direct | 1,507 | 2.856 | 0.005180 |
| 19 | Science China-Life Sciences | 1,621 | 2.781 | 0.004510 |
| 20 | Interface Focus | 1,272 | 2.693 | 0.005870 |
| 21 | ASTROBIOLOGY | 2,414 | 2.603 | 0.005550 |
| 22 | SAUDI JOURNAL OF BIOLOGICAL SCIENCES | 1,095 | 2.564 | 0.002480 |



| | | | | |
|----|--|--------|-------|----------|
| 23 | CHRONOBIOLOGY INTERNATIONAL | 4,215 | 2.562 | 0.007100 |
| 24 | RADIATION RESEARCH | 8,394 | 2.539 | 0.007920 |
| 25 | RADIATION AND ENVIRONMENTAL BIOPHYSICS | 1,468 | 2.398 | 0.002460 |
| 26 | AEROBIOLOGIA | 1,166 | 2.202 | 0.001060 |
| 27 | JOURNAL OF THERMAL BIOLOGY | 2,614 | 2.157 | 0.003440 |
| 28 | JOURNAL OF THEORETICAL BIOLOGY | 19,308 | 2.113 | 0.022280 |
| 29 | Biology Open | 1,457 | 2.095 | 0.008260 |
| 30 | CRYOBIOLOGY | 3,949 | 1.996 | 0.003490 |
| 31 | INTERNATIONAL JOURNAL OF RADIATION BIOLOGY | 4,417 | 1.992 | 0.004350 |
| 32 | BIOLOGICAL BULLETIN | 4,803 | 1.950 | 0.002280 |
| 33 | BIOELECTROMA GNETICS | 2,760 | 1.933 | 0.002570 |
| 34 | Advances in Experimental Medicine and Biology | 14,818 | 1.881 | 0.029570 |
| 35 | COMPUTERS IN BIOLOGY AND MEDICINE | 3,760 | 1.836 | 0.006550 |
| 36 | JOURNAL OF RADIATION RESEARCH | 2,270 | 1.788 | 0.004620 |
| 37 | AMERICAN JOURNAL OF HUMAN BIOLOGY | 2,959 | 1.780 | 0.005110 |
| 38 | BIOLOGICAL RESEARCH | 1,230 | 1.692 | 0.001630 |
| 39 | BIOSYSTEMS | 2,471 | 1.652 | 0.002850 |
| 40 | MATHEMATICAL MEDICINE AND BIOLOGY-A JOURNAL OF THE IMA | 469 | 1.610 | 0.000920 |
| 41 | International Journal of Astrobiology | 475 | 1.598 | 0.001340 |
| 42 | BRAZILIAN JOURNAL OF MEDICAL AND BIOLOGICAL RESEARCH | 4,568 | 1.578 | 0.004000 |

| | | | | |
|----|---|--------|-------|----------|
| 43 | JOURNAL OF MATHEMATICAL BIOLOGY | 4,399 | 1.566 | 0.006540 |
| 44 | BioScience Trends | 614 | 1.545 | 0.001230 |
| 45 | EXCLI Journal | 514 | 1.462 | 0.001120 |
| 46 | BIOMETRIKA | 18,608 | 1.448 | 0.015690 |
| 47 | JOURNAL OF BIOSCIENCES | 2,495 | 1.422 | 0.003410 |
| 48 | COMPUTATIONAL BIOLOGY AND CHEMISTRY | 1,025 | 1.331 | 0.001780 |
| 49 | BIOMETRICS | 19,952 | 1.329 | 0.014450 |
| 50 | ELECTROMAGNETIC BIOLOGY AND MEDICINE | 573 | 1.272 | 0.000960 |
| 51 | BULLETIN OF MATHEMATICAL BIOLOGY | 3,434 | 1.263 | 0.004440 |
| 52 | MATHEMATICAL BIOSCIENCES | 4,736 | 1.246 | 0.004820 |
| 53 | ANNALS OF HUMAN BIOLOGY | 1,865 | 1.240 | 0.002600 |
| 54 | JOURNAL OF ETHNOBIOLOGY | 422 | 1.217 | 0.000490 |
| 55 | Journal of Biological Research-Thessaloniki | 198 | 1.200 | 0.000330 |
| 56 | MICROSCOPY RESEARCH AND TECHNIQUE | 4,509 | 1.147 | 0.003980 |
| 57 | COMPTESE RENDUS BIOLOGIES | 2,111 | 1.100 | 0.002640 |
| 58 | TURKISH JOURNAL OF BIOLOGY | 1,049 | 1.038 | 0.001480 |
| 59 | Central European Journal of Biology | 744 | 1.016 | 0.001670 |
| 60 | ORIGINS OF LIFE AND EVOLUTION OF BIOSPHERES | 1,224 | 1.000 | 0.001260 |
| 61 | JOURNAL OF BIOLOGICAL EDUCATION | 519 | 0.946 | 0.000420 |
| 62 | FOLIA BIOLOGICA | 529 | 0.939 | 0.000630 |
| 63 | JOURNAL OF AGRICULTURAL BIOLOGICAL AND ENVIRONMENTAL STATISTICS | 809 | 0.852 | 0.001780 |
| 64 | HUMAN BIOLOGY | 1,809 | 0.825 | 0.001080 |
| 65 | THEORY IN BIOSCIENCES | 338 | 0.778 | 0.000800 |
| 66 | BIOLOGIA | 1,756 | 0.759 | 0.002500 |

| | | | | |
|----|---|-------|-------|----------|
| 67 | JOURNAL OF THE HISTORY OF BIOLOGY | 482 | 0.737 | 0.000640 |
| 68 | PROCEEDINGS OF THE BIOLOGICAL SOCIETY OF WASHINGTON | 1,269 | 0.667 | 0.000520 |
| 69 | BRAZILIAN ARCHIVES OF BIOLOGY AND TECHNOLOGY | 1,826 | 0.644 | 0.001650 |
| 70 | CRYOLETTERS | 890 | 0.628 | 0.000780 |
| 71 | BIOLOGICAL RHYTHM RESEARCH | 494 | 0.624 | 0.000590 |
| 72 | FOLIA BIOLOGICA-KRAKOW | 333 | 0.581 | 0.000400 |
| 73 | ACTA BIOLOGICA HUNGARICA | 448 | 0.506 | 0.000420 |
| 74 | REVISTA DE BIOLOGIA TROPICAL | 2,017 | 0.495 | 0.002090 |
| 75 | BRAZILIAN JOURNAL OF BIOLOGY | 1,932 | 0.479 | 0.002330 |
| 76 | Open Life Sciences | 35 | 0.448 | 0.000080 |
| 77 | Theoretical Biology Forum | 23 | 0.421 | 0.000070 |
| 78 | JOURNAL OF BIOLOGICAL SYSTEMS | 395 | 0.390 | 0.000540 |
| 79 | Archives of Biological Sciences | 725 | 0.352 | 0.001370 |
| 80 | AMERICAN BIOLOGY TEACHER | 552 | 0.318 | 0.000290 |
| 81 | ZHURNAL OBSHCHEI BIOLOGII | 206 | 0.313 | 0.000150 |
| 82 | BIOLOGY BULLETIN | 421 | 0.299 | 0.000560 |
| 83 | Bioscience Journal | 544 | 0.267 | 0.001410 |
| 84 | PERIODICUM BIOLOGORUM | 289 | 0.184 | 0.000250 |

Copyright © 2017 Thomson Reuters

Decision Letter

The decision letter does not appear in the electronic version of my paper for reasons of data protection.

Characterization in humans of *in vitro* leukocyte maximal telomerase activity capacity (mTAC) and association with stress.

Karin de Punder¹, Christine Heim^{1,2}, Ingo Przesdzing³, Pathik D. Wadhwa^{4,5,6,7,8} & Sonja Entringer^{1,6,8}

¹Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health (BIH), Institute of Medical Psychology, Berlin, Germany

²Department of Biobehavioral Health, College of Health and Human Development, Pennsylvania State University, Pennsylvania, USA

³Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health (BIH), Department of Experimental Neurology and Center for Stroke Research Berlin (CSB), Berlin, Germany

Departments of ⁴Psychiatry & Human Behavior, ⁵Obstetrics & Gynecology, ⁶Pediatrics, ⁷Epidemiology, and the ⁸Development, Health and Disease Research Program, University of California, Irvine, School of Medicine, Irvine, CA, USA

Keywords: Telomerase activity, Leukocytes, Stress, Phytohaemagglutinin (PHA)

Address correspondence to:

Sonja Entringer
Institute for Medical Psychology
Charité University Medicine
Luisenstr. 57
10117 Berlin
Germany
Tel: +49 30 450 529 222
Fax: +49 30 450 529 990
Email: sonja.entringer@charite.de

OR

Pathik D. Wadhwa, MD, PhD.

Professor of Psychiatry & Human Behavior, Obstetrics & Gynecology, Pediatrics, and Epidemiology
Director, UC Irvine Development, Health and Disease Research Program

University of California, Irvine, School of Medicine
3117 Gillespie Neuroscience Research Facility (GNRF)
837 Health Sciences Road
Irvine, CA 92697
Tel: (949) 824-8238
Fax: (949) 824-8218
E-mail: pwadhwa@uci.edu

Summary

The goal of the present study was to develop and validate a measure of *maximal telomerase activity capacity (mTAC)* for use in human studies of telomere biology and, to determine its association with measures of stress and stress responsivity. The study was conducted in a population of 28 healthy young women and men who were assessed serially across two separate days, at multiple time points, and in response to a standardized laboratory stressor. Venous blood was collected at each of these multiple assessments and an *in vitro* mitogen challenge (PHA supplemented with interleukin(IL)-2) was used to stimulate telomerase activity in leukocytes. After first establishing the optimal post-stimulation time course to characterize mTAC, we determined the within-subject stability and the between-subject variability of mTAC. The major findings of our study are as follow: 1) The optimal time point to quantify human leukocyte mTAC appears to be at 72 hours after mitogen stimulation; 2) mTAC exhibits substantial within-subject stability (correlations ranged between r 's 0.68 – 0.82) and between-subject variability, with a high intra-class coefficient (ICC = 0.70), indicating greater between-subject relative to within-subject variability; 3) mTAC is not influenced by situational factors including time of day, cortisol, acute stress exposure, and immune cell distribution in the pre-stimulation blood sample; and 4) a significant proportion of the between-subject variability in mTAC is associated with measures of stress and stress responsivity (mTAC is lower in subjects reporting higher levels of perceived (chronic) stress and exhibiting higher psychophysiological stress reactivity). Based collectively on these findings, it appears that mTAC, as proposed and operationalized, empirically meets the key criteria to represent a potentially useful individual difference measure of telomerase activity capacity of human leukocytes.

1. Introduction

1.1. Overview

A substantial and converging body of epidemiological, clinical and experimental evidence supports a fundamental role for the telomere biology system in the maintenance of DNA and cellular integrity, with important implications for health and disease risk across a wide range of age-related disorders [1-4]. The telomere biology system comprises of two closely interlinked components – the length of telomeres (TL; non-coding double-stranded repeats of guanine-rich tandem DNA sequences and shelterin protein structures that cap the ends of linear chromosomes), and the activity of telomerase (the reverse transcriptase enzyme that adds telomeric DNA to telomeres) [5, 6]. The majority of human epidemiological and clinical studies of the role of telomere biology in health and disease risk have focused largely on the telomere length component of this system. Relatively few studies have considered the role of telomerase. Because the expression and

activity of telomerase constitutes a *critical* and *complementary* (i.e., non-redundant) component of the functional integrity of the telomere biology system, we suggest that it may be important to incorporate telomerase-related measures in studies. This, then, leads to the question of how to optimally quantify this component of the system in human epidemiological or clinical studies. The relatively few studies that have included measures of telomerase have typically measured telomerase expression or activity under basal conditions or in terms of its acute (short-term) response to systemic challenges (see, e.g., [7]). It is, however, challenging to reliably quantify or interpret these data for several reasons. Firstly, telomerase is typically not expressed, or expressed only at very low levels, in most resting cells (including immune cells) [8]. Secondly, telomerase levels may vary as a function of cell cycle stage and other factors [9]. And thirdly, differences or changes in telomerase may reflect either the direct effects of states or conditions that stimulate telomerase expression (e.g., infection), or the secondary (compensatory/counter-regulatory) adaptations to states or conditions that reduce telomere length [10]. Thus, in contrast to assessment of basal telomerase, we suggest it may be more informative to assess telomerase expression or activity in cells in response to a standardized stimulus, such as a mitogen challenge, in well controlled *ex vivo* conditions. The potential advantage of this approach is that it may bypass the above-mentioned limitations to provide an indicator of individual differences in the *capacity* of the telomere biology system to respond to an ecologically relevant challenge. This, in principle, is similar to the information provided about the integrity of the glucose homeostatic system by serial measures of blood glucose and insulin in response to ingestion of a standardized glucose load. Accordingly, we propose that maximal telomerase activity capacity (hereinafter referred to as mTAC) may represent an individual difference measure that, by itself, or in combination with measures of telomere length, could prove to be potentially and particularly informative in studies of telomere biology, health and disease risk. In order for a construct such as mTAC to serve as a potentially useful individual difference measure, it should meet at least two criteria: high within-subject stability, and substantial between-subject variability. Other considerations in this specific context include selection of the optimal cell population, challenge, its dose, and the time course that best captures telomerase expression/activity.

Given the importance of stress and stress physiology as a likely regulator of the telomere biology system, the observation that many of the previous studies of human telomerase have been conducted in the context of stress, and the specific interest of our own research program in the effects of stress and stress biology on health and disease risk, we additionally elected to determine within-subject stability and between-subject variability of mTAC in the context of stress. Accordingly, we used an extensively validated acute psychosocial stress challenge (the Trier Social Stress Test; TSST) to examine the within-subject stability of mTAC, and we used reliable and previously-validated measures of perceived (chronic) psychological stress and individual differences in physiological stress reactivity to examine the between-subject variability of mTAC. Since several physiological systems (including stress biology) exhibit chronobiological regulation,

we also considered the influences of situational factors such as time of day of blood sample collection and cortisol concentration on the within-subject stability of mTAC, and of factors such as sleep quality, chronotype and the slope of the diurnal change in cortisol on the between-subject variability of mTAC. Last, we determined the extent to which mTAC may be driven by the composition of immune cell subpopulations before mitogen stimulation.

1.2. *The telomere biology system*

The telomere biology system is a highly evolutionarily conserved system that plays a central role in maintaining the integrity of the genome and cell. As mentioned above, telomere biology refers to the structure and function of two closely interlinked entities—*telomeres*, non-coding double-stranded repeats of guanine-rich tandem DNA sequences and shelterin protein structures that cap the ends of linear chromosomes [11, 12], and *telomerase*, the reverse transcriptase enzyme that adds telomeric DNA to telomeres [1, 5, 6].

Because DNA polymerase is unable to fully replicate the 3' end of the DNA strand, telomeres lose approximately 30-150 base pairs (bp) with each cell division and eventually reach a critical short length, resulting in decreased recruitment of shelterin proteins to form the protective internal nucleotide loops, which, in turn, leads to cellular senescence. Once cells become senescent, they exhibit a variety of (epi)genetic and morphological changes that result in loss of cell and tissue function. Shortened telomeres have been linked to several age-related disease risk factors, disease prevalence and progression [1-4, 13-18] and early mortality [19, 20]. Moreover, some recent reports have suggested a *causal* role for telomeres in the etiology of many of these adverse health outcomes [21].

The enzyme telomerase is a ribonucleoprotein consisting of a RNA component (TR or TERC) and a catalytic protein domain (TERT). Conventional DNA polymerase machinery is unable to fully replicate the ends of linear chromosomes. The enzyme telomerase utilizes its own template to add short TG-rich repeats to chromosome ends, thus reversing or attenuating their gradual erosion at each round of replication [6, 22]. Telomerase is regulated by epigenetic, translational and posttranslational mechanisms [23]. Its expression varies during development, cell cycle stage, and across cell types [9, 24]. Typically, telomerase activity is diminished or absent in most adult somatic cells, with the exception of cells with a strong potential for division, such as germ cells, stem cells of proliferating tissues, and activated immune cells [24]. Of particular relevance here, it is well established that activated lymphocytes express high telomerase levels [8, 9, 25, 26]. This up-regulation of telomerase is believed to prevent immune cell senescence and facilitate a fast and profound clonal cell expansion. Very occasionally cells bypass the cellular senescence and DNA damage signaling pathways described above to constitutively express high levels of telomerase, which is a characteristic feature in about 90% of all malignancies [27]. This feature of telomerase biology is beyond the scope of the current study, and is therefore not addressed here.

Telomerase not only maintains telomere length but also preserves healthy cell function. Telomerase promotes proliferation of resting stem cells, modulates signaling pathways during embryogenesis and normal adult tissue genesis, protects cellular proliferation capacity and survival under conditions of cellular stress [28], and gets excluded from the nucleus to co-localize with mitochondria to protect mitochondrial DNA and function [29, 30]. Thus, because the effects of telomerase on cellular function extend beyond, and are uncoupled from, those of telomere lengthening [31], measures of telomerase expression and activity could, in addition to telomere length, provide valuable insight regarding the role of the telomere biology system in health and disease risk.

1.3. Basal leukocyte telomerase

Studies of the association of basal leukocyte telomerase with health, disease risk (other than cancer) and conditions such as stress have largely yielded inconsistent results. Several but not all studies suggest that basal telomerase expression/activity appears to be up-regulated in autoimmune disorders, with differences between active and inactive disease [32], and between early- and advanced-stage disease [33]. Findings in the context of psychiatric disorders such as depression, schizophrenia, and posttraumatic stress disorder (PTSD) appear to be more heterogeneous [34-38]. The same pattern of mixed findings is evident in the context of chronic stress exposure, with some suggesting suppression [7, 39-41], and others suggesting stimulatory effects of stress on telomerase activity [10, 42]. This heterogeneity may be a consequence of the fact that telomerase is regulated in response to various factors such as cell cycle stage [9], stress hormones [7], and inflammation [43, 44], which, in turn, may introduce bias based on time of day of blood sample collection [45], current infections [43, 44], physical exercise [46], or acute stress exposure status [7]. In addition, elevations in basal telomerase may reflect the counter-regulatory (compensatory, secondary) adaptations to states/conditions that reduce telomere length. For example, high telomerase in conjunction with shorter TL may be indicative of a physiologically stressed system [10]. Lastly, because telomerase is normally expressed at very low levels in resting cells, the lower limit of detection of many telomerase assays may place constraints in terms of reliable quantification [47].

1.4. Stimulated leukocyte telomerase

Hiyama *et al* [8] were the first to describe the up-regulation of telomerase in leukocytes after *in vitro* mitogen stimulation. Telomerase was detectable in very low levels in isolated peripheral blood mononuclear cells (PBMCs), but increased up to 300-1000 fold over a 1 week-period in cultured T-cells stimulated with PHA and IL-2, and increased up to 30 fold over a 1 week-period in cultured B-cells stimulated with Pokeweed mitogen. Similarly, Yamada *et al* [26] used PHA and IL-2 to stimulate PBMCs and observed over a 96 h period that telomerase activity started to increase after 24 h and peaked at 72 h.

The observation by Son *et al* [48] that the capacity for induction of telomerase activity in T- or B-cells after *in vitro* stimulation varied significantly across subjects but did not change as a function of subject's age (the age of study participants ranged between birth and 94 years age) provides the first indication that this measure may reflect a stable individual difference characteristic. This observation has since been replicated by other studies [49, 50]. However, to date, only few studies have characterized the association of measures of stimulation-induced telomerase with health and disease states [50-54]. Broadly, these studies have reported that the mitogen stimulated leukocyte telomerase response appears to be attenuated in subjects with autoimmune conditions such as systemic lupus erythematosus [52], rheumatoid arthritis [50, 53], and unchanged in conditions such as atopic dermatitis [54] and chronic hepatitis B infection [51].

Two studies have examined the association of stimulated human leukocyte telomerase with chronic stress. An *in vitro* experiment by Choi *et al* [55] modeled the effect of chronic stress exposure and found that co-exposure of human T-cells stimulated with PHA and IL-2 to exogenous cortisol reduced hTERT transcription and inhibited telomerase production across a 3-day period. In a study of caregivers of patients with Alzheimer's disease (stress exposure group) and age-matched controls, basal PBMC and T-cell telomerase levels were increased in caregivers compared to controls, while no differences were observed across a 3-day period in antigen-stimulated telomerase levels (with anti-CD3/CD28 monoclonal antibody) [42].

1.5. Maximal Telomerase Activity Capacity (mTAC) measure: key considerations

Based on the findings and considerations discussed above, we propose that mTAC may represent an individual difference measure that, by itself, or in combination with measures of telomere length, could prove to be potentially and particularly informative in studies of telomere biology, health and disease risk. Several considerations guided our development of this measure for possible use in human epidemiological and clinical studies, including the following questions: What is the optimal cell population, optimal challenge, its optimal dose, and the optimal time course that captures cell capacity for telomerase expression/activity?

We selected PBMCs as the cell population of choice because it is relatively easy and convenient to obtain blood samples; telomere length in human studies is most commonly measured in peripheral leukocytes; and telomere length and telomerase activity are closely associated with immune function due to their important role in lymphocyte development, differentiation and replicative capacity. We, therefore, reasoned that quantifying mTAC in PBMCs could be particularly informative, and this also would increase the feasibility of the use of this measure in clinical studies (as opposed to a single more specific immune cell types such as T-cells).

PBMCs were stimulated with phytohemagglutinin (PHA, 10 µg/ml) supplemented with IL-2 (50 units/ml), because it induces lymphocyte (especially T-cell) proliferation. In humans, telomerase is typically

expressed in lymphocytes by induction of activating and proliferating pathways (e.g., T-cell clonal expansion during a viral infection). We selected the 10 µg/ml PHA dose because maximal lymphocyte proliferation is known to occur at this dose [56], and we added IL-2 because of its importance in the maintenance of immune cells, including T-cells, B-cells and NK cells, and because it has a synergistic effect on the proliferation of T cells and on leukocyte telomerase expression [26, 57].

To establish the optimal time course for characterization of mTAC we considered 3 investigational parameters: *i*) telomerase activity, as quantified by the TeloTAGGG Telomerase PCR ELISA plus assay; *ii*) changes in immune cell populations; and *iii*) changes in cell viability. Previous studies have reported that across a time period spanning 8 consecutive days after mitogen stimulation, peak leukocyte hTERT mRNA expression occurs between 3 to 5 days after stimulation [50]. Most studies to date have used a 3-day incubation protocol with and without the addition of IL-2, and have used PHA, [49, 58] CD3-monoclonal antibodies [42, 50, 53, 54], or PMA/Ionomycin [48, 52] as a stimulant to induce telomerase activity or hTERT mRNA expression in leukocytes or leukocyte subsets. Our criteria for characterizing the ‘optimal’ time was to determine the balance between the time required to enable quantification of the maximal telomerase activity capacity without producing major alterations in immune cell subpopulations and cell viability (as these changes could, in and of themselves, influence telomerase production and activity [48, 59]).

1.6. Goals

Thus, to summarize, the primary goals of our study were to establish the optimal time course to characterize human leukocyte maximal telomerase activity capacity (mTAC), determine the within-subject stability and between-subject variability of mTAC, and determine its association with measures of stress and stress responsivity. To determine the within-subject stability of mTAC, we considered the possible influences of situational factors such as time of day of blood sample collection, cortisol concentration, and acute social stress exposure. To determine the between-subject variability of mTAC, we considered the possible influence of factors including sleep quality, chronotype, the slope of the diurnal change in cortisol, perceived (chronic) stress, and psychophysiological stress responsivity. Last, we also determined the extent to which changes in mTAC may be driven by (or a reflection of) changes in immune cell subpopulations following mitogen stimulation.

2. Materials and Methods

2.1. Participants

The study population comprised 28 young adults (14 women and 14 men, mean age 23.8 ± 3.3 (SD) years) recruited through announcements at universities in Berlin, Germany, and 5 additional adults who

donated a single blood sample for the mTAC optimal time course assessment. All subjects were healthy (no acute or chronic health problems), as ascertained by self-report and confirmed by a brief clinical examination. All subjects also were non-smokers and medication free, except for oral contraceptives (all female subjects were oral contraceptives users). Mean body mass index (BMI) was 20.9 ± 1.7 (SD). The study was approved by the medical ethics committee of Charité Universitätsmedizin Berlin and was conducted in accordance with the Declaration of Helsinki. All subjects provided written informed consent before participation.

2.2. Study protocol

Subjects reported to our research laboratory on two occasions: Test Day 1 – the “chronobiology day”; and Test Day 2 – the “Trier Social Stress Test (TSST) day” (see **Figure 1**), with an approximately one week interval between the first and second study visits. On both study days, participants were asked to refrain from strenuous physical exercise and consumption of alcoholic beverages 24 h prior the study assessments.

On Test Day 1, participants were asked to come to the laboratory after an overnight fast. Venous blood and saliva samples were collected at three time points across the day (at 8 a.m., 1 p.m., and 7 p.m.). Participants were asked to refrain from eating at least 2.5 h before the 1 p.m. and 7 p.m. blood draws. On Test Day 2, participants were asked to come to the laboratory at 2 p.m. and to refrain from eating at least 2.5 h before that time. Participants were then exposed to the Trier Social Stress Test (TSST). This laboratory-based protocol consists of a free speech task and a mental arithmetic task of 15 min duration performed in front of an audience and a camera [60]. This protocol is among the most extensively validated tasks for the induction and assessment of acute psychophysiological stress responses, and has been found to induce significant endocrine (cortisol, ACTH), and autonomic nervous system (as indexed by heart rate and blood pressure) responses in the vast majority of subjects [61]. Blood and saliva samples were obtained before (-10 min) and after (+30 min, +90 min) the TSST, whereas heart rate was measured continuously (Actiheart, CamNtech) during this period. Saliva was collected by placing cotton swabs (Salivettes, Sarstedt, Nümbrecht, Germany) in the participant’s mouth for 2 min, and salivettes were immediately frozen at -80° C. Blood was collected in citrate containing vacutainers (BD Vacutainer) and processed as described below for various assays.

2.3. Questionnaires

Participants completed several standardized and previously-validated questionnaires. Perceived (chronic) psychological stress over the past month was quantified using the 10-item version of the *Perceived Stress Scale (PSS)* [62]. Chronotype was assessed using the *Morningness Eveningness Questionnaire (MEQ)* [63]. Sleep quality (global sleep quality score) was assessed using the *Pittsburgh Sleep Quality Index (PSQI)* [64].

2.4. Cortisol assays

Salivary cortisol was determined using a commercial ELISA kit (Salimetrics, 1-3002), with a sensitivity of 0.007 µg/dL. Intra-assay and inter-assay coefficients of variability were 5.9% and 7.3%, respectively.

2.5. Maximal telomerase activity capacity (mTAC)

Peripheral blood mononuclear cell (PBMC) isolation: PBMCs were isolated from peripheral blood collected in citrate tubes through a standard Ficoll protocol with SepMate tubes (Stemcell Technologies), and live cells were counted with a hemocytometer. Cells were then frozen using a Mister Frosty freezing chamber at a concentration of 1×10^7 cells/ml in freezing media of fetal bovine serum (HyClone, Thermo Scientific) and 10% DMSO (Sigma-Aldrich) until subsequent stimulation experiments (for more details see Supplemental material)

Mitogen stimulation protocol: Cells were thawed and washed in RPMI 1640 medium (Gibco®) containing 10% fetal bovine serum (HyClone, Thermo Scientific). Exactly 1×10^6 PBMCs were stimulated with phytohemagglutinin (PHA) (10 µg/ml) (Sigma-Aldrich) (to induce lymphocyte proliferation) [56], supplemented with IL-2 (50 units/ml) (Sigma-Aldrich) (for the maintenance of immune cell, including T-cells, B-cells and NK cells, and its synergistic effect on lymphocyte proliferation and leukocyte telomerase expression [26, 57]). Cells were cultured at 37°C and 5% CO₂ for a period of 8 days for the time course study, and for a period of 72 h for the other studies. For each individual and each study day, cells obtained at each of the three different time points were thawed and stimulated at the same time (in the afternoon) and were cultured in the same 12-well plate (for more details see Supplemental material).

Telomerase activity: Telomerase can be quantified by determining the presence of hTERT and/or other proteins of the telomerase enzyme complex, or by measuring the activity of the telomerase enzyme. Telomerase activity can be assessed by using the telomeric repeat amplification protocol (TRAP) assay, which relies on the ability of the telomerase enzyme to add telomere repeats to specifically designed oligonucleotide primers [65]. Since the induction of telomerase activity in lymphocytes has been previously determined to occur independent of changes in mRNA or protein expression [23], we elected to quantify activity (instead of expression), and did so using the TeloTAGGG Telomerase PCR ELISA plus kit (Roche), which combines a TRAP assay with detection by ELISA. The linear range of this assay was first assessed using extracts of various cell numbers (stimulated PBMCs). Cell pellets were lysed in ice-cold CHAPS lysis buffer at a concentration of 1000 cells/µl and incubated for 30 min on ice. After incubation, the lysates were centrifuged at 16,000 x g for 20 min at 4°C. An extract corresponding to 1,000 cells (1 µl) was added to each

PCR reaction. Amplification products were hybridized to a DIG-labelled telomeric repeat-specific probe bound to a streptavidin-coated 96-well plate. The binding reaction was detected with an anti-DIG-peroxidase antibody, visualized by a color reaction product and quantified photometrically. The absorbance of each sample was measured at 450 nm reading against the blank (reference wavelength 620 nm). Each negative sample was obtained by heat treatment (30 min at 94 °C). The relative telomerase activities of the samples were determined by comparing the signal from the sample to the signal obtained using a known amount of control template.

In order to determine the optimal stimulation time to characterize human leukocyte mTAC, we conducted an *in vitro* time course study using isolated PBMCs from 5 volunteers. PBMCs were stimulated with PHA (10 µg/ml) and IL-2 (50 units/ml) and cultured in 8 separate wells for 8 days. On each day, PBMC telomerase activity was assessed and (stimulated) PBMCs were stained with fluorescent antibodies against different immune cell subpopulations and analyzed using flow cytometry (see section 2.6.).

2.6. FACS Flow Cytometry

In a subset of 13 participants, PBMCs from each of the three assessment time points across Test Day 1 were stained with fluorescent antibodies against different immune cell subpopulations before *in vitro* stimulation. PBMCs were incubated for 15 min at 4°C, with antibodies against CD45 (Vioblue) CD3 (APC), CD4 (PerCp), CD8 (APC-*io770*), CD14 (FITC), CD19 (PE-*Vio770*), CD16 (PE), and CD56 (PE) (Miltenyl Biotec). Cells were washed in PBS containing 0.5% bovine serum albumin, fixed in 2% formaldehyde, and analyzed using a BD FACSCanto II with FACSDiva 6.1.3. The data analysis was performed with FlowJo 10.1r5.

2.7. Statistics

Repeated measures ANOVAS were computed to assess the effects of time (day) on change in mTAC. Greenhouse–Geisser corrections were applied, and adjusted results are reported. Pearson’s correlations were used to determine the within-subject variability of mTAC. In addition, the intra-class coefficient (ICC) of mTAC was computed to determine the proportion of total variation in mTAC that is attributable to variation between subjects relative to variation within subjects. The area-under-the-curve with respect to ground (AUCg) was used to quantify total mTAC and cortisol output (see formula 2 in [66]). The slope of the diurnal change in cortisol occurring on Test Day 1 was calculated by fitting a linear regression line, which predicted the cortisol values from morning to evening. Repeated measures ANOVAS were computed to assess the effect of time on change in cortisol and heart rate. Heart rate was measured prior to the TSST (-5 min, with an average value of 5 min duration), during the TSST (average value of 15 min duration), and +40 min after the beginning of the TSST (average value of 5 min duration). Measures of psychophysiological stress reactivity

were determined by calculating the percent change in cortisol from before to 30 min after the TSST, and the percent heart rate increase from resting state to average heart rate measured during the TSST phase. mTAC may be influenced by the distribution of immune cell subpopulations present in the sample before stimulation. Thus, we examined whether mTAC varies as a function of variation in immune cell subpopulations. Immune cell subpopulations in pre-mitogen stimulated blood and mTAC (i.e., following mitogen stimulation) were characterized in samples from each of the three time points across Test Day 1. Linear mixed models were used to predict telomerase activity as a function of each immune cell type percent by including the repeatedly measured mTAC as the outcome and each immune cell type percent as varying covariates. This model assessed whether total PBMC telomerase activity was, in part, a function of certain cell subpopulation frequencies while accounting for within-subject autocorrelations. For all analyses, p -values $< .05$ were considered significant. Data analysis was performed using SPSS statistical software (SPSS 23.0, Inc., Chicago, IL, USA).

3. Results

3.1. Aim 1: Optimal time course to characterize the maximal telomerase activity capacity (mTAC) response

As shown in **Figure 2a**, maximal telomerase activity was observed on day 3, 4 and 5 of the time course study. Concurrently, the flow cytometry data indicate that changes in immune cell subtypes (specifically percentage of CD8+ T cells) started occurring on and after day 4 (**Figure 2b**), and that the amount of cell debris/lysis (as determined by the FCS/SSC signal) also started to increase from day 4 onwards. Based collectively on these 3 sets of results (maximal telomerase activity, minimal change in immune cell subtypes, and minimal cell degradation) we ascertained that a 3-day (72 h) period represents the optimal time point at which to determine PBMC mTAC in response to PHA/IL-2 mitogen stimulation challenge.

3.2. Aim 2: Within-subject stability of mTAC

3.2.1. Chronobiological influences on mTAC

mTAC (i.e., PBMC telomerase activity 72 h after mitogen stimulation) was not significantly different across participants' blood samples collected serially in the morning, afternoon and evening (8 a.m., 1 p.m., 7 p.m.; main effect time: $F_{1.5, 40.3} = .48$, $p = .58$, $\eta^2 = .018$, see **Figure 3a**), suggesting no influence of chronobiology on mTAC. As expected, cortisol concentrations in these serially-collected blood samples exhibited a diurnal rhythm and declined significantly from morning to evening ($F_{1.1, 29.8} = 44.1$, $p < .001$, $\eta^2 =$

.62). Cortisol concentration at these three time points was not associated with any of the corresponding mTAC measures. Finally, the within-subject correlations of mTAC levels from these serially-obtained blood samples was strong and ranged between r 's 0.68 – 0.82 (see **Table 1**). The intra-class coefficient of mTAC across the day (proportion of total variation in mTAC that is attributable to variation between subjects relative to variation within subjects) was 0.7, indicating substantially greater between-subject variability relative to within-subject across time variability.

3.2.2. Acute stress exposure and mTAC

As expected, exposure to the TSST produced significant changes in cortisol (-10, +30, +90 min, main effect time: $F_{1.5, 40.3} = 35.9$, $p < .001$, $\eta^2 = .57$) and heart rate (average heart rate before, during, and after TSST, main effect time: $F_{1.6, 36.6} = 80.7$, $p < .001$, $\eta^2 = .78$). There was, however, no difference in the mTAC levels between the serially-collected blood samples obtained *before* (1 sample) and *after* TSST exposure (2 samples) (main effect time: $F_{2, 48} = 1.4$, $p = .26$, $\eta^2 = .054$, see **Figure 3b**), suggesting that on average, mTAC was not influenced by acute stress exposure.

3.3. Aim 3: Between-subject variability of mTAC: age, sex, BMI, sleep quality and chronotype

The average of the mTAC (AUCg) measures from the serially-collected blood samples across the course of the day was not significantly associated with participant age ($r = -.27$, $p = .17$, $R^2 = .073$), BMI ($r = -.11$, $p = .60$, $R^2 = .012$) or sex ($r = -.047$, $p = .81$, $R^2 = .002$), and also was not related with either sleep quality (PSQI global sleep quality score, $r = -.21$, $p = .29$, $R^2 = .044$), chronotype (MEQ score, $r = -.10$, $p = .63$, $R^2 = .01$) or the slope of the diurnal change in cortisol ($r = -.19$, $p = .35$, $R^2 = .036$).

3.4. Aim 3: Between-subject variability of mTAC: Perceived stress and individual differences in physiological stress reactivity

3.4.1. Perceived psychological stress and mTAC

Participants' perceived (chronic) stress score (PSS) was inversely associated with average mTAC (AUCg) (Figure 4a, ($r = -.34$, $p = .08$, $R^2 = .12$). Since the PSS is not a diagnostic instrument and there are no clinical cut-off values, we used our study population-based tertiles to categorize our participants into low, moderate and high stress groups. When we compared the high perceived (chronic) stress group (upper tertile PSS, mean score = 23 ± 2.8) with the low/medium perceived (chronic) stress group (lower 2 tertiles PSS, mean score = 12 ± 3.2) there was a significant group effect of PSS on mTAC ($F_{1, 25} = 5.6$, $p = .026$, $\eta^2 = .18$), with individuals exposed to high perceived stress exhibiting significantly lower mTAC (**Figure 4b**).

3.4.2. Biological stress-reactivity and mTAC

Next, we examine the association of individual differences in physiological stress reactivity (endocrine (cortisol) and autonomic (heart rate) responses to the TSST) with average mTAC (AUCg). One subject had a cortisol response (percent change from before to 30 min after the TSST) more than 3 SD above the mean and was therefore excluded from the analysis. The cortisol and heart rate responses to the TSST were moderately inter-correlated ($r = .60$, $p = .002$, $R^2 = .36$).

The cortisol response to the TSST was negatively related to mTAC (AUCg) (**Figure 5a**, $r = -.57$, $p = .004$, $R^2 = .32$). The heart rate response also was negatively – although not statistically significantly – associated with mTAC (AUCg) (**Figure 5b**, $r = -.34$, $p = .12$, $R^2 = .12$).

3.5. Immune cell composition and mTAC

Some studies have suggested that basal [47] and stimulated [59] telomerase activities can vary by immune cell subtype. We, therefore, determined whether our measure of PBMC mTAC was influenced by the distribution of immune cell subpopulations present in the blood sample before mitogen stimulation. On Test Day 1, before *in vitro* stimulation, immune cell type frequencies were obtained from the three serially-collected blood samples across the day in a subgroup of study participants ($n=13$) [7]. The *B*-coefficients of the linear mixed models presented in **Table 2** reflect the change in mTAC (change in relative telomerase activity, RTA) for each percent change in cell type of the different immune cell subtypes modeled simultaneously for all three time points. None of the relationships were significant, suggestion that this measure of mTAC is not altered as a function of immune cell type distribution present in the sample before stimulation.

4. Discussion

Based on the results of the time course study (*i.e.*, determination of maximal telomerase activity with minimal change in immune cell subtypes and minimal cellular degradation), we determined that a *72h post mitogen stimulation* period appears to represent the *optimal time point* at which to quantify human leukocyte mTAC. Using this optimal time point, our next set of results determined that the mTAC measure exhibits *substantial within-subject stability* across time and does not appear to be influenced by age, sex, BMI, and situational factors including time of day, cortisol concentration, acute stress exposure, and immune cell distribution prior to mitogen stimulation. The mTAC measure also exhibits *substantial between-subject variability*. Particularly, participants' perceived (chronic) stress level over the past 1-month period accounted for 12% of the variation in the mTAC measure, with a 25% difference in mTAC between subjects reporting high compared to medium or low levels of perceived (chronic) stress. Moreover, individual differences in a

key stress-related trait – psychophysiological stress responsivity (cortisol response to TSST exposure) – accounted for as much as 32% of the variation in mTAC. Finally, the mTAC intra-class coefficient (ICC) indicated substantially greater between-subject relative to within-subject variability. Collectively, these findings support our premise that the maximal leukocyte telomerase activity capacity construct (mTAC) empirically meets the criteria to represent a potentially useful individual difference measure.

Our finding that telomerase activity peaks between 3 to 5 days after mitogen stimulation replicates previous studies [26, 59]. However, given the changes we observed after day 4 in immune cell subtypes and cell debris/lysis, we suggest that a 3-day period represents the optimal time point at which to characterize human leukocyte mTAC in response to PHA/IL-2 mitogen stimulation challenge.

In our study, mTAC was not associated with participants' age. While our study was limited in exploring this relationship given the relatively restricted age range in our cohort, we note that most previous studies of stimulated immune cell telomerase activity responses also did *not* find a significant effect of age on this measure [48-50], suggesting that mTAC may reflect a stable individual difference characteristic. Discrepancies between these findings and those of the 2 studies that did find an effect of age on stimulated telomerase activity [51, 58] may perhaps be accounted for by differences in the mitogen stimulations protocols and the immune cell subpopulations that were assessed. In our study there was also no significant effect on mTAC of time of day, basal cortisol concentrations assessed across the day, sex, BMI, sleep quality, chronotype and the slope of the diurnal change in cortisol. To the best of our knowledge these factors have not been investigated/reported in previous studies using stimulated measures of telomerase expression/activity.

Our findings suggest that individual differences in mTAC are independent of percentages of different immune cell subsets present before *in vitro* stimulation. However, we cannot completely rule out the contribution of differences in immune cell composition. For example, here, we analyzed the total fraction of CD8+ T-cells and did not discriminate between naïve, effector and memory CD8+ cytotoxic T-cells. It has been shown in mice that effector and memory CD8+ T-cells express higher levels of telomerase activity after acute viral infection [67], indicating that the percentage of naïve and virus-specific CD8+ T-cells can influence mTAC after stimulation *in vitro*. In addition, mTAC could be influenced by the percentage of senescent cells. Lin *et al* [47] showed that in total PBMC, senescent CD28- T-cells had lowest telomerase activity and shortest telomere length. CD28 signaling is required for optimal telomerase up-regulation, indicated by the paralleled loss of telomerase activity and CD28 expression in T-cells after chronic antigen stimulation *in vitro* [59].

Previous studies have reported that chronically stressed individuals exhibit lower levels of *unstimulated* telomerase activity (e.g., [7, 39-41]). Our results indicate that individuals reporting high levels of perceived (chronic) psychological stress display lower stimulated telomerase activity capacity (mTAC). A

previous study by Damjanovic *et al* [42], showed no differences in induced PBMC or T-cell telomerase activity levels (anti-CD3/CD28 monoclonal antibody stimulation for 72 h) between caregivers of Alzheimer's patients (stress exposure group) and controls. This discrepancy with our findings may perhaps be explained by differences in the mitogen stimulation protocols that were employed, or by the fact that stress exposure in the caregivers was operationalized using a depression scale, whereas we assessed perceived stress over the past month using the perceived stress scale.

Our protocol particularly stimulates cell types involved in cellular immunity, the part of the immune system that is initially affected by chronic stress exposure [68]. Chronic stress induces a shift in the production of type 1 cytokines towards type 2 cytokines, and it has been associated with blunted mitogen-induced lymphocyte proliferation and mitogen induced IL-2 production [69], both activators of signaling pathways stimulating telomerase activity [9, 57]. Therefore, the observation that higher levels of perceived (chronic) psychological stress are associated with lower mTAC may be a reflection of impairment in cell-mediated immunity. In addition, chronic stress exposure is associated with a higher level of oxidative stress [40, 70]. Oxidative stress decreases telomerase activity and induces senescence (or apoptosis) via DNA damage-induced activation of the p53 pathway, probably by causing erosion of telomeres or perhaps by inducing nuclear exclusion of hTERT into mitochondria [29, 30]. Senescent CD28- T-cells show impaired up-regulation of telomerase activity after antigen stimulation [59], and could therefore inhibit mTAC.

In our study population, a larger cortisol increase in response to an acute psychosocial stressor was related to lower mTAC. An individual's biological stress reactivity measure can be considered a stress-related trait and has been studied in the context of several stress-related pathologies and conditions [71]. Previous findings showed that greater cortisol responses to an acute stressor was associated with shorter telomere length, as was higher overnight urinary free cortisol levels and flatter daytime cortisol slope [72]. In line with our finding, exposure to exogenous cortisol has been shown to inhibit telomerase production in human T-cells stimulated with PHA and IL-2 [55]. Glucocorticoids place a limit on the maximal activity of the immune system, modulate inflammatory gene transcription [73], and can either through direct action or through the modulation of cytokine release (type 1 cytokines towards type 2 cytokines) inhibit lymphocyte proliferation and NK cell activity [69], thereby influencing telomerase regulatory pathways [9, 57]. Cortisol levels measured during the chronobiology test day (test day 1) were not associated to mTAC. However, mTAC was negatively related to the cortisol response to acute stress. It is likely that individuals experiencing persistently exaggerated stress responses are exposed to greater levels of cortisol over longer periods of time, which may have important consequences on immune system function. In a study by Epel *et al*, in which basal/unstimulated telomerase activity was measured soon after a standardized stress test, the cortisol response was positively associated with telomerase activity [7]. As discussed by these authors, telomerase activity levels may change dynamically in response to stress, likely as a protective functional response to

protect telomeric regions from stress-induced acute increases in biological stress mediators such as cortisol or oxidative stress. Two key differences between this study and ours is that the Epel *et al* study [7] did not use a mitogen challenge to stimulate telomerase expression and did not characterize *maximal* capacity of cells to express telomerase (*i.e.*, mTAC) but measured only the short-term response (up to 90 min) to acute social stress, whereas we assessed telomerase in response to a mitogen challenge and across a 3-day period of time.

In the present study, individuals displaying higher cortisol responses also showed higher heart rate responses, indicating a coupling between these stress-related endocrine and autonomic systems.

Catecholamines, which are normally produced in response to acute stress by the adrenal medulla and postganglionic sympathetic nerve fibers, activate β 2-receptors present on immune cells. *In vitro* studies suggest that β 2-receptor stimulation by catecholamines activates the cAMP signaling pathway, inhibiting mitogen-stimulated T-cell proliferation and NK-cells activation [74]. Hence, catecholamines could also inhibit leukocyte mTAC after stimulation *in vitro*.

Strengths of the present study include the following: We first established the optimal mTAC time course and stimulation conditions before the development and use of the *in vitro* study stimulation protocol to address the study aims. Furthermore, to the best of our knowledge, our study is the first to systematically investigate the within- and between-subject stability of stimulated telomerase activity responses (mTAC) in women and men with respect to chronobiological and stress-related influences.

Limitations of our study include the unavailability of measures of telomere length in our study subjects, the relatively modest sample size, and the restricted range of age and BMI in our study subjects. Any interpretation of sex effects is limited by the fact that all the women in our study were using oral contraceptives. While the absence of chronobiological or acute stress effects on mTAC in our study may provide a justification for future studies against requiring multiple mTAC measures over the course of a day or in response to an acute challenge, we note, based on the within-subject correlations of serial measures of mTAC indicating a shared variance of approximately 50-65%, that measurement issues such as single *vs* average of multiple samples remain an important consideration. Lastly, because mTAC was measured in isolated circulating immune cells (PBMCs) in the context of immune cell activation and clonal T-cell expansion, it is unclear whether this measure also reflects the capacity of hematopoietic stem and progenitor cells to induce a telomerase activation response.

As discussed earlier, only a few studies to date have examined the characteristics of stimulation-induced telomerase responses in the context of health and disease states [50-54]. Based on our findings in support of the premise that the maximal leukocyte telomerase activity capacity construct (mTAC) may represent a potentially useful individual difference measure, one future research direction will be to determine the prognostic value of mTAC, independently, and in combination with measures of telomere length, with respect to health, disease risk and mortality. Also, further mechanistic studies are warranted to better

understand the contribution of specific immune cell populations such as memory and senescent T-cells, circulating levels of other stress-related biological mediators such as catecholamines, oxidative stress and pro-inflammatory cytokines, on mTAC. Future studies should also explore the characteristics of mTAC in hematopoietic stem cells [75], such as those obtained from umbilical cord blood.

In conclusion, because mTAC exhibits high within-subject stability, is not influenced by situational circumstances, and shows substantial between-subject variability in relation to stress-related traits and states, this *in vitro* measure of telomerase activity may represent a potentially useful individual difference measure in studies of telomere biology and health and disease.

5. Author's contributions

KdP coordinated the study, participated in the design of the study, carried out the laboratory assays, participated in the data analyses, and drafted the manuscript. CH participated in the data analysis and provided editorial assistance. IP performed the flow cytometry measurements and data analysis. PDW participated in the analysis and interpretation of study findings, drafted portions of the manuscript, and provided editorial assistance. SE conceived of and designed the study, participated in the analysis and interpretation of study findings, drafted portions of the manuscript, and provided final editorial oversight.

6. Competing interests

We have no competing interests to declare.

7. Funding

Funding for this study was provided by grants from Neurocure (Innovation Projects) and the Federal Ministry of Education and Research (01KR1301A). PDW and SE's efforts were also supported, in part, by US PHS (NIH) grants R01 HD-065825, R01 HD-060628 and R01 AG-050455.

8. References

- [1] Blackburn, E. H., Epel, E. S. & Lin, J. 2015 Human telomere biology: A contributory and interactive factor in aging, disease risks, and protection. *Science* **350**, 1193-1198. (DOI:10.1126/science.aab3389).
- [2] D'Mello, M. J., Ross, S. A., Briel, M., Anand, S. S., Gerstein, H. & Pare, G. 2015 Association between shortened leukocyte telomere length and cardiometabolic outcomes: systematic review and meta-analysis. *Circ Cardiovasc Genet* **8**, 82-90. (DOI:10.1161/CIRCGENETICS.113.000485).

- [3] Haycock, P. C., Heydon, E. E., Kaptoge, S., Butterworth, A. S., Thompson, A. & Willeit, P. 2014 Leucocyte telomere length and risk of cardiovascular disease: systematic review and meta-analysis. *Bmj* **349**, g4227. (DOI:10.1136/bmj.g4227).
- [4] Zhao, J., Miao, K., Wang, H., Ding, H. & Wang, D. W. 2013 Association between telomere length and type 2 diabetes mellitus: a meta-analysis. *PLoS one* **8**, e79993. (DOI:10.1371/journal.pone.0079993).
- [5] Blackburn, E. H. 2000 Telomere states and cell fates. *Nature* **408**, 53-56. (DOI:10.1038/35040500).
- [6] Blackburn, E. H. 2005 Telomeres and telomerase: their mechanisms of action and the effects of altering their functions. *FEBS letters* **579**, 859-862. (DOI:10.1016/j.febslet.2004.11.036).
- [7] Epel, E. S., Lin, J., Dhabhar, F. S., Wolkowitz, O. M., Puterman, E., Karan, L. & Blackburn, E. H. 2010 Dynamics of telomerase activity in response to acute psychological stress. *Brain, behavior, and immunity* **24**, 531-539. (DOI:10.1016/j.bbi.2009.11.018).
- [8] Hiyama, K., Hirai, Y., Kyoizumi, S., Akiyama, M., Hiyama, E., Piatyszek, M. A., Shay, J. W., Ishioka, S. & Yamakido, M. 1995 Activation of telomerase in human lymphocytes and hematopoietic progenitor cells. *Journal of immunology* **155**, 3711-3715.
- [9] Buchkovich, K. J. & Greider, C. W. 1996 Telomerase regulation during entry into the cell cycle in normal human T cells. *Mol Biol Cell* **7**, 1443-1454.
- [10] Zalli, A., Carvalho, L. A., Lin, J., Hamer, M., Erusalimsky, J. D., Blackburn, E. H. & Steptoe, A. 2014 Shorter telomeres with high telomerase activity are associated with raised allostatic load and impoverished psychosocial resources. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 4519-4524. (DOI:10.1073/pnas.1322145111).
- [11] Blackburn, E. H. & Gall, J. G. 1978 A tandemly repeated sequence at the termini of the extrachromosomal ribosomal RNA genes in Tetrahymena. *J Mol Biol* **120**, 33-53.
- [12] Moyzis, R. K., Buckingham, J. M., Cram, L. S., Dani, M., Deaven, L. L., Jones, M. D., Meyne, J., Ratliff, R. L. & Wu, J. R. 1988 A highly conserved repetitive DNA sequence, (TTAGGG)_n, present at the telomeres of human chromosomes. *Proceedings of the National Academy of Sciences of the United States of America* **85**, 6622-6626.
- [13] Coimbra, B. M., Carvalho, C. M., Moretti, P. N., Mello, M. F. & Belangero, S. I. 2017 Stress-related telomere length in children: A systematic review. *J Psychiatr Res* **92**, 47-54. (DOI:10.1016/j.jpsychires.2017.03.023).
- [14] Entringer, S., Buss, C. & Wadhwa, P. D. 2012 Prenatal stress, telomere biology, and fetal programming of health and disease risk. *Science signaling* **5**, pt12. (DOI:10.1126/scisignal.2003580).
- [15] Kordinas, V., Ioannidis, A. & Chatzipanagiotou, S. 2016 The Telomere/Telomerase System in Chronic Inflammatory Diseases. Cause or Effect? *Genes (Basel)* **7**. (DOI:10.3390/genes7090060).
- [16] Mathur, M. B., Epel, E., Kind, S., Desai, M., Parks, C. G., Sandler, D. P. & Khazeni, N. 2016 Perceived stress and telomere length: A systematic review, meta-analysis, and methodologic considerations for advancing the field. *Brain, behavior, and immunity* **54**, 158-169. (DOI:10.1016/j.bbi.2016.02.002).
- [17] Shalev, I., Entringer, S., Wadhwa, P. D., Wolkowitz, O. M., Puterman, E., Lin, J. & Epel, E. S. 2013 Stress and telomere biology: a lifespan perspective. *Psychoneuroendocrinology* **38**, 1835-1842. (DOI:10.1016/j.psyneuen.2013.03.010).
- [18] Stone, R. C., Horvath, K., Kark, J. D., Susser, E., Tishkoff, S. A. & Aviv, A. 2016 Telomere Length and the Cancer-Atherosclerosis Trade-Off. *PLoS Genet* **12**, e1006144. (DOI:10.1371/journal.pgen.1006144).
- [19] Cawthon, R. M., Smith, K. R., O'Brien, E., Sivatchenko, A. & Kerber, R. A. 2003 Association between telomere length in blood and mortality in people aged 60 years or older. *Lancet* **361**, 393-395. (DOI:10.1016/S0140-6736(03)12384-7).
- [20] Kimura, M., Hjelmborg, J. V., Gardner, J. P., Bathum, L., Brimacombe, M., Lu, X., Christiansen, L., Vaupel, J. W., Aviv, A. & Christensen, K. 2008 Telomere length and mortality: a study of leukocytes in elderly Danish twins. *American journal of epidemiology* **167**, 799-806. (DOI:10.1093/aje/kwm380).
- [21] Gadalla, S. M., Cawthon, R., Giri, N., Alter, B. P. & Savage, S. A. 2010 Telomere length in blood, buccal cells, and fibroblasts from patients with inherited bone marrow failure syndromes. *Aging (Albany NY)* **2**, 867-874. (DOI:10.18632/aging.100235).

- [22] Shore, D. & Bianchi, A. 2009 Telomere length regulation: coupling DNA end processing to feedback regulation of telomerase. *EMBO J* **28**, 2309-2322. (DOI:10.1038/emboj.2009.195).
- [23] Liu, K., Hodes, R. J. & Weng, N. 2001 Cutting edge: telomerase activation in human T lymphocytes does not require increase in telomerase reverse transcriptase (hTERT) protein but is associated with hTERT phosphorylation and nuclear translocation. *Journal of immunology* **166**, 4826-4830.
- [24] Forsyth, N. R., Wright, W. E. & Shay, J. W. 2002 Telomerase and differentiation in multicellular organisms: turn it off, turn it on, and turn it off again. *Differentiation; research in biological diversity* **69**, 188-197. (DOI:10.1046/j.1432-0436.2002.690412.x).
- [25] Bodnar, A. G., Kim, N. W., Effros, R. B. & Chiu, C. P. 1996 Mechanism of telomerase induction during T cell activation. *Experimental cell research* **228**, 58-64. (DOI:10.1006/excr.1996.0299).
- [26] Yamada, O., Motoji, T. & Mizoguchi, H. 1996 Up-regulation of telomerase activity in human lymphocytes. *Biochim Biophys Acta* **1314**, 260-266.
- [27] Jafri, M. A., Ansari, S. A., Alqahtani, M. H. & Shay, J. W. 2016 Roles of telomeres and telomerase in cancer, and advances in telomerase-targeted therapies. *Genome Med* **8**, 69. (DOI:10.1186/s13073-016-0324-x).
- [28] Sahin, E. & Depinho, R. A. 2010 Linking functional decline of telomeres, mitochondria and stem cells during ageing. *Nature* **464**, 520-528. (DOI:10.1038/nature08982).
- [29] Haendeler, J., Hoffmann, J., Diehl, J. F., Vasa, M., Spyridopoulos, I., Zeiher, A. M. & Dimmeler, S. 2004 Antioxidants inhibit nuclear export of telomerase reverse transcriptase and delay replicative senescence of endothelial cells. *Circulation research* **94**, 768-775. (DOI:10.1161/01.RES.0000121104.05977.F3).
- [30] Saretzki, G. 2009 Telomerase, mitochondria and oxidative stress. *Experimental gerontology* **44**, 485-492. (DOI:10.1016/j.exger.2009.05.004).
- [31] Blasco, M. A. 2002 Telomerase beyond telomeres. *Nat Rev Cancer* **2**, 627-633. (DOI:10.1038/nrc862).
- [32] Katayama, Y. & Kohriyama, K. 2001 Telomerase activity in peripheral blood mononuclear cells of systemic connective tissue diseases. *The Journal of rheumatology* **28**, 288-291.
- [33] Invernizzi, P., Bernuzzi, F., Lleo, A., Pozzoli, V., Bignotto, M., Zermiani, P., Crosignani, A., Battezzati, P. M., Zuin, M., Podda, M., et al. 2014 Telomere dysfunction in peripheral blood mononuclear cells from patients with primary biliary cirrhosis. *Digestive and liver disease : official journal of the Italian Society of Gastroenterology and the Italian Association for the Study of the Liver* **46**, 363-368. (DOI:10.1016/j.dld.2013.11.008).
- [34] Jergovic, M., Tomicevic, M., Vidovic, A., Bendelja, K., Savic, A., Vojvoda, V., Rac, D., Lovric-Cavar, D., Rabatic, S., Jovanovic, T., et al. 2014 Telomere shortening and immune activity in war veterans with posttraumatic stress disorder. *Progress in neuro-psychopharmacology & biological psychiatry* **54**, 275-283. (DOI:10.1016/j.pnpbp.2014.06.010).
- [35] Lindqvist, D., Epel, E. S., Mellon, S. H., Penninx, B. W., Revesz, D., Verhoeven, J. E., Reus, V. I., Lin, J., Mahan, L., Hough, C. M., et al. 2015 Psychiatric disorders and leukocyte telomere length: Underlying mechanisms linking mental illness with cellular aging. *Neuroscience and biobehavioral reviews* **55**, 333-364. (DOI:10.1016/j.neubiorev.2015.05.007).
- [36] Porton, B., Delisi, L. E., Bertisch, H. C., Ji, F., Gordon, D., Li, P., Benedict, M. M., Greenberg, W. M. & Kao, H. T. 2008 Telomerase levels in schizophrenia: a preliminary study. *Schizophrenia research* **106**, 242-247. (DOI:10.1016/j.schres.2008.08.028).
- [37] Simon, N. M., Smoller, J. W., McNamara, K. L., Maser, R. S., Zalta, A. K., Pollack, M. H., Nierenberg, A. A., Fava, M. & Wong, K. K. 2006 Telomere shortening and mood disorders: preliminary support for a chronic stress model of accelerated aging. *Biol Psychiatry* **60**, 432-435. (DOI:10.1016/j.biopsych.2006.02.004).
- [38] Wolkowitz, O. M., Mellon, S. H., Epel, E. S., Lin, J., Reus, V. I., Rosser, R., Burke, H., Compagnone, M., Nelson, J. C., Dhabhar, F. S., et al. 2012 Resting leukocyte telomerase activity is elevated in major depression and predicts treatment response. *Mol Psychiatry* **17**, 164-172. (DOI:10.1038/mp.2010.133).
- [39] Daubenmier, J., Lin, J., Blackburn, E., Hecht, F. M., Kristeller, J., Maninger, N., Kuwata, M., Bacchetti, P., Havel, P. J. & Epel, E. 2012 Changes in stress, eating, and metabolic factors are related to changes in

- telomerase activity in a randomized mindfulness intervention pilot study. *Psychoneuroendocrinology* **37**, 917-928. (DOI:10.1016/j.psyneuen.2011.10.008).
- [40] Epel, E. S., Blackburn, E. H., Lin, J., Dhabhar, F. S., Adler, N. E., Morrow, J. D. & Cawthon, R. M. 2004 Accelerated telomere shortening in response to life stress. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 17312-17315. (DOI:10.1073/pnas.0407162101).
- [41] Epel, E. S., Lin, J., Wilhelm, F. H., Wolkowitz, O. M., Cawthon, R., Adler, N. E., Dolbier, C., Mendes, W. B. & Blackburn, E. H. 2006 Cell aging in relation to stress arousal and cardiovascular disease risk factors. *Psychoneuroendocrinology* **31**, 277-287. (DOI:10.1016/j.psyneuen.2005.08.011).
- [42] Damjanovic, A. K., Yang, Y., Glaser, R., Kiecolt-Glaser, J. K., Nguyen, H., Laskowski, B., Zou, Y., Beversdorf, D. Q. & Weng, N. P. 2007 Accelerated telomere erosion is associated with a declining immune function of caregivers of Alzheimer's disease patients. *Journal of immunology* **179**, 4249-4254.
- [43] Akbar, A. N. & Vukmanovic-Stejić, M. 2007 Telomerase in T lymphocytes: use it and lose it? *Journal of immunology* **178**, 6689-6694.
- [44] Akiyama, M., Yamada, O., Hideshima, T., Yanagisawa, T., Yokoi, K., Fujisawa, K., Eto, Y., Yamada, H. & Anderson, K. C. 2004 TNF α induces rapid activation and nuclear translocation of telomerase in human lymphocytes. *Biochemical and biophysical research communications* **316**, 528-532. (DOI:10.1016/j.bbrc.2004.02.080).
- [45] Chen, W. D., Wen, M. S., Shie, S. S., Lo, Y. L., Wo, H. T., Wang, C. C., Hsieh, I. C., Lee, T. H. & Wang, C. Y. 2014 The circadian rhythm controls telomeres and telomerase activity. *Biochemical and biophysical research communications* **451**, 408-414. (DOI:10.1016/j.bbrc.2014.07.138).
- [46] Zietzer, A., Buschmann, E. E., Janke, D., Li, L., Brix, M., Meyborg, H., Stawowy, P., Jungk, C., Buschmann, I. & Hillmeister, P. 2016 Acute physical exercise and long-term individual shear rate therapy increase telomerase activity in human peripheral blood mononuclear cells. *Acta Physiol (Oxf)*. (DOI:10.1111/apha.12820).
- [47] Lin, J., Epel, E., Cheon, J., Kroenke, C., Sinclair, E., Bigos, M., Wolkowitz, O., Mellon, S. & Blackburn, E. 2010 Analyses and comparisons of telomerase activity and telomere length in human T and B cells: insights for epidemiology of telomere maintenance. *Journal of immunological methods* **352**, 71-80. (DOI:10.1016/j.jim.2009.09.012).
- [48] Son, N. H., Murray, S., Yanovski, J., Hodes, R. J. & Weng, N. 2000 Lineage-specific telomere shortening and unaltered capacity for telomerase expression in human T and B lymphocytes with age. *Journal of immunology* **165**, 1191-1196.
- [49] Benko, A. L., Olsen, N. J. & Kovacs, W. J. 2012 Estrogen and telomerase in human peripheral blood mononuclear cells. *Molecular and cellular endocrinology* **364**, 83-88. (DOI:10.1016/j.mce.2012.08.012).
- [50] Thewissen, M., Linsen, L., Geusens, P., Raus, J. & Stinissen, P. 2005 Impaired activation-induced telomerase activity in PBMC of early but not chronic rheumatoid arthritis patients. *Immunol Lett* **100**, 205-210. (DOI:10.1016/j.imlet.2005.03.007).
- [51] Fan, X. G., Huang, Y., Tang, F. Q. & Yi, H. 2000 Telomerase activity of peripheral blood lymphocytes in patients with chronic hepatitis B. *Immunol Lett* **73**, 7-11.
- [52] Fritsch, R. D., Shen, X., Illei, G. G., Yarboro, C. H., Prussin, C., Hathcock, K. S., Hodes, R. J. & Lipsky, P. E. 2006 Abnormal differentiation of memory T cells in systemic lupus erythematosus. *Arthritis and rheumatism* **54**, 2184-2197. (DOI:10.1002/art.21943).
- [53] Fujii, H., Shao, L., Colmegna, I., Goronzy, J. J. & Weyand, C. M. 2009 Telomerase insufficiency in rheumatoid arthritis. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 4360-4365. (DOI:10.1073/pnas.0811332106).
- [54] Wu, K., Volke, A., Lund, M., Bang, K. & Thestrup-Pedersen, K. 1999 Telomerase activity is spontaneously increased in lymphocytes from patients with atopic dermatitis and correlates with cellular proliferation. *Journal of dermatological science* **22**, 24-30.
- [55] Choi, J., Fauce, S. R. & Effros, R. B. 2008 Reduced telomerase activity in human T lymphocytes exposed to cortisol. *Brain, behavior, and immunity* **22**, 600-605. (DOI:10.1016/j.bbi.2007.12.004).

- [56] Darko, D. F., Gillin, J. C., Risch, S. C., Bulloch, K., Golshan, S., Tasevska, Z. & Hamburger, R. N. 1989 Mitogen-stimulated lymphocyte proliferation and pituitary hormones in major depression. *Biol Psychiatry* **26**, 145-155.
- [57] Kawauchi, K., Ihjima, K. & Yamada, O. 2005 IL-2 increases human telomerase reverse transcriptase activity transcriptionally and posttranslationally through phosphatidylinositol 3'-kinase/Akt, heat shock protein 90, and mammalian target of rapamycin in transformed NK cells. *Journal of immunology* **174**, 5261-5269.
- [58] Lin, Y., Damjanovic, A., Metter, E. J., Nguyen, H., Truong, T., Najarro, K., Morris, C., Longo, D. L., Zhan, M., Ferrucci, L., et al. 2015 Age-associated telomere attrition of lymphocytes in vivo is co-ordinated with changes in telomerase activity, composition of lymphocyte subsets and health conditions. *Clinical science* **128**, 367-377. (DOI:10.1042/CS20140481).
- [59] Valenzuela, H. F. & Effros, R. B. 2002 Divergent telomerase and CD28 expression patterns in human CD4 and CD8 T cells following repeated encounters with the same antigenic stimulus. *Clin Immunol* **105**, 117-125.
- [60] Kirschbaum, C., Pirke, K. M. & Hellhammer, D. H. 1993 The 'Trier Social Stress Test'--a tool for investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology* **28**, 76-81. (DOI:119004).
- [61] Dickerson, S. S. & Kemeny, M. E. 2004 Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research. *Psychol Bull* **130**, 355-391. (DOI:10.1037/0033-2909.130.3.355).
- [62] Cohen, S. & Williamson, G. 1988 Perceived stress in a probability sample of the United States. In *The social psychology of health: Claremont Symposium on applied social psychology* (eds. S. Spacapan & S. Oskamp). Newbury Park, CA, Sage.
- [63] Horne, J. A. & Ostberg, O. 1976 A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol* **4**, 97-110.
- [64] Buysse, D. J., Reynolds, C. F., 3rd, Monk, T. H., Berman, S. R. & Kupfer, D. J. 1989 The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* **28**, 193-213.
- [65] Greider, C. W. & Blackburn, E. H. 1989 A telomeric sequence in the RNA of Tetrahymena telomerase required for telomere repeat synthesis. *Nature* **337**, 331-337. (DOI:10.1038/337331a0).
- [66] Pruessner, J. C., Kirschbaum, C., Meinlschmid, G. & Hellhammer, D. H. 2003 Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology* **28**, 916-931.
- [67] Hathcock, K. S., Kaech, S. M., Ahmed, R. & Hodes, R. J. 2003 Induction of telomerase activity and maintenance of telomere length in virus-specific effector and memory CD8+ T cells. *Journal of immunology* **170**, 147-152.
- [68] Segerstrom, S. C. & Miller, G. E. 2004 Psychological stress and the human immune system: a meta-analytic study of 30 years of inquiry. *Psychol Bull* **130**, 601-630. (DOI:10.1037/0033-2909.130.4.601).
- [69] Bauer, M. E., Vedhara, K., Perks, P., Wilcock, G. K., Lightman, S. L. & Shanks, N. 2000 Chronic stress in caregivers of dementia patients is associated with reduced lymphocyte sensitivity to glucocorticoids. *J Neuroimmunol* **103**, 84-92.
- [70] Colaianna, M., Schiavone, S., Zotti, M., Tucci, P., Morgese, M. G., Backdahl, L., Holmdahl, R., Krause, K. H., Cuomo, V. & Trabace, L. 2013 Neuroendocrine profile in a rat model of psychosocial stress: relation to oxidative stress. *Antioxid Redox Signal* **18**, 1385-1399. (DOI:10.1089/ars.2012.4569).
- [71] Miller, G. E., Chen, E. & Zhou, E. S. 2007 If it goes up, must it come down? Chronic stress and the hypothalamic-pituitary-adrenocortical axis in humans. *Psychol Bull* **133**, 25-45. (DOI:10.1037/0033-2909.133.1.25).
- [72] Tomiyama, A. J., O'Donovan, A., Lin, J., Puterman, E., Lazaro, A., Chan, J., Dhabhar, F. S., Wolkowitz, O., Kirschbaum, C., Blackburn, E., et al. 2012 Does cellular aging relate to patterns of allostasis? An examination of basal and stress reactive HPA axis activity and telomere length. *Physiol Behav* **106**, 40-45. (DOI:10.1016/j.physbeh.2011.11.016).
- [73] Barnes, P. J. & Adcock, I. M. 2009 Glucocorticoid resistance in inflammatory diseases. *Lancet* **373**, 1905-1917. (DOI:10.1016/S0140-6736(09)60326-3).

- [74] Elenkov, I. J., Wilder, R. L., Chrousos, G. P. & Vizi, E. S. 2000 The sympathetic nerve--an integrative interface between two supersystems: the brain and the immune system. *Pharmacol Rev* **52**, 595-638.
- [75] Hordyjewska, A., Popiolek, L. & Horecka, A. 2015 Characteristics of hematopoietic stem cells of umbilical cord blood. *Cytotechnology* **67**, 387-396. (DOI:10.1007/s10616-014-9796-y).

Tables

Table 1. Within-subject correlations (Pearson’s correlations) between mTAC levels from samples obtained at the three different time points during Test Day 1.

| | mTAC 8 a.m. | mTAC 1 p.m. | mTAC 7 p.m. |
|------------------------|------------------------|------------------------|------------------------|
| mTAC 8 a.m. | 1 | .82** | .73** |
| mTAC 1 p.m. | .82** | 1 | .68** |
| mTAC 7 p.m. | .73** | .68** | 1 |

**indicates p-value < .001

Table 2. Results of mixed models predicting mTAC by immune cell subtype percentages (%) on Test Day 1.

| mTAC on Test Day 1 | | |
|---------------------------|---------------|----------------|
| <i>Cell type (%)</i> | <i>B (SE)</i> | <i>p-value</i> |
| CD4+ T-cells | -.64 (.42) | .14 |
| CD8+ T-cells | .92 (.80) | .26 |
| B-cells | .28 (1.65) | .87 |
| NK-cells | .15 (.82) | .86 |
| NKT-cells | -.10 (.31) | .98 |
| Monocytes | .52 (.53) | .34 |

Figure captions

Figure 1. Study protocol: All subjects reported to the laboratory on two occasions (Test Day 1: “Chronobiology day”; Test Day 2: “TSST day”); mTAC: maximal telomerase activity capacity; FACS: fluorescence-activated cell scanning (flow cytometry); TSST: Trier Social Stress Test.

Figure 2. Time course experiment: a) mean relative PBMC telomerase activity (\pm SD) of 5 individual measured at day 1-8 after *in vitro* PHA/IL-2 stimulation and b) mean percentages (\pm SD) of immune cell subpopulations (calculated as percentages of CD45+ cells) of 5 individuals measured at day 1-5 after PHA/IL-2 stimulation *in vitro*; RTA: relative telomerase activity.

Figure 3. mTAC levels measured on both test days: a) mean mTAC levels (\pm SE) in stimulated PBMCs isolated at 8 p.m., 1 p.m. and 7 p.m. during Test Day 1, and b) before (-10 min) and after (+30 min, +90 min) the TSST (Test Day 2); RTA: relative telomerase activity.

Figure 4. Perceived (chronic) stress score and mTAC: a) Correlation of mTAC (AUCg) with the perceived stress score (PSS), and b) mTAC levels in individuals reporting high and low/moderate stress. RTA: relative telomerase activity.

Figure 5. Correlation of mTAC (AUCg) with a) the cortisol response, and b) the heart rate response (%) to the TSST. RTA: relative telomerase activity.

Figure 1




Figure 2




Figure 3




Figure 4






Figure 5



Supplemental material

Supplemental material: Protocol for isolation, stimulation and lysis of PBMC

Materials (listed or equivalent)

BD Vacutainer 9NC (sodium citrate) 6ml (2x), Becton Dickinson (Franklin Lakes, USA), 366575
Falcon 50mL conical centrifuge tubes, Corning (Corning, USA), 352070
Falcon 15mL conical centrifuge tubes, Corning (Corning, USA), 352097
Sepmate-50 (or Sepmate-15 for smaller volumes of blood), Stemcell Technologies (Cologne, Germany), 15450
Ficoll, GE Healthcare Life Sciences (Little Chalfont, UK), 17-144-002
PBS ready to use (sterile), VWR International (Radnor, USA), K812-500 ml
Fetal bovine serum (FBS), HyClone (South Logan, USA), SH3007102
Dimethyl sulfoxide (DMSO, sterile), Sigma-Aldrich (Saint Louis, USA), D2438-5X10ML
Hemocytometer, Sigma-Aldrich (Saint Louis, USA), Z359629-1EA
Trypan blue, Sigma-Aldrich (Saint Louis, USA), T8154
RPMI-1640 medium, Gibco, Life Science Technologies (Carlsbad, USA), 21875-034
Phytohaemagglutinin (PHA), Sigma-Aldrich (Saint Louis, USA), L8754-1MG
Interleukin(IL)-2, Sigma-Aldrich (Saint Louis, USA), I7908-10KU
TeloTAGGG Telomerase PCR ELISA plus kit, Roche (Basel, Switzerland), 12013789001

Equipment

Biosafety cabinet
Centrifuge
Mr. Frosty freezing container with isopropanol
-80 °C freezer and/or liquid nitrogen freezer

Solutions

Freezing medium: FBS containing 10% DMSO
Culture medium: RPMI-1640 medium containing 10% FBS

PBMC isolation protocol

1. Procedures should be performed in a biosafety cabinet under sterile conditions.
2. Retrieve whole blood.

3. Dilute 1:1 with sterile PBS.
4. Fill Sepmate-50 tube with 15 ml Ficoll.
5. Carefully pipet the diluted blood on the Ficoll layer.
6. Spin down for 10 min at 1200 x g with the brake on.
7. Poor the supernatant within 2 sec in new 50 ml falcon tube and add PBS until 40 ml.
8. Spin down for 10 min at 300 x g (brake on), remove the supernatant and tap the bottom of tube to loosen the cell pellet.
9. Fill up the tube with 10 ml of PBS.
10. Spin down at 200 x g for 10 min (brake on), remove the supernatant and tap the bottom of tube to loosen the cell pellet.
11. Dissolve the pellet in exactly 1 ml of sterile PBS (mix thoroughly by resuspending).
12. Count cells and calculate the amount of cells/ml using a hemacytometer.
13. Spin down cells at 250 x g for 10 min (brake on), remove the supernatant and tap the bottom of tube to loosen the cell pellet.
14. Dissolve pellet in freezing medium (1 ml of freezing medium / 1×10^7 cells).
15. Divide the cells dissolved in the freezing medium over 2 (or more) sterile cryovials and freeze using a Mister Frosty at $-80\text{ }^{\circ}\text{C}$. Optional: Cryovials can be transferred to liquid nitrogen tank after 1 day.

PBMC stimulation protocol

1. Procedures should be performed in a biosafety cabinet under sterile conditions.
2. Thaw cells (quickly) in a $37\text{ }^{\circ}\text{C}$ water bath and transfer each thawed cell suspension to an appropriately labeled 15 ml tube and add pre-warmed ($37\text{ }^{\circ}\text{C}$) culture medium diluting 10 x the original volume.
3. Centrifuge at 250 x g for 10 minutes at RT, remove the supernatant and tap the bottom of tube to loosen the cell pellet.
4. Resuspend cells in exactly 1 ml of culture medium.
5. Count live cells using a hemacytometer.
6. Dilute the cells in the appropriate amount of culture medium in order to obtain a cell concentration of exactly 1×10^6 cells/ml.
7. Pipet 1 ml of cell suspension (1×10^6 cells/ml) in the well of a 12-well plate; so each well contains 1×10^6 cells.
8. Add 5 μl of IL-2 (10 KU/ml) per ml/well (final concentration = 50 units/ml).
9. Dissolve 1 mg of PHA (Sigma-Aldrich) in 1 ml of PBS (stock concentration = 1 mg/ml).

10. Add 10 μ l of PHA (1 mg/ml) per ml/well (final concentration = 10 μ g/ml) and incubate 72 hours at 37°C and 5% CO₂.

Lysis protocol

1. Procedures should be performed under nuclease free conditions.
2. Take out lysis reagent (Roche TeloTAGGG Telomerase PCR ELISA plus kit) from freezer and let it thaw to 4 °C. Keep on ice.
3. Label the appropriate amount of sterile (RNase free) 1.5 ml Eppendorf tubes.
4. Remove cells from the wells with a 1000 μ l pipet (resuspend a few times to remove aggregates) and place into Eppendorf tubes.
5. Spin down tubes at 300 x g for 10 min, RT, remove the supernatant and tap the bottom of tube to loosen the cell pellet.
6. Dissolve pellet in exactly 1 ml of PBS and carefully resuspend and/or shortly vortex.
7. Count live cells with hemacytometer.
8. Transfer a volume corresponding to exactly 200,000 cells to the corresponding Eppendorf tube:

$$\text{required cell volume } (\mu\text{l}) = \text{total volume } (1000 \mu\text{l}) * 200,000 / \text{total amount of cells}$$

9. Cool centrifuge and spin down the 200.000 cells for 5 min, 3000 x g at 4 °C, remove supernatant and keep pellet on ice.
10. Dissolve pellet in 200 μ l of cold lysis buffer (Roche TeloTAGGG Telomerase PCR ELISA plus kit).
11. Resuspend pellet and shortly vortex, keep on ice for 30 min.
12. Centrifuge for 20 min, 16.000 x g at 4 °C.
13. Carefully remove 175 μ l of the supernatant, transfer in a sterile labeled (RNase free) 1.5 ml Eppendorf tube and store at -80 °C.
14. Telomerase activity is measured in the lysates following the instructions of the Roche TeloTAGGG Telomerase PCR ELISA plus kit.

Curriculum Vitae

My curriculum vitae does not appear in the electronic version of my paper for reasons of data protection.

List of Publications

Original and Review Articles

Karin de Punder, Christine Heim, Ingo Przesdzing, Pathik D. Wadhwa, Sonja Entringer (2018). Characterization in humans of *in vitro* leukocyte maximal telomerase activity capacity (mTAC) and association with stress. *Philos Trans R Soc Lond B Biol Sci.* 5;373(1741) pii: 20160441. Impact factor: 5,846

Sonja Entringer, **Karin de Punder**, Claudia Buss, Pathik D. Wadhwa (2018). The fetal programming of telomere biology hypothesis: An update. *Philos Trans R Soc Lond B Biol Sci.* 5;373(1741) pii: 20170151. Impact factor: 5,846

Henner F. Farin, Ingrid Jordens, Mohammed H. Mosa, Onur Basak1, Jeroen Korving, Daniele V. F. Tauriello, **Karin de Punder**, Stephane Angers, Peter J. Peters, Madelon M. Maurice, Hans Clevers (2015). Visualization of the short-range Wnt gradient in the intestinal stem cell niche. *Nature.* 18;530(7590):340-3. Impact factor: 40,137

Karin de Punder, Leo Pruimboom (2015). Stress induces endotoxemia and low-grade inflammation by increasing barrier permeability. *Front. Immunol.* 6:223. Impact factor: 6,429

Leo Pruimboom, **Karin de Punder** (2015). The opioid effects of gluten exorphins: asymptomatic celiac disease. *J Health Popul Nutr.* 33:24. Impact factor: 1,187

Daleke-Schermerhorn MH, Felix T, Soprova Z, Ten Hagen-Jongman CM, Vikström D, Majlessi L, Beskers J, Follmann F, **de Punder K**, van der Wel NN, Baumgarten T, Pham TV, Piersma SR, Jiménez CR, van Ulsen P, de Gier JW, Leclerc C, Jong WS, Luirink J (2014). Decoration of outer membrane vesicles with multiple antigens by using an autotransporter approach. *Appl Environ Microbiol.* 80(18):5854-65. Impact factor: 3,807

Jong WS, Daleke-Schermerhorn MH, Vikström D, Ten Hagen-Jongman CM, de **Punder K**, van der Wel NN, van de Sandt CE, Rimmelzwaan GF, Follmann F, Agger EM, Andersen P, de Gier JW, Luirink J (2014). An autotransporter display platform for the development of multivalent recombinant bacterial vector vaccines. *Microb Cell Fact.* 25;13(1):162. Impact factor: 3,681

Karin de Punder, Leo Pruijboom (2013). The dietary intake of wheat and other cereal grains and their role in inflammation. *Nutrients.* 12;5(3):771-87. Impact factor: 3,550

van der Woude AD, Mahendran KR, Ummels R, Piersma SR, Pham TV, Jiménez CR, **de Punder K**, van der Wel NN, Winterhalter M, Luirink J, Bitter W, Houben EN (2013). Differential detergent extraction of *Mycobacterium marinum* cell envelope proteins identifies an extensively modified threonine-rich outer membrane protein with channel activity. *J Bacteriol.* 195(9):2050-9. Impact factor: 3,143

Wouter SP Jong, Zora Soprova, **Karin de Punder**, Corinne M ten Hagen-Jongman, Samuel Wagner, David Wickström, Jan-Willem de Gier, Peter Andersen, Nicole N van der Wel and Joen Luirink (2012). A structurally informed autotransporter platform for the efficient secretion and display of heterologous proteins. *Microb Cell Fact.* 18;11:85. Impact factor: 3,681

Diane Houben, Caroline Demangel, Jakko van Ingen, Jorge Perez, Lucy Baldeón, Abdallah M. Abdallah, Laxmee Caleechurn, Daria Bottai, Maaïke van Zon, **Karin de Punder**, Tridia van der Laan, Arie Kant, Ruth Bossers-de Vries, Peter Willemsen, Wilbert Bitter, Dick van Soolingen, Roland Brosch, Nicole van der Wel and Peter J. Peters (2012). ESX-1 mediated translocation to the cytosol controls virulence of mycobacteria. *Cell Microbiol.* 14(8):1287-98. Impact factor: 4,554

Eveline M. Weerdenburg, Abdallah M. Abdallah, Suman Mitra, **Karin de Punder**, Nicole N. van der Wel, Steve Bird, Ben J. Appelmelk, Wilbert Bitter, Astrid M. van der Sar (2011). ESX-5-deficient *Mycobacterium marinum* is hypervirulent in adult zebrafish. *Cell Microbiol.* 14(5):728-39. Impact factor: 4,554

Abdallah M. Abdallah, Jovanka Bestebroer, Nigel D. L. Savage, **Karin de Punder**, Maaïke van Zon, Louis Wilson, Cees J. Korbee, Astrid M. van der Sar, Tom H. M. Ottenhoff, Nicole N. van der Wel, Wilbert Bitter, and Peter J. Peters (2011). Mycobacterial secretion systems ESX-1 and ESX-5 play distinct roles in host cell death and inflammasome activation. *J Immunol.* 1;187(9):4744-53. Impact factor: 4,856

Maria H. Daleke, Alessandro Cascioferro, **Karin de Punder**, Roy Ummels, Nicole van der Wel, Peter J. Peters, Joen Luijck, Riccardo Manganelli, Wilbert Bitter (2011). Homologues of the mycobacterial lipase LipY have evolved to ESX-5 substrates by acquiring AN N-terminal PE or PPE domain. *J Biol Chem.* 27;286(21):19024-34. Impact factor: 4,125

Musa Sani, Edith N.G. Houben, Jeroen Geurtsen, Jason Pierson, **Karin de Punder**, Maaïke van Zon, Brigitte Wever, Mamadou Daffe, Ben J. Appelmelk, Wilbert Bitter, Nicole van der Wel, Peter J. Peters (2010). Direct visualization by cryo-EM of the mycobacterial capsular layer: a labile structure containing ESX-1-secreted proteins. *PLoS Pathog.* 5;6(3): e1000794. Impact factor: 6,608

de Witte MA, Jorritsma A, Swart E, Straathof KC, **de Punder K**, Haanen JB, Rooney CM, Schumacher TN (2008). An inducible caspase 9 safety switch can halt cell therapy-induced autoimmune disease. *J Immunol.* 1;180(9):6365-73. Impact factor: 4,856

Boekhoorn K, Sarabdjitsingh A, Kommerie H, **de Punder K**, Schouten T, Lucassen PJ, Vreugdenhil E (2008). Doublecortin (DCX) and doublecortin-like (DCL) are differentially expressed in the early but not late stages of murine neocortical development. *J comp neur.* 1;507(4):1639-52. Impact factor: 3,266

Book Chapters

Sonja Entringer, **Karin de Punder**, Glenn Verner, Pathik D. Wadhwa (2017). Fetal programming of telomere biology: Role of maternal nutrition, obstetric risk factors and suboptimal birth outcomes in Diet, nutrition, and fetal programming, Nutrition and Health pp.569-593. *Springer International Publishing AG.*

Published Abstracts

Karin de Punder, Christine Heim, Ingo Przesdzing, Pathik D. Wadhwa, Sonja Entringer (2017). Characterization of *in vitro* leukocyte maximal telomerase activity capacity (mTAC) as a stress-related measure for human studies. *Psychoneuroendocrinology*. 83S:1-89

Karin de Punder, Judith Overfeld, Peggy Dörr, Katja Dittrich, Sibylle M. Winter, Nina Kubiak, Gergana Karaboycheva, Christine Heim (2017). Maltreatment is associated with elevated C-reactive protein levels in 3 to 5 year-old children. *Brain Behav Immun*. 66:e17-e18

Karin de Punder, Christine Heim, Pathik D. Wadhwa, Sonja Entringer (2017). Characterization of *in vitro* leukocyte maximal telomerase activity capacity (mTAC) in humans: Association with chronic stress exposure and stress-reactivity. *Brain Behav Immun*. 66:e17

Maaïke van Zon, **Karin de Punder**, Astrid van der Sar, Roland Brosch, Rogelio Hernandez Pando, Anita Grootemaat, Wikky Tigchelaar-Gutter, Nicole van der Wel (2017). Subcellular localization of *M. tuberculosis in vivo* and effect of the adaptive immunity. *Ultrastruct Pathol*. 41(1):133

Karin de Punder, Judith Overfeld, Peggy Dorr, Katja Dittrich, Sibylle M. Winter, Nina Kubiak, Gergana Karaboycheva, Christine Heim (2016). Exposure to child maltreatment is associated with elevated stress and immune mediators in children aged 3–5 years. *Psychoneuroendocrinology*. 71S:1–78

Karin de Punder, Christine Heim, Pathik D. Wadhwa, Sonja Entringer (2016). *In vitro* stimulated leukocyte telomerase activity is associated with chronic stress exposure. *Psychoneuroendocrinology*. 71S:1–78

Judith Overfeld, Karin Nadig-Haynes, **Karin de Punder**, Peggy Dorr, Katja Dittrich, Sibylle M. Winter, John-Dylan Haynes, Claudia Buss, Christine Heim (2016). Hippocampal volume in children aged 3–5 years with verified maltreatment exposure. *Psychoneuroendocrinology*. 71S:1–78

Karin de Punder, Christine Heim, Sonja Entringer (2016). Plasma levels of lipopolysaccharide-binding protein in response to psychosocial stress induction: Association with sympathetic nervous system response. *Brain Behav Immun.* 57:e25

Karin de Punder, Christine Heim, Sonja Entringer (2015). Validation of a measure of in-vitro stimulated telomerase expression as a stress-related biomarker for human studies. *Psychoneuroendocrinology.* 61:1–78

Acknowledgements

My acknowledgements do not appear in the electronic version of my paper for reasons of data protection.