

Erratum: “Multi-reference approach to the calculation of photoelectron spectra including spin-orbit coupling” [J. Chem. Phys. 143, 074104 (2015)]

Gilbert Grell, Sergey I. Bokarev, Bernd Winter, Robert Seidel, Emad F. Aziz, Saadullah G. Aziz, and Oliver Kühn

Citation: *The Journal of Chemical Physics* **145**, 089901 (2016); doi: 10.1063/1.4961314

View online: <http://dx.doi.org/10.1063/1.4961314>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/145/8?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Multi-reference approach to the calculation of photoelectron spectra including spin-orbit coupling](#)
J. Chem. Phys. **143**, 074104 (2015); 10.1063/1.4928511

[Perturbational treatment of spin-orbit coupling for generally applicable high-level multi-reference methods](#)
J. Chem. Phys. **141**, 074105 (2014); 10.1063/1.4892060

[On the R -dependence of the spin-orbit coupling constant: Potential energy functions of Xe 2 + by high-resolution photoelectron spectroscopy and ab initio quantum chemistry](#)
J. Chem. Phys. **128**, 234306 (2008); 10.1063/1.2937133

[On the simulation of photoelectron spectra in molecules with conical intersections and spin-orbit coupling: The vibronic spectrum of C H 3 S](#)
J. Chem. Phys. **127**, 104309 (2007); 10.1063/1.2764052

[Ab initio based study of the ArO – photoelectron spectra: Selectivity of spin-orbit transitions](#)
J. Chem. Phys. **112**, 5852 (2000); 10.1063/1.481186



NEW Special Topic Sections

NOW ONLINE
Lithium Niobate Properties and Applications:
Reviews of Emerging Trends

AIP Applied Physics Reviews

Erratum: “Multi-reference approach to the calculation of photoelectron spectra including spin-orbit coupling” [J. Chem. Phys. 143, 074104 (2015)]

Gilbert Grell,¹ Sergey I. Bokarev,^{1,a)} Bernd Winter,² Robert Seidel,² Emad F. Aziz,^{2,3} Saadullah G. Aziz,⁴ and Oliver Kühn¹

¹Institut für Physik, Universität Rostock, Albert-Einstein-Str. 23-24, 18059 Rostock, Germany

²Helmholtz-Zentrum Berlin für Materialien und Energie, Methods for Material Development, Albert-Einstein-Strasse 15, 12489 Berlin, Germany

³Department of Physics, Freie Universität zu Berlin, Arnimalle 14, 14159 Berlin, Germany

⁴Chemistry Department, Faculty of Science, King Abdulaziz University, 21589 Jeddah, Saudi Arabia

(Received 27 July 2016; accepted 5 August 2016; published online 24 August 2016)

[<http://dx.doi.org/10.1063/1.4961314>]

In Ref. 1 we have presented a novel first principles approach to the calculation of photoelectron spectra (PES) and a comparison with the widely used sudden approximation (SA)²⁻⁴ has been performed. Two mistakes have been made in Ref. 1 which are corrected in this erratum. First, within the SA the intensity of a transition is proportional to the absolute square of the Dyson orbital (DO) norm

$$\sigma(\mathcal{E}_k) \propto |\langle \Psi_F^{N-1} | \hat{a} \Psi_I^N \rangle|^2. \quad (1)$$

However, in Figs. 2 and 5 of Ref. 1, the DO norm was not squared. Second, Eq. (10) from the original publication contained a typo and should read

$$|D_{FI}^{\text{SA}}|^2 = |\langle \Psi_F^{N-1} \psi^{\text{el}}(\mathbf{k}) | \hat{d} | \Psi_I^N \rangle|^2 \approx |\langle \Psi_F^{N-1} | \hat{a} \Psi_I^N \rangle|^2. \quad (2)$$

Taking the square alters the discussion on the applicability of the SA to the calculation of X-ray PES of transition metal compounds. We emphasize that the main results of Ref. 1, i.e., the PES obtained via the numerical integration of the DO with the ionized electron wave function, are not affected.

The corrected SA PES are presented in Fig. 1 together with the full calculation results. In contrast to what is claimed in Ref. 1, Fig. 1 shows that the SA predicts the intensities of all peaks with an accuracy that is comparable to the full calculation.




FIG. 1. (Left panel) Calculated and experimental PES of the gas-phase water for 180 eV photon energy. (Right panel) Experimental (2M FeCl₂ aqueous solution) and calculated (for the [Fe(H₂O)₆]²⁺ cluster) core PES for 925 eV photon energy.

^{a)}Electronic mail: sergey.bokarev@uni-rostock.de

Thus, the SA provides a good approximation to the full calculation of L-edge core PES of $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ where the kinetic energy of the outgoing electron is quite high and all transitions have the same character, i.e., being ionizations from the 2p orbitals.

Although the results obtained within the SA for the L-edge PES of $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$ are quite accurate, the SA should be used with caution. If the spectrum contains transitions involving DOs of different character, e.g., localized on metal and ligands, or the kinetic energy of the outgoing electron is low, the applicability of the SA should be investigated on a case to case basis. We would like to stress that the protocol proposed in Ref. 1 that utilizes the numerical integration for the matrix elements allows us to obtain reliable intensities also in cases, where the widely used SA cannot be applied.

We acknowledge financial support by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah (Grant No. D-003-435), Deutsche Forschungsgemeinschaft (Grant No. KU952/10-1), and Collaborative Research Center 1109.

¹G. Grell, S. I. Bokarev, B. Winter, R. Seidel, E. F. Aziz, S. G. Aziz, and O. Kühn, *J. Chem. Phys.* **143**, 074104 (2015).

²R. Manne and T. Åberg, *Chem. Phys. Lett.* **7**, 282 (1970).

³B. T. Pickup, *Chem. Phys.* **19**, 193 (1977).

⁴T. Åberg, *Phys. Rev.* **156**, 35 (1967).