

The solute transport and binding profile of a novel nucleobase cation symporter 2 from the honeybee pathogen *Paenibacillus larvae*

Amanda J. Stoffer-Bittner¹, Candace R. Alexander¹, Douglas W. Dingman², George S. Mourad¹ and Neil P. Schultes³

¹ Department of Biology, Indiana University-Purdue University Fort Wayne, IN, USA

² Department of Entomology, The Connecticut Agricultural Experiment Station, New Haven, CT, USA

³ Department of Plant Pathology & Ecology, The Connecticut Agricultural Experiment Station, New Haven, CT, USA

Keywords

heterologous complementation; nucleobase; nucleobase cation symporter; *Paenibacillus larvae*; PIUacP; transporter

Correspondence

N. P. Schultes, Department of Plant Pathology & Ecology, The Connecticut Agricultural Experiment Station, 123 Huntington St., New Haven, CT 0651, USA
Fax: 203-974-8502
Tel: 203-974-8464
E-mail: Neil.Schultes@ct.gov

(Received 20 February 2018, revised 22 June 2018, accepted 24 June 2018)

doi:10.1002/2211-5463.12488

Here, we report that a novel nucleobase cation symporter 2 encoded in the genome of the honeybee bacterial pathogen *Paenibacillus larvae* reveals high levels of amino acid sequence similarity to the *Escherichia coli* and *Bacillus subtilis* uric acid and xanthine transporters. This transporter is named *P. larvae* uric acid permease-like protein (PIUacP). Even though PIUacP displays overall amino acid sequence similarities, has common secondary structures, and shares functional motifs and functionally important amino acids with *E. coli* xanthine and uric acid transporters, these commonalities are insufficient to assign transport function to PIUacP. The solute transport and binding profile of PIUacP was determined by radiolabeled uptake experiments via heterologous expression in nucleobase transporter-deficient *Saccharomyces cerevisiae* strains. PIUacP transports the purines adenine and guanine and the pyrimidine uracil. Hypoxanthine, xanthine, and cytosine are not transported by PIUacP, but, along with uric acid, bind in a competitive manner. PIUacP has strong affinity for adenine K_m $7.04 \pm 0.18 \mu\text{M}$, and as with other bacterial and plant NCS2 proteins, PIUacP function is inhibited by the proton disruptor carbonyl cyanide *m*-chlorophenylhydrazone. The solute transport and binding profile identifies PIUacP as a novel nucleobase transporter.

Paenibacillus larvae is a Gram-positive bacterium that is the causal agent of American foulbrood of *Apis mellifera* Linnaeus [1,2]. *P. larvae* causes disease on larvae but not adult bees [3,4]. Larvae ingest *P. larvae* endospores (the infectious agent) from contaminated royal jelly and are most susceptible at twelve to thirty-six hours after hatching [4]. Spores germinate and greatly proliferate in the larval midgut prior to infection across the midgut epithelium [5]. Recently, it was shown that spore germination can be triggered by

L-tyrosine and uric acid *in vitro* [6]. Uric acid, a waste product from purine degradation and metabolized proteins, accumulates to high levels in the midgut since the mid and hindgut are not yet connected in young larvae [7]. Once the midgut epithelium is breached, bacterial enter and proliferate in the hemocoel, producing extracellular proteases [8] and toxins [9] leading to cell death and forming a ropey mass of cellular degradation. Depletion of food supplies causes the bacteria to produce highly resistant spores. Worker

Abbreviations

AzG, aza-guanine purine transporter A; COG, cluster of orthologous groups; NCS1, nucleobase cation symporter 1; NCS2, nucleobase cation symporter 2; PIUacP, *P. larvae* uric acid permease-like protein; UapA, uric acid/xanthine transporters A.

bees sense the decay, open the brood chamber, remove the carcass, and become coated with spores that are available to infect subsequent larvae [3,4]. The abundance of cellular metabolites, including nitrogen-rich nucleobases, nucleosides, and nucleotides, in the hemocoel of infected larvae offers *P. larvae* ample opportunity to import valuable nutrients for rapid proliferation.

Nucleobase transporters, ubiquitous among plants, fungi, and bacteria, belong to five independent families. The families include nucleobase cation symporter 1 (NCS1), found in bacteria, fungi, and plants; nucleobase cation symporter 2 (NCS2) family, also known as the nucleobase ascorbate transporter (NAT) family, present in bacteria, fungi, and plants; the closely related aza-guanine-like transporters (AzgA) found in bacteria, fungi, and plants and plant-specific purine permease and ureide permease families [10–16]. The NCS2 family has been the most extensively studied [14]. In plants, NCS2 tend to have broad and mixed solute transport and binding profiles including combinations of xanthine, uric acid, uracil, adenine, guanine, cytosine, and hypoxanthine [17,18 & Schultes and Mourad pers comm]. Plant cells have complex nucleobase biochemistry including *de novo* synthesis, salvage, and catabolic pathways that rely upon different intercellular compartments with diverse environments. In contrast, single-cell microbes primarily use nucleobase transporters for acquisition of these nitrogen-rich compounds from the external environment. Unlike plants, microbes have NCS2 with very restricted solute transport and binding profiles. The fungus *Aspergillus nidulans* contains two NCS2—UapA and UapC—that transport uric acid and xanthine [14] and the adenine–hypoxanthine–guanine transporter AzgA [19,20]. Both UapA and AzgA have been extensively studied with dozens of mutations characterized at the biochemical level [21,22]. *Escherichia coli* contains ten NCS2 with narrow solute transport profiles: the cluster of orthologous group COG2252 contains EcGHXQ (YgfQ) and EcGHXP (YjcD) that transport guanine and hypoxanthine [23] and EcAdeP (YieG) and EcAdeQ (YicO) that transport adenine [23]; the cluster of orthologous group COG2233 contains EcUraA that transports uracil [24], EcXanP(YicE) and EcXan Q (YgfO) that transport xanthine [25], EcUacT that transports uric acid [26], while EcRutG [27] and EcYbbY are of unknown function [28]. The three-dimensional structures of EcUraA with uracil in the inward-open and occluded states [24,29] and that of an inward-facing AnUapA with xanthine [30] have recently been solved. In addition, a number of

EcNCS2 proteins have undergone extensive mutagenesis and biochemical characterization [24–26,31,32]. With this impressive backdrop, we functionally characterize a novel NCS2 transporter from *Paenibacillus larvae* which we named PIUacP.

Materials and methods

Microbial strains and growth conditions

P. larvae subspecies *larvae* strain NRRL B-3650 was grown on MYPGP media [33]. *Escherichia coli* strain DH5 α (*fhuA2 lac Δ U169 phoA glmV44 Φ 80' lacZ Δ M15 gyrA96 recA1 relA1 endA1 thi-1 hsdR17*) was grown on LB media with 50 mg·mL⁻¹ carbenicillin and used for molecular cloning. *S. cerevisiae* strains RG191 [*MAT α , fyc2A::kanMX4, his3 Δ 1, leu2 Δ 0, met15 Δ 0, ura3 Δ 0*] [34] and NC122-sp6 [*MAT α leu2 fur4 Δ*] [35] were grown in synthetic complete (SC) medium—leucine (Sigma-Aldrich, St. Louis, MO) at 30 °C. Yeast transformations were performed by the lithium acetate method [36].

Nucleic acid manipulations

Genomic DNA was isolated from *P. larvae* strain B-3650 using QIAamp Tissue kit (Qiagen Inc. Santa Clarita, CA). The coding region for PIUacP was PCR-amplified from genomic DNA using oligonucleotides *PIXPa* 5' cccaagctctcgcagatgcgtaaaagcaagactactacc 3' and *PIXPb* 5' ataagaatcggccgcttaagcggcatgaacgttttctgc 3' using conditions 94 °C 2 min; 94 °C 15 s; 55 °C 30 s; 70 °C 2 min; repeat 30X; 72 °C 10 min. The resulting PCR product was purified using QIAquick PCR Purification Kit (Qiagen Inc. Santa Clarita, CA) and, along with yeast expression vector pRG399, cut with restriction endonucleases *Xho* I and *Not* I, ligated and transformed into *E. coli* strain DH5 α to generate plasmid pNS515. DNA sequence analysis was performed at W.M. Keck Biotechnology Resource Laboratory at Yale School of Medicine (New Haven CT, USA) to verify sequence integrity. pNS515 was transformed into yeast strains RG191 and NC122-Sp6 and transformants selected on SD—leucine media. RNA was isolated from *P. larvae* strain B-3650 culture grown in MYPGP to an OD₆₀₀ of 0.64 using the RNeasy Protection Kit/RNA protect Bacterial Reagent (Qiagen Inc. Santa Clarita, CA, USA) and treated with RNase free DNaseI (Roche, Indianapolis, IN, USA). Genomic DNA, DNaseI-treated total RNA, and DNaseI-treated total RNA with Invitrogen Super-Script One-Step RT-PCR treatment (ThermoFisher, Waltham, Massachusetts, USA) were amplified with primers *PIXPa* and *PIXPb* to generate a 1.3-kb fragment using amplification conditions of 94 °C 2 min; 94 °C 15 s; 55 °C 30 s; 70 °C 2 min; repeat 33X; and 72 °C 10 min. DNA products were visualized by agarose gel electrophoresis. Phylogenetic analysis employed Phylogeny.fr

[37] using MUSCLE alignment [38] and either maximum likelihood PhyML 3.1 [39] or Bayesian inference Mr. Bayes 3.2.3 [40].

Radiolabel uptake by yeast expressing PIUacP

Time course for the uptake of 0.5 μM [2,8- ^3H]-adenine was performed in RG191 cells containing pRH399 or pNS515 and concentrated to a density proportional to $\text{OD}_{600} = 4$. Samplings were harvested at time intervals of 0, 5, 10, 20, 40, 60, and 90 min. Yeast grown for 24 h at 30 $^{\circ}\text{C}$ was concentrated to $\text{OD}_{600} = 4$ and incubated for 0 and 5 min with 0.5 μM [8- ^3H]-guanine, [2,8- ^3H]-adenine, [8- ^3H]-xanthine, [5- ^3H]-cytosine, [8- ^3H]-hypoxanthine, or 1.0 μM [5,6- ^3H]-uracil (Moravek, Brea, CA), in 100 mM citrate buffer (pH 3.5) with 1% glucose. 25 μL aliquots were added to 4 mL of ice-cold water and filtered through a 0.45 μm Metrical membrane filter (Gelman Sciences, Ann Arbor, MI). Filters were then washed with 8 mL of water, and radioactivity was measured by a Beckman LS 6000IC liquid scintillation counter (Beckman Coulter, Fullerton, CA). Experiments were performed in triplicate. Statistical analysis used an independent paired *t*-test. Significance was measured at $P = 0.05$ (*), and the error bars represent the standard error of mean.

Radiolabeled substrate competition study of PIUacP

A substrate competition experiment was performed to further refine the solute binding capacity of PIUacP. Radiolabeled uptake of 1 μM [2, 8- ^3H]-adenine was performed in the presence of various heterologous unlabeled competitors including adenine, guanine, cytosine, hypoxanthine, xanthine, uric acid, and uracil. Yeast cells concentrated to $\text{OD}_{600} \approx 4$ expressing pNS515 were incubated with 1 μM [2, 8- ^3H]-adenine in the presence of 1 mM unlabeled heterologous competitors. The samplings were carried out at 5 min for both experiments, and radiolabeled substrate was measured as described above. Experiments were performed in triplicate. Statistical analysis used an independent paired *t*-test. Significance was measured at $P = 0.05$ (*), and the error bars represent the standard error of mean.

Transport kinetics and inhibitor studies of PIUacP

The K_m value of PIUacP for adenine was measured through traditional Michaelis–Menten kinetics experiments. RG191 cells ($\text{OD}_{600} = 4$) containing *PIUacP* were incubated with increasing concentrations of [2,8- ^3H]-adenine (0.1, 0.25, 0.5, 1.0, 10.0, 20.0, and 30.0 μM) for 10 min, and the amount of radiolabel taken up by cells was measured as mentioned above. Three independent replicas were used for each substrate concentration. Substrate saturation data were fitted

by nonlinear regression. Estimates of K_m were calculated from the double reciprocal Lineweaver–Burk plot transformation [41]. Radiolabeled uptake was performed as previously described at 1 μM [2,8- ^3H]-adenine. RG191 cells containing *PIUacP* were incubated with 1 μM [2,8- ^3H]-adenine alone or in the presence of 100 μM carbonyl cyanide *m*-chlorophenylhydrazone (CCCP) or 1 mM ouabain octahydrate. The reaction was incubated at 30 $^{\circ}\text{C}$ with samples taken at 0 and 5 min, and radioactivity was measured as above. The error bars represent the standard error of the mean of three independent experiments. An independent paired *t*-test was used to measure statistical significance. Statistical significance was measured at $P = 0.05$ (*).




Fig. 1. Phylogenetic relationships of PIUacP with select *Escherichia coli* and *Bacillus subtilis* nucleobase transporters. Amino acid sequence accession numbers are given as PIUacP (EFX44190.1); *B. subtilis* PbuX (P42086), PucJ (O32139), PucK (O32140); *E. coli* UraA (BAA16385.1), RutG (P75892.2), XanQ (P67444.2), XanP (P0AGM9.1), UacT (Q46821.1), YbbY (P77328.2), GHXQ (Q46817.2), GHXP (P0AF52.1), AdeP (P31466.2), and AdeQ (P31440.3). Phylogenetic trees were constructed using Phylogeny.fr [37], 2008) using MUSCLE alignment [38] and employing maximum likelihood (A) PhyML 3.1 [39] or Bayesian inference (B) Mr. Bayes 3.2.3 [40].

Results and Discussion

P. larvae genome contains a locus with similarities to bacterial uric acid and xanthine transporters

A search of the *P. larvae* B-3560 genome through the National Center for Biotechnology Information

database with the amino acid sequence of the *E. coli* uric acid transporter UacT (also known as YgfU) using sequence comparison software (tblastn) [42] matched with a presumptive protein (Gb# EFX44190.1) [43]. Phylogenetic trees derived from maximum likelihood or Bayesian inference detailing the amino acid sequence comparisons with other characterized nucleobase cation symporter 2 from Gram-positive *Bacillus subtilis* and




Fig. 2. Alignment of PIUacP with select *Escherichia coli* and *Bacillus subtilis* xanthine and uric acid transporters. Alignment of amino acid sequences of *Paenibacillus larvae* UacP with select bacterial xanthine and uric acid transporters as given by ClustalW [55]. Amino acid sequences are as designated in Fig 1. The NAT motif conserved among NCS2 proteins is designated by black bar above amino acid sequences [52,56]. Amino acid residues identified by site-directed mutagenesis in the EcUacT that affect function (bold black text with gray background) are identified [26]. Amino acid residues identified by site-directed mutagenesis in EcXanQ that affect function (bold italic underlined text) are identified [31,47]. Transmembrane spanning domains for PIUacP are indicated by + above amino acid sequence, as predicted by TMHMM v2.0 [46] (TM1–TM8 and TM11–TM14) or as identified in EcUacT (TM9 and TM10) [26]. Transmembrane spanning domains for EcUacT are identified as underlined text [22].

Gram-negative *E. coli* reveal a close association between *P. larvae* protein (Gb# EFX44190.1) and *B. subtilis* and *E. coli* uric acid and xanthine transporters in COG2233 (Fig. 1A,B). Based upon this similarity, the *P. larvae* protein (Gb# EFX44190.1) is now referred to as *P. larvae* uric acid permease-like protein (PIUacP). PIUacP shares a high level of amino acid similarity with *B. subtilis* uric acid transporters BsPucK (58.9% amino acid identity/87.4% amino acid similarity) and BsPucJ (53.7%/86.2%) [44], xanthine transporter BsPbuX (59.6%/90.5%) [45], with lesser, but still substantial, similarity to *E. coli* uric acid transporter EcUacT (38.6%/72.2%) [26], and the xanthine transporters EcXanQ (33.6%/68.9%) and EcXanP (27.1%/62.5%) [25]. PIUacP also shares secondary structures, such as fourteen transmembrane spanning (TM) domains, and key signature motifs that are common to uric acid and xanthine transporters as well as other NCS2 (NAT) proteins (Fig. 2). PIUacP contains twelve common overlapping TM domains (TM1–TM8; TM11–TM14) as predicted by secondary structure prediction program TMHMM v2.0 [46] when aligned with EcUacT, while the shorter TM domains 9 and 10 present in the crystal structures of EcUraA and AnUapA [24,30] are not predicted by most secondary structure prediction programs. However, the overall levels of amino acid sequence similarity, common motifs, and transmembrane spanning domain positions are insufficient to assign PIUacP as either a low-affinity uric acid transporter or a high-affinity xanthine transporter.

Extensive site-directed mutagenesis combined with functional analysis of both EcUacT and EcXanQ has identified residues important for transport and solute selectivity. Some of these residues are conserved or present as similar amino acids in PIUacP at the corresponding sites with EcUacT or EcXanQ, while other identified residues are not conserved. Six of nine amino acids previously identified by site-directed mutagenesis for function in EcUacT are shared in PIUacP (Fig. 2). In EcUacT, aspartic acid (298) essential for function is present as a similar glutamic acid in PIUacP, while serine (426), involved in uric acid affinity in EcUacT, is present as the like amino acid asparagine in PIUacP [26]. Similarly, PIUacP shares identity with ten of eighteen amino acids identified as important for function in EcXanQ as determined by site-directed mutagenesis (Fig. 2) [31,32,47]. A number of key amino acid residues affecting EcXanQ function in the presumptive solute binding cavity are not conserved in PIUacP: EcXanQ alanine (323) serves to sense xanthine binding and glycine (333) forms part of the purine permeation pathway [47] are, however, present as unlike serine and lysine residues in PIUacP, respectively; phenylalanine

(322) and asparagine (326) essential for high-affinity xanthine binding in EcXanQ [31] are present as unlike tyrosine and valine in PIUacP, respectively; alteration of residue tyrosine (425) in EcXanQ leads to depressed xanthine accumulation [32], but is present as a lysine in PIUacP (Fig. 2). Unfortunately, neither overall amino acid similarities nor the commonality of functionally important residues with other characterized NCS2 proteins are sufficient to assign PIUacP as either a uric acid transporter or a xanthine transporter. An experimental approach to determine function is required to address this question. Endpoint reverse transcription polymerase chain reaction of *P. larvae* RNA from vegetative cells reveals that the *PIUacP* locus is expressed (Fig. 3).




Fig. 3. Expression of *PIUacP* by *Paenibacillus larvae*. Expression of *PIUacP* in *P. larvae* by endpoint RT-PCR. Lane 1, exACTGene 1kb plus DNA ladder (Fisher); lane 2, genomic DNA; lane 3, DNaseI-treated total RNA without reverse transcription; and lane 4, RT-PCR of DNaseI-treated total RNA all amplified with *PIUacP*-specific primers.




Fig. 4. Time-dependent uptake of [³H]-adenine by yeast containing *PIUacP*. Yeast strain RG191 containing pNS515 (Δ *fcy2/PIUacP*) was monitored for the uptake of [2,8-³H]-adenine over a ninety-minute time period. Significance was measured at $P = 0.05$ (*). All error bars represent standard error of the mean. Experiments were performed in triplicate.

Heterologous expression of PIUacP in nucleobase transport-deficient *S. cerevisiae* strains defines solute transport and binding profile

Plasmid pNS515 was transformed into *S. cerevisiae* strains RG191 and NC122-Sp6. Strain RG191 carries a deletion of the *Fcy2* locus that encodes for the guanine–adenine–hypoxanthine transporter and is deficient in transport of these nucleobases. Strain NC122-Sp6 carries a deletion of the *Fur4* locus that encodes for the uracil transporter and is deficient for transport of uracil. Yeast genomes do not contain loci for xanthine or uric acid transport. These yeast strains and expression plasmids have been successfully employed in heterologous expression experiments monitoring for the uptake of radiolabeled purines and pyrimidines for

a variety of plant NCS1 and NCS2 loci for guanine, adenine, cytosine, hypoxanthine, xanthine, and uracil [48–52]. A time course monitoring the uptake of [³H]-adenine for strain RG191 harboring *PIUacP* (RG191/pNS515) displays a saturation after 60 min (Fig. 4). Uptake of [³H]-adenine at 5-min incubation is within the linear phase and was used as incubation time for subsequent experiments. Figure 5A–C reveals that yeast strains containing pNS515 took up significantly more [³H]-adenine, [³H]-guanine, and [³H]-uracil than strains containing the empty vector. However, no significant difference in uptake was observed for [³H]-cytosine, [³H]-hypoxanthine, or [³H]-xanthine (Fig. 5D–F). *PIUacP* is an adenine, guanine, and uracil transporter, but not a xanthine transporter. *PIUacP* stands in contrast to canonical bacterial NCS2 transporters from




Fig. 5. Uptake of radiolabeled nucleobases by yeast containing *PIUacP*. Yeast strain RG191 alone ($\Delta fcy2$) or containing pNS515 ($\Delta fcy2/PIUacP$) was monitored at 0 and 5 min for the uptake of (A) [³H]-guanine, (B) [2,8-³H]-adenine, (D) [5-³H]-cytosine, or (E) [8-³H]-hypoxanthine. Strain NC122-Sp6 alone or with pNS515 was monitored at 0 and 5 min for the uptake of (C) [5,6-³H]-uracil or (F) [8-³H]-xanthine. Statistical analysis used an independent paired *t*-test. Significance was measured at $P = 0.05$ (*). All error bars represent standard error of the mean. Experiments were performed in triplicate.

E. coli that tend to be specific for the transport of one or two nucleobases (e.g., *E. coli* uracil transporter [24], xanthine transporters [25], adenine and guanine/hypoxanthine transporters [23]).

A series of substrate competition experiments were performed to determine the binding—but not transport—ability of nucleobases on PIUacP. Here, yeast-harboring *PIUacP* were incubated with radiolabeled adenine alone or with an excess of unlabeled competitors, and the uptake of adenine determined. If competitor molecules bind to the same site in the transporter as does adenine, then adenine transport can be affected—regardless of whether the competitor molecule is actually transported. It is clear that guanine, cytosine, hypoxanthine, xanthine, uric acid, and uracil all compete with adenine and reduce the amount of radiolabel uptake (Fig. 6). The solute transport and binding profile for PIUacP is thus defined as the ability to transport adenine, guanine, and uracil, and to bind cytosine, hypoxanthine, xanthine, and uric acid. This broad nucleobase transport and binding profile differentiates PIUacP from other characterized bacterial NCS2/COG2233 transporters that have both narrow transport and binding profiles (e.g., EcXanQ transports xanthine, but neither transports nor competitively binds adenine, cytosine, hypoxanthine, uric acid,




Fig. 6. Substrate competition studies of yeast containing *PIUacP*. Yeast strain RG191 containing pNS515 ($\Delta fcy2/PIUacP$) was monitored for the uptake of [2,8- 3H]-adenine in the presence of excess of unlabeled competitors. Measurements were scored against the uptake of [2,3- 3H]-adenine without competitors as 100% (not shown). Significance was measured at $P = 0.05$ (*). All error bars represent standard error of the mean. Experiments were performed in triplicate.

or uracil and displays only weak competitive binding with guanine [25]).

The broad solute recognition of PIUacP is reminiscent of some bacterial NCS2 COG2252 and some eukaryotic NCS2 transporters. Recently, it was shown that two AzgA-like transporters of *P. larvae*, *PIAzg1* and *PIAzg2*, belonging to COG2252, display a wide solute transport profile including adenine, guanine, xanthine, uracil, and cytosine [53]. Eukaryotic NCS2 transporters have similarly broad transport profiles including *A. nidulans* AzgA transporting adenine, guanine, and hypoxanthine [19]; *Arabidopsis thaliana* Azg1




Fig. 7. Kinetic values of select nucleobases for PIUacP. Substrate saturation assay in yeast strain RG191 containing pNS515 ($\Delta fcy2/PIUacP$) with [2,8- 3H]-adenine (A). Double reciprocal Lineweaver-Burk plot (B) for PIUacP was constructed from the substrate saturation curve. All error bars represent standard error of the mean. Experiments were performed in triplicate.

Table 1. Effects of inhibitors on the function of PIUacP.

	%[2,8- 3H]-adenine uptake
RG191/ <i>PIUacP</i>	100 ± 35.3
RG191/ <i>PIUacP</i> + CCCP*	41.8 ± 1.5
RG191/ <i>PIUacP</i> + Ouabain	114.9 ± 11.3

Significance was measured at $P = 0.05$.

*indicates that the uptake of radio labeled adenine for RG191/*PIUacP* + CCCP* is significantly different from the uptake of radio labeled adenine for the control RG191/*PIUacP*.

and Azg2 transporting adenine and guanine; *Zea mays* Azg2–Azg3 transporters transporting adenine, guanine, cytosine, hypoxanthine, and xanthine (Mourad and Schultes, pers. comm.); the *A. thaliana* AtNAT1-8 and AtNAT12 that transport xanthine, and a combination of adenine, guanine, uracil, and cytosine [18; Mourad and Schultes, pers. comm.]. Nonetheless, the solute transport and binding profile for PIUacP is distinct from other NCS2 [54]. It is clear that despite the close amino acid similarity to other bacterial xanthine or uric acid transporters that have narrow solute transport and binding profiles, PIUacP functions as neither.

Biochemical and kinetic properties PIUacP differ from the uric acid transporter in *E. coli*

The affinity of PIUacP for adenine was determined using Michaelis–Menten kinetics experiments in which RG191 harboring *PIUacP* was incubated with increasing concentrations of [³H]-adenine and then resulting radiolabel uptake was monitored. Data reveal that PIUacP has a high affinity for adenine with a K_m $7.04 \pm 0.18 \mu\text{M}$ (Fig. 7A,B), a similar affinity as those determined for other NCS2 adenine transporters EcAdeP ($K_m = 1.0 \mu\text{M}$) [23] or AtNAT3 and AtNAT12 ($K_m = 10.12 \mu\text{M}$ and $1.74 \mu\text{M}$, respectively) [18].

Yeast strain RG191 containing *PIUacP* takes up significantly less [³H]-adenine in the presence of the proton uncoupler carbonyl cyanide *m*-chlorophenylhydrazone, but not the Na⁺ gradient disruptor ouabain (Table 1). As with other bacterial, fungal, and plant NCS2, PIUacP appears to be a proton–nucleobase symporter, in contrast to mammalian NCS2 that rely upon a Na⁺ gradient for function [14]. During rapid cell growth in infected honeybee larvae, a rich plethora of compounds released from disrupted cells are available for uptake by the pathogen. Here, it is shown that *P. larvae* has the capacity to import adenine, guanine, and uracil from external sources. This obligate pathogen grows in a nutrient-rich environment and would benefit from transporters that move a wide range of available nutrients.

Conclusions

Analysis of a nucleobase cation symporter 2 from *P. larvae* reveals a close amino acid sequence similarity to other functionally characterized bacterial xanthine and uric acid transporters. However, functional analysis of PIUacP in a yeast heterologous system reveals it has a broad solute transport and binding profile moving adenine, guanine, and uracil and binding hypoxanthine, xanthine, cytosine, and uric acid, unlike other bacterial xanthine and uric acid transporters. This

work reveals that close amino acid sequence similarity by itself cannot be used as a surrogate for assigning function.

Acknowledgements

We thank Regan Huntley and Carol Clark at CAES for expert technical assistance. This work was funded by research funds from IPFW to GSM and Hatch Fund CONH00253 to NPS.

Author contributions

NPS and GSM designed the experiments. AJS and CRA performed the experiments, analyzed the data, and prepared figures. NPS, DWD, and GSM wrote the manuscript.

References

- White GF (1906) *The bacteria of the apiary, with special reference to bee diseases*. Bureau of Entomology Technical Series no 14. US Department Agriculture, Washington DC.
- Katznelson H (1950) *Bacillus pulvifaciens* (n. sp.), an organism associated with powdery scale of honey bee larvae. *J Bacteriol* **59**, 153–155.
- Genersch E (2010) American Foulbrood in honey bees and its causative agent, *Paenibacillus larvae*. *J Invertebr Pathol* **103**, S10–S19.
- Ebeling J, Knispel H, Hertlein G, Fünfhaus A and Genersch E (2015) Biology of *Paenibacillus larvae*, a deadly pathogen of honey bee larvae. *Appl Microbiol Biotechnol* **17**, 7387–7395.
- Yue D, Nordhoff M, Wieler LH and Genersch E (2008) Fluorescence *in situ*-hybridization (FISH) analysis of the interactions between honey bee larvae and *Paenibacillus larvae*, the causal agent of American Foulbrood of honey bees (*Apis mellifera*). *Environ Microbiol* **10**, 1612–1620.
- Alvarado I, Phui A, Elekonich MM and Abel-Santos E (2013) Requirements for *in vitro* germination of *Paenibacillus larvae* spores. *J Bacteriol* **195**, 1005–1011.
- Winston ML (1987) *The Biology of the Honey Bee*. Harvard University Press, Cambridge, MA.
- Antúnez K, Anido M, Schlapp G, Evans JD and Zunino P (2009) Characterization of secreted proteases of *Paenibacillus larvae*, potential virulence factors involved in honey bee larval infection. *J Invertebr Pathol* **102**, 129–132.
- Poppinga L and Genersch E (2015) Molecular pathogenesis of American foulbrood: how *Paenibacillus larvae* kills honey bee larvae. *Curr Opin Insect Sci* **10**, 29–36.
- Desimone M, Catoni E, Ludewig U, Hilpert M, Schneider A, Kunze R, Tegeder M, Frommer WB and

- Schumacher K (2002) A novel superfamily of transporters for allantoin and other oxo-derivatives of nitrogen heterocyclic compounds in *Arabidopsis*. *Plant Cell* **14**, 847–856.
- 11 Schmidt A, Su Y-H, Kunze R, Warner S, Hewitt M, Slocum RD, Ludewig U, Frommer WB and Desimone M (2004) UPS1 and UPS2 from *Arabidopsis* mediate high affinity transport of uracil and 5-fluorouracil. *J Biol Chem* **279**, 44817–44824.
 - 12 Schmidt A, Baumann N, Schwarzkopf A, Frommer WB and Desimone M (2006) Comparative studies on Ureide Permeases in *Arabidopsis thaliana* and analysis of two alternative splice variants of AtUPS5. *Planta* **224**, 1329–1340.
 - 13 Pantazopoulou A and Diallinas G (2007) Fungal nucleobase transporters. *FEMS Microbiol Rev* **31**, 657–675.
 - 14 Gournas C, Papageorgiou I and Diallinas G (2008) The nucleobase-ascorbate transporter (NAT) family: genomics, evolution, structure-function relationships and physiological role. *Mol BioSyst* **4**, 404–416.
 - 15 Jelesko JG (2012) An expanding role for purine uptake permease-like transporters in plant secondary metabolism. *Front Plant Sci* **3**, 1–5.
 - 16 Ma P, Baldwin JM, Baldwin SA and Henderson PJF (2013) Membrane transport proteins: the nucleobase cation symporter 1 family. In *Encyclopedia of Biophysics* (Roberts GCK, ed), pp. 1485–1489. Springer, European Biophysical Societies' Association.
 - 17 Argyrou E, Sophianopoulou V, Schultes N and Diallinas G (2001) Functional characterization of a maize purine transporter by expression in *Aspergillus nidulans*. *Plant Cell* **13**, 953–996.
 - 18 Niopek-Witz S, Deppe J, Lemieux MJ and Möhlmann T (2014) Biochemical characterization and structure-function relationship of two plant NCS2 proteins, the nucleobase transporters NAT3 and NAT12 from *Arabidopsis thaliana*. *Biochim Biophys Acta* **1838**, 3025–3035.
 - 19 Cecchetto G, Amillis S, Diallinas G, Scazzocchio C and Drevet C (2004) The AzgA purine transporter of *Aspergillus nidulans*. Characterization of a protein belonging to a new phylogenetic cluster. *J Biol Chem* **279**, 3132–3141.
 - 20 Goudela S, Tsilivi H and Diallinas G (2006) Comparative kinetic analysis of AzgA and Fcy21p, prototypes of the two major fungal hypoxanthine-adenine-guanine transporter families. *Mol Membr Biol* **23**, 291–303.
 - 21 Kosti V, Lambrinidis G, Myriantopoulos V, Diallinas G and Mikros E (2012) Identification of the substrate recognition and transport pathway in a eukaryotic member of the nucleobase-ascorbate transporter (NAT) family. *PLoS One* **7**, e41939.
 - 22 Kryptou E, Lambrinidis G, Evangelidis T, Mikros E and Diallinas G (2014) Modelling, substrate docking and mutational analysis identify residues essential for function and specificity of the major fungal purine transporter AzgA. *Mol Microbiol* **93**, 129–145.
 - 23 Papakostas K, Botou M and Frillingos S (2013) Functional identification of the hypoxanthine/guanine transporters YjcD and YgfQ and the adenine transporters PurP and YicO of *Escherichia coli* K-12. *J Biol Chem* **288**, 36827–36840.
 - 24 Lu F, Li S, Jiang Y, Fan H, Lu G, Deng D, Dang S, Zhang X, Wang J and Yan N (2011) Structure and mechanism of the uracil transporter UraA. *Nature* **472**, 243–246.
 - 25 Karatza P and Frillingos S (2005) Cloning and functional characterization of two bacterial members of the NAT/NCS2 family in *Escherichia coli*. *Mol Membr Biol* **22**, 251–261.
 - 26 Papakostas K and Frillingos S (2012) Substrate selectivity of YgfU, a uric acid transporter from *Escherichia coli*. *J Biol Chem* **287**, 15684–15695.
 - 27 Kim KS, Pelton JG, Inwood WB, Andersen U, Kustu S and Wemmer DE (2010) The Rut pathway for pyrimidine degradation: novel chemistry and toxicity problems. *J Bacteriol* **192**, 4089–4102.
 - 28 Cusa E, Obradors N, Baldoma L, Badia J and Aguilar J (1999) Genetic analysis of a chromosomal region containing genes required for assimilation of allantoin nitrogen and linked glycolate metabolism in *Escherichia coli*. *J Bacteriol* **181**, 7479–7484.
 - 29 Yu X, Yang G, Yan C, Baylon JL, Jiang J, Fan H, Lu G, Hasegawa K, Okumura H, Wang T *et al.* (2017) Dimeric structure of the uracil: proton symporter UraA provides mechanistic insights into the SLC4/23/26 transporters. *Cell Res* **27**, 1020–1033.
 - 30 Alguet T, Amillis S, Leung J, Lambrinidis G, Capaldi S, Scull NJ, Craven G, Iwata S, Armstrong A, Mikros E *et al.* (2016) Structure of eukaryotic purine/H⁺ symporter UapA suggests a role for homodimerization in transport activity. *Nature Commun* **7**, 11336.
 - 31 Karatza P, Panos P, Georgopoulou E and Frillingos S (2006) Cysteine-scanning analysis of the nucleobase-ascorbate transporter signature motif in YgfO permease of *Escherichia coli*. *J Biol Chem* **281**, 39881–39890.
 - 32 Papakostas K, Georgopoulou E and Frillingos S (2008) Cysteine-scanning analysis of putative helix XII in the YgfO Xanthine permease. *J Biol Chem* **283**, 13666–13678.
 - 33 Dingman DW and Stahly DP (1983) Medium promoting sporulation of *Bacillus larvae* and metabolism of medium components. *Appl Environ Microbiol* **46**, 860–869.
 - 34 Winzler EA, Shoemaker DD, Astromoff A, Liang H, Anderson K, Andre B, Bangham R, Benito R, Boeke JD, Bussey H *et al.* (1999) Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* **285**, 901–906.

- 35 Jund R, Chevallier MR and Lacroute F (1988) Primary structure of the uracil transport protein of *Saccharomyces cerevisiae*. *Eur J Biochem* **171**, 417–424.
- 36 Gietz DR and Woods RA (2002) Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol* **350**, 87–96.
- 37 Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard JF, Guindon S, Lefort V, Lescot M *et al.* (2008) Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res* **36**, W465–W469.
- 38 Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**, 1792–1797.
- 39 Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W and Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**, 307–321.
- 40 Ronquist F and Huelsenbeck JP (2003) MrBayes 3. Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574.
- 41 Ritchie RJ and Prvan T (1996) Current Statistical Methods for estimating the K_m and V_{max} of Michaelis-Menten Kinetics. *Biochem Educ* **24**, 16–206.
- 42 Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* **5**, 401–410.
- 43 Chan QWT, Cornman RS, Birol I, Liao NY, Chan SK, Docking TR, Jackman SD, Taylor GA, Jones SJ, de Graaf DC *et al.* (2011) Updated genome assembly and annotation of *Paenibacillus larvae*, the agent of American foulbrood disease of honey bees. *BMC Genom* **12**, 450.
- 44 Schultz AC, Nygaard P and Saxild HH (2001) Functional analysis of 14 genes that constitute the purine catabolic pathway in *Bacillus subtilis* and evidence for a novel regulon controlled by the PucR transcription activator. *J Bacteriol* **183**, 3293–3302.
- 45 Christiansen LC, Schou S, Nygaard P and Saxild HH (1997) Xanthine metabolism in *Bacillus subtilis*: characterization of the xpt-pbuX operon and evidence for purine- and nitrogen-controlled expression of genes involved in xanthine salvage and catabolism. *J Bacteriol* **179**, 2540–2550.
- 46 Krogh A, Larsson B, von Heijne G and Sonnhammer ELL (2001) Predicting transmembrane protein topology with a hidden markov model: application to complete genomes. *J Mol Bio* **305**, 567–580.
- 47 Georgopoulou E, Mermelekas G, Karena E and Frillingos S (2010) Purine substrate recognition by the nucleobase-ascorbate transporter signature motif in the YgfO xanthine permease: Asn-325 binds and Ala-323 senses substrate. *J Biol Chem* **285**, 19422–19433.
- 48 Mansfield TA, Schultes NP and Mourad GS (2009) AtAzg1 and AtAzg2 comprise a novel family of purine transporters in Arabidopsis. *FEBS Lett* **583**, 481–486.
- 49 Mourad GS, Tippmann-Crosby J, Hunt KA, Gicheru Y, Bade K, Mansfield TA and Schultes NP (2012) Genetic and molecular characterization reveals a unique nucleobase cation symporter 1 in Arabidopsis. *FEBS Lett* **586**, 1370–1378.
- 50 Schein J, Hunt KA, Minton J, Schultes NP and Mourad GS (2013) The nucleobase cation symporter 1 from *Chlamydomonas reinhardtii* and the evolutionary distant *Arabidopsis thaliana* share function and establish a plant-specific solute transport profile. *Plant Physiol Biochem* **70**, 52–60.
- 51 Minton JA, Rapp M, Stoffer AJ, Schultes NP and Mourad GS (2016) Heterologous complementation studies reveal the solute transport profiles of a two-member nucleobase cation symporter 1 (NCS1) family of *Physcomitrella patens*. *Plant Physiol Biochem* **100**, 12–17.
- 52 Rapp M, Schein J, Hunt KA, Nalam V, Mourad GS and Schultes NP (2016) The solute specificity profiles of nucleobase cation symporter 1 (NCS1) from *Zea mays* and *Setaria viridis* illustrate functional flexibility. *Protoplasma* **253**, 611–623.
- 53 Alexander CR, Dingman DW, Schultes NP and Mourad GS (2018) The solute transport profile of two Aza-guanine transporters from the Honey bee pathogen *Paenibacillus larvae*. *FEMS Microbiol Lett* **365**, 1–8. <https://doi.org/10.1093/femsle/fny018>
- 54 Goudela S, Karatza P, Koukaki M, Frillingos S and Diallinas G (2005) Comparative substrate recognition by bacterial and fungal purine transporters of the NAT/NCS2 family. *Mol Membr Biol* **22**, 263–275.
- 55 Thompson JD, Higgins DG and Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**, 4673–4680.
- 56 Koukaki M, Vlanti A, Goudela S, Pantazopoulou A, Gioule H, Tournaviti S and Diallinas G (2005) The nucleobase ascorbate transporter (NAT) signature motif in UapA defines the function of the purine translocation pathway. *J Mol Biol* **350**, 499–513.