

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

**IDENTIFICATION OF METABOLITES AND THERMAL TRANSFORMATION
PRODUCTS OF QUINOLONES IN RAW COW MILK BY LIQUID
CHROMATOGRAPHY COUPLED TO HIGH RESOLUTION MASS
SPECTROMETRY**

A. Junza¹, S. Barbosa², R. Codony³, A. Jubert³, J. Barbosa^{1,2}, D. Barrón^{1*}

¹Universitat de Barcelona, Departament de Química Analítica, Campus de l'Alimentació de Torribera.

Avda. Prat de la Riba, 171, 08921 Sta. Coloma de Gramanet, Barcelona, Spain

²Universitat de Bracelona, Departament de Química Analítica, Martí i Franquès, 1-11, 08028 Barcelona, Spain

³Laboratori Interprofessional Lleter de Catalunya (ALLIC) Ctra. Vilassar a Cabrils s/n, 08348 Cabrils, Barcelona, Spain

* To whom correspondence should be sent.

Phone: +34-934033797

E-mail: dolores.barron@ub.edu

30 **ABSTRACT**

31 The presence of residues of antibiotics, metabolites and thermal transformation products
32 (TPs), produced during thermal treatment to eliminate pathogenic microorganisms in milk,
33 could be represent a risk for people.

34

35 Cow milk samples spiked with enrofloxacin (ENR), ciprofloxacin (CIP), difloxacin (DIF)
36 and sarafloxacin (SAR) and milk samples from cows medicated with ENR were submitted
37 to several thermal treatments. The milk samples were analysed by liquid chromatography
38 mass spectrometry (LC-MS) to find and identify TPs and metabolites. In this work, 27 TPs
39 of four quinolones and 24 metabolites of ENR were found. Some of these compounds had
40 been reported previously, but others were characterized for the first time, including lactose-
41 conjugated CIP, the formamidation reaction for CIP and SAR, and hydroxylation or ketone
42 formation to produce three different isomers for all quinolones studied.

43

44

45 Keywords: antibiotics, milk, quinolones, thermal treatments, transformation products,
46 metabolites, elucidation, ToF, LTQ-Orbitrap

47

48

49 **1. INTRODUCTION**

50 Quinolones are one of the classes of antibiotics that are most widely used in veterinary
51 practice to treat bacterial infections of animals in livestock farming and bovine milk
52 production. Antibiotics are also used as growth promoters, although this practice has been
53 forbidden by the European Union (EU) since 2006¹.

54 The presence of antibiotic residues in food is a potential risk to consumers because of
55 direct toxic effects such as allergic reactions or the induction of resistant strains of
56 bacteria^{2,3}. To ensure human health, the EU has established maximum residue limits
57 (MRL) for some antibiotics in foodstuffs of animal origin (37/2010/EC)⁴. Active
58 substances are usually included in this regulation. However, their metabolites such as
59 ciprofloxacin, the main metabolite of enrofloxacin, are only considered in a few cases.

60

61 Milk is one of the most commonly consumed foods in the world. Drug residues and
62 metabolites can be found in milk if the time between drug administration and milking is
63 too short. Before consumption, cow milk has to be subjected to thermal treatment to
64 eliminate pathogenic microorganisms. Two common thermal processes are pasteurization,
65 which consists of heating at 60°C for 30 min or at 72°C for 15 s, and sterilization, by which
66 milk is heated at 120°C for 20 min^{5,6}. During thermal processes, the antibiotic residues and
67 their metabolites could undergo transformations, depending on the temperature and the
68 duration of the process.

69 The study of metabolites and their thermal degradation products is of interest because these
70 new compounds may be a health risk to consumers and could be more persistent and
71 harmful than the drug administered⁷.

72 Numerous papers have been published on analytical methods to determine target
73 substances and their main metabolites in several matrices⁸⁻²¹. However, there are few
74 studies on the identification and determination of metabolites, degradation, and
75 transformation products (TPs). The analytical methods consist of chromatographic
76 separation prior to detection with a mass spectrometer. Several strategies are used to
77 identify non-targets, including accurate mass measurement to determine the elemental
78 composition and distinguish isobaric molecular ions, multiple mass fragmentations (MSⁿ),
79 and complementary techniques such as NMR to confirm structure^{7,22-34}. Some metabolites
80 of quinolones are described in the bibliography^{25,35-41}, but to our knowledge no study has
81 been carried out on metabolites in milk which has been exposed to thermal treatment.

82

83 The aim of this study was to characterize the metabolites and TPs of four quinolones ENR,
84 CIP, DIF and SAR (Figure 1) in cow's milk subjected to thermal treatment. In order to test
85 the new metabolites and TPs found in spiked milk samples, milk samples from animals
86 medicated with ENR were submitted to the same thermal treatment. Samples were
87 analysed by liquid chromatography (LC) coupled to high resolution mass spectrometry
88 (ToF and LTQ-Orbitrap).

89

90 **2. EXPERIMENTAL PROCEDURES**

91 **2.1. Chemicals and apparatus**

92 The standards were purchased from several pharmaceutical firms: ENR from Cenavisa
93 (Reus, Spain), CIP from Ipsen Pharma (Barcelona, Spain), DIF from Abbott (Madrid,
94 Spain) and SAR from AK Scientific Inc. (CA,USA).

95

96 All reagents were of analytical grade unless indicated. Formic acid (HOCHO), acetic acid
97 (HOAc), acetonitrile (MeCN), methanol (MeOH), sodium dihydrogenphosphate and
98 sodium hydroxide (NaOH) were supplied by Merck (Darmstadt, Germany). Water was
99 ultrapurified by a Milli-Q system (Millipore, MA, USA).

100

101 The quinolone individual stock solutions (ENR, CIP, DIF and SAR) were prepared at a
102 concentration of $500 \mu\text{g}\cdot\text{mL}^{-1}$ in 50 mM HOAc aqueous solution. The individual working
103 solutions that were used to spike the milk samples were prepared by dilution of the
104 individual stock solution at a concentration of $20 \mu\text{g}\cdot\text{mL}^{-1}$ in water.

105

106 Sodium dihydrogenphosphate solution 0.1 M adjusted with NaOH 5 M to pH 10 was added
107 to milk samples prior to the clean-up step.

108

109 The SPE cartridges used in this study were Oasis HLB ($3 \text{ cm}^3/60 \text{ mg}$), obtained from
110 Waters (Milford, MA, USA).

111

112 To simulate the thermal treatment of milk, a laboratory oven (Mettler) was used to warm
113 the samples. A Rotanta 460RS (Hettich Zentrifugen, Germany) centrifuge was used to
114 separate precipitated proteins and fat of supernatant after the extraction. The SPE step was
115 carried out in a Supelco vacuum manifold with disposable liners for 24 cartridges
116 (Bellefonte, PA, USA), connected to a Supelco vacuum tank. Evaporation to dryness was
117 performed in a TurboVap LV (Caliper Life Science, Hopkinton, MA, USA). A Crison
118 2002 potentiometer ($\pm 0.1 \text{ mV}$) (Crison, Barcelona, Spain) with a Crison 5203 combined
119 pH electrode from Orion Research (Boston, MA, USA) was used to measure the pH of the
120 phosphate solution and the mobile phase.

121 **2.2. Milk samples**

122 Milk samples used in the work have been supplied by the Laboratori Interprofessional
123 Lleter de Catalunya (ALLIC), control laboratory of milk in Catalonia. The samples were
124 analysed previously in ALLIC using a screening method to determine if the milk contained
125 residues of antibiotics. The milk samples that were negative in antibiotics were used as
126 blank samples in this work. Four positive samples in ENR were used to study the
127 metabolites.

128

129 **2.3. LC-MS instrumentation and working conditions**

130 **2.3.1. LC conditions**

131 Quinolones, their metabolites and degradation products were separated in a Zorbax Eclipse
132 XDB-C8 column (5 μm , 4.6 \times 150 mm) from Agilent Technologies (Santa Clara, CA,
133 USA), using a pre-column Kromasil C8 (5 μm , 4.6 \times 15 mm) supplied by Akady
134 (Barcelona, Spain) when samples were analysed by LC-ToF. A Pursuit UPS C18 column
135 (2.4 μm , 2 \times 50 mm) from Varian (Harbor City, CA, USA) was used when samples were
136 injected into the LC-LTQ-Orbitrap.

137

138 For both columns, the mobile phase was composed of an aqueous solution of 0.1%
139 HOCHO (solvent A) and MeCN with 0.1% HOCHO (solvent B) at a constant flow rate of
140 1 $\text{mL}\cdot\text{min}^{-1}$ and 0.3 $\text{mL}\cdot\text{min}^{-1}$, respectively. The injection volume was 20 μL in LC-ToF
141 and 10 μL in LC-LTQ-Orbitrap. Separations were carried out in a Zorbax Eclipse XDB-C8
142 column for 11 min under the following gradients: from 0 to 1 min, 15% B; 4 min, 45% B;
143 7 min, 56% B; 8.5 min, 15% B; and 11 min, 15% B. Separations were carried out in a
144 Pursuit UPS C18 column under the following conditions: from 0 to 3.5 min, 1% B; 4.5

145 min, 25% B; 5 min, 50% B; 6.5 min, 50% B; 7.5 min, 25% B; 8.5 min, 10% B; and 11
146 min, 1% B.

147

148 **2.3.2. LC-MS (ToF)**

149 A HP Agilent Technologies 1100 LC system was equipped with an autosampler and
150 coupled to a 6220 oa-ToF LC/MS mass spectrometer with an electrospray ionisation
151 source (ESI) (Agilent Technologies, Santa Clara, CA, USA). The system was controlled by
152 Mass Hunter workstation software (Agilent Technologies, Santa Clara, CA, USA) for the
153 acquisition and processing of data from the ToF mass spectrometer.

154

155 The optimum parameters of ToF in positive mode were as follows: capillary voltage 4000
156 V, drying gas (N₂) temperature 300°C, drying gas (N₂) flow rate 9 L·min⁻¹, nebulizer gas
157 (N₂) 40 psi, fragmentor voltage 150 V, skimmer voltage 60 V, and OCT 1 RF voltage 250
158 V. The ToF-MS mass resolving power was approximately 10,000 FWHM (Full width at
159 half maximum) at *m/z* 922. Spectra were acquired over the *m/z* 50-1100 range. Data
160 storage was in profile and centroid modes.

161

162

163 **2.3.3. LC-MS (LTQ-Orbitrap)**

164 An Accela LC system (Thermo Scientific, Hemel Hempstead, UK) was equipped with a
165 thermostatic autosampler and coupled to an LTQ Orbitrap Velos mass spectrometer
166 (Thermo Scientific, Hemel Hempstead, UK), with an ESI source. XCalibur software was
167 used for the data analyses.

168

169 The operation parameters used were source voltage, 3.5 kV; sheath gas (N₂): 40 (arbitrary
170 units); auxiliary gas (N₂): 10 (arbitrary units); sweep gas (N₂): 10 (arbitrary units); and
171 capillary temperature, 275°C. Default values were used for most other acquisition
172 parameters (Fourier transform (FT) automatic gain control (AGC) target 1·10⁶ for MS
173 mode and 5·10⁴ for MSⁿ mode). Milk samples were first analysed in full MS mode with the
174 Orbitrap mass resolving power set at 30,000 FWHM at *m/z* 400. The following analyses
175 were carried out in MSⁿ mode with the Orbitrap mass resolving power set at 15,000
176 FWHM at *m/z* 400. The maximum injection time was set to 100 ms with one micro scan
177 for MS mode and to 500 ms with one micro scan for MSⁿ mode. TPs were fragmented in
178 the HCD Collision Cell because some metabolites were not fragmented in the trap. The
179 selected HCD voltage was 45 V, although in some cases it was raised to 50 and 60 V. The
180 mass range was from *m/z* 100 to 1000.

181

182 **2.4. Procedures**

183 **2.4.1 Thermal study**

184 In this study, individual standards of quinolones (ENR, CIP, DIF and SAR) at 1 µg·mL⁻¹ in
185 water were warmed to degrade the antibiotics. All quinolones were heat-treated under
186 conditions of temperature and time corresponding to three thermal processes of milk:
187 pasteurization at 60°C for 30 min (T1) or 72°C for 15 s (T2) and sterilization at 120°C for
188 20 min (T3). Subsequently, individual milk samples spiked with ENR, CIP, DIF and SAR
189 were warmed at 120°C for 20 and 60 min (T3.20 and T3.60). The samples heated at
190 different temperatures were compared with spiked non-heated samples (T0) and with blank
191 milk samples at same thermal condition, ie, for example samples at T3.20 were compared
192 with blank at T3.20. Milk samples from animals medicated with ENR were also thermally

193 treated at T3.20 and T3.60 and compared with the same sample but non-heated. All
194 experiments were made by triplicate.

195

196 **2.4.2. Sample treatment and clean-up (SPE)**

197 The method for analyzing the milk sample was used previously to determine the antibiotics
198 in the milk⁹⁻¹². The extraction method consist of the addition of 0.5 mL of phosphate
199 solution 0.1 M at pH 10 and 2 mL of water to 2 g of milk, centrifugation of samples, and an
200 SPE process. The SPE cartridges used were Oasis HLB and these were preconditioned
201 with 1 mL of MeOH, 1 mL of water and 1 mL of 0.1 M phosphate solution at pH 10. After
202 sample loading, the cartridge was washed with 3 mL of water and the analytes were eluted
203 with 2 mL of MeOH.

204

205 The methanolic fraction eluted from the SPE was evaporated to dryness under a nitrogen
206 stream by TurboVap LV and a water temperature of 35°C. The extract was then
207 reconstituted with 200 µL of water and filtered through membrane filters (Ultra free
208 Durapore PVDF 0.22 µm from Millipore), before injection into the LC-LTQ-Orbitrap.

209

210 **2.4.2. Data treatment**

211 The LC-ToF data were treated using two procedures to find metabolites and TPs in milk
212 samples from medicated cow with ENR. The first corresponds to the manual comparison
213 of mass spectra of blanks and samples. This procedure was used initially to detect the TPs
214 in standards and spiked cow milk. The other option consists of the use of Mzmine2^{42,43} free
215 software, which provides a list of all m/z on mass spectra. The list can be shortened by
216 removing all the ions that only appear in the blank spectra and those that have a similar or
217 higher intensity in the blanks than in the samples. Only the ions which appeared between 1

218 to 9 min and from 150 to 800 Da were considered. Also, a mass defect filter (MDF) of 120
219 mDa was applied.

220

221

222 **3. RESULTS AND DISCUSSION**

223 **3.1. Determination of thermal TPs by LC-ToF-MS**

224 Blank and spiked water samples were subjected to the thermal conditions explained in
225 Section 2.3.1 and analysed by LC-ToF-MS. The total ion current chromatograms (TIC) of
226 spiked water samples were compared with the TIC of blank water samples to examine for
227 possible TPs. The two chromatograms were nearly identical, except for the spiked
228 antibiotic peak.

229

230 The mass spectra of blank and spiked water samples were compared by overlapping. The
231 only difference between the spiked sample and the blanks was the peak of parent
232 compound. However, a decrease in the area of the protonated molecule of quinolones was
233 noticed when the temperature rose. The highest decrease was seen when the samples were
234 heated at 120°C for 20 min (T3.20) where in some cases up to 20%. A new condition of
235 120°C for 60 min (T3.60) was also applied to ensure that TPs were obtained in sufficient
236 high concentrations to be detected.

237

238 Raw cow milk samples free of quinolones were spiked and heated under the selected
239 conditions. To assess whether any TPs had formed, the same procedure as that used for the
240 water standards was followed. In this case, some ions were identified in the mass spectra of
241 the spiked milk samples that were not observed in the blank milk samples. Some ions were
242 identified as doubly charged protonated molecules, with characteristic losses of water and

243 carbon dioxide. These ions were also detected in the water samples. The doubly protonated
244 identity was confirmed by the isotopic pattern. The difference between $[M+2H]^{2+}$ and
245 $[M+1+2H]^{2+}$ was m/z 0.5. In addition, the same retention time (t_R) of the compounds in the
246 extracted ion chromatogram (EIC) was observed⁴⁴. The losses of H₂O (18.0105) and CO₂
247 (43.9898) are not considered as TPs because these were also observed in the source by
248 CID-in-source fragmentation^{40,45-48}. The ions m/z 332.1405 and 386.1311 were detected in
249 samples that were spiked with ENR and DIF respectively. These m/z values are consistent
250 with the m/z of CIP and SAR. The EIC were obtained for blank and spiked samples to
251 check the presence of TPs. Table 1 shows all the ions found in milk and their
252 identification, sorted by the original antibiotic. As can be observed, the errors between
253 experimental mass and assigned structure were lower than 3.0 ppm.

254

255 To assign a structure to the remaining of the ions that were observed, an in-house database
256 was created using a list of 106 metabolic reactions⁴⁹ that can take place in milk, and
257 another list of milk components (including carbohydrates, amino acids, vitamins and lipids;
258 a total of 188 components)⁵⁰. The exact mass of the starting antibiotic or a known fragment
259 is introduced into the database and a series of combinations of reactions and/or possible
260 additions are obtained that give rise to a molecule with an exact mass similar to that
261 observed in the mass spectra of ToF.

262

263

264 **3.2. Identification of thermal TPs by LC-LTQ-Orbitrap-MS**

265 As a ToF spectrometer cannot fragment molecules, we used an LTQ-Orbitrap to elucidate
266 the TPs, as it has a higher mass resolving power than ToF and can fragment the compound
267 to obtain structural information.

268

269 For each quinolone, Table 2 shows the retention time (t_R), m/z values, the gains and losses
270 of mass compared to the parent quinolone, the m/z of the TPs observed, and the fragments
271 observed in mass spectra. Table 2 also shows the proposed reactions, with the assigned
272 molecular formula, the double bond equivalents (DBE) of neutral molecules, the
273 theoretical mass, and the error in ppm calculated for the formula. The information is
274 provided as common reactions that occurred in all the studied quinolones, and other
275 reactions that were only observed in some quinolones.

276

277 *Common reactions of the four quinolones*

278

279 Several gains and losses of mass were common among the four quinolones studied. Our
280 database shows that the gain of 15.9949 m/z (Table 2) could correspond to two different
281 reactions: hydroxylation and N-oxide formation. An evaluation of MS/MS spectra could be
282 used to differentiate between both structures. As an example, Figure 2A shows the MS/MS
283 spectrum of ions m/z 416.1416 in samples of DIF and discussion about elucidation is given
284 in supporting information. The position of OH could not be assigned with certainty, but our
285 proposal is hydroxylation in position 3 of the piperazine ring, as in oxo-ciprofloxacin, the
286 metabolite described in the bibliography^{36,37,40}. For the four quinolones, the hydroxylated-
287 TPs were observed, but the concentration of TPs from ENR and DIF were higher than CIP
288 and SAR ones. N-oxide formation only occurred in ENR and DIF in N⁴-alkylpiperazinyl.
289 This metabolite has been described previously^{38,51}. For this structure, a loss of OH is
290 characteristic in fragmentation and was experimentally observed in Figure 2B of the
291 MS/MS spectrum of ions m/z 416.1416 for DIF. In this case, the DBEs of TPs undergo no
292 alteration of values in comparison to the DBE of parent quinolones. Different

293 fragmentation patterns were observed for the hydroxylated and N-oxide compounds that
294 were found, as can be observed in Figure 2A and B. In Figure 2B, corresponding to the N-
295 oxide TP, the peak of the precursor ion disappeared almost completely when the same
296 voltage was applied (HCD: 45). This finding suggests that hydroxylated compounds are
297 more stable than N-Oxide TP. The structures assigned to the rest of the ions are shown in
298 Figure 2. Table 2 shows the fragmentation ions, the molecular formula, the DBE, and the
299 calculated error in the assignment of the different structures.

300

301 A gain of 29.9740 m/z was also observed for the four quinolones, as shown in Table 2.
302 Three peaks were found for ENR and DIF, whereas only one peak was observed for their
303 main metabolites (CIP and SAR), at a concentration too low to be isolated and fragmented.
304 The mass spectra of all peaks of ENR and DIF were studied. Figure 3 shows the mass
305 spectra of ions with m/z 390.1460 of ENR. In Figure 3A, the separation by LC-LTQ-
306 Orbitrap is shown with peaks at 5.71, 7.57 and 7.99 min. Although the retention time of
307 ENR matches peak 1, the mass spectra are different, as can be observed in Figures 3B and
308 C. A comparison of both mass spectra revealed only two ions (m/z 245.1077 and 72.0805)
309 in common. The m/z 245.1077 indicates that the differences between the TP of peak 1 and
310 ENR are in the loss fragment (part of the piperazinyl ring). The accurate mass of the
311 fragment m/z 128.0706 corresponds to molecular formula $C_6H_{10}NO_2$ with a DBE value of
312 3 and is the key to the elucidation as is shown in Figure 3C and in the supporting
313 information⁵².

314

315 Unlike the first molecule, the other two ions of m/z 390.1460 (peaks 2 and 3) had low
316 fragmentation as can be observed in Figure 3. Nevertheless, their structures were
317 elucidated. The retention time of these peaks are different, but their mass spectra are

318 identical, which indicates that they are isomers (Figure 3D). The m/z 245.1085 observed in
319 the mass spectrum of ENR (Figure 3B) could not be found in the spectra of peaks 2 and 3.
320 The molecules were broken up from the piperazinyl ring, resulting in the ions 100.0756
321 and 291.0774. The fragment 100.0756 is the same as that observed in Figure 3C, which
322 indicates that there was only a ketone in the ring. The fragmentation of 291.0774 makes us
323 to conclude that OH is on the aromatic ring. Two different vacant positions in the aromatic
324 ring led to the two different isomers. For DIF, the fragmentations were similar to those
325 shown in ENR, as can be observed in Table 2.

326
327 TPs corresponding to net loss of 26.0146 were observed for all compounds (ENR, CIP,
328 DIF and SAR). For the TPs of ENR and DIF the losses of N^4 -alkylamine from piperazinyl
329 ring were observed. However, as CIP and SAR do not have an alkyl group in N^4 , the losses
330 of ammonia (17.0275) were seen. Moreover, the TPs had DBE one unit lower than the
331 parent compound, due to the break in the piperazinyl ring. Therefore, the peaks at m/z
332 334.1560, 306.1246, 374.1307 and 360.1156 have been assigned to desethylenequinolone
333 TPs³⁴⁻³⁷. As Table 2 shows, when Orbitrap was used to elucidate the structures, the
334 calculation errors (ppm) were lower than 2 ppm. Slightly higher error values were obtained
335 when ToF was used.

336

337 *Other reactions*

338 The major metabolites, CIP and SAR, underwent other transformations, as can be observed
339 in Table 2. Due to the absence of the alkyl chain in the N^4 -piperazinyl ring, two reactions
340 acetylation (Acetyl-CIP (m/z 374.1509) and Acetyl-SAR (m/z 428.1416)) and
341 formamidation (Formamide-CIP (m/z 375.1463) and Formamide-SAR (m/z 429.1416))
342 took place. The corresponding explanation of elucidation is in supplementary material.

343

344 In addition to the above-mentioned reactions, CIP gave three more TPs (m/z : 274.0986,
345 573.1429 and 656.2457). The ion 573.1429 was observed at a very low concentration and
346 could not be fragmented. For the mass 274.0986, the ion 292.1091 could also be observed,
347 at less intensity, corresponding to the gain of a water molecule. The main ion obtained by
348 MS² of the ion 274.0986 was 231.0564, and this was isolated and fragmented again. In the
349 MS³ spectrum, the ion m/z 249.0668 was also observed with m/z 231.0564 which
350 corresponds to gain of a water molecule. This gain of water was only observed in the ions
351 that kept the carbonyl group of acid, which is capable of hydrating the initial carboxylic
352 acid again. The rest of the ions were mainly due to fragmentation of the piperazine ring.
353 The structure assigned to m/z 292.1091 was CIP, with loss of the cyclopropyl group.

354

355 The last TP (656.2457) was obtained by heating the spiked milk sample with CIP at T3.60.
356 When a HCD voltage of 45 V was used, only fragments with low m/z were found. To
357 obtain structural information, the voltage was reduced to 35 V. The CIP ion (m/z 332.1400)
358 could be identified among all the fragments as can be seen in Figure 4. As can be followed
359 in supporting information, the molecule CIP (332.1405) is combined with the lactose
360 (galactose + glucose) present in milk at high concentration. This new product was favoured
361 when samples were heated at a high temperature for a long time (60 min).

362

363 **3.3. Effect of temperature on the stability of TPs**

364 For each TP ion, the areas obtained from milk samples were plotted vs. the heating
365 conditions (T0, T3.20 and T3.60) (Figure 5). For ENR and CIP (Figure 5A and 5B), all the
366 TPs were observed at T0 except for hydroxylated quinolones (m/z 376.1656_1 and
367 348.1334, respectively). The concentration of these ions was raised by heating at T3.20,
368 and disappeared at T3.60. In the case of ENR (Figure 5A), the concentration of some TPs,

369 including CIP (332.1402), desethylene-ENR (334.1569) and two isomers of hydroxylation
370 and ketone formation (390.1461_1 and 390.1461_3), increased slightly by warming at
371 T3.20, and decreased at T3.60. For CIP, T3.60 was required for conjugation with lactose
372 (656.2477) to occur (Figure 5B). The acetylated (374.1535) and formamidated (375.1463)
373 were present at T0 and their concentrations increased by heating. The 362.1148 ion
374 (corresponding to gain $[M+H+29.974]^+$) was initially present (T0), but was destroyed by
375 heating.

376

377 In contrast, the concentration of most of the ions in DIF samples increased slightly when
378 the temperature was increased (Figure 5C). A general downward trend of DIF ions was
379 seen at T3.60. Completely different behaviour was observed for the 430.1185_3 ion, which
380 underwent sharp decrease at T3.20 and a sharp rise in formation when the warming time
381 was increased, possibly due to the ion coming from different sources, and for the
382 430.1185_1 ion, which disappeared completely at T3.20. The acetylated (428.1407) and
383 formamidated (429.1391) TPs of SAR (Figure 5D) behaved in the same way as those of
384 CIP, *i.e.*, their concentration increased with temperature.

385

386 **3.4 Determination of metabolites and TPs in medicated cow milk**

387 Four raw milk samples from cows medicated with ENR were submitted to the same
388 thermal treatments as the spiked milk samples. The extracts were analysed by LC-ToF and
389 the metabolites and TPs were fragmented by LC-LTQ-Orbitrap.

390

391 In this case, metabolites and TPs were found using the two data treatments (Section 2.3.3).
392 When exact mass, retention time and MS/MS spectra were compared, some of the
393 compounds in milk spiked with ENR and CIP were also present in medicated cow milk.

394

395 Table 3 shows the retention time (t_R) of metabolites in samples analysed by LC-ToF, the
396 accurate m/z of metabolites determined by ToF and LTQ-Orbitrap, the assigned molecular
397 formula, the theoretical mass, the error in ppm calculated for the assigned formula, the
398 proposed structures, and the double bond equivalents (DBE) of the neutral molecule.

399

400 CIP (m/z 332.1405), the main metabolite of ENR, was one of the common compounds. In
401 addition, desethylene metabolites of ENR (m/z 334.1561)(ENR-6) and CIP (m/z
402 306.1248)(ENR-3) were observed. The TPs of ENR corresponded to hydroxylation (ENR-
403 16) and N-Oxide formation (ENR-17)(m/z 376.1667) and two isomers of compound m/z
404 390.1460 (ENR-18 and ENR-19) (hydroxylation and ketone formation, Figure 3D) were
405 also found. Four TPs of CIP were observed in medicated cow milk. At 3.6 min was found
406 the hydroxy-CIP (m/z 348.1354)(ENR-8) and at 6.1 min the m/z 362.1147(ENR-11),
407 although this time the MS² spectrum could not be obtained due to its low concentration.
408 The other metabolites were from acetylation (m/z 374.1511)(ENR-14) and formamidation
409 (m/z 375.1463)(ENR-15) of CIP.

410

411 Other compounds were found in milk from medicated cows, but not in spiked milk. These
412 compounds were produced by metabolism of the antibiotic in the cow, and some of them
413 were identified for the first time.

414

415 The metabolite with the highest intensity after CIP (332.1405) and desethylene-ENR
416 (334.1561)(ENR-6) had m/z 263.0834 (ENR-1). The m/z of the metabolite was lower than
417 the m/z of ENR. Therefore, the MS/MS spectrum was analysed to find which part of the
418 parent compound had been modified. Figure 6A shows MS/MS spectrum of ENR-1. It

419 was concluded that the metabolite structure corresponded to CIP without a piperazine ring
420 (Supporting information). Only the amino group was maintained, as shown in Table 3.

421

422 In the MS/MS spectrum of metabolite ENR-2 with m/z 291.0783 (Figure 6B), the ion m/z
423 263.0834 (ENR-1), discussed above in supporting information and in Figure 6A, was
424 observed. The difference between both ions corresponded to a carbonyl group (27.9949).
425 The assigned structure for this metabolite is given in Table 3.

426

427 Another metabolite elucidated was m/z 307.1095 (ENR-4). Some explanations of the
428 fragments are shown in Figure 6C and supporting information. Its structure can be
429 observed in Table 3.

430

431 The ion m/z 334.1189 (ENR-5) was identified as formyl-desethyleno-CIP²⁵. Its structure
432 was checked by its mass spectrum. The molecule was broken up from the piperazinyl ring,
433 resulting in the ions m/z 72.0445 and 263.0834 cited before. Other important fragments
434 were m/z 245.0730 [263.0834-H₂O], 230.0489 [334.1189-H₂O-Cyclopropyl-Formamide]
435 and 217.0411 [230.0489-CH].

436

437 A compound (ENR-9) with the same m/z as hydroxy-CIP (m/z 348.1354)(ENR-8) was also
438 studied, because the retention time differed by about 4 minutes from the peak observed in
439 spiked samples. The MS/MS spectrum (Figure 6D) was very similar to the spectrum of
440 formyl-desethyleno-CIP (m/z 334.1189)(ENR-5). The most striking difference between both
441 spectra was that instead of fragment m/z 72.0444 was observed a fragment of m/z 86.0601.
442 This fragment corresponds to the addition of a methyl group to fragment m/z 72.0444. In

443 conclusion, a structure of acetyldesethylene-CIP (ENR-9) is suggested, which could be
444 formed from the oxidation of the side chain of desethylene-ENR (m/z 334.1561)(ENR-6).

445

446 The EIC of ion m/z 362.1137 (ENR-12 and ENR-13) from a full-scan of ToF was very
447 similar to that obtained for the m/z 390.1460 (ENR-18 and ENR-19) in spiked samples
448 with ENR (Figure 3). Three peaks were observed in the EIC. The first one was at 6.1 min,
449 which coincides with the peak observed in samples fortified with CIP. Also were seen two
450 peaks at 6.8 and 7.1 min, with the same shape as the two peaks at 7.6 and 7.99 in Figure 3.
451 Unfortunately, when the sample was analysed by LC-LTQ-Orbitrap, it was degraded and
452 only one peak was observed and fragmented. In the MS/MS spectrum, the ions m/z
453 291.0776 and 259.0820 were present, as elucidated previously in Figure 3D. Another
454 fragment that helped to determine the structure corresponded to m/z 72.0444, which is like
455 fragment 100.0757 of metabolite 390.1460 without the alkyl chain of ENR. Due to the
456 resemblances between the EICs and mass spectra of ions m/z 362.1137 (ENR-12 and ENR-
457 13) and 390.1460 (ENR-18 and ENR-19), can be assume that these compounds are the
458 same, but one is from CIP and the other from ENR, respectively.

459

460 The last two elucidated metabolites corresponded to oxo-CIP (m/z 346.1198)(ENR-7) and
461 formyl-CIP (360.1354)(ENR-10), which have been described previously in literature^{36,40}.
462 Their structures were confirmed by their MS/MS spectra.

463

464 The rest of the metabolites found in medicated cow milk could not be elucidated. In any
465 case, Table 3 shows their m/z , retention time (t_R), hypothetical molecular formula, double
466 bond equivalents (DBE) and errors in ppm.

467

468 **AUTHOR INFORMATION**

469 **Funding source**

470 The authors are grateful to the Ministerio de Ciencia e Innovación (MICINN) (Project
471 CTQ2010-19044/BQU) for financial support. A.J. received support from MICINN (BES-
472 2011-045849) for the training of researchers.

473

474 **Supporting Information Available**

475 Explanation about MS elucidation of some TPs and metabolites of quinolones (ENR, CIP,
476 DIF and SAR) has been included. This material is available, free of charge, via the Internet
477 at <http://pubs.acs.org>.

478

479

480

481 **REFERENCES**

482

483 1. Council Regulation No 1831/2003 of the European Parliament and of the Council of 22
484 September 2003 on additives for use in animal nutrition.

485 2. Hammerum, A.M.; Heuer, O.E.; Lester, C.H.; Agers, Y.; Seyfarth, A.M.; Emborg, H.D.;
486 Frimodt-Moller, N.; Monnet, D.L. Comment on: withdrawal of growth-promoting
487 antibiotics in Europe and its effects in relation to human health. *Int J Antimicrob Agents*
488 **2007**, 30, 466-468.

489 3. Fábrega, A.; Sánchez-Céspedes, J.; Soto, S.; Vila, J. Quinolone resistance in the food
490 chain. *Int J Antimicrob Agents* **2008**, 31, 307-315.

491 4. Commission of the European Communities, Official Journal of the European
492 Communities No. 37/2010 L15/1, December 22, 2009.

493 5. Zorraquino, M.A.; Althaus, R.L.; Roca, M.; Molina, M.P. Effect of heat treatments on
494 aminoglycosides in milk. *J Food Prot.* **2009**, 72, 1338-1341.

495 6. Early, R. The technology of dairy products. Edition 2.; Ed. Aspen Publishers Inc.;
496 Gaithersburg, Maryland, USA, **1998**.

497 7. Picó, Y.; Barceló, D. The expanding role of LC/MS in analyzing metabolites and
498 degradation products of food contaminants. *Trends Anal Chem.* **2008**, 27, 821-835.

499 8. Becker, M.; Zittlan, E.; Petz, M. Residue analysis of 15 penicillins and cephalosporins in
500 bovine muscle, kidney and milk by liquid chromatography–tandem mass spectrometry.
501 *Anal Chim Acta* **2004**, 520, 19-32.

502 9. Junza, A.; Amatya, R.; Barrón, D.; Barbosa, J. Comparative study of the LC–MS/MS
503 and UPLC–MS/MS for the multi-residue analysis of quinolones, penicillins and

504 cephalosporins in cow milk, and validation according to the regulation 2002/657/EC. *J*
505 *Chromatogr B* **2011**, 879, 2601-2610.

506 10. Reyns, T.; Cherlet, M.; De Baere, S.; De Backer, P.; Croubels, S. Rapid method for the
507 quantification of amoxicillin and its major metabolites in pig tissues by liquid
508 chromatography-tandem mass spectrometry with emphasis on stability issues. *J*
509 *Chromatogr B* **2008**, 861, 108-116.

510 11. Granelli, K.; Elgerud, C.; Lundström, A.; Ohlsson, A.; Sjöberg, P. Rapid multi-residue
511 analysis of antibiotics in muscle by liquid chromatography–tandem mass spectrometry.
512 *Anal Chim Acta* **2009**, 637, 87-91.

513 12. Martínez-Huélamo, M.; Jiménez-Gámez, E.; Hermo, M.P.; Barrón, D.; Barbosa, J.
514 Determination of penicillins in milk using LC-UV, LC-MS and LC-MS/MS. *J Chromatogr*
515 *A* **2009**, 32, 2385-2393.

516 13. Nemetlu, E.; Kir, S.; Katlan, D.; Beksaç, M.S. Simultaneous multiresponse
517 optimization of an HPLC method to separate seven cephalosporins in plasma and amniotic
518 fluid: Application to validation and quantification of cefepime, cefixime and cefoperazone.
519 *Talanta* **2009**, 80, 117-126.

520 14. Mol, H.G.J.; Plaza-Bolaños, P.; Zomer, P.; de Rijk, T.C.; Stolker, A.A.M.; Mulder,
521 P.P.J. Toward a Generic Extraction Method for Simultaneous Determination of Pesticides,
522 Mycotoxins, Plant Toxins, and Veterinary Drugs in Feed and Food Matrixes. *Anal Chem.*
523 **2008**, 80, 9450-9459.

524 15. Marazuela, M.D.; Bogialli, S. A review of novel strategies of sample preparation for
525 the determination of antibacterial residues in foodstuffs using liquid chromatography-based
526 analytical methods. *Anal Chim Acta* **2009**, 645, 5-17.

- 527 16. Moreno-Bondi, M. C.; Marazuela, M.D.; Herranz, S.; Rodríguez, E. An overview of
528 sample preparation procedures for LC-MS multiclass antibiotic determination in
529 environmental and food samples. *Anal Bioanal Chem.* **2009**, 395, 921-946.
- 530 17. Blasco, C.; Torres, C.M.; Picó, Y. Progress in analysis of residual antibacterials in
531 food. *Trends Anal Chem.* **2007**, 26, 895-913.
- 532 18. Andreu, V.; Blasco, C.; Picó, Y. Analytical strategies to determine quinolone residues
533 in food and the environment. *Trends Anal Chem.* **2007**, 26, 534-556.
- 534 19. Bogialli, S.; Di Corcia, A. Recent applications of liquid chromatography–mass
535 spectrometry to residue analysis of antimicrobials in food of animal origin. *Anal Bioanal*
536 *Chem.* **2009**, 395, 947-966.
- 537 20. Tang, Q.; Yang, T.; Tan, X.; Luo, J. Simultaneous Determination of Fluoroquinolone
538 Antibiotic Residues in Milk Sample by Solid-Phase Extraction - Liquid Chromatography -
539 Tandem Mass Spectrometry. *J.Agric. Food Chem.* **2009**, 57, 4535-4539.
- 540 21. Romero-González, R.; Aguilera-Luiz, M.M.; Plaza-Bolaños, P.; Garrido Frenich, A.;
541 Martínez Vidal, J.L. Food contaminant analysis at high resolution mass spectrometry:
542 Application for the determination of veterinary drugs in milk. *J Chromatogr A* **2011**, 1218,
543 9353-9365.
- 544 22. Kaufmann, A. Strategy for the elucidation of elemental compositions of trace analytes
545 based on a mass resolution of 100 000 full width at half maximum. *Rapid Commun Mass*
546 *Spectrom.* **2010**, 24, 2035-2045.
- 547 23. Zhu, M.; Ma, L.; Zhang, D.; Ray, K.; Zhao, W.; Humphreys, W.G.; Skiles, G.;
548 Sanders, M.; Zhang, H. Detection and Characterization of Metabolites in Biological
549 Matrices Using Mass Defect Filtering of Liquid Chromatography/High Resolution Mass
550 Spectrometry Data. *Drug Metab Dispos* **2006**, 34, 1722-1733.

- 551 24. Nägele, E.; Moritz, R. Structure Elucidation of Degradation Products of the Antibiotic
552 Amoxicillin with Ion Trap MSⁿ and Accurate Mass Determination by ESI TOF. *J Am Soc*
553 *Mass Spectr.* **2005**, 16, 1670-1676.
- 554 25. Calza, P.; Medana, C.; Carbone, F.; Giancotti, V.; Baiocchi, C. Characterization of
555 intermediate compounds formed upon photoinduced degradation of quinolones by high-
556 performance liquid chromatography/high-resolution multiple-stage mass spectrometry.
557 *Rapid Commun Mass Spectrom.* **2008**, 22, 1533-1552.
- 558 26. Gómez, M.J; Gómez-Ramos, M.M.; Malato, O.; Mezcuca, M.; Fernández-Alba, A.R.
559 Rapid automated screening, identification and quantification of organic micro-
560 contaminants and their main transformation products in wastewater and river waters using
561 liquid chromatography–quadrupole-time-of-flight mass spectrometry with an accurate-
562 mass database. *J Chromatogr A* **2010**, 1217, 7038-7054.
- 563 27. Kaufmann, A. Determination of the elemental composition of trace analytes in complex
564 matrices using exact masses of product ions and corresponding neutral losses. *Rapid*
565 *Commun Mass Spectrom.* **2007**, 21, 2003-2013.
- 566 28. Berendsen, B.J.A.; Essers, M.L.; Mulder, P.P.J.; van Bruchem Lommen, A.; van
567 Overbeek, W.M., Stolker, L.A.M. Newly identified degradation products of ceftiofur and
568 cephapirin impact the analytical approach for quantitative analysis of kidney. *J*
569 *Chromatogr A* **2009**, 1216, 8177-8186.
- 570 29. Krauss, M.; Singer, H.; Hollonder, J. LC–high resolution MS in environmental
571 analysis: from target screening to the identification of unknowns. *Anal Bioanal Chem.*
572 **2010**, 397, 943-951.
- 573 30. Prasad, B.; Garg, A.; Takwani, H.; Singh, S. Metabolite identification by liquid
574 chromatography-mass spectrometry. *Trends Anal Chem.* **2011**, 30, 360-387.

- 575 31. Lacorte, S.; Fernández-Alba, A.R. Time of flight mass spectrometry applied to the
576 liquid chromatography analysis of pesticides in water and food. *Mass Spectrom Rev.* **2006**,
577 25, 866-880.
- 578 32. Malik, A.K.; Blasco, C.; Picó, Y. Liquid chromatography–mass spectrometry in food
579 safety. *J Chromatogr A* **2010**, 1217, 4018-4040.
- 580 33. Koulman, A.; Lane, G.A.; Harrison, S.J.; Volmer, D.A. From differentiating
581 metabolites to biomarkers. *Anal Bioanal Chem.* **2009**, 394, 663-670.
- 582 34. Jacobs, D.M.; Spiesser, L.; Garnier, M.; de Roo, N.; van Dorsten Hollebrands, B.; van
583 Velzen, E.; Draijer, R.; van Duynhoven, J. SPE–NMR metabolite sub-profiling of urine.
584 *Anal Bioanal Chem.* **2012**, 404, 2349-2361.
- 585 35. Sturini, M.; Speltini, A.; Maraschi, F.; Profumo, A.; Pretali, L.; Fasani, E.; Albini, A.
586 Photochemical Degradation of Marbofloxacin and Enrofloxacin in Natural Waters. *Envir*
587 *Sci Thecno.* **2010**, 144, 4564-4569.
- 588 36. Anadón, A.; Martínez-Larrañaga, M.R.; Iturbe, J.; Martínez, M.A.; Díaz, M.J.; Frejo,
589 M.T.; Martínez, M. Pharmacokinetics and residues of ciprofloxacin and its metabolites in
590 broiler chickens. *Res Vet Sci.* **2001**, 71, 101-109.
- 591 37. Turnipseed, S. B.; Storey, J.M.; Clark, S.B.; Miller, K.E. Analysis of Veterinary Drugs
592 and Metabolites in Milk Using Quadrupole Time-of-Flight Liquid Chromatography–Mass
593 Spectrometry. *J Agric Food Chem.* **2011**, 59, 7569-7581.
- 594 38. Sukul, P.; Lamshöft, M.; Kusari, S.; Zühlke, S.; Spiteller, M. Metabolism and excretion
595 kinetics of ¹⁴C-labeled and non-labeled difloxacin in pigs after oral administration, and
596 antimicrobial activity of manure containing difloxacin and its metabolites. *Environ Res.*
597 **2009**, 109, 225-231.

- 598 39. Turiel, E.; Bordin, G.; Rodríguez, A.R. Study of the evolution and degradation
599 products of ciprofloxacin and oxolinic acid in river water samples by HPLC-UV/MS/MS-
600 MS. *J Environ Monit.* **2005**, 7, 189-195.
- 601 40. Volmer, D.A.; Mansoori, B.; Locke, S.J. Study of 4-Quinolone Antibiotics in
602 Biological Samples by Short-Column Liquid Chromatography Coupled with Electrospray
603 Ionization Tandem Mass Spectrometry. *Anal Chem.* **1997**, 69, 4143-4155.
- 604 41. Terzic, S.; Senta, I.; Matosic, M.; Ahel, M. Identification of biotransformation products
605 of macrolide and fluoroquinolone antimicrobials in membrane bioreactor treatment by
606 ultrahigh-performance liquid chromatography/quadrupole time-of-flight mass
607 spectrometry.) *Anal Bioanal Chem.* **2011**, 401, 353-363.
- 608 42. <http://mzmine.sourceforge.net> (17/01/2014)
- 609 43. Pluskal, T.; Castillo, S.; Villar-Briones, A.; Oresic, M. MZmine 2: Modular framework
610 for processing, visualizing, and analyzing mass spectrometry-based molecular profile data.
611 *Bioinformatics* **2010**, 11, 395-405
- 612 44. Wu, L.; Liu, D.Q. Data-dependent tandem mass spectrometry of doubly protonated
613 molecules in pharmaceutical analysis. *Rapid Commun Mass Spectrom.* **2007**, 21, 455-458.
- 614 45. Clemente, M.; Hermo, M.P.; Barrón, D.; Barbosa, J. Confirmatory and quantitative
615 analysis using experimental design for the extraction and liquid chromatography–UV,
616 liquid chromatography–mass spectrometry and liquid chromatography–mass
617 spectrometry/mass spectrometry determination of quinolones in turkey muscle. *J*
618 *Chromatogr A* **2006**, 1135, 170-178
- 619 46. Picó, Y.; Andreu, V. Fluoroquinolones in soil—risks and challenges. *Anal Bioanal*
620 *Chem.* **2007**, 387, 1287-1299.
- 621 47. Hermo, M.P.; Barrón, D.; Barbosa, J. Multiresidue determination of quinolones
622 regulated by the European Union in bovine and porcine plasma. Application of

623 chromatographic and capillary electrophoretic methodologies. *J Chromatogr A* **2008**, 1201,
624 1-14.

625 48. Kaufmann, A.; Butcher, P.; Maden, K.; Widmer, M.; Giles, K.; Uría, D. Are liquid
626 chromatography/electrospray tandem quadrupole fragmentation ratios unequivocal
627 confirmation criteria? *Rapid Commun Mass Spectrom.* **2009**, 23, 985-998.

628 49. Thermo Scientific Analytical Tools for Metabolism. Biotransformation Reference
629 Guide 2009

630 50. <http://foodbs.org/examples/milk>. (17/01/2014)

631 51. Rust, H.; Scheer, M.; Hallenbach, W. BAYER AG, Preparation of quinolone and
632 naphthylidinone N-oxides as antibacterials. Patent ES 2 161 361 T3.

633 52. Badertscher, M.; Bischofberger, K.; Munk, M.E.; Pretsch, E. A Novel Formalism to
634 Characterize the Degree of Unsaturation of Organic Molecules. *J Chem Inf Comput Sci.*
635 **2001**, 41, 889-893.

636

637

638 **FIGURE CAPTIONS**

639

640 **Fig 1**

641 Structures of the four quinolones studied (ENR, CIP, DIF and SAR).

642

643 **Fig 2**

644 A) MS² spectrum for the hydroxylated TP of DIF. B) MS² spectrum for the N-oxide TP of
645 DIF.

646

647 **Fig 3**

648 A) EIC of ion m/z 390.1460 in the milk sample spiked with ENR. B) MS² spectrum for
649 ENR. C) MS² spectrum for the first peak (t_R 5.71 min). D) MS² spectrum for the remaining
650 peaks (t_R 7.57 and 7.99 min).

651

652 **Fig 4**

653 Product ion scan (MS²) of CIP lactose conjugate (precursor ion m/z 656.2457). Top left,
654 extended mass spectrum where losses of water molecules are shown.

655

656 **Fig 5**

657 Influence of the different warming conditions in areas of different TPs of ENR, CIP, DIF
658 and SAR.

659


660 **Fig 6**

661 MS² spectra for the metabolites of ENR. A) m/z 263.0828 (ENR-1). B) m/z 291.0776
662 (ENR-2). C) m/z 307.1089 (ENR-4). D) m/z 348.1354 (ENR-9).

663

664 **TABLE OF CONTENTS GRAPHIC**

665



666

Table 1. Accurate masses of ions observed in milk spiked samples but not in blank milk samples by LC-ToF. The errors (ppm) are calculated from the theoretical and proposed structures.

ENR		CIP				SAR					
<i>m/z</i> (exp)	<i>m/z</i> (theoretical)	Structure	Error (ppm)	<i>m/z</i> (exp)	<i>m/z</i> (theoretical)	Structure	Error (ppm)	<i>m/z</i> (exp)	<i>m/z</i> (theoretical)	Structure	Error (ppm)
157.5685	157.5686	[CIP+2H-H ₂ O] ²⁺	0.6	157.5690	157.5686	[CIP+2H-H ₂ O] ²⁺	2.5	184.5637	184.5639	[SAR+2H-H ₂ O] ²⁺	1.1
166.5740	166.5739	[CIP+2H] ²⁺	0.6	166.5742	166.5739	[CIP+2H] ²⁺	1.8	193.5692	193.5692	[SAR+2H] ²⁺	0.1
167.5820	n.a.*	[334.1569] ²⁺	-	274.0989	n.a.	n.a.	-	342.1422	342.1412	[SAR+H-CO ₂] ⁺	2.9
171.5845	171.5843	[ENR+2H-H ₂ O] ²⁺	1.2	288.1511	288.1507	[CIP+H-CO ₂] ⁺	1.4	360.1181	n.a.	n.a.	-
180.5897	180.5897	[ENR+2H] ²⁺	0.1	314.1293	314.1288	[CIP+H-H ₂ O] ⁺	1.6	368.1193	368.1205	[SAR+H-H ₂ O] ⁺	3.3
316.1817	316.1820	[ENR+H-CO ₂] ⁺	0.9	332.1400	332.1405	[CIP+H] ⁺	1.5	386.1311	386.1311	[SAR+H] ⁺	0.1
332.1402	332.1405	[CIP+H] ⁺	0.9	348.1334	n.a.	n.a.	-	402.1241	n.a.	n.a.	-
334.1569	n.a.	n.a.	-	362.1148 ^b	n.a.	n.a.	-	416.1069	n.a.	n.a.	-
342.1603	342.1612	[ENR+H-H ₂ O] ⁺	2.6	374.1509	n.a.	n.a.	-	428.1410	n.a.	n.a.	-
360.1719	360.1718	[ENR+H] ⁺	0.3	375.1563	n.a.	n.a.	-	429.1395	n.a.	n.a.	-
376.1656 ^a	n.a.	n.a.	-	573.1431	n.a.	n.a.	-				
390.1461 ^b	n.a.	n.a.	-	656.2477	n.a.	n.a.	-				
719.3349	719.3363	[2ENR+H] ⁺	1.9								
DIF		CIP				SAR					
<i>m/z</i> (exp)	<i>m/z</i> (theoretical)	Structure	Error (ppm)	<i>m/z</i> (exp)	<i>m/z</i> (theoretical)	Structure	Error (ppm)	<i>m/z</i> (exp)	<i>m/z</i> (theoretical)	Structure	Error (ppm)
191.5714	191.5717	[DIF+2H-H ₂ O] ²⁺	0.5	184.5637	184.5639	[SAR+2H-H ₂ O] ²⁺	1.1	193.5692	193.5692	[SAR+2H] ²⁺	0.1
200.5761	200.5770	[DIF+2H] ²⁺	3.0	342.1422	342.1412	[SAR+H-CO ₂] ⁺	2.9	360.1181	n.a.	n.a.	-
299.0980	n.a.	n.a.	-	360.1181	n.a.	n.a.	-	368.1193	368.1205	[SAR+H-H ₂ O] ⁺	3.3
374.1284	n.a.	n.a.	-	368.1193	368.1205	[SAR+H-H ₂ O] ⁺	3.3	386.1311	386.1311	[SAR+H] ⁺	0.1
382.1367	382.1362	[DIF+H-H ₂ O] ⁺	1.3	386.1311	386.1311	[SAR+H] ⁺	0.1	402.1241	n.a.	n.a.	-
386.2209	386.1311	[SAR+H] ⁺	0.3	402.1241	n.a.	n.a.	-	416.1069	n.a.	n.a.	-
400.1446	400.1467	[DIF+H] ⁺	2.0	416.1069	n.a.	n.a.	-	428.1410	n.a.	n.a.	-
416.1396 ^a	n.a.	n.a.	-	428.1410	n.a.	n.a.	-	429.1395	n.a.	n.a.	-
430.1185 ^b	n.a.	n.a.	-	429.1395	n.a.	n.a.	-				

(*) n.a: Non-assigned.

(a) 2 chromatographic peaks with the same *m/z* (at different retention time).

(b) 3 chromatographic peaks with the same *m/z* (at different retention time).

Table 2. Accurate masses of parent quinolones, given reactions, all TPs observed and the fragments obtained by LC-LTQ-Orbitrap. The DBE calculated with the deprotonated molecular formula and the calculated error (ppm) of the assignment of structure.

	t_R^a (min)	m/z	Gains or losses	Reactions (*)	m/z TPs	Fragmentation	Molecular formula	DBE	Theoretical mass	Error (ppm)
Common reactions of 4 quinolones										
ENR	4.7	360.1718	+15.9949	HX	376.1666	358.1560, 340.1121, 286.1018, 244.0912, 243.0877	$C_{19}H_{23}FN_3O_4$	10	376.1667	-0.3
ENR	5.8	360.1718	+15.9949	N-O	376.1665	359.1639, 330.1611, 315.1741, 300.1506, 257.1085, 231.0928, 84.0807	$C_{19}H_{23}FN_3O_4$	10	376.1667	-0.5
CIP	3.6	332.1405	+15.9949	HX	348.1355	330.1248, 310.1184, 136.0617	$C_{17}H_{19}FN_3O_4$	10	348.1354	0.3
DIF	6.0	400.1467	+15.9949	HX	416.1416	398.1307, 343.0880, 329.0725, 316.0647, 256.0801	$C_{21}H_{20}F_2N_3O_4$	13	416.1416	0.0
DIF	6.3	400.1467	+15.9949	N-O	416.1418	399.1388, 370.1359, 355.1489, 299.09899	$C_{21}H_{20}F_2N_3O_4$	13	416.1416	0.5
SAR	5.8	386.1311	+15.9949	HX	402.1261	384.1156, 382.1197, 364.1089, 343.0882, 316.0653, 136.0758	$C_{20}H_{18}F_2N_3O_4$	13	402.1260	0.2
ENR	5.8	360.1718	+29.9740	HX + K	390.1458	372.1353, 362.1509, 346.1560, 318.1246, 301.0982, 289.09813, 245.1083, 128.0706, 100.0756, 72.0808	$C_{19}H_{21}FN_3O_5$	11	390.1460	-0.5
ENR	6.7	360.1718	+29.9740	HX + K	390.1462	372.1356, 362.1510, 291.07754, 273.0670, 259.0514, 100.0757, 72.0808	$C_{19}H_{21}FN_3O_5$	11	390.1460	0.5
ENR	6.8	360.1718	+29.9740	HX + K	390.1461	372.1355, 362.1509, 291.0774, 273.0669, 259.0513, 100.0756, 72.0809	$C_{19}H_{21}FN_3O_5$	11	390.1460	0.3
CIP	6.1	332.1405	+29.9740	HX + K	362.1141 ^b		$C_{17}H_{17}FN_3O_5$	11	362.1147	-1.7
DIF	6.0	400.1467	+29.9740	HX + K	430.1209	412.1102, 402.1259, 386.1309, 368.1204, 329.0730, 299.0988, 197.1029, 114.0549, 70.0650	$C_{21}H_{18}F_2N_3O_5$	14	430.1209	0.0
DIF	6.9	400.1467	+29.9740	HX + K	430.1207	412.1101, 402.1258, 384.1151, 345.0678, 327.0572, 86.0599	$C_{21}H_{18}F_2N_3O_5$	14	430.1209	-0.5
DIF	7.1	400.1467	+29.9740	HX + K	430.1207	412.1101, 402.1258, 384.1151, 345.0678, 327.0572, 86.0599	$C_{21}H_{18}F_2N_3O_5$	14	430.1209	-0.5
SAR	6.7	386.1311	+29.9740	HX + K	416.1053 ^b		$C_{20}H_{16}F_2N_3O_5$	14	416.1053	0.0
ENR	4.4	360.1718	-26.0146	DE + H	334.1560	316.1464, 311.3906, 296.1393, 289.098, 263.0825, 219.0927, 164.1069, 86.0600, 72.0807	$C_{17}H_{21}FN_3O_3$	9	334.1561	-0.3
CIP	3.6	332.1405	-26.0146	DE + H	306.1246	288.1141, 268.108, 263.0824, 245.1084	$C_{15}H_{17}FN_3O_3$	9	306.1248	-0.7
DIF	5.8	400.1467	-26.0146	DE + H	374.1307	354.1246, 343.0884, 336.1138, 317.0728, 299.0988, 280.1242, 273.0830	$C_{19}H_{18}F_2N_3O_3$	12	374.1311	-1.1
SAR	5.9	386.1311	-26.0146	DE + H	360.1156	343.0883, 342.1048, 340.1089, 322.0986, 317.07310, 299.0991	$C_{18}H_{16}F_2N_3O_3$	12	360.1154	0.6





Other reactions											
ENR	5.0	360.1718	- 28.0303		DE	332.1404		C ₁₇ H ₁₉ FN ₃ O ₃	10	332.1405	-0.3
DIF	6.0	400.1467	- 14.0157		DM	386.1309		C ₂₀ H ₁₈ F ₂ N ₃ O ₃	13	386.1311	-0.5
CIP	7.0	332.1405	+ 42.0103		A	374.1509	356.1402, 314.1297, 300.0777, 272.0827, 249.0669, 243.0563, 231.0562, 215.0249	C ₁₉ H ₂₁ FN ₃ O ₄	11	374.1511	-0.5
SAR	8.0	386.1311	+ 42.0103		A	428.1416	410.1310, 382.1361, 368.1203	C ₂₂ H ₂₀ F ₂ N ₃ O ₄	14	428.1416	0.0
CIP	6.6	332.1405	+ 43.0056		F	375.1463	357.1356, 340.1096, 332.1402, 314.1297, 270.0671, 243.0563, 231.0563, 216.0692	C ₁₈ H ₂₀ FN ₄ O ₄	11	375.1463	0.0
SAR	7.2	386.1311	+ 43.0056		F	429.1369	411.1264, 386.1312, 368.1205, 348.1141	C ₂₁ H ₁₉ F ₂ N ₄ O ₄	14	429.1369	0.0
CIP	5.0	332.1405	- 58.0419		DY +DCP	274.0986 ^c	231.0562 ^c , 217.0404 ^c , 203.0611, 187.0663, 175.0664, 163.0665, 148.0556, 136.557 ^c	C ₁₄ H ₁₃ FN ₃ O ₂	10	274.0986	0.0
CIP	5.0	332.1405	+ 241.0024			573.1429 ^b					
CIP	4.0	332.1405	+ 324.1056		LC	656.2457	638.2450, 620.2246, 494.1929, 476.1825, 404.1616, 344.1400, 332.1401	C ₂₉ H ₃₉ FN ₃ O ₁₃	12	656.2461	-0.6






^a Retention time when LC-ToF is used.

^b Concentration too low to be fragmented; ^c These masses were also observed with a molecule of water (+ 18.0105) but at low concentration.





(*) HX: hydroxylation; N-O: N-Oxide; K: ketone formation; DE: deethylation; H: Hydrogenation; CC: cysteine conjugation; DC: Decarboxylation; DF: Reductive defluorination; DM: Demethylation; A: Acetylation; F: Formamidation; DY: Dehydration; DCP: Decyclopropylation; LC: Lactose conjugation.

Table 3. Metabolites and TPs found in medicated cow milk with ENR sorted by accurate mass. Accurate masses analysed by ToF and Orbitrap and their theoretical mass assigned to molecular formula are given with calculated error (ppm) for the assignment of suggested structure for the two analysers.

m/z (ToF)	m/z (Orbitrap)	m/z theoretical	t_R^a (min)	Molecular formula ^b	Error (ppm) (ToF)	Error (ppm) (Orbitrap)	Suggested Structure	DBE	
263.0828	263.0834	263.0826	6.85	$C_{13}H_{12}FN_2O_3^+$	0.8	3.0		9	ENR-1
288.1349	288.1345	288.1343	2.9	$C_{15}H_{18}O_3N_3^+$	2.1	0.7	n.a		
291.0770	291.0783	291.0776	6.7	$C_{14}H_{12}FN_2O_4^+$	-2.1	2.4		10	ENR-2
302.1501	302.1506	302.1499	5.7	$C_{16}H_{20}N_3O_3^+$	0.7	2.3	n.a		
306.1239	306.1256	306.1248	3.6	$C_{15}H_{17}FN_3O_3^+$	-2.9	2.6		9	ENR-3
307.1083	307.1097	307.1089	6.5	$C_{15}H_{16}FN_2O_4^+$	-1.9	1.9		9	ENR-4
329.1485	329.1498	329.1496	3.4	$C_{18}H_{21}N_2O_4^+$	-3.3	0.6	n.a		
329.1485	329.1498	329.1496	4.9	$C_{18}H_{21}N_2O_4^+$	-3.3	0.6	n.a		

332.1408	332.1405	332.1405	5.0	$C_{17}H_{19}FN_3O_3^+$	0.9	0		10	CIP
334.1189	334.1207	334.1198	6.4	$C_{16}H_{17}FN_3O_4^+$	-2.7	2.7		10	ENR-5
334.1539	334.1566	334.1561	4.35	$C_{17}H_{21}FN_3O_3^+$	-6.5	1.5		9	ENR-6
346.1190	346.1204	346.1198	6.5	$C_{17}H_{17}FN_3O_4^+$	-2.3	1.7		11	ENR-7
348.1343	348.1350	348.1354	3.6	$C_{17}H_{19}FN_3O_4^+$	-3.1	-1.1		10	ENR-8


348.1343	348.1364	348.1354	6.5	$C_{17}H_{19}FN_3O_4^+$	-3.1	2.9		10	ENR-9
360.1357	360.1360	360.1354	6.9	$C_{18}H_{19}FN_3O_4^+$	0.8	1.7		11	ENR-10
362.1137	362.1152	362.1147	6.1	$C_{17}H_{17}FN_3O_5^+$	-2.8	1.4		11	ENR-11
362.1137	362.1152	362.1147	6.8	$C_{17}H_{17}FN_3O_5^+$	-2.8	1.4		11	ENR-12
362.1137	362.1152	362.1147	7.1	$C_{17}H_{17}FN_3O_5^+$	-2.8	1.4		11	ENR-13
374.1510	374.1523	374.1511	7.0	$C_{19}H_{21}FN_3O_4^+$	-0.3	2.7		11	ENR-14

375.1462	375.1464	375.1463	6.6	$C_{18}H_{20}FN_4O_4^+$	-0.3	0.3		11	ENR-15
376.1656	376.1671	376.1667	4.6	$C_{19}H_{23}FN_3O_3^+$	-2.9	1.1		10	ENR-16
376.1656	376.1673	376.1667	5.8	$C_{19}H_{23}FN_3O_3^+$	-2.9	1.6		10	ENR-17
390.1459	390.1461	390.1460	6.65	$C_{19}H_{21}FN_3O_5^+$	-0.3	0.3		11	ENR-18
390.1459	390.1461	390.1460	6.85	$C_{19}H_{21}FN_3O_5^+$	-0.3	0.3		11	ENR-19


^a Retention time when LC-ToF is used.

^b These molecular formulas are hypothetical in case of metabolites without assigned structure.


n.a : Non-assigned




CIP
m/z 332.1405



SAR
m/z 386.1311



ENR
m/z 360.1718



DIF
m/z 400.1467

Fig 1





Fig 2










Fig 4

Fig 5









Fig 6

Electronic Supporting Information for the research article:

**IDENTIFICATION OF METABOLITES AND THERMAL
TRANSFORMATION PRODUCTS OF QUINOLONES IN RAW COW MILK
BY LIQUID CHROMATOGRAPHY COUPLED TO HIGH RESOLUTION
MASS SPECTROMETRY**

A. Junza¹, S. Barbosa², R. Codony³, A. Jubert³, J. Barbosa^{1,2}, D. Barrón^{1*}

¹Departament de Química Analítica, Campus de l'Alimentació de Torribera
Avda. Prat de la Riba, 171, 08921 Sta. Coloma de Gramanet, Barcelona, Spain

²Departament de Química Analítica
Martí Franquès, 1-11, 08028 Barcelona, Spain

³Laboratori Interprofessional Lleter de Catalunya (ALLIC)
Ctra. Vilassar a Cabrils s/n, 08348 Cabrils, Barcelona,

* Email: dolores.barron@ub.edu

Content:

- Explanations about MS elucidation of some TPs and metabolites of quinolones (ENR, CIP, DIF and SAR).

Section 3.2: Common reactions of the four quinolones

1. TPs of DIF (m/z 416.1416)

In the mass spectrum of the ion m/z 416.1416 (Figure 2A), two losses of water (18.0105 m/z) were observed. The first was due to carboxylic acid and the same loss was observed for the rest of the quinolones. The second loss of water observed in the spectra is only possible if an OH group is in position 2 or 3 of the piperazine ring.

2. TPs of ENR (m/z 390.1460) Peak 1

The accurate mass of the fragment m/z 128.0706 corresponds to molecular formula $C_6H_{10}NO_2$ with a DBE value of 3 and is the key to the elucidation (Figure 3C). This was followed by two consecutive fragmentations, corresponding to losses of 27.9949 (CO) [$128.0706 - CO \rightarrow 100.0756$; $100.0756 - CO \rightarrow 72.0808$] and the reduction of one unit of DBE. These fragments indicate that a ketone and an alcohol are present in the piperazinyl ring, as observed in the suggested fragmentation shown in Figure 3C. The rest of the fragments are due to losses of CO_2 , H_2O and pieces of the piperazine ring. It should be noted that the DBE of some fragments such as m/z 301.0982 and 328.1454 are 10.5. The structures represented in the mass spectrum (Figure 3C) have 11 unsaturations (double bonds and rings). The DBE calculated is the average of the resonance structures of drawn structure (DBE: 11) and the carbocation (DBE:10)⁵².

Section 3.2: Other reactions

1. TPs of CIP and SAR (acetylation and formamidation)

Two different reactions took place on CIP and SAR. The first was acetylation, which could be justified by a loss of acetyl group (42.0103) (Acetyl-CIP(-H₂O): 356.1402 → 314.1297 and Acetyl-SAR(-H₂O): 410.1310 → 368.1203), as observed in the MS spectra (not shown). The second reaction corresponds to formamidation. For CIP and SAR, the loss of formamide and subsequent gain of 2H occurred in one step (Formamide-CIP(-H₂O): 357.1356 → 314.1296 and Formamide-SAR(-H₂O): 411.1264 → 368.1205). Moreover, in the case of CIP, the loss of NH₃ (357.1356 → 340.1096), followed by the loss of a carbonyl and gain of 2H (340.1096 → 314.1296), were also observed. Due to the absence of the alkyl chain in the N⁴-piperazinyl ring, both reactions (acetylation and formamidation) took place.

2. TPs of CIP (*m/z* 656.2457)

Figure 4 shows the mass spectrum of this TP with its fragmentations. The two main characteristic losses were the loss of -180.0633 (656.2457 → 476.1825), which corresponds to the loss of a glucose, and the loss of a galactose without a water molecule (476.1825 → 344.1400). In the spectrum, steady losses of six water molecules were observed due to the OH present on the sugars, as can be seen in the smaller mass spectrum in Figure 4. All these losses could be explained if the molecule CIP (332.1405) is combined with the lactose (galactose + glucose) present in milk at high concentration. This new product was favoured when samples were heated at a high temperature for a long time (60 min).

Section 3.4: Determination of metabolites and TPs in medicated cow milk

1. Metabolite ENR-1 (m/z 263.0834)

In the MS/MS spectrum of ENR-1 (Figure 6A), the loss of a water molecule ($263.0834 \rightarrow 245.0729$) and CO ($245.0729 \rightarrow 217.0778$) from carboxylic acid can be observed. The presence of cyclopropyl can also be justified by ion m/z 222.0441, which is the result of the loss of cyclopropyl from the parent metabolite ($263.0834 - 41.0391 \rightarrow 222.0441$).

2. Metabolite ENR-4 (m/z 307.1095)

In this case, characteristic losses of the carboxylic group and cyclopropyl could be observed in the spectrum of ENR-4 (Figure 6C). The modifications of structure with respect to ENR, were again found over the piperazine ring. As can be seen, the ion m/z 248.0596 loses the methyl group, which results in the ion m/z 235.0520 that later loses a molecule of water ($235.0520 \rightarrow 217.0411$).