

1  
2  
3  
4 **1 Functional and immunogenic characterization of diverse HCV glycoprotein E2 variants**

5  
6 2  
7 3 Tanvi Khera<sup>1#</sup>, Patrick Behrendt<sup>1,2,11#</sup>, Dorothea Bankwitz<sup>1</sup>, Richard J.P. Brown<sup>1</sup>, Daniel Todt<sup>1, 12</sup>, Mandy  
8  
9 4 Doepke<sup>1</sup>, Abdul Ghafoor Khan<sup>3</sup>, Kai Schulze<sup>4</sup>, John Law<sup>5</sup>, Michael Logan<sup>5</sup>, Darren Hockman<sup>5</sup>, Jason  
10  
11 5 Alexander Ji-Xhin Wong<sup>5</sup>, Leona Dold<sup>6,7</sup>, Victor Gonzalez-Motos<sup>8</sup>, Ulrich Spengler<sup>9</sup>, Abel Viejo-  
12  
13 6 Borbolla<sup>8</sup>, Luisa Ströh<sup>8</sup>, Thomas Krey<sup>8,11</sup>, Alexander W. Tarr<sup>10</sup>, Eike Steinmann<sup>1,11,12</sup>, Michael P.  
14  
15 7 Manns<sup>2,11</sup>, Florian Klein<sup>6,7</sup>, Carlos A. Guzman<sup>4</sup>, Joseph Marcotrigiano<sup>3</sup>, Michael Houghton<sup>5</sup>, Thomas  
16  
17 8 Pietschmann<sup>1, 11\*</sup>

18  
19 9 <sup>1</sup>Institute of Experimental Virology, TWINCORE, Centre for Experimental and Clinical Infection  
20  
21 10 Research; a joint venture between the Medical School Hannover (MHH) and the Helmholtz Centre for  
22  
23 11 Infection Research (HZI), Hannover, Germany

24  
25 12 <sup>2</sup>Department of Gastroenterology, Hepatology and Endocrinology, Hannover Medical School, Hannover,  
26  
27 13 Germany

28  
29 14 <sup>3</sup>National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892-  
30  
31 15 9806, USA

32  
33 16 <sup>4</sup>Department of Vaccinology and Applied Microbiology, Helmholtz Centre for Infection Research,  
34  
35 17 Braunschweig, Germany

36  
37 18 <sup>5</sup>Li Ka Shing Institute of Virology, Department of Medical Microbiology & Immunology, University of  
38  
39 19 Alberta, Edmonton, Canada

40  
41 20 <sup>6</sup>Institute of Virology, Center for Molecular Medicine Cologne (CMMC), University of Cologne,  
42  
43 21 Cologne, Germany.

44  
45 22 <sup>7</sup>German Centre for Infection Research (DZIF), partner site Cologne, Germany,

46  
47 23 <sup>8</sup>Institute of Virology, Hannover Medical School, 30625 Hannover, Germany

48  
49 24 <sup>9</sup> Department of Internal Medicine 1, Rheinische Friedrich-Wilhelms-University Bonn, Bonn, Germany

50  
51 25 <sup>10</sup>NIHR Nottingham Digestive Diseases Biomedical Research Centre and School of Life Sciences, The  
52  
53 26 University of Nottingham, Nottingham, UK

54  
55 27 <sup>11</sup>German Centre for Infection Research (DZIF), partner site Hannover-Braunschweig, Germany

56  
57 28 <sup>12</sup>Department of Molecular and Medical Virology, Ruhr-University Bochum, Bochum, Germany

58  
59 29 # equally contributing authors

60  
61 30 \*Correspondence Address: Prof. Dr. rer. nat. Thomas Pietschmann, Institute of Experimental Virology,  
62  
63 31 Twincore, Centre for Experimental and Clinical Infection Research, Feodor-Lynen Str 7-9, 30625  
64  
65 32 Hannover, thomas.pietschmann@twincore.de, Phone: +49-511-220027130, FAX +49-511-220027139.

66  
67 33 **Keywords:** HCV, Glycoproteins, immunogen, antibodies, recombinant proteins **Word count:** 5678

68  
69 34 **Conflict of interest:** The authors declare no conflicts of interest that pertain to this work.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 35 Abstract

36 **Background and Aims** Induction of cross-reactive antibodies targeting conserved epitopes of the  
37 envelope proteins E1E2 is a key requirement for an HCV vaccine. Conserved epitopes like the viral  
38 CD81-binding site are targeted by rare broadly neutralizing antibodies. However, these viral segments are  
39 occluded by variable regions and glycans. We aimed to identify antigens exposing conserved epitopes and  
40 to characterize their immunogenicity.

41 **Methods** We created HCV variants with mutated glycosylation sites and/or hypervariable region 1  
42 (HVR1). Exposure of the CD81 binding site and conserved epitopes was quantified by soluble CD81 and  
43 antibody interaction and neutralization assays. E2 or E1-E2 heterodimers with mutations causing epitope  
44 exposure were used to immunize mice. Vaccine-induced antibodies were examined and compared with  
45 patient-derived antibodies.

46 **Results** Mutant viruses bound soluble CD81 and antibodies targeting the CD81 binding site with  
47 enhanced efficacy. Mice immunized with E2 or E1E2 heterodimers incorporating these modifications  
48 mounted strong, cross-binding, and non-interfering antibodies. E2-induced antibodies neutralized the  
49 autologous virus but they were not cross-neutralizing.

50 **Conclusions** Viruses lacking the HVR1 and selected glycosylation sites expose the CD81 binding site and  
51 cross-neutralization antibody epitopes. Recombinant E2 proteins carrying these modifications induce  
52 strong cross-binding but not cross-neutralizing antibodies.

## 53 Lay summary

54 Conserved viral epitopes can be made considerably more accessible for binding of potently neutralizing  
55 antibodies by deletion of HVR1 and selected glycosylation sites. Recombinant E2 proteins carrying these  
56 mutations are unable to elicit cross-neutralizing antibodies suggesting that exposure of conserved epitopes  
57 is not sufficient to focus antibody responses on production of cross-neutralizing antibodies.

## 58 Highlights

- 59 • High resolution mapping of the impact of HVR1 and glycosylation sites on CD81 binding,  
60 antibody binding and virus neutralization
- 61 • Viral mutants lacking HVR1 and selected glycosylation sites are functional and they expose the  
62 viral CD81 binding site and conserved cross-neutralization epitopes
- 63 • E2 proteins with these mutations induce cross-binding and non-interfering antibodies in mice

## 64 Introduction

65 Hepatitis C virus (HCV) is a global health burden affecting approximately 71 million people worldwide  
66 [1]. Infection often leads to chronic hepatitis, with the subsequent risk for liver cirrhosis and  
67 hepatocellular carcinoma. Persistent HCV infection is now curable with the introduction of direct-acting  
68 antivirals (DAAs). However, a prophylactic HCV vaccine is not available. Since viral re-infection is  
69 possible and as many HCV infected individuals are not diagnosed, a vaccine against HCV would facilitate  
70 global HCV eradication programs.

71 The extreme diversity of HCV is a major obstacle for vaccine development [2]. The HCV E1E2 proteins  
72 are essential for viral cell entry, they bind the HCV receptor CD81 and they are targets for neutralizing  
73 antibodies. Hence, immunogens based on E1E2 represent one major branch of vaccine development [3, 4]  
74 and numerous approaches to induce E1E2-targeting broadly neutralizing antibodies (bNabs) have been  
75 explored [5]. For recombinant E1E2, the most advanced HCV subunit vaccine candidate, induction of  
76 robust cross-binding and cross-neutralizing antibody responses was observed in multiple animal models  
77 and in humans [4, 6, 7, 8].

78 HCV has evolved mechanisms to evade humoral immune responses including high functional flexibility  
79 and variability of immunogenic portions of its envelope proteins [9]. The highest sequence variability  
80 occurs in the first 27 amino acids of the N-terminus of E2, which is referred to as the hypervariable region  
81 1 (HVR1), and which is dispensable for HCV infection *in vitro* [10, 11]. The HVR1 is immunogenic and  
82 most patients mount antibodies targeting this region [12]. However, these antibodies rapidly select  
83 resistant viral variants [13]. Deletion of the HVR1 renders HCV more susceptible to antibody  
84 neutralization, and it increases virus binding to soluble CD81, suggesting that this region occludes key  
85 neutralization epitopes and the viral CD81 binding site [10, 11]. HCV E1E2 heterodimers are also heavily  
86 glycosylated at multiple sites both within E1 and E2 and glycans modulate glycoprotein function and  
87 antibody neutralization [14, 15, 16]. Structural analyses of the E2 core domain show that the conserved  
88 CD81 binding site is surrounded by several glycosylation sites and that it overlaps with the epitopes of  
89 bNabs isolated from patients like for instance HC1 and HC11 [17, 18, 19, 20] (Fig. 1). Strikingly, *in vitro*  
90 HCV is unable to escape antibody pressure by HC-1 and HC-11 suggesting that immune responses  
91 targeting these epitopes may confer robust protection [21, 22].

92 We hypothesized that the HVR1 and protein glycosylation limits access to these conserved viral epitopes.  
93 In turn, immunogenicity of these epitopes may be low and antibodies targeting these viral regions may  
94 arise only very infrequently. To overcome this limitation and to focus immune responses to conserved

1  
2  
3  
4 95 viral epitopes we combined targeted protein deglycosylation with deletion of HVR1 to create viruses and  
5  
6 96 proteins with increased exposure of the crucial CD81 binding site and possibly superior immunogenicity.  
7

## 8 9 97 **Results**

### 10 11 98 **Inactivation of glycosylation sites and deletion of HVR1 increase exposure of conserved** 12 13 99 **neutralization epitopes and the CD81 binding site**

14  
15  
16 100 To generate an improved HCV vaccine antigen, we aimed to increase the exposure of the CD81-binding  
17  
18 101 site by combined mutation of the HVR1 and selected glycosylation sites. Jc1 reporter viruses expressing  
19  
20 102 the J6 (GT2a) E1E2 proteins [23, 24] were constructed harboring N to A substitutions at glycosylation  
21  
22 103 sites N417 (417) (numbering according to the H77 reference stain; J6 numbering in brackets), N423 (423),  
23  
24 104 N448 (448), N532 (534) and N645 (649) (Fig. 1A). A previous report had shown that mutations at these  
25  
26 105 residues still allowed viral entry of JFH1 and can therefore be studied and further evaluated *in vitro* [15].  
27  
28 106 Molecular modelling of the glycans associated with the HCV E2 core structure revealed that some were  
29  
30 107 proximal to the CD81 binding site and conserved neutralization epitopes (Fig. 1B). Therefore, to further  
31  
32 108 increase exposure of the CD81 binding site, we combined these mutations with the deletion of the HVR1.  
33  
34 109 Fitness of mutant viruses was examined by transfection of Huh7-Lunet N#3 cells that do not express  
35  
36 110 CD81 [25], and therefore do not permit reinfection of the cells. Replication of all viral mutants was  
37  
38 111 comparable to wild type (WT) Jc1 (Fig. 1C). Huh-7.5 cells were inoculated with virus stocks after  
39  
40 112 normalization to viral core protein that acts as a marker for virus particle release. In line with our previous  
41  
42 113 report [10], deletion of HVR1 resulted in a 5-fold reduction of specific infectivity of Jc1 (Fig. 1D).  
43  
44 114 Likewise, point mutations of individual glycosylation sites were well tolerated as specific infectivity was  
45  
46 115 unaffected or at most reduced by ca. 5-fold.

47  
48 116 To examine the impact of these mutations on the exposure of conserved neutralization epitopes within the  
49  
50 117 CD81 binding site, we used the soluble, large-extracellular loop of CD81 (CD81-LEL) and two potent  
51  
52 118 bNabs (HC-1 and HC-11) that target the CD81 binding site [17, 18, 19, 20, 21], for precipitation of the  
53  
54 119 indicated viral mutants (Fig. 2A and B). Deglycosylation of specific residues in the parental Jc1 virus  
55  
56 120 slightly increased binding of HC-1 in particular for mutant N448A and N534A as is evidenced by  
57  
58 121 enhanced core protein precipitation compared to Jc1, without being statistical significant (Fig. 2A).  
59  
60 122 However, the deletion of HVR1 significantly enhanced precipitation by this antibody, which did not  
61  
62 123 further increase by additional point mutations of glycosylation sites at indicated positions (Fig. 2A).  
63  
64 124 Similar results were observed when the HC-11 antibody was used: mutation of individual glycosylation  
65  
66 125 sites increased antibody binding (e.g. N423A, N448A). Deletion of HVR1 had the greatest effect, which  
67  
68 126 was not further boosted by mutation of the glycosylation sites (Fig. 2B). Comparable results were obtained

1  
2  
3  
4 127 when we used CD81-LEL for precipitation. In the context of parental Jc1, inactivation of individual  
5  
6 128 glycosylation sites modestly increased precipitation (Fig. 2C). As may be expected from the available  
7  
8 129 structural information (Fig. 1B) [17, 18], the effect was greatest for mutants N423A and N534A, as these  
9  
10 130 glycosylation sites are directly adjacent to the CD81 binding site. Deletion of HVR1 had the greatest  
11  
12 131 effect and precipitation of viruses lacking HVR1 was not further boosted by deglycosylation of specific  
13  
14 132 residues.

### 15 133 **Deletion of HVR1 and glycosylation sites enhances virus neutralization**

16  
17  
18 134 To characterize the accessibility of HCV antibody epitopes, we performed neutralization assays with a  
19  
20 135 panel of bNAbs (Fig. 3A and C and supplementary table 1). Moreover, we quantified virus neutralization  
21  
22 136 by CD81-LEL (Fig. 3B and C and supplementary table 1). Dose dependence of neutralization for selected  
23  
24 137 viruses is displayed in figure 3C. Radar plots indicating the inhibitory concentration 90% (IC<sub>90</sub>) of  
25  
26 138 different antibodies or the CD81-LEL against parental HCV and all mutant viruses is displayed in figure  
27  
28 139 3A and 3B. Each plot shows a specific virus and its sensitivity towards given antibodies (Fig. 3A),  
29  
30 140 whereas figure panel 3B highlights the susceptibility of each virus to competition by CD81-LEL. In the  
31  
32 141 context of parental Jc1, deletion of glycans at position N423 and N448 increased the susceptibility to all  
33  
34 142 tested antibodies, whereas removal of glycosylation at residues N417, N534, and N649 had little effect  
35  
36 143 (Fig. 3A upper panels). As expected, deletion of HVR1 enhanced neutralization by all antibodies tested.  
37  
38 144 Combination of deletion of HVR1 with deletion of specific N-glycosylation sites modified neutralization  
39  
40 145 by these antibodies, and the Jc1-ΔHVR1-N534A mutant exhibited a further enhanced susceptibility to  
41  
42 146 neutralization by HC-1 and HC-11 (Fig. 3A and 3C). Removal of glycosylation at the N423, N534, and  
43  
44 147 N417 site in the context of WT Jc1 enhanced inhibition of infection by CD81-LEL, whereas mutation of  
45  
46 148 the N417 and N649 sites had no effect (Fig. 3B). Deletion of HVR1 increased neutralization by CD81-  
47  
48 149 LEL and this was further enhanced ca. 5-10-fold by addition of the N534 mutation. Therefore, combined  
49  
50 150 deletion of HVR1 and inactivation of glycosylation site N534 enhanced antibody and CD81 binding to  
51  
52 151 virus particles and it had the most drastic effect on neutralization by bNabs and soluble CD81.

53  
54  
55 152 Next, we evaluated neutralization of Jc1, Jc1-N534A, Jc1-ΔHVR1 and Jc1-ΔHVR1-N534A by sera from  
56  
57 153 individuals chronically infected with HCV GT 1 and 2 (Fig. 4A-C). Similar to the results of neutralization  
58  
59 154 with monoclonal antibodies and CD81-LEL (Fig. 3), the sera poorly neutralized Jc1. Neutralization was  
60  
61 155 enhanced by elimination of the N534 site in Jc1, further boosted by deletion of HVR1, and it was maximal  
62  
63 156 when the deletion of HVR1 was combined with the inactivation of the N534 glycosylation site with minor  
64  
65 157 interpatient variation, independent of the patients` HCV genotype (Fig. 4A-C). Thus, combined deletion of  
66  
67 158 HVR1 and inactivation of glycosylation site N534 enhanced antibody and CD81 binding to virus particles  
68  
69 159 and it had the most drastic effect on neutralization by bNabs and soluble CD81 and patient sera. As these

1  
2  
3  
4 160 viruses are infectious they present well-folded and functional viral envelope proteins that display  
5  
6 161 conserved epitopes in a more accessible manner. Thus, envelope proteins carrying these modifications,  
7  
8 162 particularly the deletion of HVR1 combined with the N534A mutation, may constitute good candidates for  
9  
10 163 focusing antibody responses on conserved viral epitopes.

### 11 12 164 **Immunogenicity of recombinant E2 proteins with and without HVR1 and glycosylation**

13  
14 165 To test this, we prepared soluble, truncated HCV J6 E2 protein variants in 293T cells and immunized  
15  
16 166 BALB/c mice. These proteins have been previously reported to inhibit virus infection indicating that they  
17  
18 167 are properly folded [26]. Supplementary figure 1 shows the immunization scheme. Briefly, animals  
19  
20 168 received 30 µg of soluble E2 injected together with bis-(3',5')-cyclic dimeric adenosine monophosphate  
21  
22 169 (c-di-AMP) as adjuvant and were boosted thrice with the same formulation. As control, animals were  
23  
24 170 vaccinated with PBS and adjuvant only. We also used two vaccination series where we changed antigens  
25  
26 171 in each boost in order to promote the stimulation of antibodies specific for the conserved epitopes. In  
27  
28 172 sequence A animals were primed using WT J6 E2 (GT2a), followed by three boosters with E2ΔHVR1 (J6  
29  
30 173 GT2a derivative), E2ΔHVR1/N534A (J6 GT2a derivative), and WT Con1 E2 (GT1b), respectively. In  
31  
32 174 sequence B mice were primed with E2ΔHVR1/N534A, followed by three booster immunizations with  
33  
34 175 E2ΔHVR1 (J6 GT2a derivative), WT Con1 (GT1b) E2, and WT J6 (GT2a) E2, respectively. Finally, to  
35  
36 176 examine the relevance of global glycosylation for antigenicity, we vaccinated with WT J6 (GT2a) or with  
37  
38 177 J6 (GT2a) E2ΔHVR1 that had been deglycosylated by PNGaseF treatment prior to vaccination.

39 178 The binding of vaccine-induced antibodies to 293T cell derived recombinant E2 proteins or E1-E2  
40  
41 179 heterodimers from HCV infected cells was determined by ELISA (Fig. 5 and supplementary figure S2).  
42  
43 180 Fig 5A shows end point titers of sera from vaccinated animals reacting with H77 (GT1a), Con1 (GT1b),  
44  
45 181 J6 (GT2a) or J6ΔHVR1 (GT2a) recombinant proteins. High binding titers across these GT1 and GT2  
46  
47 182 antigens were observed in all vaccinated groups up to a maximal end point dilution approaching 10E6  
48  
49 183 (Fig. 5A). Moreover, irrespectively of which protein was used for vaccination, most of these antibodies  
50  
51 184 targeted binding sites outside of the HVR1, as end point titers were comparable between J6 (GT2a) and  
52  
53 185 J6ΔHVR1 (GT2a) ELISA antigens. Cross-binding activity to these recombinant E2 proteins was clearly  
54  
55 186 enhanced for vaccination with E2ΔHVR compared to wildtype recombinant E2 (Fig.5C left panel).  
56  
57 187 Moreover, vaccination according to sequence A and B resulted in greater cross-binding to recombinant E2  
58  
59 188 proteins. To evaluate the binding of these antibodies to heterodimeric E1-E2 protein complexes from  
60  
61 189 different HCV strains, we used plates coated with galanthus nivalis lectin (GNA) to capture HCV E1E2  
62  
63 190 complexes expressed in Huh-7.5 cells transfected with infectious HCVcc chimeras. Subsequently, these  
64  
65 191 plates were incubated with sera from our vaccinated animals and end point binding titers were calculated

1  
2  
3  
4 192 (Fig. 5B). In each group of vaccinated animals, anti-E2 antibodies binding to all examined E1-E2 protein  
5  
6 193 complexes were detectable. Thus, each vaccination approach induced broadly cross-binding antibodies  
7  
8 194 that recognize E2 proteins of all major HCV GTs. With exception of the vaccination with J6 $\Delta$ HVR-534  
9  
10 195 E2 protein, however, all vaccination approaches induced antibodies with cross-binding activity to these  
11 196 E1-E2 heterodimers indistinguishable from the ones induced by parental J6 E2 (Fig. 5B and 5C right  
12  
13 197 panel). Taken together, all vaccination approaches induced strong cross-binding antibodies to E2 and E1-  
14  
15 198 E2 heterodimers. Vaccination with different E2 proteins, most notably with E2 $\Delta$ HVR1, enhanced cross-  
16 199 binding when judged by ELISAs involving recombinant E2. However, this difference was not detected  
17  
18 200 when cross-binding was quantified with ELISAs based on E1-E2 heterodimers from cells replicating  
19  
20 201 infectious virus.

### 22 202 **Evaluation of neutralizing activity of vaccine induced antibodies**

23  
24  
25 203 To examine autologous virus and virus cross-neutralizing responses we incubated Jc1 and GT5a reporter  
26  
27 204 viruses, respectively, with increasing doses of purified IgG from vaccinated animals or from patient sera  
28  
29 205 (Fig. 6A). The autologous virus Jc1 was neutralized by antibodies from the animals vaccinated with  
30  
31 206 deglycosylated E2, with deglycosylated  $\Delta$ HVR1 or the series A antigen combinations. The strength of  
32  
33 207 these autologous neutralizing antibody responses was at least as strong as the neutralization by polyclonal  
34  
35 208 human antibodies collected from a panel of GT1, 2 and 3 chronically infected individuals. This correlated  
36  
37 209 with comparable binding of patient-derived and vaccine-induced antibodies to recombinant E2 and virus  
38  
39 210 particle associated E2, quantified by surface plasmon resonance and virus particle precipitation  
40  
41 211 (supplementary figure S4). However, none of the vaccine-induced antibodies were able to cross-neutralize  
42  
43 212 the GT5a reporter virus which was at least partially neutralized by the majority of patient derived  
44  
45 213 antibodies.

46  
47 214 It has been reported that HCV induces antibodies that bind to E2 and that interfere with the activity of  
48  
49 215 neutralizing antibodies, thereby masking vaccine-induced antibody neutralization [27, 28]. To explore  
50  
51 216 whether our vaccine candidates induced such interfering antibodies, we used pooled polyclonal mouse  
52  
53 217 IgGs from each vaccinated group and mixed them with polyclonal IgGs from a patient that displays strong  
54  
55 218 neutralizing activity. The neutralization efficacy was not decreased by addition of IgG vaccinated mice,  
56  
57 219 suggesting that the vaccines did not induce antibodies interfering with neutralization (Fig. 6B).

58  
59 220 The role of glycans and the HVR1 for immunogenicity was also assessed using protein from a different  
60  
61 221 genotype (GT1a H77), as well as in the context of E1E2 heterodimers purified from CHO cell extracts. A  
62  
63 222 detailed description of the purification procedure is provided in the materials and methods section and the  
64  
65 223 quality control of the purification is given in supplementary figure S5A. The binding of E1E2 (WT), E1E2

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

224 (N417Q) and E1E2 (N532Q) antigens to CD81-LEL was examined by ELISA and revealed enhanced  
225 binding for E1E2 (N532Q) compared to parental E1E2 and the N417Q mutant (Fig. 7A and  
226 supplementary figure S5B). These GT1a antigens were used to immunize mice resulting in the induction  
227 of E2 specific antibody responses comparable between all groups (Fig. 7B). As shown in figure 7C all  
228 three vaccine candidates induced antibodies competing with binding of the known monoclonal antibodies  
229 AR3B, 1:7, HC33.4 and AP33 as assessed by a competition assay. Finally, we used GT1a H77 HCVpp  
230 assays to examine neutralization efficacy of vaccine induced antibodies. Each vaccine candidate induced  
231 HCVpp neutralizing antibodies whereby no significant differences between the vaccine candidates were  
232 observed (Fig. 7D).



## Discussion

Here we provide a high resolution map of viral determinants that govern exposure of the CD81 binding site and of conserved neutralization epitopes. We show that these changes are well tolerated indicating that they do not abrogate the functioning of E1/E2. The most global change in antibody binding/neutralization and CD81 binding/neutralization was accomplished by deletion of HVR1 combined with inactivation of glycosylation site 534. Thus, we chose this combined modification for vaccination approaches involving recombinant proteins. To rule out strain-specific differences between immunogens, we created these variants in the background of J6 (GT2a) E2 and in the context of the H77 (GT1a) strain. We prepared H77 E1E2 heterodimers from CHO cell extracts and recombinant E2 protein secreted from 293T cells to examine the importance of producer cells and the relevance of E1. To globally assess the relevance of glycosylation in our vaccine candidates we also examined the immunogenicity of 293T cell-derived proteins that had been enzymatically deglycosylated prior to vaccination. Finally, we included a vaccination protocol involving variable immunogens and in each case at least one immunogen from an alternative viral GT.

All vaccination approaches induced robust cross-binding antibody responses as determined by ELISA assays involving recombinant proteins or cell extracts expressing E1-E2 heterodimers from all major HCV GTs (Fig. 5). Vaccination with E1-E2 heterodimers induced antibodies that competed with the binding of previously described bNAbs AR3B, 1:7, HC33.4 and AP33 and we detected virus neutralization in HCVpp assays (Fig. 7). However, inactivation of N417 or N532 glycosylation sites in the context of H77 E1E2 heterodimers did not grossly affect these responses although the mutation N532Q mediated increased binding of the recombinant E1E2 complex to CD81. Among vaccinations with 293T-derived E2 proteins immunization with J6-E2 $\Delta$ HVR1 and the two immunization series A and B mounted superior cross-binding antibodies compared with J6-E2 and the other examined approaches. However, this was only detected when recombinant E2 proteins from 293T cells were used as ELISA antigens and not when E1-E2 heterodimers partially purified from HCV replicating cells were employed. Thus, some of our E2-protein based vaccination approaches successfully improved production of cross-binding antibodies to the recombinant immunogens but not to the E1-E2 heterodimer expressed in infected cells. Thus, apparently there are structural differences between these proteins and E1-E2 heterodimers. These differences could impact on vaccine efficacy and they should be considered when standardizing tests between laboratories and when estimating cross-binding between different studies. It is also possible that such structural differences were responsible that vaccination with 293T-derived E2 proteins triggered only modest autologous neutralizing antibodies and no cross-neutralization. It has been shown that purified, UV-inactivated HCVcc particles induce cross-neutralizing antibodies in mice and non-human primates [29,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

30]. Thus, it will be interesting to explore if vaccination with the HCVcc variants characterized here induces a focused and enhanced neutralizing antibody response.

It is surprising that vaccination with E2 from 293T cells induced very robust, cross-binding antibodies that, however, did not cross-neutralize. One possible explanation was that these immunogens induced antibodies that interfere with cross-neutralization as has been described previously [27, 28]. However, the vaccine induced antibodies did not interfere with cross-neutralization of HCV patient derived polyclonal antibodies (Fig. 6). Thus, we believe that the 293T-derived E2 vaccines examined here efficiently induced antibodies that target conserved binding epitopes, but that targeting these epitopes is not cross-neutralizing. It is possible that this reflects in part a species-specific limitation of immunogenicity of recombinant E2 proteins in mice as recent results by other groups and us suggest that anti-HCV neutralizing antibody responses are more readily induced in guinea pigs [31, 32]. Although neutralizing antibodies were elicited, those were primarily strain-specific, thus explaining the neutralization of autologous Jc1 particles. Notably, the conserved CD81 binding site seems to be structurally highly flexible, and it was proposed that this flexibility may favor induction of non-neutralizing antibodies [33]. In line with this, it is possible that the recombinant E2 proteins used here preferentially adopt a conformation that induces binding, but non-neutralizing antibodies and that the conformation inducing neutralizing antibodies may be under represented in these recombinant proteins. Thus, epitope exposure alone may be insufficient and should be complemented by vaccine design aiming at rigidifying this region.

The lack of robust immunocompetent animal models permissive for HCV makes it hard to assess if the quantity and quality of vaccine-induced immunity is sufficient to confer protection. Thus, it is difficult to predict to which extent antibody responses induced by current vaccination approaches, including the ones presented here, contribute to protection. *In vivo*, antibody functions independent of direct virus particle neutralization could facilitate protection. For instance antibody dependent cytotoxicity (ADCC) may contribute to clearance of HCV infected cells. In this regard, binding, but non-neutralizing antibodies may contribute to HCV protection, as has been recently reported for an HIV infection *in vivo* [34]. Therefore, development of vaccination approaches inducing strong T cell responses as well as nNAb and bNAb and involving all antibody effector functions is likely of key importance for induction of robust protection.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

294 **Acknowledgements**

295 We are grateful to Takaji Wakita and Jens Bukh for JFH1 and J6CF isolates and HCV chimeras,  
296 respectively, and to Charles Rice for Huh-7.5 cells and the 9E10 antibody. We thank Steven Fong for  
297 generous provision of CBH-7, HC-1, HC-11, HC84-20, HC84-21, and HC84-22 monoclonal antibodies,  
298 and Arvind Patel for providing AP33, and Mansun Law for gift of AR3B. We are particularly thankful to  
299 Thomas Schulz and the members of his diagnostics team for help in the quantification of HCV core  
300 protein levels. We would also like to thank all members of the Institute for Experimental Virology for  
301 helpful suggestions and discussions.

302 Twincore is a joint venture of Hannover Medical School and the Helmholtz-Centre for infection research.  
303 T.P. was supported by a grant from the DZIF (HCV-vaccine), and T.P. and C.A.G. were supported by the  
304 Helmholtz-Alberta Initiative for Infectious Disease Research (HAI-IDR).

## 306 **Materials and Methods**

### 307 **Plasmid constructs**

308 The renilla-luciferase harboring Jc1 [23] and Jc1-ΔHVR1 [10] viral cDNA clones with or without alanine  
309 substitutions within N-linked glycosylation-sites of E2 (N417A, N423A, N448A, N534A and N649A)  
310 were constructed by standard PCR-based techniques and verified by sequencing.

### 311 **HCVcc neutralization assay**

312 For inhibition of HCV infection, 200 µl of a Huh7.5 cell suspension ( $5 \times 10^4$  cells per ml) was seeded into  
313 each well of a 96-well plate 24 h prior to inoculation. Luciferase reporter viruses were mixed with serial  
314 dilutions of indicated serum/antibody concentrations and pre incubated for 1 h. This mixture was used to  
315 inoculate cells for 4 h in triplicates per dilution. Thereafter, 170 µl of DMEM was added onto the cells.  
316 Viral infection was determined 48 or 72 h after infection by removing the supernatant and lysing of the  
317 cells by addition of passive lysis buffer or water and measurement of RLU using a 96-well plate reader  
318 (Berthold).

### 319 **HCV pseudotyped viruses (HCVpp) neutralization assay**

320 HCV pseudotyped viruses (HCVpp) expressing a luciferase reporter were generated as described [35]. For  
321 neutralization assays, Huh7.5 cells were plated on poly-lysine coated 96-well plates 1 day prior to  
322 infection. HCVpp were diluted 1:10 and premixed with heat inactivated diluted sera (1:100) for 1 h at  
323 37°C followed by addition to Huh 7.5 cells. Six hours post-infection, the antibody-virus inoculum was  
324 replaced with fresh culture medium. Cells were processed 48 h post-infection using the Bright-glo  
325 luciferase assay system (Promega, Madison, WI, USA). Luminescence was measured using an Enspire  
326 plate reader (Perkin-Elmer, Waltham, MA, USA). The neutralization activity was calculated using the  
327 following formula: % neutralization = (pre-post)/pre  $\times$  100 where pre/post represent the luciferase activity  
328 done after incubating with either the pre- or post-vaccination sera.

### 329 **Molecular modeling**

330 Modelling of the glycosylated core structure (PDB 4MWF) was performed using Glycoprotein Builder  
331 ([www.glycam.org](http://www.glycam.org)). Glycans are depicted as wireframes; protein as a surface. The predicted CD81 binding  
332 site and mAb HC1/HC11 epitopes (highlighted in green) were annotated using UCSF Chimera  
333 ([www.cgl.ucsf.edu/chimera](http://www.cgl.ucsf.edu/chimera)) using a space fill model. Other amino acid side chains were coloured by  
334 hydrophobicity (red = hydrophobic; blue = hydrophilic as determined by the Kyte-Doolittle scale).

1  
2  
3  
4 335 **References**  
5  
6

- 7 336 1 WHO. 2017. Hepatitis C. WHO fact sheet 164. WHO G, Switzerland:.
- 8 337 2 Tarr AW, Khera T, Hueging K, Sheldon J, Steinmann E, Pietschmann T, *et al.* Genetic diversity  
9 338 underlying the envelope glycoproteins of hepatitis C virus: Structural and functional consequences and  
10 339 the implications for vaccine design. *Viruses* 2015;**7**:3995-4046.
- 11 340 3 Walker CM. Designing an HCV vaccine: a unique convergence of prevention and therapy? *Curr*  
12 341 *Opin Virol* 2017;**23**:113-9.
- 13 342 4 Houghton M. Prospects for prophylactic and therapeutic vaccines against the hepatitis C viruses.  
14 343 *Immunol Rev* 2011;**239**:99-108.
- 15 344 5 Fauvelle C, Colpitts CC, Keck ZY, Pierce BG, Fong SK, Baumert TF. Hepatitis C virus vaccine  
16 345 candidates inducing protective neutralizing antibodies. *Expert Rev Vaccines* 2016;**15**:1535-44.
- 17 346 6 Law JL, Chen C, Wong J, Hockman D, Santer DM, Frey SE, *et al.* A hepatitis C virus (HCV) vaccine  
18 347 comprising envelope glycoproteins gpE1/gpE2 derived from a single isolate elicits broad cross-genotype  
19 348 neutralizing antibodies in humans. *PLoS One* 2013;**8**:e59776.
- 20 349 7 Frey SE, Houghton M, Coates S, Abrignani S, Chien D, Rosa D, *et al.* Safety and immunogenicity of  
21 350 HCV E1E2 vaccine adjuvanted with MF59 administered to healthy adults. *Vaccine* 2010;**28**:6367-73.
- 22 351 8 Landi A, Law J, Hockman D, Logan M, Crawford K, Chen C, *et al.* Superior immunogenicity of HCV  
23 352 envelope glycoproteins when adjuvanted with cyclic-di-AMP, a STING activator or archaeosomes.  
24 353 *Vaccine* 2017;**35**:6949-56.
- 25 354 9 Cashman SB, Marsden BD, Dustin LB. The Humoral Immune Response to HCV: Understanding is  
26 355 Key to Vaccine Development. *Front Immunol* 2014;**5**:550.
- 27 356 10 Bankwitz D, Steinmann E, Bitzegeio J, Ciesek S, Friesland M, Herrmann E, *et al.* Hepatitis C virus  
28 357 hypervariable region 1 modulates receptor interactions, conceals the CD81 binding site, and protects  
29 358 conserved neutralizing epitopes. *J Virol* 2010;**84**:5751-63.
- 30 359 11 Prentoe J, Jensen TB, Meuleman P, Serre SB, Scheel TK, Leroux-Roels G, *et al.* Hypervariable  
31 360 region 1 differentially impacts viability of hepatitis C virus strains of genotypes 1 to 6 and impairs virus  
32 361 neutralization. *J Virol* 2011;**85**:2224-34.
- 33 362 12 Zibert A, Schreier E, Roggendorf M. Antibodies in human sera specific to hypervariable region 1  
34 363 of hepatitis C virus can block viral attachment. *Virology* 1995;**208**:653-61.
- 35 364 13 Weiner AJ, Geysen HM, Christopherson C, Hall JE, Mason TJ, Saracco G, *et al.* Evidence for  
36 365 immune selection of hepatitis C virus (HCV) putative envelope glycoprotein variants: potential role in  
37 366 chronic HCV infections. *Proc Natl Acad Sci U S A* 1992;**89**:3468-72.
- 38 367 14 Helle F, Goffard A, Morel V, Duverlie G, McKeating J, Keck ZY, *et al.* The neutralizing activity of  
39 368 anti-hepatitis C virus antibodies is modulated by specific glycans on the E2 envelope protein. *J Virol*  
40 369 2007;**81**:8101-11.
- 41 370 15 Helle F, Vieyres G, Elkrief L, Popescu CI, Wychowski C, Descamps V, *et al.* Role of N-linked glycans  
42 371 in the functions of hepatitis C virus envelope proteins incorporated into infectious virions. *J Virol*  
43 372 2010;**84**:11905-15.
- 44 373 16 Goffard A, Callens N, Bartosch B, Wychowski C, Cosset FL, Montpellier C, *et al.* Role of N-linked  
45 374 glycans in the functions of hepatitis C virus envelope glycoproteins. *J Virol* 2005;**79**:8400-9.
- 46 375 17 Kong L, Giang E, Nieusma T, Kadam RU, Cogburn KE, Hua Y, *et al.* Hepatitis C virus E2 envelope  
47 376 glycoprotein core structure. *Science* 2013;**342**:1090-4.
- 48 377 18 Khan AG, Whidby J, Miller MT, Scarborough H, Zatorski AV, Cygan A, *et al.* Structure of the core  
49 378 ectodomain of the hepatitis C virus envelope glycoprotein 2. *Nature* 2014;**509**:381-4.
- 50 379 19 Keck ZY, Li TK, Xia J, Gal-Tanamy M, Olson O, Li SH, *et al.* Definition of a conserved  
51 380 immunodominant domain on hepatitis C virus E2 glycoprotein by neutralizing human monoclonal  
52 381 antibodies. *J Virol* 2008;**82**:6061-6.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

20 Pierce BG, Keck ZY, Lau P, Fauvelle C, Gowthaman R, Baumert TF, *et al.* Global mapping of antibody recognition of the hepatitis C virus E2 glycoprotein: Implications for vaccine design. *Proc Natl Acad Sci U S A* 2016.

21 Keck ZY, Xia J, Wang Y, Wang W, Krey T, Prentoe J, *et al.* Human monoclonal antibodies to a novel cluster of conformational epitopes on HCV E2 with resistance to neutralization escape in a genotype 2a isolate. *PLoS Pathog* 2012;**8**:e1002653.

22 Keck ZY, Saha A, Xia J, Wang Y, Lau P, Krey T, *et al.* Mapping a region of hepatitis C virus E2 that is responsible for escape from neutralizing antibodies and a core CD81-binding region that does not tolerate neutralization escape mutations. *J Virol* 2011;**85**:10451-63.

23 Pietschmann T, Kaul A, Koutsoudakis G, Shavinskaya A, Kallis S, Steinmann E, *et al.* Construction and characterization of infectious intragenotypic and intergenotypic hepatitis C virus chimeras. *Proc Natl Acad Sci U S A* 2006;**103**:7408-13.

24 Reiss S, Rebhan I, Backes P, Romero-Brey I, Erfle H, Matula P, *et al.* Recruitment and activation of a lipid kinase by hepatitis C virus NS5A is essential for integrity of the membranous replication compartment. *Cell Host Microbe* 2011;**9**:32-45.

25 Bitzegeio J, Bankwitz D, Hueging K, Haid S, Brohm C, Zeisel MB, *et al.* Adaptation of hepatitis C virus to mouse CD81 permits infection of mouse cells in the absence of human entry factors. *PLoS Pathog* 2010;**6**:e1000978.

26 Whidby J, Mateu G, Scarborough H, Demeler B, Grakoui A, Marcotrigiano J. Blocking hepatitis C virus infection with recombinant form of envelope protein 2 ectodomain. *J Virol* 2009;**83**:11078-89.

27 Kachko A, Frey SE, Sirota L, Ray R, Wells F, Zubkova I, *et al.* Antibodies to an interfering epitope in hepatitis C virus E2 can mask vaccine-induced neutralizing activity. *Hepatology* 2015;**62**:1670-82.

28 Zhang P, Zhong L, Struble EB, Watanabe H, Kachko A, Mihalik K, *et al.* Depletion of interfering antibodies in chronic hepatitis C patients and vaccinated chimpanzees reveals broad cross-genotype neutralizing activity. *Proc Natl Acad Sci U S A* 2009;**106**:7537-41.

29 Akazawa D, Moriyama M, Yokokawa H, Omi N, Watanabe N, Date T, *et al.* Neutralizing antibodies induced by cell culture-derived hepatitis C virus protect against infection in mice. *Gastroenterology* 2013;**145**:447-55 e1-4.

30 Yokokawa H, Higashino A, Suzuki S, Moriyama M, Nakamura N, Suzuki T, *et al.* Induction of humoral and cellular immunity by immunisation with HCV particle vaccine in a non-human primate model. *Gut* 2016.

31 Stamatakis Z, Coates S, Evans MJ, Winer M, Crawford K, Dong C, *et al.* Hepatitis C virus envelope glycoprotein immunization of rodents elicits cross-reactive neutralizing antibodies. *Vaccine* 2007;**25**:7773-84.

32 Vietheer PT, Boo I, Gu J, McCaffrey K, Edwards S, Owczarek C, *et al.* The core domain of hepatitis C virus glycoprotein E2 generates potent cross-neutralizing antibodies in guinea pigs. *Hepatology* 2017;**65**:1117-31.

33 Kong L, Lee DE, Kadam RU, Liu T, Giang E, Nieuwma T, *et al.* Structural flexibility at a major conserved antibody target on hepatitis C virus E2 antigen. *Proc Natl Acad Sci U S A* 2016.

34 Horwitz JA, Bar-On Y, Lu CL, Fera D, Lockhart AAK, Lorenzi JCC, *et al.* Non-neutralizing Antibodies Alter the Course of HIV-1 Infection In Vivo. *Cell* 2017;**170**:637-48 e10.

35 Hsu M, Zhang J, Flint M, Logvinoff C, Cheng-Mayer C, Rice CM, *et al.* Hepatitis C virus glycoproteins mediate pH-dependent cell entry of pseudotyped retroviral particles. *Proc Natl Acad Sci U S A* 2003;**100**:7271-6.

36 Owsianka AM, Timms JM, Tarr AW, Brown RJ, Hickling TP, Szwejk A, *et al.* Identification of conserved residues in the E2 envelope glycoprotein of the hepatitis C virus that are critical for CD81 binding. *J Virol* 2006;**80**:8695-704.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

37 Drummer HE, Boo I, Maerz AL, Poumbourios P. A conserved Gly436-Trp-Leu-Ala-Gly-Leu-Phe-Tyr  
430 motif in hepatitis C virus glycoprotein E2 is a determinant of CD81 binding and viral entry. J Virol  
431 2006;**80**:7844-53.

432

## Figure Legends

**Figure 1: Schematic and structure of HCV E2 and characterization of mutant viruses.** (A) N-linked glycosylation sites (NGS) in the HCV E2 ectodomain are indicated by inverted open triangles. For positional referencing, a graphic chart of the E2 protein is located directly below, with two hypervariable regions and the intergenic variable region (HVR1, HVR2 and igVR) highlighted, and the stem region of E2 marked in light grey. All numbering is relative to the full-length ORF position in the H77 (GT1a) reference strain (accession number NC\_004102) while numbers in brackets indicate the homologous position in the J6 (GT2a) E2 ectodomain. The HVR1 region and glycan sites, which were deleted, are indicated in red. Zoomed in region highlights the viral CD81 binding site with key contact residues displayed in blue [36, 37]. Amino acid important for binding of bNAbs HC1, HC11, and HC84-like antibodies are highlighted with colored inverted triangles (pink, HC1; green HC11, orange HC84-series). \*\* indicates indels responsible for length differences between H77 (GT1a) and J6 E2 (GT2a). Note that HC84.20 also binds to Y613 and W616 and HC84.22 also binds to W616. (B) Structural modelling of the E2 core domain with the CD81, HC1 and HC11 binding sites highlighted in green. Glycosylation was modelled using Glycoprotein Builder and are shown and numbered according to the H77 (GT1a) reference strain. The E2 protein is colored by hydrophobicity of aa side chains (red and blue represent hydrophobic and hydrophilic residues, respectively) and green shows antibody or CD81 contact residues. (C) Replication of mutant viruses with alanine substitutions at indicated positions with or without deletion of the HVR1 within the E2 protein. Viral replication was determined by renilla-luciferase activity of cell lysates 48 h after transfection (n=3, duplicates each; SD). (D) Specific infectivity of particles. Released virus was normalized to equal amounts of core and was used to infect Huh-7.5 cells. Infection is displayed as RLU per well (n=3, measured in triplicates each; SD). The dotted line depicts the background of the assay as determined by measurements of uninfected cell lysates.

**Figure 2: Precipitation of mutant viruses with monoclonal antibodies and with GST-CD81-LEL.** Mutant viruses were normalized to equal amounts of core and incubated with either HC-1 (A), HC-11 (B) or with GST-C81-LEL (C), respectively. Virus-antibody or virus-CD81-LEL-GST complexes were precipitated by addition of protein G or glutathione coated beads. Viral core protein was determined to measure bound viral particles. Data are expressed as fold over control (antibody RO4 or GST) and compared to WT virus (n=4, ANOVA, Friedman, Dunn's multiple comparison, p<0.05 (\*), p<0.01 (\*\*), p<0.001 (\*\*\*)). Symbols represent individual IPs.



1  
2  
3  
4 464 **Figure 3: Neutralization of Jc1 and mutant viruses by antibodies or by receptor competition with**  
5 **GST-CD81LEL.** Viruses were incubated with serial dilutions of given antibodies (A) or GST-CD81-LEL  
6 465 (B) and infection efficiency was quantified by luciferase assays. Selected dose-response curves are plotted  
7 466 in panel (C). Radar plots in (A) highlight IC<sub>90</sub> values (µg/ml) for antibody neutralization, and IC<sub>90</sub> values  
8 467 of GST-CD81-LEL receptor competition in panel (B), respectively. The IC<sub>90</sub> values were calculated using  
9 468 non-linear regression method on graph pad version 6. Abbreviations: RLU, Relative light units.  
10  
11  
12  
13  
14

15 470 **Figure 4: Neutralization of viruses by GT1 and GT2 patient sera.** Indicated viruses were mixed with  
16 471 serial dilutions of sera of uninfected (A, #1-3), GT1 (B, #1-3) or GT2 infected (C, #1-3) individuals.  
17 472 Infection efficiency was quantified by inoculation of Huh-7.5 cells and subsequent luciferase assays. It is  
18 473 plotted relative to control infections in the absence of human serum. Means ± SD (n=2(A/B/C #1-2) or  
19 474 3(A/B/C #3)).  
20  
21  
22  
23

24 475 **Figure 5: Cross-binding of vaccine-induced antibodies to HCV GT1 to 6 glycoproteins.** (A)  
25 476 Recombinant E2 proteins (2 µg/ml) from different HCV GTs were immobilized on ELISA plates. (B)  
26 477 Alternatively, E1-E2 heterodimers prepared from virus transfected cells were used to determine antibody  
27 478 cross binding. Plates were incubated with serial dilutions of mouse sera from vaccinated and mock treated  
28 479 animals, and bound antibodies were detected with a secondary anti-mouse antibody coupled to horseradish  
29 480 peroxidase (HRP). The background detected upon incubation with sera from mock vaccinated animals  
30 481 was subtracted and the end point binding titers were calculated based on the median effect method as  
31 482 described in the materials and methods section. Raw data are provided in the supplementary figure 2A and  
32 483 B. (C) Antibody cross-binding expressed relative to vaccination with the parental J6-E2 protein and  
33 484 measured by ELISAs using recombinant E2 (left panel) or E1-E2 heterodimers extracted from cells with  
34 485 infectious HCV (right panel). Horizontal bars represent the median value of cross binding relative to the  
35 486 wildtype J6-E2 vaccination.  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 487 **Figure 6: Quantification of vaccine-induced neutralizing antibody responses.** (A) Neutralization of  
46 488 autologous Jc1 and of heterologous GT5A reporter virus by vaccine induced mouse antibodies and human  
47 489 antibodies from HCV patients. Viruses were incubated with given doses of antibodies and infection  
48 490 efficiency was determined and expressed relative to infections conducted in the absence of antibodies.  
49 491 Means ± SD of three technical replicates are shown. (B) Influence of vaccine-induced mouse IgG on  
50 492 neutralization by human neutralizing antibodies. Purified mouse IgGs (100µg/ml; pooled from all animals  
51 493 in one group) were mixed with human IgG (10µg/ml) from an HCV patient, pre-incubated with GT5a  
52 494 reporter virus for 1h, and then used to inoculate Huh-7.5 cells. Infection efficiency was determined as  
53 495 above and is expressed relative to infections conducted in the absence of any IgGs. Means ± SD (n=3).  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 496 **Figure 7: Vaccination with E1E2 heterodimers with or without glycosylation site mutation induces**  
5 **comparable neutralizing antibody responses. (A)** Effect of N-glycosylation mutations on E1E2  
6 497 **interaction with CD81-LEL.** Purified WT or N-glycosylation mutant (N417Q and N532Q) E1E2 proteins  
7 498 (25, 50 and 100 ng) were added to CD81-LEL coated wells or BSA coated wells (single 100 ng E1E2  
8 499 dose) as a negative control. Bound E1E2 antigens were detected by anti-E2 (H53) mAb. Values are shown  
9 500 as a percentage of WT E1E2 binding at the highest dose tested (100 ng). (\*) Designates  $p < 0.05$  respective  
10 501 to WT (100 ng) by One way ANOVA; Tukey post-hoc test. **(B)** HCV E2 binding antibodies induced by  
11 502 vaccination. WT recombinant E2 (384-661) H77c antigens were coated to ELISA plates in triplicate and  
12 503 probed with post-vaccinated mice sera. Binding of E2-specific antibodies from WT, N417Q, and N532Q  
13 504 (1000, 2000, 4000-fold dilutions) vaccinated animals compared to control **(C)** sera (1000-fold dilution)  
14 505 were detected by anti-mouse HRP conjugated secondary antibody and peroxidase substrate. The OD450-  
15 506 570 nm values (mean and SEM) plotted vs serum dilution. Shown is one representative of three  
16 507 independent experiments. **(C)** Competition of mice antisera with HCV cross-neutralizing monoclonal  
17 508 antibodies (mAb) to E1E2. Microtiter wells containing GNA-purified E1E2 H77c were incubated with  
18 509 diluted terminal antiserum (1:100) from control (C) or E1E2 (WT, N417Q or N532Q antigen) vaccinated  
19 510 mice. After washing, plates were incubated with anti-HCV mAbs. Bound AR3B, 1:7 and HC33.4 were  
20 511 detected with anti-human alkaline phosphatase-conjugated secondary antibody and biotin-AP33 detected  
21 512 using neutravidin-alkaline phosphatase. The percentages of mAb binding were calculated relative to the  
22 513 amount of mAb bound in the absence of antiserum. Shown are mean values for each group  $\pm$  range from  
23 514 three independent experiments. (\*) designates  $p < 0.05$  respective to the control group by One way  
24 515 ANOVA; Tukey post-hoc test. **(D)** Comparison of the HCVpp (H77) neutralization response from WT  
25 516 and N-glycosylation mutant E1E2 vaccinated mice. Neutralization assay using homologous HCVpp H77  
26 517 (1a) were performed using pre- and post-vaccinated sera (1:100) and the group means with SEMs plotted  
27 518 from representatives of three independent experiments. Positive control: Anti-CD81 mAb [1  $\mu\text{g/ml}$ ].  
28 519  
29 520 Abbreviations: OD, optical density.







**Figure 1****A****B****C****D**

Figure 2


A



B




C




**Figure 3**

**A**



**B**



**C**




Figure 4






Figure 5


A



B




C




**Figure 6**

**A**


**Jc1 (GT2a)**










**GT 5a**



**B**






**Figure 7****A****B****C****D**


## Parental HCV

Conserved epitopes/CD81 binding site partially occluded




## Mutant HCV

Conserved epitopes/CD81 binding site more exposed



## Recombinant E2

HVR1-deleted  
deglycosylated



## Antibody response

Strong binding to recombinant protein  
Enhanced cross-binding to E2  
Comparable binding to E1-E2  
Strain-specific neutralization  
No cross-neutralization

## Highlights

- High resolution mapping of the impact of HVR1 and glycosylation sites on CD81 binding, antibody binding and virus neutralization
- Viral mutants lacking HVR1 and selected glycosylation sites are functional and they expose the viral CD81 binding site and conserved cross-neutralization epitopes
- E2 proteins with these mutations induce cross-binding and non-interfering antibodies in mice