



You have downloaded a document from
RE-BUŚ
repository of the University of Silesia in Katowice

Title: Assets and threats to Molinia meadows (Molinion caeruleae Alliance) on chosen NATURA 2000 areas in the eastern part of the Silesian Upland

Author: Teresa Nowak, Beata Węgrzynek, Barbara Tokarska-Guzik

Citation style: Nowak Teresa, Węgrzynek Beata, Tokarska-Guzik Barbara. (2015). Assets and threats to Molinia meadows (Molinion caeruleae Alliance) on chosen NATURA 2000 areas in the eastern part of the Silesian Upland. "Acta Scientiarum Polonorum. Agricultura" (Vol. 14, nr 4 (2015), s. 49-61).



Uznanie autorstwa - Użycie niekomercyjne - Licencja ta pozwala na kopowanie, zmienianie, remiksowanie, rozpowszechnianie, przedstawianie i wykonywanie utworu jedynie w celach niekomercyjnych. Warunek ten nie obejmuje jednak utworów zależnych (mogą zostać objęte inną licencją).

ASSETS AND THREATS TO *Molinia* MEADOWS (*Molinion caeruleae* ALLIANCE) ON CHOSEN NATURA 2000 AREAS IN THE EASTERN PART OF THE SILESIAN UPLAND

Teresa Nowak, Beata Węgrzynek, Barbara Tokarska-Guzik
University of Silesia in Katowice

Abstract. The studies on diversity and conservation status of *Molinia* meadows as significantly endangered natural habitats are very important in the context of nature protection on the European scale. The collected basic data for exemplary protected objects in the Silesian Upland, whose main subject of protection is the already mentioned natural habitat, provide basis for continuing monitoring and for drawing conclusions concerning their protection. This was the main aim of the undertaken research. Occurrence of the *Galio veri-Molinietum* and the *Selino carvifoliae-Molinietum caeruleae* associations from the *Molinion caeruleae* alliance were observed on the analyzed areas. The most important floristic assets of these areas include occurrence of a significant number of 5 plant species under strict protection as well as occurrence of *Laserpitium prutenicum* regarded as critically endangered on the areas of the Silesian Voivodeship. Communities on the designated Natura 2000 areas of "Meadows in Jaworzno" above all are endangered by natural processes (plant succession), while the area of "Meadows in Sławków" by human activities connected with amelioration and development as well as devastation of the area by off-road vehicles. Significance of the areas designated for protection in preserving valuable phytocenotic structures has been highlighted, especially in the aspect of their location on urban areas.

Key words: endangered habitats, meadow communities, "Meadows in Jaworzno", "Meadows in Sławków", non-forest vegetation, Silesian Voivodeship

INTRODUCTION

In Europe, after the end of the II World War, shift in agricultural management from extensive into intensive, caused significant transformation of habitats and depletion of

Corresponding author: dr Teresa Nowak, Department of Botany and Nature Protection of University of Silesia in Katowice, Jagiellońska 28, 40-032 Katowice, e-mail:
teresa.nowak@us.edu.pl

© Copyright by Wydawnictwa Uczelniane Uniwersytetu Technologiczno-Przyrodniczego w Bydgoszczy, Bydgoszcz 2015

agrocenoses [Robinson and Sutherland 2002, Poptcheva *et al.* 2009]. Therefore, a very important task of biodiversity protection is on one hand determination of the current status of habitats and plant communities on areas utilized in agriculture, and on the other hand monitoring them. Owing to this, type of changes may be precisely determined, as well as their scale and trends. Significance of monitoring species and natural habitats is highlighted as a fundamental activity providing important information for efficient realization of nature protection [Schmeller 2008]. The critically endangered ones, both in terms of the continent and of Poland, include semi-natural meadow ecosystems. Intensification in agriculture and abandoning less productive types of meadows result, among other processes, in the succession of trees and shrubs with a simultaneous decrease in floristic diversity and disappearance of many protected and endangered species [Balátová-Tuláčková 1985, Huhta *et al.* 2001, Huhta and Rautio 2005, Poschlod *et al.* 2005, Michalska-Hejduk and Kopeć 2012a]. The most naturally valuable habitats include extensively utilized meadows of various moisture level, on which there develop phytocenoses from the *Molinion caeruleae* alliance. In Poland, over the past few decades, their diminishing is observed [Kornaś and Dubiel 1990, Barabasz 1994, Denisiuk *et al.* 1995, Kucharski 1999, 2009, Brzeg and Wojterska 2001, Michalska-Hejduk 2001, Barabasz-Krasny 2002, Kącki 2007]. In some regions of Poland, additional factors affecting transformation of habitats and plant cover are intensive industrialization and urbanization. These areas include most of the Silesian Upland (south Poland). However, despite heavy degradation of natural environment occurring for ages, there are some vegetation enclaves which are relatively well developed in terms of phytocenosis, also including meadow communities. However, they were not the subject of a complex study. Data occurring in literature is often general or concerns only some regions, selected types of meadows or occurrence of protected and rare species in meadow communities [Ciepał and Jędrzejko 1977, Babczyńska-Sendek *et al.* 1994, Babczyńska-Sendek 1998, 2009, Bernacki and Nowak 1994, Nowak and Bernacki 1997, Tokarska-Guzik 1997, Zalewska 1997, Bula and Nowak 2000, Czylok and Baryła 2003, Malewski 2006, Suder 2008, Babczyńska-Sendek and Henel 2009, Tokarska-Guzik *et al.* 2012b]. Even such fragmentary documentation confirms unique abundance and floristic diversity of meadows occurring here in the past. In the industrial landscape of the Upper Silesia (the southeastern part of the historical and geographical region of Silesia), meadow communities have mosaic distribution and usually cover small areas. Especially in such degraded areas these are valuable centers of biodiversity, which should be under special protection.

In 2008, parts of meadows of the best preserved structure were suggested to be included in the protection as Natura 2000 areas. In the present study, special attention has been paid to two of them located on urban areas: "Meadows in Jaworzno" and "Meadows in Slawków". One of the types of natural habitats protected here are *Molinia* meadows from the order *Molinion caeruleae* (code 6410), and they are the main subject of this study.

The main aims of the study included presentation of the current diversity of the analyzed type of meadow communities, preliminary assessment of their conservation status, as well as determination of the most important floristic assets, as well as indicating trends in management on areas of the studied protected localities.

MATERIAL AND METHODS

The research included meadows of two Natura 2000 areas: "Meadows in Jaworzno" and "Meadows in Sławków". According to physical and geographical division they are located in the Silesian Upland [Kondracki 2009], in two towns, at the eastern border of the Upper Silesia, in the central part of the Silesian Voivodeship. The area of "Meadows in Jaworzno" ($50^{\circ}12' N$; $19^{\circ}19' E$) consists of 4 parts of the total area of 36.45 ha [<http://natura2000.gdos.gov.pl/datafiles>] situated in the district of Ciężkowice, in the north-eastern part of town. The area of "Meadows in Sławków" ($50^{\circ}18' N$; $19^{\circ}20' E$) covers 3 parts of a total area of 50.97 ha [<http://natura2000.gdos.gov.pl/datafiles>], located in the following districts: Ciołkowizna, Kozioł and Korzeniec, in the central part of town. On each of the mentioned areas, only their small parts are still utilized in agriculture, no mowing nor grazing has been conducted on majority of these areas for at least 10 years. However, burning of meadows can be observed there in spring. Moreover, in the neighboring areas, a progressing single-detached dwelling is observed along with an accompanying infrastructure.

Field studies were carried out in July 2012. Based on the observations conducted on the whole area of the analyzed localities, borders of the occurrence of the analyzed habitat were established on these areas. In order to determine phytosociological diversity of the habitats from the *Molinion caeruleae* alliance, as well as its current conservation status and threats, some elements of methodology were applied to monitor species and habitats of Natura 2000 [Kącki and Zaluski 2004, Michalska-Hejduk and Kopeć 2012b]. In localities where the authors considered *Molinion coeruleae* patches as developed in a representative way for particular areas, the following research areas were designated: 4 in case of "Meadows in Jaworzno" (J I-J IV) and 3 in case of "Meadows in Sławków" (S I-S III). They were designated arbitrarily, taking into consideration, among other things, habitat diversity and current stage of ecological transformations. In three cases (the area J I and J III as well as S III) they had a form of a transect of 10 m \times 200 m, in other cases they had a different shape, though they maintained a similar size, i.e. 2000 m². Within each of the areas, in the typically developed habitat patches, 3 phytosociological relevés were taken in each case with Braun-Blanquet method of an area of 100 m². If possible they were located in a similar distance from each other. In case of transects, efforts were made to localize relevés at the beginning, in the middle and at the end of it. Geographical coordinates of their centers were given for each of them. In total, 21 relevés were taken (12 for "Meadows in Jaworzno" and 9 for "Meadows in Sławków") and they were juxtaposed in a phytosociological table determining diversity of their vegetation communities. Nomenclature of plant species was adopted based on the up-to-date study [Mirek *et al.* 2002], while taxonomy and nomenclature of communities according to Matuszkiewicz [2008]. Within the whole research areas, some selected parameters of natural habitat were evaluated [Michalska-Hejduk and Kopeć 2012b]: (i) typical species, i.e. characteristic and differential for the *Molinion caeruleae* alliance [Matuszkiewicz 2008], (ii) invasive alien species [Tokarska-Guzik *et al.* 2012a], (iii) expansive species of herbaceous plants, (iv) expansion of shrubs and saplings. All these parameters were used to determine conservation status of the studied natural habitat, and to indicate threats to it. Evaluation of flora was carried out based on the list of protected species in Poland (Regulation of the Ministry of Environment of 9 October 2014) as well as on "The red list of vascular plants of Silesian Voivodeship" [Parusel and Urbisz 2012]. Moreover, abundance of

population was estimated for these species. Depending on the type of growth of particular species, either the area that they covered was given, or the shoots were counted (data exclusively for generative shoots). In case of counting, precise resources are given for species having up to 30 shoots. However, for others, they were estimated while determining first the total area taken, and then multiplying it by the number of shoots on the randomly selected 1 m².

RESULTS AND DISCUSSION

Conducted field experiments confirmed compliance of the current proportion of area of the studied natural habitat with the one given in the documentation of both analyzed Natura 2000 areas [<http://natura2000.gdos.gov.pl/datafiles>]. In case of "Meadows in Jaworzno", *Molinia* meadows covered app. 60% of the area, while on the area of "Meadows in Sławków" their proportion was significantly lower, app. 15%.

Despite a small area of the analyzed localities and strong pressure on the side of human, communities on the meadows from the *Molinion caeruleae* alliance are quite diversified (Table 1) and present significant floristic and phytocenotic assets. They may be classified as the *Galio veri-Molinietum* and the *Selino carvifoliae-Molinietum caeruleae* association. Moreover, small part of communities developed fragmentarily with a slight proportion of species characteristic for the association. More often and more abundantly represented there are species from the *Molinion caeruleae* alliance and from the order *Molinietalia*. Among them there are: *Deschampsia caespitosa*, *Molinia caerulea*, *Sanguisorba officinalis* and *Angelica sylvestris*. In impoverished phytocenoses, the dominant species are often: *Rubus caesius* (with coverage sometimes reaching 80%), *Calamagrostis epigejos* and *Solidago canadensis*. Patches of the association *Galio veri-Molinietum*, for the first time described by Kącki [2007], developed on drier parts of meadows with an organic bedding containing calcium carbonate, characterized by a significant proportion of thermophilous species. Communities of a similar character, namely *Galio borealis-Molinietum*, are given for Greater Poland [Brzeg and Wojterska 2001], while in the Silesian Upland they were observed in Dąbrowa Górnica and Trzebyczka [Suder 2008]. Phytocenoses from this area were floristically abundant, on average there occurred 45 species per relevé (from 24 to 65). Moreover, occurrence of 10 protected species was observed on them, and 8 species considered as rare. Patches of associations on the studied Natura 2000 areas do not indicate such floristic diversity. From 20 to 42 species were observed there, on average app. 30 species per relevé (Table 1), which in turn makes them closer to the mean value for phytocenoses of the association for Poland [Kącki 2012]. In total, in the patches of the discussed communities, occurrence of 138 species of vascular plants was observed. Apart from characteristic and determining species for the association such as *Galium boreale* (at significant cover), *Betonica officinalis*, *Serratula tinctoria*, *Inula salicina*, *Silaum silaus*, *Galium verum* in analysed patches *Dianthus deltoides*, *Genista tinctoria*, *Knautia arvensis*, *Libanotis pyrenaica*, *Thymus pulegioides* were recorded. In the area studied, the meadows are often located in close proximity to xerothermic grasslands and thermophilous fringes from where thermophilous plant species can penetrate into the meadow communities. Another characteristic trait of this community is the dominance of dicotyledonous over monocotyledonous plants including grasses, which confirmed the observation of other authors [Suder 2008, Kącki 2012].

Also occurrence of phytocenoses of *Selino carvifoliae-Molinietum caeruleae* was observed. On the studied areas, the most numerous and most frequent characteristic species of this association is *Selinum carvifolia*, while of the differential one: *Carex panicea*. Moreover, there occur: *Iris sibirica*, *Carex acutiformis*, *Gentiana pneumonanthe* and *Laserpitium prutenicum*. The latter species was observed on the area of "Meadows in Jaworzno" for the first time. It belongs to the rarest elements of vascular plant flora in the Silesian Upland. Significant proportion, greater than in case of the previous association, had species from the order *Arrhenatheretalia* and class *Molinio-Arrhenatheretea*, including representatives of the family Poaceae: *Deschampsia caespitosa*, *Arrhenatherum elatius*, *Holcus lanatus*, *Phleum pratense*. Greater coverage than in the previously described communities was also achieved here by *Molinia caerulea*. Number of species per relevé oscillated from 25 to 35, being on average 30. These values are similar to the ones given for the eastern part of the Silesian Upland: 22, 47 and 33 [Suder 2008], respectively, as well as for other regions in Poland: on average 34 species per relevé [Kącki 2012]. Patches of this community were observed mainly on the lower parts of the terrain, with a higher level of groundwater, and their physiognomy had a valley-tussock structure, mainly because of quite numerously occurring *Molinia caerulea* and *Deschampsia caespitosa*.

While analyzing the basic parameter of the conservation status of a natural habitat, i.e. occurrence of typical species (characteristic of the *Molinion caeruleae* alliance), it was found that on each of the analyzed areas, there were observed 10 out of 15 such species [Matuszkiewicz 2008]. *Laserpitium prutenicum* was observed only in Jaworzno. Another characteristic species, *Carex tomentosa*, although occurred on meadows in Ślawków, it was outside the analyzed areas. Compared with the described patches of meadows from other parts of the Silesian Upland [Suder 2008, Babczyńska-Sendek and Henel 2009], the number of typical species in the analyzed phytocenoses was slightly lower. However, except *Molinia caerulea*, sometimes *Galium boreale* and *Serratula tinctoria*, they occurred here in lower numbers.

Other parameters of the conservation status are: presence of invasive alien species and expansion of trees and shrubs. In the analyzed patches 3 species of invasive plants of American origin were observed. These are: *Solidago canadensis*, *Padus serotina* and *Quercus rubra*. Their proportion was insignificant, they did not reach 5% of coverage on the analyzed areas. Apart from the two latter species on meadows there were also observed other species of trees and shrubs of different age: from seedlings to a few-year-old plants. These were: *Rubus caesius*, *Crataegus monogyna*, *Frangula alnus*, *Corylus avellana*, *Cornus sanguinea*, *Viburnum opulus*, *Betula pendula*, and *Tilia cordata*. Their total coverage also did not reach 5%. The symptom of the occurring secondary succession of vegetation connected with abandoning utilization may also be presence of herbaceous species such as: *Filipendula ulmaria*, *Lysimachia vulgaris*, *Valeriana officinalis*, as well as other herbaceous species like: *Calamagrostis epigejos*, *Cirsium arvense*, *Phragmites australis*, covering usually up to 20%.

Conservation status of the described patches of meadows may be considered, in the period of observation, as good (appropriate). However, visible symptoms indicate change in this assessment in a short time, unless particular actions are taken.

Table 1. Meadow communities recorded on chosen Natura 2000 areas in the eastern part of the Silesian Upland
 Tabela 1. Zbiorowiska łąkowe odnotowane na terenach wybranych obszarów Natura 2000 wschodniej części Wyżyny Śląskiej

		Cover coefficient – Współczynnik pokrycia																				
		Constancy – Stałość																				
Succesive number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	
Numer porządkowy																						
Field number of relevé – Numer zdjęcia w terenie	10	11	15	13	9	7	17	12	18	14	8	2	1	20	3	21	19	16	6	5	4	
Locality Stanowisko	J IV 1	J IV 2	S I 3	S I 1	J III 3	J III 1	S II 2	J IV 3	S II 3	S I 2	J III 2	J I 2	J I 1	S III 2	J I 3	S III 3	S III 1	S II 1	J II 3	J II 2	J II 1	
Date of relevé	27.07	27.07	03.08	03.08	27.07	27.07	03.08	27.07	03.08	03.08	27.07	27.07	03.08	27.07	03.08	03.08	03.08	03.08	27.07	27.07	27.07	
Data	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	2012	
Cover of shrub layer b																						
Pokrycie warstwy krzewowej b, %	3	3	2	–	10	–	10	3	2	1	10	3	5	1	–	1	2	5	–	1	5	
Cover of herb layer c																						
Pokrycie warstwy zielnej c, %	95	100	95	100	95	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	
Cover of moss layer d																						
Pokrycie warstwy mchowej d, %	–	5	5	5	–	–	5	–	–	5	5	10	5	10	10	5	–	5	5	20	–	
Species number in relevé – Liczba gatunków w zdjiku	30	30	26	42	27	24	20	29	30	39	25	30	31	29	28	25	35	24	21	27	23	
Ch. Ass. et D.* Galio veri-Molinietum																						
<i>Galium boreale</i>	2	2	3	3	1	1	2	2	2	1	1	3	3	1	1	1	1	1	1	1	IV	1250
<i>Beonica officinalis</i>	3	2	2	1	2	2	2	2	1	1	1	1	1	1	1	1	1	1	1	+	IV	822
<i>Gladiolus imbricatus</i>	+	1	r	+	1	r	2	2	1	1	1	1	1	1	1	1	1	1	1	1	III	324
<i>Serratula tinctoria</i>	1	2	4	3	4	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	II	513
<i>Galium verum*</i>	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	II	73
<i>Inula salicina</i> 2 (1, 2), <i>Silium silaus</i> + (8), 1 (3), 2 (10), 3 (4)																						
Ch. Ass. et D.* Selino carvifoliae-Molinietum caeruleae																						
<i>Selinum carvifolia</i>	3	2	+	+	2	+	+	+	2	+	2	3	3	2	2	2	2	+	+	+	V	1300
<i>Carex panicoides*</i>	–	–	–	–	–	–	–	–	–	–	–	3	2	1	2	2	1	2	+	+	III	478
<i>Iris sibirica</i>	–	1	2	–	–	–	–	–	–	3	2	–	2	–	–	2	–	–	–	–	II	619
<i>Carex acutiformis</i> * 1 (12, 19), <i>Gentiana pneumonanthe</i> + (9), 1 (1, 6), <i>Laserpitium prutenicum</i> + (4, 10), 2 (11)																						

Table 1 continued – cd. tabelli 1

Ch. All. et D.* Molinion caeruleae	2	2	3	3	2	2	3	4	2	3	3	4	3	+	+	V	3001
<i>Molinia caerulea</i>	3	2	.	3	.	1	2	.	1	2	.	1	1	.	.	2	III
<i>Succisa pratensis</i>	2	1	.	3	.	r	2	.	r	+	1	1	1	.	.	2	III
<i>Potentilla erecta*</i>	2	.	.	r	.	.	2	+	+	1	1	1	1	.	.	+	632
<i>Salix repens</i> subsp. <i>rosmarinifolia</i> * C	1	.	+	.	2	.	2	.	.	.	+	.	+	.	.	+	III
<i>Briza media</i> * + (1), 1 (2, 4), 2 (14), <i>Carex flava</i> * + (1), <i>Parnassia palustris</i> * + (4, 10), <i>Pimpinella saxifraga</i> * + (6)	264
Ch. All. Filipendulion	+	2	1	3	1	1	+	1	+	1	1	1	1	+	1	3	538
<i>Filipendula ulmaria</i>	.	+	2	1	2	1	1	2	+	1	1	1	1	1	1	.	III
<i>Lysimachia vulgaris</i>	.	+	2	1	2	1	1	1	+	1	3	3	+	.	.	1	370
<i>Valeriana officinalis</i>	.	2	+	.	2	1	1	1	1	+	1	1	1	.	.	1	369
<i>Hypericum tenuifolium</i>	+	.	.	.	+	.	.	2	.	.	1	1	.	.	.	+	II
<i>Lythrum salicaria</i>	+	.	.	1	1	1	1	1	1	1	.	.	109
<i>Geranium palustre</i> + (19), <i>Stachys palustris</i> + (2), <i>Thalictrum lucidum</i> + (18)	.	.	.	+	.	.	+	.	+	1	1	1	1	.	.	.	II
Ch All. et D.* Calthion	1	2	3	1	2	3	.	2	2	4	3	2	2	1	1	2	IV
<i>Cirsium rivulare</i>	.	.	.	2	.	2	.	2	2	+	3	2	1	1	3	1	522
<i>Polygonum bistorta</i>	.	.	.	2	.	2	.	2	2	+	3	2	1	1	1	1	631
<i>Caltha palustris</i> + (2), 2 (6), <i>Cirsium oleraceum</i> + (14), 1 (16), <i>Geum rivale</i> * + (11, 14, 16), 1 (13), <i>Juncus conglomeratus</i> + (17), 1 (14), <i>J. effusus</i> + (17), 1 (14)	
Ch. O. Molinetalia	
<i>Sanguisorba officinalis</i>	1	1	3	1	2	2	3	3	3	2	3	2	2	1	1	2	V
<i>Deschampsia caespitosa</i>	.	1	2	1	1	1	1	3	1	1	2	2	3	2	2	2	130
<i>Angelica sylvestris</i>	+	+	2	+	1	1	1	1	1	1	2	2	2	1	2	3	646
<i>Lotus uliginosus</i>	.	.	1	+	1	1	1	1	1	1	1	1	1	1	1	1	144
<i>Equisetum palustre</i>	.	.	1	2	1	1	1	1	1	1	1	1	1	1	1	1	132
<i>Galium uliginosum</i>	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	120
<i>Colchicum autumnale</i>	+	+	1	1	1	1	1	1	1	1	1	1	1	1	1	1	97
<i>Cirsium palustre</i> + (1, 8), 2 (14, 17), <i>Gaultheria mollugo</i> 1 (17), 2 (9, 16), <i>Lychnis flos-cuculi</i> (8), <i>Poa angustifolia</i> + (8, 12, 17), 2 (14)	
Ch. O. Arrhenatheretalia	
<i>Arrhenatherum elatius</i>	+	2	3	.	2	1	1	1	454	
<i>Achillea millefolium</i>	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	120
<i>Dactylis glomerata</i>	.	.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	72
<i>Centaura oxylopis</i>	+	1	1	1	1	1	1	1	1	1	1	1	3
<i>Carum carvi</i> + (6, 20), 1 (2), 2 (13), <i>Crepis biennis</i> r (9, 21), + (14), <i>Daucus carota</i> r (15), <i>Heracleum sphondylium</i> + (13, 16, 20), <i>Juncus inflexus</i> 1 (4), <i>Knautia arvensis</i> 1 (4), <i>Leucanthemum vulgare</i> + (8), <i>Lotus corniculatus</i> + (7), <i>Lysimachia nummularia</i> + (2, 15)	

abbreviations – objaśnienia skrótów: J – Jaworzno, S – Śląsków, I, II, IV – successive number of transects studied – numer kolejny analizowanego transekta, 1, 2, 3 – number of relevé taken within transect – numer zbiioru w transektie

The studied areas are also characterized by great floristic assets. A particular asset here is presence of 5 strictly protected species: *Epipactis palustris*, *Gentiana pneumonanthe*, *Gladiolus imbricatus*, *Gymnadenia conopsea* subsp. *conopsea*, *Iris sibirica*, as well as 4 species under partial protection: *Centaurium erythraea* subsp. *erythraea*, *Colchicum autumnale*, *Listera ovata* and *Phyteuma orbiculare*. The latter species represents the mountain element in the flora of the Polish lowlands and belongs to a very rare species (2 localities) in the Silesian Upland [Tokarska-Guzik 1997, Nowak 1999]. Most of the mentioned species in the growing season 2012 were represented quite numerously (Table 2). Moreover, outside the patches of the discussed communities, occurrence of two more species was observed under strict protection: *Liparis loeselii*, *Orchis morio*, and two under partial protection: *Dactylorhiza majalis* and *Pedicularis palustris*. In addition, 16 from the observed plant species (including 7 among the protected ones) have been classified as the endangered ones in the Silesian Voivodeship [Parusel and Urbisz 2012] (Table 2). *Laserpitium prutenicum* belongs to the particularly valuable ones in this group and it got category CR (critically endangered).

Table 2. Species threatened and protected by law recorded in the areas studied

Tabela 2. Gatunki zagrożone i prawnie chronione odnotowane na badanych obszarach

Taxon name Nazwa taksonu	Protection by law Ochrona prawnna	Categories of threat Kategorie zagrożenia	Abundance/size of population Liczliwość/wielkość populacji	
			„Meadows in Jaworzno” „Łąki w Jaworznie”	„Meadows in Śląsków” „Łąki w Śląskowie”
<i>Centaurium erythraea</i> subsp. <i>erythraea</i>	§	–	*15	20
<i>Colchicum autumnale</i>	§	VU	300	500
<i>Epipactis palustris</i>	§§	NT	50	100
<i>Equisetum pratense</i>	–	LC	**5 m ²	1 m ²
<i>Gentiana pneumonanthe</i>	§§	VU	200	400
<i>Gladiolus imbricatus</i>	§§	NT	400	200
<i>Gymnadenia conopsea</i> subsp. <i>conopsea</i>	§§	NT	–	15
<i>Inula salicina</i>	–	VU	25 m ²	–
<i>Iris sibirica</i>	§§	VU	20 m ²	15 m ²
<i>Laserpitium prutenicum</i>	–	CR	800	
<i>Listera ovata</i>	§	LC	50	80
<i>Parnassia palustris</i>	–	NT	30	20
<i>Phyteuma orbiculare</i>				
<i>Serratula tinctoria</i>	–	NT	40 m ²	50 m ²
<i>Silaum silaus</i>	–	VU	100 m ²	80 m ²
<i>Thalictrum flavum</i>	–	VU	2 m ²	–
<i>Thalictrum lucidum</i>	–	NT	5 m ²	10 m ²

§§ – strictly protected species – gatunki ścisłe chronione, § – partly protected species (according to the Regulation of the Ministry of Environment [2014]) – gatunki częściowo chronione (zgodnie z Rozporządzeniem Ministra Środowiska [2014])

categories of threat in the Silesian Voivodeship (according to Parusel and Urbisz [2012]) – kategorie zagrożenia w województwie śląskim (zgodnie z opracowaniem Parusel i Urbisz [2012]): CR – critically endangered – krytycznie zagrożony, VU – vulnerable – narażony, LC – least concern – najmniejszej troski, NT – near of threat – bliski zagrożenia

* number of flowering shoots – liczba pędów kwitnących, ** total area of clump-forming plant species – całkowita powierzchnia zajmowana przez gatunki o wzroście kępowym

According to Kącki [2012], development of particular types of communities on *Molinia* meadows and their conservation, beside habitat conditions is also affected by the method of agricultural management. Mowing, poorly tolerated by *Molinia caerulea*, reduces its occurrence, while grazing is favorable for the development of dicotyledonous plant species. A similar balance was observed in the analyzed areas. Interactions resulting from natural or anthropogenic processes threaten preservation of *Molinia* meadows in the appropriate condition [Załuski 2007, 2011, Kącki 2012, Michalska-Hejduk and Kopeć 2012b]. Based on the conducted observations, threats from both these categories have been identified. Apart from native species, both herbaceous and woody ones mentioned above, among plants entering meadow communities there were observed the following invasive species of North American origin: *Padus serotina*, *Quercus rubra*, and *Solidago canadensis*. These processes are more visible on the area of „Meadows in Jaworzno”. On the other hand, main anthropogenic factors threatening the described phytocenoses are: the cessation of mowing and grazing, improvement in drainage, periodic burning off, expansion of urbanisation, proximity of many communication routes together with the presence of the ditches accompanying them and the devastation of land by cross-country vehicles. Influence of this group of factors is more visible on „Meadows in Sławków”.

CONCLUSIONS

1. Floristic and phytocenotic assets of the discussed Natura 2000 areas fully justify bringing these forms of nature protection into existence, especially in the aspect of preserving them in the form relatively slightly transformed on urban areas.
2. Protection of the characterized areas will contribute to preservation of local diversity, enabling at the same time conservation of genetic resources of many rare and protected species of vascular plants.
3. There is a real possibility to protect designated habitats through active treatments and reduction in external factors, e.g. building development.
4. To preserve characteristic structure of the analyzed meadows, it is recommended to mow them in late summer, once a year or once every two years, which should be included in the plan of protection tasks.
5. It is necessary to monitor the analyzed natural habitat in order to determine tendencies of changes, which will help modify methods of its management.

REFERENCES

- Babczyńska-Sendek, B. (1998). Półnaturalne i naturalne zbiorowiska nieleśne Górnego Śląska na tle zróżnicowania przestrzeni tego regionu. *Studia i materiały waloryzacji przestrzeni Górnego Śląska*, 2, 37-47.
- Babczyńska-Sendek, B. (2009). Significance of protection of the meadow and grassland communities for maintenance the floristic diversity in the area of the south-eastern Silesian Upland (Poland). *Biodiv. Res. Conserv.*, 13, 49-60.
- Babczyńska-Sendek, B., Cabała, S., Wilczek, Z. (1994). Ochrona łąk w okolicach Trzebyszczki na Wyżynie Śląskiej. *Chrońmy Przyr. Ojcz.*, 50(2), 74-79.

- Babczyńska-Sendek, B., Henel, A. (2009). Mid-forest meadows of Mokrznia and Poręba Forests – an important feature of the Silesian Upland biodiversity. [In:] J. Holeksa, B. Babczyńska-Sendek, S. Wika (eds), The role of geobotany in biodiversity conservation, University of Silesia, 45-55.
- Balátová-Tuláčková, E. (1985). Chorological phenomena of the *Molinietalia* communities in Czechoslovakia. *Vegetatio*, 59, 111-117.
- Barabasz, B. (1994). Wpływ modyfikacji tradycyjnych sposobów gospodarowania na przemiany roślinności łąk z klasy *Molinio-Arrhenatheretea*. *Wiad. Bot.*, 38(1-2), 85-94.
- Barabasz-Krasny, B. (2002). Sukcesja roślinności na łąkach, pastwiskach i nieużytkach porolnych Pogórza Przemyskiego. *Fragm. Flor. Geobot. Polonica, Suppl.* 4, 3-81.
- Bernacki, L., Nowak, T. (1994). Materiały do rozmieszczenia i poznania zasobów chronionych gatunków centralnej części Wyżyny Śląsko-Krakowskiej. *Acta Biol. Sil.*, 25(42), 24-42.
- Brzeg, A., Wojterska, M. (2001). Zespoły roślinne Wielkopolski, ich stan poznania i zagrożenia. [W:] M. Wojterska (ed), Szata roślinna Wielkopolski i Pojezierza Południowo-Pomorskiego, Wyd. Nauk. Bogucki Poznań, 39-110.
- Bula, R., Nowak, T. (2000). Kosaciec syberyjski (*Iris sibirica L.*) w zbiorowiskach łąkowych na Wyżynie Śląskiej. *Nat. Sil. Super.*, 4, 45-56.
- Ciepał, R., Jędrzejko, K. (1977). Nowe stanowisko pełnika europejskiego *Trollius europaeus* i kosaćca syberyjskiego *Iris sibirica* na Wyżynie Śląskiej. *Chrońmy Przyr. Ojcz.*, 33(2), 50-53.
- Czylok, A., Baryła, J. (2003). Notatki florystyczne i ekologiczne z okolic Dąbrowy Górnictwnej i Sławkowa (Wyżyna Śląska). *Nat. Sil. Super.*, 7, 11-17.
- Denisiuk, Z., Korzeniak, J., Płecha, R. (1995). Godne ochrony łąki w Opatkowicach pod Krakowem. *Chrońmy Przyr. Ojcz.*, 51(4), 30-35.
- Huhta, A.-P., Rautio, P. (2005). Condition of semi-natural meadows in northern Finland today do the classical vegetation types still exist? *Ann. Bot. Fenn.*, 42, 81-93.
- Huhta, A.-P., Rautio, P., Juha, T., Laine, K. (2001). Restorative mowing on an abandoned semi-natural meadow: short-term and predicted long-term effects. *J. Veg. Sci.*, 12, 677-686.
- <http://natura2000.gdos.gov.pl/datafiles> (dostęp 15.01.2015)
- Kącki, Z. (2007). Comprehensive syntaxonomy of *Molinion* meadows in southwestern Poland. *Acta Bot. Sil. Monographiae*, 2, 1-134.
- Kącki, Z. (2012). Variability and long-term changes in the composition of *Molinia* meadows in Poland: a case study using a large data set from the Polish Vegetation Database. *Acta Bot. Sil. Monographiae*, 7, 1-131.
- Kącki, Z., Załuski, T. (2004). Zmiennowilgotne łąki trzęślicowe (*Molinion*). [Purple-moor grass meadows (*Molinion*)]. [W:] J. Herbich (ed), Poradniki ochrony siedlisk i gatunków Natura 2000. Murawy, łąki, ziołośla, wrzosowiska, zarośla, Ministerstwo Środowiska Warszawa, 159-170.
- Kondracki, J. (2009). Geografia regionalna Polski. Wyd. 2. Wyd. Nauk. PWN Warszawa.
- Kornaś, J., Dubiel, E. (1990). Przemiany zbiorowisk łąkowych w Ojcowskim Parku Narodowym w ostatnim trzydziestoleciu. *Prądnik. Prace Muz. Szaf.*, 2, 97-106.
- Kucharski, L. (1999). Szata roślinna łąk Środkowej Polski i jej zmiany w XX stuleciu. Wyd. Uniw. Łódzkiego.
- Kucharski, L. (2009). Grassland vegetation of central Poland – classification and conservation problems. [In:] J. Holeksa, B. Babczyńska-Sendek, S. Wika (eds), The role of geobotany in biodiversity conservation. University of Silesia, 101-109.
- Malewski, K. (2006). Zbiorowiska łąkowe dolin rzecznych zlewni Białej Przemszy. *Nat. Sil. Sup.*, 9, 41-61.
- Matuszkiewicz, W. (2008). Przewodnik do oznaczania zbiorowisk roślinnych Polski. Ser. *Vademecum Geobotanicum*, Wyd. Nauk. PWN Warszawa.
- Michalska-Hejduk, D. (2001). Stan obecny i kierunki zmian zbiorowisk nieleśnych Kampinoskiego Parku Narodowego. *Monogr. Bot.*, 89, 1-134.

- Michalska-Hejduk, D., Kopeć, D. (2012a). Dynamics of semi-natural vegetation with a focus on *Molinion* meadows after 50 years of strict protection. *Pol. J. Environ. Stud.*, 21(6), 1731-1741.
- Michalska-Hejduk, D., Kopeć, D. (2012b). Zmiennowilgotne łąki trzęślicowe (*Molinion*). [W:] W. Mróz (ed), Monitoring siedlisk przyrodniczych. Przewodnik metodyczny. Część III. GIOŚ Warszawa, 40-52.
- Mirek, Z., Piekoś-Mirek, H., Zajac, A., Zajac, M. (2002). Flowering plants and pteridophytes of Poland. A checklist. W. Szafer Institute of Botany, Polish Academy of Sciences Kraków.
- Nowak T. 1999. Atlas rozmieszczenia roślin naczyniowych na terenie wschodniej części Garbu Tarnogórskiego (Wyżyna Śląska). Materiały, Opracowania 2. Centrum Dziedzictwa Przyrody Górnego Śląska Katowice.
- Nowak, T., Bernacki, L. (1997). Materiały do poznania flory oraz zasobów wybranych płatów łąk wschodnich obrzeży aglomeracji górnośląskiej. *Acta Biol. Sil.*, (30)47, 139-152.
- Parusel, J.B., Urbisz, A. (red.) (2012). Czerwona lista roślin naczyniowych województwa śląskiego. Raporty i Opinie, 6, 105-177.
- Poptcheva, K., Schwartz, P., Vogel, A., Kleinebecker, T., Höllzel, N. (2009). Changes in wet meadow vegetation after 20 years of different management in a field experiment (North-West Germany). *Agric. Ecosyst. and Environ.*, 134, 108-114.
- Poschlod, P., Bakker, J.P., Kahmen, S. (2005). Changing land use and its impact on biodiversity. *Basic Appl. Ecol.*, 6(2), 93-98.
- Robinson, R.A., Sutherland, W.J. (2002). Post-war changes in arable farming and biodiversity in Great Britain. *J. Appl. Ecol.*, 39, 157-176.
- Shmeller, D.S. (2008). European species and habitat monitoring: where are we now? *Biodivers. Conserv.*, 17, 3321-3326.
- Suder, A. (2008). Purple-moor grass meadows (Alliance *Molinion caeruleae* Koch 1926) in the eastern part of Silesia Upland: phytosociological diversity and aspects of protection. *Nat. Conserv.*, 65, 63-77.
- Tokarska-Guzik, B. (1997). Rozmieszczenie i zasoby roślin chronionych na terenie miasta Jaworzno. *Acta Biol. Sil.*, 30(47), 106-124.
- Tokarska-Guzik, B., Dajdok, Z., Zajac, M., Zajac, A., Urbisz, A., Danielewicz, W., Hołyński, C. (2012a). Rośliny obcego pochodzenia w Polsce ze szczególnym uwzględnieniem gatunków invazyjnych. GDOŚ Warszawa.
- Tokarska-Guzik B., Rostański A., Gorczyca J., Herczek A., Dulias R. (2012b). Przyroda miasta Jaworzna. Urząd Miejski w Jaworznie, Jaworzno.
- Zalewska, J. (1997). Łąki trzęślicowe w dolinie Przemszy. *Ochr. Przyr.*, 54, 73-79.
- Załuski, T. (2007). Zagrożenie i ochrona zespołów trawiastych. [W:] F. Frey (red.), Księga Polskich traw, W. Szafer Institute of Botany Kraków, 283-316.
- Załuski, T. (2011). Vegetation transformations of Kujawy-Pomerania Region in the last twenty years period. *Acta Univ. Lodz.*, 7, 75-98.

WALORY I ZAGROŻENIA ZMIENNOWILGOTNYCH ŁĄK TRZĘŚLICOWYCH (ZWIĄZEK *Molinion caeruleae*) NA TERENACH WYBRANYCH OBSZARÓW NATURA 2000 WSCHODNIEJ CZĘŚCI WYZYNY ŚLĄSKIEJ

Streszczenie. Badania zróżnicowania i stanu zachowania zmiennowilgotnych łąk trzęślicowych jako istotnie zagrożonych siedlisk przyrodniczych są bardzo ważne w kontekście ochrony przyrody w skali Europy. Zebrane dane podstawowe dla przykładowych obiektów chronionych na Wyżynie Śląskiej, których głównym przedmiotem ochrony jest wymienione siedlisko przyrodnicze, dają podstawę do kontynuowania monitoringu i wysuwanego wniosków dotyczących ich ochrony. Było to głównym celem podjętych

badań. Na analizowanych obszarach odnotowano występowanie płatów zespołu *Galio veri-Molinietum* i *Selino carvifoliae-Molinietum caeruleae* ze związką *Molinion caeruleae*. Do najważniejszych walorów florystycznych tych obszarów należy obecność 5 gatunków roślin objętych ochroną ścisłą o znacznej liczebności oraz występowanie *Laserpitium prutenicum*, uznanego za krytycznie zagrożony na terenie województwa śląskiego. Zbiorowiska na terenie projektowanego obszaru Natura 2000 „Łąki w Jaworznie” zagrozone są przede wszystkim przez procesy naturalne (sukcesja roślinności), natomiast na obszarze „Łąki w Sławkowie” – przez działania człowieka związane z melioracją i zabudową oraz dewastację terenu przez pojazdy terenowe. Podkreślono znaczenie proponowanych do ochrony obszarów dla zachowania cennych układów fitocenotycznych, szczególnie w aspekcie ich lokalizacji na terenach miejskich.

Key words: „Łąki w Jaworznie”, „Łąki w Sławkowie”, roślinność nieleśna, województwo śląskie, zagrożone siedliska, zbiorowiska łąkowe

Accepted for print – Zaakceptowano do druku: 24.08.2015

For citation – Do cytowania:

Nowak, T., Węgrzynek, B., Tokarska-Guzik, B. (2015). Assets and threats to *Molinia* meadows (*Molinion caeruleae* alliance) on chosen Natura 2000 areas in the eastern part of the Silesian Upland. *Acta Sci. Pol. Agricultura*, 14(3), 49-61.