

## Article

# Surface EMG signal normalisation and filtering improves sensitivity of equine gait analysis

St George, Lindsay Blair, Roy, SH, Richards, James, Sinclair, Jonathan Kenneth and Hobbs, Sarah Jane

Available at <http://clock.uclan.ac.uk/28496/>

*St George, Lindsay Blair ORCID: 0000-0002-5531-1207, Roy, SH, Richards, James ORCID: 0000-0002-4004-3115, Sinclair, Jonathan Kenneth ORCID: 0000-0002-2231-3732 and Hobbs, Sarah Jane ORCID: 0000-0002-1552-8647 (2019) Surface EMG signal normalisation and filtering improves sensitivity of equine gait analysis. Equine and Comparative Exercise Physiology . ISSN 1478-0615*

It is advisable to refer to the publisher's version if you intend to cite from the work.

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>

# Surface EMG signal normalisation and filtering improves sensitivity of equine gait analysis

St. George, L.<sup>1</sup>, Roy S.H.<sup>3</sup>, Richards, J.<sup>2</sup>, Sinclair, J.<sup>1</sup>, Hobbs, S. J.<sup>1</sup>.

<sup>1</sup>University of Central Lancashire, Centre for Applied Sport and Exercise Sciences, Preston, Lancashire PR1 2HE, United Kingdom;

<sup>2</sup>University of Central Lancashire, Allied Health Research Unit, Preston, Lancashire PR1 2HE, United Kingdom;

<sup>3</sup>Delsys Inc., Natick, MA 01760

**Corresponding author:** Dr. Lindsay St. George,  
University of Central Lancashire,  
Preston, PR1 2HE, United Kingdom.  
Tel: 01772 893775 Email: [lbstgeorge@uclan.ac.uk](mailto:lbstgeorge@uclan.ac.uk)

**Running Header:** Effect of normalisation and filtering on equine sEMG outcome measures.

## Abstract

Low-frequency noise attenuation and normalisation are fundamental signal processing (SP) methods for surface electromyography (sEMG), but are absent, or not consistently applied, in equine biomechanics. The purpose of this study was to examine the effect of different band-pass filtering and normalisation conventions on sensitivity for identifying differences in sEMG amplitude-related measures, calculated from leading (LdH) and trailing hindlimb (TrH) during canter, where between-limb differences in vertical loading are known. sEMG and 3D-kinematic data were collected from the right Biceps Femoris in 10 horses during both canter leads. Peak hip and stifle joint angle and angular velocity were calculated during stance to verify between-limb biomechanical differences. Four SP methods, with and without normalisation and high-pass filtering, were applied to raw sEMG data. Methods 1 (M1) to 4 (M4) included DC-offset removal and full-wave rectification. Method 2 (M2) included additional normalisation relative to maximum sEMG across all strides. Method 3 (M3) included additional high-pass filtering (Butterworth 4<sup>th</sup> order, 40Hz cut-off), for artefact attenuation. M4 included the addition of high-pass filtering and normalisation. Integrated EMG (iEMG) and average rectified value (ARV) were calculated using processed sEMG data from M1 – M4, with stride duration as the temporal domain. sEMG parameters, within M1 – M4, and kinematic parameters were grouped by LdH and TrH and compared using repeated measures ANOVA. Significant between-limb differences for hip and stifle joint kinematics were found, indicating functional differences in hindlimb movement. M2 and M4, revealed significantly greater iEMG and ARV for LdH than TrH ( $p < 0.01$ ), with M4 producing the lowest p values and **largest** effects sizes. Significant between-limb differences in sEMG parameters were not observed with M1 and M3. The results indicate that equine sEMG SP should include normalisation and high-pass filtering to improve sensitivity for identifying differences in muscle function associated with biomechanical changes during equine gait.

**Keywords:** surface electromyography, horse, signal processing, high-pass filter, normalise

**Conflict of Interest:** none

## 1 **1. Introduction**

2 Surface electromyography (sEMG) has long been used as a non-invasive tool for investigating  
3 the human neuromuscular system. Within the equine biomechanics field, the use of sEMG is  
4 relatively scarce compared to human studies but has gained **popularity** in the past 10 years.  
5 sEMG has proven to be a useful method for understanding equine muscle function during  
6 normal locomotion (Harrison *et al.*, 2012; Jansen *et al.*, 1992; Robert *et al.*, 1999; Zsoldos *et*  
7 *al.*, 2010a; Zsoldos *et al.*, 2010b), but also for differentiating the biomechanical effects of speed  
8 (Robert *et al.*, 2001a; Robert *et al.*, 2002), incline (Crook *et al.*, 2010; Hodson-Tole, 2006;  
9 Robert *et al.*, 2000; Robert *et al.*, 2001b), fatigue (Cheung *et al.*, 1998; Colborne *et al.*, 2001;  
10 Williams *et al.*, 2013) and lameness (Zaneb *et al.*, 2009) on equine gait. The relative ease of  
11 sEMG signal acquisition makes it an attractive tool for both human and equine researchers, but  
12 sEMG signal quality and processing techniques must be carefully considered for accurate  
13 analysis and interpretation of muscle function in response to changes in biomechanics.

14 Some of the factors that influence signal quality can be mitigated by technological advances in  
15 sensor design and complying with best practice for locating and adhering the sensor to the skin  
16 (Clancy *et al.*, 2002; De Luca, 1997; De Luca *et al.*, 2010; Roy *et al.*, 2007). There are  
17 applications however when contamination of the low-frequency sEMG spectra, from baseline  
18 and movement artefact noise, and intra/ inter-individual subject characteristics, such as  
19 subcutaneous fat thickness, are unavoidable and of particular concern when interpreting the  
20 sEMG signal (De Luca *et al.*, 2010; Halaki and Ginn, 2012; Kuiken *et al.*, 2003; Lehman and  
21 McGill, 1999; Nordander *et al.*, 2003). Methodological guidelines for human sEMG recording  
22 and processing have been published and describe optimal signal processing (SP) methods to  
23 mitigate these sources of error. Of these, the International Society of Electrophysiology and  
24 Kinesiology (ISEK) (Winter *et al.*, 1980) and Standards for Reporting EMG data (Merletti and  
25 Di Torino, 1999), recommend SP methods that include high-pass filtering, for attenuating low-  
26 frequency noise contamination (De Luca *et al.*, 2010; Van Boxtel, 2001; Van Boxtel *et al.*,  
27 1998), and normalisation, for reducing inter and intra-subject variability (Burden, 2010; Halaki  
28 and Ginn, 2012; Lehman and McGill, 1999).

29 Unfortunately, standards for sEMG signal detection and processing in equine subjects are not  
30 currently available and methodological variation within equine sEMG literature is particularly  
31 evident for SP methods (Valentin and Zsoldos, 2016). Furthermore, fundamental low-  
32 frequency noise attenuation and normalisation techniques are absent, or not consistently  
33 applied in the equine sEMG literature (Valentin and Zsoldos, 2016). Reliance on human  
34 subject-based sEMG guidelines for equine subjects is not recommended, as differences in size,  
35 mass, bipedal vs. quadrupedal gait and skin properties alone are sufficient to question their  
36 equivalence. Thus, the need for a best practice framework that follows human sEMG guidelines  
37 while taking into consideration the unique challenges associated with detecting and processing  
38 sEMG data from equine subjects, has been initiated (St. George *et al.*, 2018; Valentin and  
39 Zsoldos, 2016). Our recent work (St. George *et al.* 2018) demonstrated that simply adopting  
40 human sEMG guidelines for removal of motion artefact for equine gait studies is not adequate.  
41 The removal of low-frequency noise contamination within sEMG signals obtained from *Biceps*  
42 *Femoris* and *Triceps Brachii* during trot and canter was found to be more effective using a  
43 high-pass filter with a 30 - 40 Hz cut-off frequency, when compared to the standard  
44 recommendation of a 10 – 20 Hz cut-off frequency shown for human studies (De Luca *et al.*,  
45 2010). Although the need for an equine-specific high-pass filtering cut-off has been

46 demonstrated for optimal attenuation of signal noise in these studies, the practical effect of  
47 different high-pass filtering cut-offs on the sensitivity of sEMG outcome measures for equine  
48 gait analysis has not yet been investigated. Similar questions arise for SP practices involving  
49 sEMG signal normalisation. Valentin and Zsoldos (2016) reported that normalisation  
50 techniques are frequently absent in the equine sEMG literature, but no studies have  
51 demonstrated the consequences of this on the interpretation of equine sEMG data. In the human  
52 literature, Lehman and McGill (1999) investigated the effect of normalisation on the sensitivity  
53 of sEMG outcome measures for analysing the relationship between upper and lower rectus  
54 abdominus (RA) during a trunk curl exercise. When data were normalised to a maximum  
55 voluntary contraction (MVC) activity of the upper and lower RA were comparable, which was  
56 considered a clinically correct interpretation of muscle function, but when normalisation was  
57 omitted, a large asymmetry between upper and lower RA activity was observed (Lehman and  
58 McGill, 1999). High-pass filtering and normalisation have therefore been shown to improve  
59 the sensitivity of accurately interpreting human sEMG signal findings in relation to changes in  
60 biomechanics. However, no studies to date have demonstrated their effect on sensitivity for  
61 identifying differences in equine muscle function during gait analysis.

62 The purpose of this study is to test whether adopting a more rigorous SP protocol for sEMG  
63 filtering and normalisation provides greater sensitivity, **reflected by smaller P-values and larger**  
64 **effect sizes**, in identifying differences in muscle activation during equine gait, when compared  
65 to the current standard. For this example, sEMG and three-dimensional kinematic data,  
66 obtained unilaterally from right hindlimb and the vertebral head of the right *Biceps Femoris*  
67 (BF) during canter, were chosen *a priori*. The canter is a three-phase asymmetrical gait with a  
68 footfall pattern as follows: 1) trailing hindlimb (TrH); 2) leading hindlimb (LdH) and trailing  
69 forelimb (TrF) (as a diagonal pair); and 3) leading forelimb (LdF). During canter,  
70 biomechanically different demands are placed on hindlimb, depending on these phases of gait,  
71 with the LdH experiencing greater vertical loading (Merkens *et al.*, 1993) and flexion of stifle  
72 and tarsal joints (Back *et al.*, 1997) during stance than TrH. Because the BF acts to adduct the  
73 hindlimb and extend the hip and stifle joints during stance (Payne *et al.*, 2005; Robert *et al.*,  
74 1999), the differences in loading are expected to produce different levels of BF muscle  
75 activation, which are measured as differences in sEMG signal amplitudes during equine gait  
76 analysis. In this study, peak joint angle and angular velocity are calculated for the hip and stifle  
77 joint during the stance phase to characterize biomechanical differences for the equine subjects  
78 during the periods of sEMG signal measurement of the BF muscle. To evaluate the effects of  
79 band-pass filtering and normalisation, four different SP methods were applied to the BF sEMG  
80 signals. Methods 1 (M1) to 4 (M4) included DC-offset removal and full-wave rectification.  
81 Method 2 (M2) included additional normalisation relative to maximum sEMG across all  
82 strides. Method 3 (M3) included additional high-pass filtering (Butterworth 4<sup>th</sup> order, 40Hz  
83 cut-off), for artefact attenuation and M4 included the addition of both high-pass filtering and  
84 normalisation. Commonly employed amplitude-based sEMG parameters were computed from  
85 the processed sEMG signals from each method to quantify the magnitude of muscle activation.  
86 It is hypothesised that incorporating normalisation with the most recent recommendations for  
87 equine filtering at 40 Hz (St. George *et al.*, 2018) will provide the greatest sensitivity for  
88 identifying statistically significant differences in BF sEMG activation between LdH and TrH,  
89 **which correspond to between-hindlimb differences in joint kinematics** during canter.

90

## 91 2. Methods

92 Ethical approval for this study was obtained from the University of Central Lancashire's  
93 Animal Projects Committee (RE/13/04/SH). Written informed consent was obtained from all  
94 horse owners, riders and handlers prior to data collection.

95 Data were collected from 10 horses (age:  $9.7 \pm 2.6$  years, height:  $161.9 \pm 6.3$  cm, sex: 7  
96 geldings, 3 mares, breed: various). All horses were in training and free from lameness, as  
97 defined by their owner. sEMG and 3D kinematic data were collected unilaterally from the right  
98 hindlimb at 2088 Hz and 232 Hz respectively during ridden canter trials. Unilateral sEMG and  
99 kinematic data were collected during right and left canter lead trials, when the right hindlimb  
100 functioned as LdH and TrH, respectively. This was done to study how the different SP methods  
101 influence measures of muscle activation from one muscle when it is analysed under different  
102 loading conditions. sEMG data were collected from right BF using wireless sEMG sensors  
103 (Trigno™, Delsys Inc., USA), with a bi-polar parallel bar electrode configuration and an inter-  
104 electrode distance of 10 mm. Sensor sites for BF were approximately halfway between the third  
105 trochanter and patella, and approximately 9 cm cephalad to the cranial margin of  
106 *Semitendinosus* (Schuurman *et al.*, 2003). Prior to sensor adhesion, sensor sites were prepared  
107 by removing all hair and thoroughly cleaning with isopropyl alcohol wipes. A small amount of  
108 saline solution was applied to the electrode bars to act as an electrolytic solution (Clancy *et al.*,  
109 2002; Cram and Rommen, 1989). Sensors were then adhered to prepared sites using a  
110 combination of Delsys Adhesive Surface Interface strips (Delsys Inc., USA) and strips of  
111 double-sided tape, which were applied to the top and bottom of the sensor above each electrode  
112 pair. The sensor was positioned on the muscle belly, with electrode bars oriented perpendicular  
113 to the underlying muscle fibre direction (De Luca, 1997; Hermens *et al.*, 2000).

114 Three-dimensional kinematic data were collected to detect right hindlimb hoof impact and lift-  
115 off gait events for stride segmentation. In addition, kinematic data were collected to calculate  
116 peak joint angle and angular velocity for the hip and stifle joints during stance phase, for which  
117 the BF functions as an extensor (Payne *et al.*, 2005; Robert *et al.*, 1999), as a means of analysing  
118 muscle activity in relation to expected biomechanical differences in hindlimb function.  
119 Spherical retro reflective markers, (25 mm diameter) (Qualisys AB, Sweden) were positioned  
120 over the following anatomical landmarks on the right hindlimb: the most ventral part of the  
121 tuber coxae, greater trochanter, lateral epicondyle of the femur, talus and the center of rotation  
122 of the metatarsalphalangeal and distal interphalangeal joints. A marker was also attached over  
123 the croup for stride velocity calculation. Data were collected using eight Qualisys Oqus  
124 cameras (Qualisys AB, Sweden). Cameras were positioned side-by-side in a linear  
125 configuration and an extended calibration was conducted to collect data from multiple strides  
126 (Figure 1). The calibration volume was approximately 8 m in length.

127 **Figure 1. Equipment set-up for data collection showing camera configuration and instrumented**  
128 **equine subject.**

### 129 2.1 Data Collection

130 Data were collected during ridden canter trials using Qualisys Track Manager (QTM) software  
131 (Qualisys AB, Sweden). **Five different riders, with similar experience and ability, rode the**  
132 **horses during data collection. Each horse was ridden by their usual rider, who either owned or**  
133 **had experience riding them.** Kinematic and sEMG data were synchronously acquired using an

134 external trigger system (Delsys Trigger Module, Delsys Inc., USA). A static trial was initially  
135 recorded for each horse. Following the static trial, each horse progressed through the capture  
136 volume during ridden canter. Horses were permitted to travel at their preferred velocity and  
137 riders were instructed to position horses adjacent to placing poles, positioned on the ground  
138 approximately 4.5 m from the cameras to demarcate the optimal capture volume (Figure 1).  
139 Three successful trials were collected from each horse during right and left canter lead, which  
140 were randomised. A trial was successful when the horse held the canter and the correct canter  
141 lead through the calibrated volume and did not deviate from the optimal capture volume. The  
142 number of strides collected within the calibrated volume differed between horses, largely due  
143 to differences in sizes and stride lengths. Thus, the number of strides collected from each horse  
144 was not standardised in this study.

## 145 2.2 Data Processing and Analysis

146 Kinematic data were tracked in QTM and both kinematic and sEMG data were imported into  
147 Visual3D (C-Motion Inc., USA) for further analysis. Kinematic data were interpolated and  
148 low-pass filtered (Butterworth 4th order), with a cut-off frequency of 12 Hz, as determined  
149 using residual analysis. Hindlimb hoof impact and lift-off events were calculated from  
150 kinematic data using a hindlimb sagittal plane angle in accordance with Holt *et al.* (2017).  
151 Kinematic gait events were applied to sEMG signals to segment the signal into stance and  
152 swing phases. A constant delay of 20 ms between kinematic and sEMG data was corrected for  
153 by shifting sEMG signals forward by 5 frames prior to applying kinematic gait events. To  
154 calculate stride velocity, the first derivative of the croup marker was calculated in the sagittal  
155 plane, and the average velocity was calculated between consecutive hoof impact events.  
156 Kinematic markers were used to define the distal and proximal ends of the pelvis, femur, tibia  
157 and third metatarsal segments of the right hindlimb. A segment coordinate system (SCS) was  
158 defined for segment, with the X axis as mediolateral, Y axis as cranio-caudal and Z axis as  
159 axial. Joint angles were calculated in the sagittal plane, as rotation around the SCS X axis,  
160 using the proximal and distal segments for each joint. Joint angular velocity was determined  
161 by calculating the first derivative of the hip and stifle joint angle signals. Flexion was defined  
162 as positive and extension as negative. During stance phase, vertical forces are primarily  
163 absorbed by shortening of the hindlimb between the stifle joint and hoof (Hjerten *et al.*, 1994),  
164 while the distance between the stifle joint and tuber coxae increases as the hip joint undergoes  
165 extension (Back *et al.*, 1996; Back *et al.*, 1995; Hodson *et al.*, 2001). Thus, peak joint angle  
166 and angular velocity were calculated for hip joint extension and stifle joint flexion during  
167 stance phase.

168 Raw sEMG signals were differentially amplified by a factor gain of 909, a common-mode  
169 rejection ratio (CMRR) of > 80 dB and an internal Butterworth high-pass ( $20 \pm 5$  Hz cut-off, >  
170 40 dB/dec) and low-pass filter ( $450 \pm 50$  Hz cut-off, >80 dB/dec). Post-processing of signals  
171 was conducted in Visual3D, where four SP methods were applied to the raw sEMG data. M1  
172 represents the most commonly applied SP method within existing equine sEMG literature and  
173 includes DC-offset removal and full-wave rectification of signals following acquisition. M2  
174 follows the same protocol as M1 but includes additional normalisation relative to a maximal  
175 reference voluntary contraction (RVC) (Lehman and McGill, 1999; Sousa and Tavares, 2012;  
176 Yang and Winter, 1984). In this instance, the RVC represents the maximum sEMG outcome  
177 measure observed across all canter strides within each horse. The use of an RVC is based on  
178 recommendations from human studies where obtaining a maximal voluntary contraction

179 (MVC) is not possible or difficult to obtain, for example in participants experiencing pain or  
180 with neurologic disorders (Burden and Bartlett, 1999; Dankaerts *et al.*, 2004; Lehman and  
181 McGill, 1999; Yang and Winter, 1984). M3 and M4 follow the same SP protocol as M1 but  
182 include additional high-pass filtering using a Butterworth 4th order filter with a 40 Hz cut-off  
183 frequency based on recent equine recommendations (St. George *et al.*, 2018), with M4  
184 including both high-pass filtering and normalisation relative to the RVC, which have not been  
185 adopted routinely in equine studies.

186 Integrated EMG (iEMG) and average rectified value (ARV) represent commonly reported  
187 amplitude-based outcome measures in equine sEMG literature for studies examining  
188 differences in muscle function during gait (Robert *et al.*, 2001a; Robert *et al.*, 2001b; Robert  
189 *et al.*, 2000; Robert *et al.*, 2002; Zaneb *et al.*, 2009; Zsoldos *et al.*, 2010a; Zsoldos *et al.*, 2010b).  
190 **iEMG represents the area under the voltage curve, where the sEMG signal is integrated over a**  
191 **specified time interval, and ARV represents the mean value of the full-wave rectified sEMG**  
192 **signal over a specified time interval (Merletti and Di Torino, 1999; Winter *et al.*, 1980).** The  
193 effect of the different SP methods was therefore evaluated using **iEMG and ARV, which** were  
194 calculated in accordance with Merletti and Di Torino (1999) and Winter *et al.* (1980) using the  
195 full-wave rectified signal from Methods 1 - 4 and stride duration as the time interval.

### 196 2.3 Statistical analysis

197 For each sEMG outcome measure (**iEMG, ARV**), data from LdH and TrH were grouped within  
198 each SP method (M1, M2, M3, M4). Ensemble averages (mean  $\pm$  SD) were calculated for each  
199 sEMG (iEMG and ARV) and kinematic (peak joint angle and peak joint angular velocity)  
200 outcome measure to examine differences between limbs. One-way repeated measures  
201 ANOVAs were used to compare kinematic and sEMG outcome measures from LdH and TrH,  
202 with sEMG outcome measures compared within each SP method. Significant differences were  
203 identified at  $P < 0.05$  and effect sizes were established using partial  $\eta^2$  ( $pn^2$ ). Sensitivity for  
204 identifying differences were therefore based on conditions which provided the lowest p-value  
205 and largest effect size.

## 206 3. Results

207 Across all horses, 115 strides were analysed, with 62 and 53 strides analysed when the right  
208 hindlimb acted as TrH and LdH, respectively. Stride velocity was  $4.6 \pm 0.4$  m/s across all  
209 horses. Mean peak joint angle and angular velocity ( $\pm$ SD) data for the stifle and hip joints are  
210 presented in Figure 2 and Table 1 for normalized canter strides. During stance phase, the LdH  
211 exhibited significantly greater stifle joint flexion ( $p=0.001$ ) and hip joint extension ( $p=0.000$ )  
212 than the TrH. Significantly greater peak flexion velocity was observed in the LdH for the stifle  
213 joint ( $p=0.000$ ) and significantly greater peak extension velocity was observed in the TrH for  
214 the hip joint ( $p=0.037$ ) during stance phase.

215 Figure 2. Mean (bold line) and standard deviation (shaded area) joint angle ( $^\circ$ ) and joint angular  
216 velocity data ( $^\circ/s$ ) for a) hip joint angle, b) stifle joint angle, c) hip joint angular velocity, d)  
217 stifle joint angular velocity from LdH (blue) and TrH (red). Data are normalised over one canter  
218 stride, with the hoof-lift off event demarcated by the green vertical line. Flexion was defined  
219 as positive and extension as negative. Overall average peak joint angle and peak joint angular  
220 velocity events are presented on corresponding graphs as red and blue arrows for TrH and LdH,  
221 respectively.

222 Table 1. Mean ( $\pm$  sd) peak joint angle ( $^{\circ}$ ) and peak joint angular velocity data ( $^{\circ}/s$ ) from the  
 223 stifle joint and hip joint during stance phase. Data are grouped according to limb (LdH and  
 224 TrH). Differences between LdH and TrH are presented for each joint as P values and effects  
 225 sizes ( $pn^2$ ). Significant differences ( $P<0.05$ ) between limbs are **denoted by bold text**.

	TrH	LdH	P value	$pn^2$
Hip Joint Angle ( $^{\circ}$ )	-33.7 (8.7)	-44.1 (7.6)	<b>0.000</b>	0.823
Stifle Joint Angle ( $^{\circ}$ )	33.9 (6.3)	39.3 (5.5)	<b>0.001</b>	0.723
Hip Joint Angular Velocity ( $^{\circ}/s$ )	-226.6 (40.8)	-181.4 (37.9)	<b>0.037</b>	0.399
Stifle Joint Angular Velocity ( $^{\circ}/s$ )	342.6 (97.3)	481.5 (81.4)	<b>0.000</b>	0.866

226

227 Descriptive and inferential statistics for sEMG outcome measures are presented in Table 2 and  
 228 3 and show that LdH exhibited greater mean ARV and iEMG values than TrH across all SP  
 229 methods. Between limb differences for ARV and iEMG were only significant when  
 230 normalisation was applied in M2 and M4, with the addition of high-pass filtering in M4  
 231 resulting in a lower p value and higher effect size (ARV:  $p=0.002$ , iEMG:  $p=0.002$ ) than M2  
 232 (ARV:  $p=0.017$ , iEMG:  $p=0.016$ ). M1 and M3, which did not include normalisation, did not  
 233 detect significant differences between limbs for iEMG and ARV ( $p>0.05$ ). For both iEMG and  
 234 ARV, the addition of high-pass filtering in M3 was again found to produce higher effect sizes  
 235 and lower p values that approached significance (ARV:  $p=0.066$ , iEMG:  $p=0.074$ ) than Method  
 236 1 (ARV:  $p=0.101$ , iEMG:  $p=0.109$ ). Combined mean and standard deviation data from all  
 237 subjects in Tables 2 and 3 reveal that normalisation, employed in M2 and M4, resulted in  
 238 reduced standard deviation for iEMG and ARV outcome measures. Figures 3 and 4 illustrate  
 239 this finding by showing decreased intrasubject variability and more distinct between-limb  
 240 differences when normalisation is applied to ARV data from two different horses (Figure 3 and  
 241 4c, d). Figures 3 and 4 also illustrate the effect of high-pass filtering on between-limb  
 242 differences, which are in accordance with findings presented in Tables 2 and 3. When high-  
 243 pass filtering is applied to sEMG signals in Figures 3 and 4 (b, d), which represent M3 and M4,  
 244 respectively, a distinct between-limb difference is observed, with the LdH clearly showing  
 245 greater amplitude of sEMG activity than TrH. In comparison, when high-pass filtering is not  
 246 applied in Figures 3 and 4 (a, c), which represent M1 and M2, respectively, sEMG signals from  
 247 TrH often overlap with signals from LdH. Thus, the omission of high-pass filtering in Figures  
 248 3 and 4 does not result in distinct between limb differences.

249 Table 2. Mean ( $\pm$  sd) for iEMG, calculated using processed sEMG signals from Methods 1 to  
 250 4 and grouped according to limb (LdH and TrH). Differences between LdH and TrH within  
 251 Methods 1 to 4 are presented for each outcome measure as P values and effects sizes ( $pn^2$ ).  
 252 Significant differences ( $P<0.05$ ) between limbs are **denoted by bold text**.

Outcome Measure	Signal Processing Method	TrH	LdH	P value	$pn^2$
iEMG	<b>Method 1</b> <ul style="list-style-type: none"> <li>• DC offset removal</li> <li>• 20 - 450 Hz band pass filtered (<math>\mu V.s</math>)</li> </ul>	18.2 (13.7)	28.6 (30.8)	0.109	0.288



	<b>Method 2</b> <ul style="list-style-type: none"> <li>• DC offset removal</li> <li>• 20 - 450 Hz band pass filtered</li> <li>• Normalised</li> </ul> (% Maximum Value)	65.1 (17.2)	82.1 (7.1)	<b>0.016</b>	0.536
	<b>Method 3</b> <ul style="list-style-type: none"> <li>• DC offset removal</li> <li>• 20 - 450 Hz band pass filtered</li> <li>• Butterworth high-pass filtered (40 Hz cut-off)</li> </ul> (μ V.s)	10.4 (6.6)	17.7 (17.2)	0.074	0.345
	<b>Method 4</b> <ul style="list-style-type: none"> <li>• DC offset removal</li> <li>• 20 - 450 Hz band pass filtered</li> <li>• Butterworth high-pass filtered (40 Hz cut-off)</li> <li>• Normalised</li> </ul> (% Maximum Value)	57.4 (17.0)	79.1 (6.8)	<b>0.002</b>	0.720

253

254 Table 3. Mean ( $\pm$  sd) for ARV, calculated using processed sEMG signals from Methods 1 to 4  
255 and grouped according to limb (LdH and TrH). Differences between LdH and TrH within  
256 Methods 1 to 4 are presented for each outcome measure as P values and effects sizes ( $pn^2$ ).  
257 Significant differences ( $P < 0.05$ ) between limbs are **denoted by bold text**.

Outcome Measure	Signal Processing Method	TrH	LdH	P value	$pn^2$
ARV	<b>Method 1</b> <ul style="list-style-type: none"> <li>• DC offset removal</li> <li>• 20 - 450 Hz band pass filtered</li> </ul> (μ V)	30.6 (22.1)	47.8 (49.4)	0.101	0.300
	<b>Method 2</b> <ul style="list-style-type: none"> <li>• DC offset removal</li> <li>• 20 - 450 Hz band pass filtered</li> <li>• Normalised</li> </ul> (% Maximum Value)	65.2 (17.4)	82.4 (7.6)	<b>0.017</b>	0.533
	<b>Method 3</b> <ul style="list-style-type: none"> <li>• DC offset removal</li> <li>• 20 - 450 Hz band pass filtered</li> <li>• Butterworth high-pass filtered (40 Hz cut-off)</li> </ul> (μ V)	14.1 (12.6)	24.9 (30.1)	0.066	0.362
	<b>Method 4</b> <ul style="list-style-type: none"> <li>• DC offset removal</li> <li>• 20 - 450 Hz band pass filtered</li> <li>• Butterworth high-pass filtered (40 Hz cut-off)</li> <li>• Normalised</li> </ul>	57.4 (16.7)	78.9 (6.6)	<b>0.002</b>	0.710

	(% Maximum Value)				
--	-------------------	--	--	--	--

258

259 Figure 3. Mean (bold line) and standard deviation (shaded area) sEMG ARV from **Horse 2**,  
 260 obtained from right Biceps Femoris during left lead (red signals) and right lead (blue signals)  
 261 canter when the right hindlimb functions as TrH and LdH, respectively. Data are normalised  
 262 over one canter stride, with the green vertical line on the x-axis representing the hoof-lift off  
 263 event. sEMG signals are smoothed using an RMS filter (window length: 0.125 s, window  
 264 overlap: 0.121 s). The different signal processing methods are represented by a) Method 1, b)  
 265 Method 3, c) Method 2, d) Method 4.

266 Figure 4. Mean (bold line) and standard deviation (shaded area) sEMG ARV data from **Horse**  
 267 **4**, obtained from right Biceps Femoris during left lead (red signals) and right lead (blue signals)  
 268 canter when the right hindlimb functions as TrH and LdH, respectively. Data are normalised  
 269 over one canter stride, with the green vertical line on the x-axis representing the hoof-lift off  
 270 event. sEMG signals are smoothed using an RMS filter (window length: 0.125 s, window  
 271 overlap: 0.121 s). The different signal processing methods are represented by a) Method 1, b)  
 272 Method 3, c) Method 2, d) Method 4.

273 Figure 5 provides an individual example of how the application of a Butterworth high-pass  
 274 filter with a 40 Hz cut-off frequency, as applied in M3 and M4, can influence both ARV and  
 275 iEMG outcome measures and the interpretation of between-limb differences for LdH and TrH.  
 276 A comparison of band-pass filtered sEMG signals from TrH and LdH in Figure 5 (a, b) and  
 277 their corresponding full-wave rectified signals in Figure 5 (c, d) with high-pass filtered signals  
 278 in Figure 5 (g – j) illustrates how additional high-pass filtering alters the amplitude of sEMG  
 279 activation by removing low-frequency artefacts. The influence of high-pass filtering on  
 280 outcome measures are evidenced in Figure 5 (c, d), where failure to apply high-pass filtering  
 281 results in TrH exhibiting greater ARV and iEMG values than LdH. In contrast, the application  
 282 of high-pass filtering in Figure 5 (e, f) results in LdH showing greater ARV and iEMG values,  
 283 which is in accordance with overall results from this study (Table 2 and 3) and previous  
 284 biomechanical literature describing functional differences between LdH and TrH during canter.  
 285 Thus, failure to high-pass filter sEMG signals can lead to erroneous interpretation of results.  
 286 In accordance with Tables 2 and 3 and Figures 3 and 4, Figure 5 (k, l) also provides a visual  
 287 representation of how M4's combination of additional high-pass filtering, to attenuate low-  
 288 frequency noise sources, and normalisation, to reduce intra subject variability, results in the  
 289 greatest difference between LdH and TrH for sEMG outcome measures.

290 Figure 5. sEMG data obtained from right Biceps Femoris of **Horse 2** over one left lead (red  
 291 signals) and one right lead (blue signals) canter stride when the right hindlimb functions as TrH  
 292 and LdH, respectively. Signal processing steps for Methods 1 – 4 are illustrated as follows: a,  
 293 b) band-pass filtered signals (20 – 450 Hz), c, d) full-wave rectification of band-pass filtered  
 294 signals in a. and b. (Method 1), e, f) normalisation of band-pass filtered and full-wave rectified  
 295 signals in c. and d. using maximum observed value (Method 3), g, h) band-pass filtered (20 –  
 296 450 Hz) and high-pass filtered (40 Hz cut-off frequency) signals, i, j) full-wave rectification of  
 297 band-pass and high-pass filtered signals in g. and h. (Method 2), k, l) normalisation of band-  
 298 pass filtered, high-pass filtered and full-wave rectified signals in i. and j. using maximum  
 299 observed value (Method 4). iEMG and ARV data are provided for corresponding signals. The  
 300 hoof-lift off event is represented by the green tick on the x-axis.

#### 301 4. Discussion

302 In this study, sEMG data were obtained from the right BF during canter, to compare the  
303 sensitivity of four different SP methods for identifying differences in muscle activity that  
304 results from known differences in limb loading between LdH and TrH. Although it is known  
305 from the literature that the LdH experiences the greatest peak vertical loading of approximately  
306 1.2 times the horse's body weight, and the TrH experiences the smallest with peak vertical  
307 loading approximately equal to the horse's body weight (Merkens *et al.*, 1993), this study  
308 provides further kinematic evidence for these functional differences within our data set. In this  
309 study, significantly greater stifle joint flexion was accompanied by significantly greater peak  
310 flexion velocity for LdH during stance, which is indicative of an increased rate of stifle joint  
311 loading than that observed in TrH. Although significantly greater hip joint extension was found  
312 in the LdH, this coincided with significantly lower peak hip joint extension velocity than TrH,  
313 indicating that TrH experiences a greater rate of hip joint loading than LdH. Previous equine  
314 EMG studies report BF activity from late swing phase to late stance phase during trot and  
315 canter and postulate that the BF functions eccentrically during stance phase to stabilise the hip  
316 and stifle joints during limb loading (Crook *et al.*, 2010; Robert *et al.*, 1999; Tokuriki and Aoki,  
317 1995). Based on our findings from kinematic data, it is therefore argued that the BF generates  
318 eccentric muscle activity with a greater force in the LdH to stabilise the hip joint and prevent  
319 involuntary flexion of the stifle joint (Denoix, 2014; Robert *et al.*, 1999), which experiences a  
320 greater joint loading rate than TrH, during increased vertical limb loading (Merkens *et al.*,  
321 1993). Significantly higher BF muscle activity was observed in LdH than TrH when M2 and  
322 M4 were applied, which agrees with reported functional differences in LdH and TrH from the  
323 literature (Back *et al.*, 1997; Merckens *et al.*, 1993) and from kinematic data presented in this  
324 study. Thus, the significant increase in BF activity in the LdH, observed when M2 and M4 are  
325 applied, provides an accurate representation of BF activity during canter.

326 It is important to note that the SP methods employed in this study were not chosen arbitrarily.  
327 M1 was based on a review of existing equine sEMG literature and represents the most  
328 commonly employed sEMG SP method within this field. M4 was based on a combination of  
329 best practice for human sEMG SP, where the importance of low-frequency noise attenuation  
330 and normalisation techniques are well established (Burden, 2010; De Luca *et al.*, 2010; Lehman  
331 and McGill, 1999) and recent, equine-specific recommendations for high-pass filtering (St.  
332 George *et al.*, 2018). M2 and M3 provide intermediary SP methods, which were used to identify  
333 the individual contributions of normalisation and high-pass filtering for identifying differences  
334 in BF muscle activity between LdH and TrH. Following the application of all SP methods,  
335 amplitude-based outcome measures were calculated and compared, revealing significant  
336 differences in muscle activity between LdH and TrH when M2 and M4 were applied, but that  
337 M1 and M3 did not provide a sensitive enough metric to detect significant differences. Thus,  
338 the hypothesis that following recommended guidelines for sEMG SP, which includes  
339 normalisation and high-pass filtering, enables the identification of functional differences in  
340 muscle activation that would otherwise be missed was accepted. Of all methods, M4 resulted  
341 in the greatest between-limb differences in muscle activity, as evidenced by the lowest p values  
342 and highest effect sizes for iEMG and ARV. Thus, SP techniques used for M4 may serve as a  
343 basis for developing standardisation for equine sEMG SP. However, when considering why  
344 M4 produces outcome measures that best reflect biomechanical differences between hindlimbs  
345 at canter, as well as the highest magnitude of between-limb differences, it is important to

346 discuss the relative contributions of combining the 40 Hz high-pass filtering with  
347 normalisation.

#### 348 *4.1 Effect of high-pass filtering on outcome measures*

349 In human sEMG literature, movement artefact and baseline noise sources are known to  
350 contaminate the sEMG frequency spectra between 0 and 20 Hz (Clancy *et al.*, 2002; De Luca  
351 *et al.*, 2010; Van Boxtel, 2001). Such artefacts influence the shape of the sEMG frequency  
352 spectra and can dominate the total signal power, leading to erroneous interpretation of both  
353 spectral and amplitude-based sEMG signal outcome measures (De Luca *et al.*, 2010; Van  
354 Boxtel, 2001). Thus, attenuation of low-frequency noise in human studies is achieved using  
355 appropriate high-pass filtering techniques, where a cut-off frequency  $\geq 20$  Hz is recommended  
356 for maximally attenuating artefacts whilst minimising the removal of true sEMG signal content  
357 (De Luca *et al.*, 2010; Van Boxtel, 2001; Van Boxtel *et al.*, 1998). A more recent study carried  
358 out a similar approach among horses, where a high-pass filter cut-off frequency of 30 to 40 Hz  
359 was recommended for sEMG signals obtained from the BF of equine subjects during canter  
360 (St. George *et al.*, 2018). This recommended high pass filter was therefore employed for M3  
361 and M4 in the current study.

362 The beneficial effects of low-frequency noise attenuation on decreased intrasubject variability  
363 and increased between-limb differences in muscle activity when M3 and M4 are applied are  
364 illustrated in Figures 3 – 4. Furthermore, evidence for potential misinterpretation of muscle  
365 activity when high-pass filtering is omitted from SP is presented in Figure 5 (c, d, e, f), where  
366 M1 and M2 produce greater iEMG and ARV for TrH than LdH. M3, which employed high-  
367 pass filtering without normalisation, did not produce statistically significant differences in  
368 muscle activity between LdH and TrH. However, in comparison to M1, which did not employ  
369 high-pass filtering, M3 produced greater between-limb differences with lower p values, lower  
370 standard deviation and higher effects sizes for iEMG and ARV. Statistical power depends on  
371 both sample size and effect size, thus with a higher effect size it is possible to detect significant  
372 differences with a smaller sample size (Sullivan and Feinn, 2012). In equine sEMG research,  
373 the ability to employ a smaller sample size is advantageous due to the challenges associated  
374 with data acquisition, for example behavioural constraints and the time-consuming skin  
375 preparation process. Therefore, although the addition of high-pass filtering in M3 did not detect  
376 statistically significant between-limb differences in BF activity, attenuating low-frequency  
377 noise sources improves the sensitivity of SP methods by decreasing inter and intrasubject  
378 variability and increasing the magnitude of between-limb differences in muscle activity, which  
379 can lead to decreased sample size requirements for equine sEMG studies.

#### 380 *4.2 Effect of normalisation on outcome measures*

381 Normalisation converts the amplitude of an sEMG signal to a scaled value, generally the  
382 percentage of a MVC or RVC from a specific task (Burden, 2010; Lehman and McGill, 1999).  
383 This technique is fundamental for comparisons of amplitude-related sEMG outcome measures  
384 across subjects, muscles and trials/ days (Burden, 2010; Halaki and Ginn, 2012; Lehman and  
385 McGill, 1999; Mathiassen *et al.*, 1995) due to sources of variability associated with relative  
386 differences in sensor location, among other factors (De Luca, 1997). However, this is the first  
387 known study to demonstrate the effect of normalisation on sensitivity for identifying  
388 differences in muscle activity in relation to biomechanical differences in equine gait. The effect  
389 of normalisation on outcome measures in this study are clearly illustrated in Tables 2 and 3,

390 where application of M2 and M4 resulted in significantly greater BF activity in LdH compared  
391 to TrH. Standard deviation values in Tables 2 and 3 also show that the omission of  
392 normalisation in M1 and M3 resulted in increased variation in iEMG and ARV variables, which  
393 will have influenced the non-significant results in the statistical analysis. Findings from this  
394 study indicate that reduced standard deviation from normalisation represents the major  
395 contribution to significant statistical findings and is therefore recommended for equine sEMG  
396 SP. However, the contribution of high-pass filtering should not be overlooked, as it is the  
397 combination of high-pass filtering and normalisation in M4 that provided the most sensitive SP  
398 method for detecting differences in BF activity in relation to biomechanical differences  
399 between LdH and TrH during canter.

## 400 **5. Conclusion**

401 sEMG signals, obtained from BF during canter, exhibited significantly different amplitude-  
402 based outcome measures between LdH and TrH when normalisation and recommended band-  
403 pass filtering techniques for equine sEMG signals (St. George et al., 2018) were applied. The  
404 changes in muscle function that were observed were consistent with underlying biomechanical  
405 differences in hindlimb loading during canter. However, between limb differences were not  
406 observed when high-pass filtering and normalisation were omitted from SP. Therefore,  
407 functional between-limb differences may be missed depending on the SP procedures employed  
408 for equine gait analysis. More specifically, findings from this study illustrate the importance of  
409 including both appropriate band-pass filtering and normalisation techniques to facilitate  
410 accurate interpretation of the equine sEMG signal. It is our intent that these findings may  
411 accelerate further best practice guidelines and standardisation efforts within the equine sEMG  
412 field to facilitate knowledge transfer via consistent methodology.

## 413 **References**

- 414 Back, W., Schamhardt, H. and Barneveld, A., 1996. Are kinematics of the walk related to the  
415 locomotion of a warmblood horse at the trot? *Veterinary Quarterly* 18: 79-84.
- 416 Back, W., Schamhardt, H. and Barneveld, A., 1997. Kinematic comparison of the leading and  
417 trailing fore-and hindlimbs at the canter. *Equine Veterinary Journal* 29: 80-83.
- 418 Back, W., Schamhardt, H., Savelberg, H., Van Den Bogert, A., Bruin, G., Hartman, W. and  
419 Barneveld, A., 1995. How the horse moves: 2. Significance of graphical representations  
420 of equine hind limb kinematics. *Equine Veterinary Journal* 27: 39-45.
- 421 Burden, A., 2010. How should we normalize electromyograms obtained from healthy  
422 participants? What we have learned from over 25 years of research. *Journal of*  
423 *Electromyography and Kinesiology* 20: 1023-1035.
- 424 Burden, A. and Bartlett, R., 1999. Normalisation of EMG amplitude: an evaluation and  
425 comparison of old and new methods. *Medical Engineering & Physics* 21: 247-257.
- 426 Cheung, T., Warren, L., Lawrence, L. and Thompson, K., 1998. Electromyographic activity of  
427 the long digital extensor muscle in the exercising Thoroughbred horse. *Equine*  
428 *Veterinary Journal* 30: 251-255.
- 429 Clancy, E.A., Morin, E.L. and Merletti, R., 2002. Sampling, noise-reduction and amplitude  
430 estimation issues in surface electromyography. *Journal of Electromyography and*  
431 *Kinesiology* 12: 1-16.
- 432 Colborne, G., Birtles, D. and Cacchione, I., 2001. Electromyographic and kinematic indicators  
433 of fatigue in horses: a pilot study. *Equine Veterinary Journal* 33: 89-93.
- 434 Cram, J.R. and Rommen, D., 1989. Effects of skin preparation on data collected using an EMG  
435 muscle-scanning procedure. *Biofeedback and Self-regulation* 14: 75-82.

436 Crook, T., Wilson, A. and Hodson-Tole, E., 2010. The effect of treadmill speed and gradient  
437 on equine hindlimb muscle activity. *Equine Veterinary Journal* 42: 412-416.

438 Dankaerts, W., O'Sullivan, P.B., Burnett, A.F., Straker, L.M. and Danneels, L.A., 2004.  
439 Reliability of EMG measurements for trunk muscles during maximal and sub-maximal  
440 voluntary isometric contractions in healthy controls and CLBP patients. *Journal of*  
441 *Electromyography and Kinesiology* 14: 333-342.

442 De Luca, C.J., 1997. The use of surface electromyography in biomechanics. *Journal of Applied*  
443 *Biomechanics* 13: 135-163.

444 De Luca, C.J., Gilmore, L.D., Kuznetsov, M. and Roy, S.H., 2010. Filtering the surface EMG  
445 signal: Movement artifact and baseline noise contamination. *Journal of Biomechanics*  
446 43: 1573-1579.

447 Denoix, J.-M., 2014. *Biomechanics and physical training of the horse*. CRC Press.

448 Halaki, M. and Ginn, K., 2012. Normalization of EMG signals: to normalize or not to normalize  
449 and what to normalize to?, *Computational intelligence in electromyography analysis-a*  
450 *perspective on current applications and future challenges*. InTech.

451 Harrison, S.M., Whitton, R.C., King, M., Haussler, K.K., Kawcak, C.E., Stover, S.M. and  
452 Pandy, M.G., 2012. Forelimb muscle activity during equine locomotion. *Journal of*  
453 *Experimental Biology* 215: 2980-2991.

454 Hermens, H.J., Freriks, B., Disselhorst-Klug, C. and Rau, G., 2000. Development of  
455 recommendations for SEMG sensors and sensor placement procedures. *Journal of*  
456 *Electromyography and Kinesiology* 10: 361-374.

457 Hjerten, G., Drevemo, S. and Eriksson, L.E., 1994. Shortening of the hind limb in the horse  
458 during the stance phase. *Equine Veterinary Journal* 26: 48-50.

459 Hodson-Tole, E., 2006. Effects of treadmill inclination and speed on forelimb muscle activity  
460 and kinematics in the horse. *Equine and Comparative Exercise Physiology* 3: 61-72.

461 Hodson, E., Clayton, H. and Lanovaz, J., 2001. The hindlimb in walking horses: 1. Kinematics  
462 and ground reaction forces. *Equine Veterinary Journal* 33: 38-43.

463 Jansen, M., Schamhardt, H. and Hartman, W., 1992. Quantitative analysis of computer-  
464 averaged electromyographic profiles of intrinsic limb muscles in ponies at the walk.  
465 *American Journal of Veterinary Research* 53: 2343-2349.

466 Kuiken, T.A., Lowery, M. and Stoykov, N., 2003. The effect of subcutaneous fat on  
467 myoelectric signal amplitude and cross-talk. *Prosthetics and Orthotics International* 27:  
468 48-54.

469 Lehman, G.J. and McGill, S.M., 1999. The importance of normalization in the interpretation  
470 of surface electromyography: a proof of principle. *Journal of Manipulative and*  
471 *Physiological Therapeutics* 22: 444-446.

472 Mathiassen, S., Winkel, J. and Hägg, G., 1995. Normalization of surface EMG amplitude from  
473 the upper trapezius muscle in ergonomic studies—a review. *Journal of*  
474 *Electromyography and Kinesiology* 5: 197-226.

475 Merkens, H.W., Schamhardt, H.C. and Hartman, W., 1993. Ground reaction force patterns of  
476 Dutch Warmbloods at the canter. *American journal of veterinary research* 54: 670-674.

477 Merletti, R. and Di Torino, P., 1999. Standards for reporting EMG data. *Journal of*  
478 *Electromyography and Kinesiology* 9: 3-4.

479 Nordander, C., Willner, J., Hansson, G.-Å., Larsson, B., Unge, J., Granquist, L. and Skerfving,  
480 S., 2003. Influence of the subcutaneous fat layer, as measured by ultrasound, skinfold  
481 calipers and BMI, on the EMG amplitude. *European Journal of Applied Physiology* 89:  
482 514-519.

483 Payne, R., Hutchinson, J., Robilliard, J., Smith, N. and Wilson, A., 2005. Functional  
484 specialisation of pelvic limb anatomy in horses (*Equus caballus*). *Journal of Anatomy*  
485 206: 557-574.

- 486 Robert, C., Audigié, F., Valette, J., Pourcelot, P. and Denoix, J.M., 2001a. Effects of treadmill  
487 speed on the mechanics of the back in the trotting saddlehorse. *Equine Veterinary*  
488 *Journal* 33: 154-159.
- 489 Robert, C., Valette, J., Degueurce, C. and Denoix, J., 1999. Correlation between surface  
490 electromyography and kinematics of the hindlimb of horses at trot on a treadmill. *Cells*  
491 *Tissues Organs* 165: 113-122.
- 492 Robert, C., Valette, J. and Denoix, J., 2000. The effects of treadmill inclination and speed on  
493 the activity of two hindlimb muscles in the trotting horse. *Equine Veterinary Journal*  
494 32: 312-317.
- 495 Robert, C., Valette, J. and Denoix, J.M., 2001b. The effects of treadmill inclination and speed  
496 on the activity of three trunk muscles in the trotting horse. *Equine Veterinary Journal*  
497 33: 466-472.
- 498 Robert, C., Valette, J.P., Pourcelot, P., Audigie, F. and Denoix, J.M., 2002. Effects of trotting  
499 speed on muscle activity and kinematics in saddlehorses. *Equine Veterinary Journal* 34:  
500 295-301.
- 501 Roy, S.H., De Luca, G., Cheng, M.S., Johansson, A., Gilmore, L.D. and De Luca, C.J., 2007.  
502 Electro-mechanical stability of surface EMG sensors. *Medical & Biological*  
503 *Engineering & Computing* 45: 447-457.
- 504 Schuurman, S.O., Kersten, W. and Weijs, W.A., 2003. The equine hind limb is actively  
505 stabilized during standing. *Journal of Anatomy* 202: 355-362.
- 506 Sousa, A.S. and Tavares, J.M.R., 2012. Surface electromyographic amplitude normalization  
507 methods: a review. *Electromyography: new developments, procedures and*  
508 *applications*.
- 509 St. George, L., Hobbs, S.J., Richards, J., Sinclair, J., Holt, D. and Roy, S., 2018. The effect of  
510 cut-off frequency when high-pass filtering equine sEMG signals during locomotion.  
511 *Journal of Electromyography and Kinesiology* 43: 28-40.
- 512 Sullivan, G.M. and Feinn, R., 2012. Using effect size—or why the P value is not enough.  
513 *Journal of Graduate Medical Education* 4: 279-282.
- 514 Tokuriki, M. and Aoki, O., 1995. Electromyographic activity of the hindlimb muscles during  
515 the walk, trot and canter. *Equine Veterinary Journal* 27: 152-155.
- 516 Valentin, S. and Zsoldos, R.R., 2016. Surface electromyography in animal biomechanics: a  
517 systematic review. *Journal of Electromyography and Kinesiology* 28: 167-183.
- 518 Van Boxtel, A., 2001. Optimal signal bandwidth for the recording of surface EMG activity of  
519 facial, jaw, oral, and neck muscles. *Psychophysiology* 38: 22-34.
- 520 Van Boxtel, A., Boelhouwer, A. and Bos, A., 1998. Optimal EMG signal bandwidth and  
521 interelectrode distance for the recording of acoustic, electrocutaneous, and photic blink  
522 reflexes. *Psychophysiology* 35: 690-697.
- 523 Williams, J., Gundry, P., Richards, J. and Protheroe, L., 2013. A preliminary evaluation of  
524 surface electromyography as a tool to measure muscle fatigue in the National Hunt  
525 racehorse. *The Veterinary Nurse* 4: 566-572.
- 526 Winter, D., Rau, G., Kadefors, R., Broman, H. and De Luca, C., 1980. Units, Terms and  
527 Standards in the Reporting of EMG Research. A Report by the Ad Hoc Committee of  
528 the International Society of Electrophysiology and Kinesiology, Department of Medical  
529 Research. Rehabilitation Institute of Montreal, Montreal.
- 530 Yang, J.F. and Winter, D., 1984. Electromyographic amplitude normalization methods:  
531 improving their sensitivity as diagnostic tools in gait analysis. *Archives of Physical*  
532 *Medicine and Rehabilitation* 65: 517-521.
- 533 Zaneb, H., Kaufmann, V., Stanek, C., Peham, C. and Licka, T.F., 2009. Quantitative  
534 differences in activities of back and pelvic limb muscles during walking and trotting

535 between chronically lame and nonlame horses. American Journal of Veterinary  
536 Research 70: 1129-1134.

537 Zsoldos, R., Kotschwar, A., Kotschwar, A., Groesel, M., Licka, T. and Peham, C., 2010a.  
538 Electromyography activity of the equine splenius muscle and neck kinematics during  
539 walk and trot on the treadmill. Equine Veterinary Journal 42: 455-461.

540 Zsoldos, R., Kotschwar, A., Kotschwar, A., Rodriguez, C., Peham, C. and Licka, T., 2010b.  
541 Activity of the equine rectus abdominis and oblique external abdominal muscles  
542 measured by surface EMG during walk and trot on the treadmill. Equine Veterinary  
543 Journal 42: 523-529.

544

545