

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Dittrich, Sabine; Rudgard, William E; Woods, Kate L; Silisouk, Joy; Phuklia, Weerawat; Davong, Viengmon; Vongsouvath, Manivanh; Phommasone, Koukeo; Rattanaovong, Sayaphet; Knappik, Michael; +5 more... Craig, Scott B; Weier, Steven L; Tulsiani, Suhella M; Dance, David AB; Newton, Paul N; (2016) The Utility of Blood Culture Fluid for the Molecular Diagnosis of *Leptospira*: A Prospective Evaluation. AMERICAN JOURNAL OF TROPICAL MEDICINE AND HYGIENE, 94 (4). pp. 736-740. ISSN 0002-9637 DOI: <https://doi.org/10.4269/ajtmh.15-0674>

Downloaded from: <http://researchonline.lshtm.ac.uk/4652776/>

DOI: <https://doi.org/10.4269/ajtmh.15-0674>

Usage Guidelines:

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

The Utility of Blood Culture Fluid for the Molecular Diagnosis of *Leptospira*: A Prospective Evaluation

Sabine Dittrich,* William E. Rudgard, Kate L. Woods, Joy Silisouk, Weerawat Phuklia, Viengmon Davong, Manivanh Vongsouvath, Koukeo Phommason, Sayaphet Rattanavong, Michael Knappik, Scott B. Craig, Steven L. Weier, Suhella M. Tulsiani, David A. B. Dance, and Paul N. Newton

Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR; Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine Research Building, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom; Public Health England, Colindale, London, United Kingdom; Faculty of Science Health, Education and Engineering, University of the Sunshine Coast, Sippy Downs, Queensland, Australia; Queensland Health Forensic and Scientific Services, WHO Collaborating Centre for Reference and Research on Leptospirosis, Brisbane, Queensland, Australia; Faculty of Health, Queensland University of Technology, Brisbane, Queensland, Australia; Copenhagen Centre for Disaster Research, Global Health Section, Department of Public Health, University of Copenhagen, Copenhagen, Denmark

Abstract. Leptospirosis is an important zoonosis worldwide, with infections occurring after exposure to contaminated water. Despite being a global problem, laboratory diagnosis remains difficult with culture results taking up to 3 months, serology being retrospective by nature, and polymerase chain reaction showing limited sensitivity. *Leptospira* have been shown to survive and multiply in blood culture media, and we hypothesized that extracting DNA from incubated blood culture fluid (BCF), followed by quantitative real-time polymerase chain reaction (qPCR) could improve the accuracy and speed of leptospira diagnosis. We assessed this retrospectively, using preincubated BCF of *Leptospira* spp. positive ($N = 109$) and negative ($N = 63$) febrile patients in Vientiane, Lao PDR. The final method showed promising sensitivities of 66% (95% confidence interval [CI]: 55–76) and 59% (95% CI: 49–68) compared with direct or direct and indirect testing combined, as the respective reference standards (specificities > 95%). Despite these promising diagnostic parameters, a subsequent prospective evaluation in a Lao hospital population ($N = 352$) showed that the sensitivity was very low (~30%) compared with qPCR on venous blood samples. The disappointingly low sensitivity does suggest that venous blood samples are preferable for the clinical microbiology laboratory, although BCF might be an alternative if leptospirosis is only suspected postadmission after antibiotics have been used.

INTRODUCTION

Leptospirosis is a worldwide zoonotic disease, particularly common in rural tropical regions associated with contaminated water and soil exposure during heavy rainfall and occupational risks.¹ Poor living conditions in urban slums also contribute to an apparently increasing numbers of cases.² Leptospirosis infections are difficult to differentiate clinically from many other important causes of fever in the tropics, for example, rickettsial infections, typhoid, dengue, and malaria. In the United Kingdom and other temperate zones, cases may be indigenously associated with occupational or recreational risk factors or acquired overseas, predominantly in Southeast Asia or Central America.³ Leptospirosis usually presents as a nonspecific febrile illness, but may progress to serious renal, pulmonary, and central nervous system complications, with a case fatality rate of up to 40%.^{4,5} A lack of accurate diagnostic tools makes individual patient diagnosis to guide treatment very difficult.^{6–9} Serological methods are of limited use at admission, particularly in populations with regular exposure to the pathogen in whom only a 4-fold rise between admission and convalescent samples indicates recent infection. The organism is slow-growing and fastidious, requiring specialized prolonged culture methods and media¹⁰; hence for timely diagnosis, molecular tools have emerged as the methodologies of choice.¹¹ Several molecular methods have been developed mostly using venous (ethylenediaminetetraacetic acid [EDTA]) blood.^{9,11,12} We recently demonstrated the utility of molecular diagnosis of *Orientia*

tsutsugamushi, the agent of scrub typhus, from blood culture fluid (BCF) after 24-hour incubation (S. Dittrich, submitted). It has also been shown that *Leptospira* spp. survive in conventional blood culture media for up to 14 days but were not detected by conventional alert systems.^{13,14} Villumsen and others¹⁴ explored the use of BCF in place of venous blood as a sample for early leptospira diagnosis even days after antibiotic therapy initiation. Such techniques, especially if there was bacterial multiplication in blood cultures before quantitative real-time polymerase chain reaction (qPCR), could increase the sensitivity of assays and give additional sample matrices to work with. We, therefore, performed an initial pilot proof-of-principle and then a prospective study to determine whether blood culture bottles incubated for 24 hours in the Microbiology Laboratory of Mahosot Hospital (Vientiane, Lao PDR) could be used for the early diagnosis of leptospira by qPCR.

MATERIALS AND METHODS

Study population. The methodology was optimized using *Leptospira* spp. qPCR positive⁹ and negative samples, collected as part of studies in different geographical regions of the Lao PDR (Laos): Vientiane ($N = 65$), Luang Namtha ($N = 51$), and Salavan ($N = 18$). Samples had tested positive by qPCR ($N = 18$), culture ($N = 73$), or microscopic agglutination tests (MATs; $N = 18$) as part of published investigations.^{5,15}

The subsequent prospective study was conducted between May and August 2014 using consecutive patient samples ($N = 363$), submitted to the Microbiology Laboratory of Mahosot Hospital (102.6119°E, 17.9604°N). The inclusion criteria were fever and/or headache, rash, eschar, myalgia, arthralgia, meningitis, encephalitis, respiratory symptoms, jaundice, and acute

* Address correspondence to Sabine Dittrich, Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR. E-mail: sabine-dittrich@gmx.de

renal failure. Samples were collected on the day of admission for molecular diagnostics (serum, EDTA buffy coat, EDTA whole blood), leptospiral culture (blood clot), serology (serum), and two aerobic blood culture bottles. The aerobic blood culture bottles (Pharmaceutical Factory No. 2, Vientiane, Laos), which contain tryptic hydrolysate of casein and soy peptone with sodium polyanethol sulfonate (SPS), were inoculated with blood (> 15 years: 5 mL; 1–15 years: 2 mL; < 1 year: 1 mL), vented, and incubated at 37°C for 7 days.¹⁶ Bottles were manually checked for turbidity and blind subcultures (blood, chocolate, and MacConkey agars) performed after 24 hours and 7 days of incubation.¹⁶

BCF was collected from all blood cultures after 24-hour incubation. Bottles were repeatedly inverted until blood and media were homogeneously mixed, and 0.5 mL BCF from each blood culture bottle was transferred into another tube (total volume 1–1.5 mL of BCF per patient). Samples were stored at 4°C for up to 7 days and at –20°C for long term.

All study patients provided written informed consent prior to sample collection, with parents or guardians of any child participant providing informed consent on their behalf. Ethical approval for all investigations was granted by Oxford Tropical Research Ethics Committee (University of Oxford, United Kingdom) and the Faculty of Medical Sciences Committee (University of Health Sciences, Lao PDR).

Sample processing. EDTA blood samples (buffy coat, whole blood) and serum were processed using the Qiagen Blood Mini Kit (Qiagen, Germany) as described.¹⁵ BCF (0.5 mL) were spun for 10 minutes at 15,900 × *g* to pellet free bacteria and cells, but ~200 µL of the supernatant were discarded before proceeding as described.¹⁴ BCF was processed using the benzyl alcohol-based method, to remove, combined with the guanidine hydrochloride lysis/Qiagen Blood Mini Kit extraction, and obtained a final elution of 50 µL.¹⁴

Molecular detection. *Leptospira* spp. detection was performed using a published qPCR targeting the *rrs* gene (16S rRNA).⁹ In each run, 5 µL of template DNA (BCF, buffy coat, whole blood or serum) were added, and when using BCF-DNA, 40 mg/mL BSA was included to overcome residual inhibition due to SPS.¹⁷ Each run contained a standard curve of genomic *Leptospira* DNA (1 genome equivalent (GE)/µL to 10³ GE/µL; clinical isolate from Laos; assumed genome size, 4.7 Mb). Nontemplate controls were added to each run and were always negative; that is, no amplification was detected. All runs were performed using a Rotorgene 6000 (Qiagen).

Analysis. Threshold analysis was performed using RotorGene™ 6000 software (Version 1.7; Qiagen, Hilden, Germany). Receiver operator characteristics (ROC) analysis and figures were prepared using GraphPad Prism (Version 6.0c; San Diego, CA). In accordance with published guidelines, qPCR results were only considered positive with a Cq-value of ≤ 40 unless otherwise stated.¹⁸ MATs were performed by the WHO/FAO/OIE (World Health Organization/Food and Agriculture Organization of the United Nations/World Organization for Animal Health) Collaborating Center for Reference and Research on Leptospirosis, Australia.¹⁵ A 4-fold increase in titer between admission and convalescent samples was considered as an evidence of acute infection, whereas a 2-fold increase/decrease and a titer ≥ 1:400 were considered as “evidence of probable or recent infection.” Diagnostic accuracy was calculated in comparison to culture results, culture + qPCR, and with culture + qPCR + 4-fold rise in MAT titer.

RESULTS

Pilot study. The retrospective study aimed to pilot the use of 24-hour incubated BCF samples for molecular diagnosis and develop appropriate local qPCR threshold values. In total, BCFs from 109 *Leptospira* spp. qPCR, culture, or MAT positive and 63 *Leptospira* spp. qPCR and MAT negative patients were used. Each qPCR run was analyzed using three fluorescent threshold values (0.05, 0.075, 0.1), to include or exclude low positive samples and optimize the assay sensitivity and specificity accordingly (Figure 1). Using ROC analysis, a threshold level of 0.1 gave the best balance between sensitivity and specificity with an estimated sensitivity of 65.9% (95% confidence interval [CI]: 55.3–75.6) and specificity of 95.2% (95% CI: 86.7–99.0) when compared with culture and qPCR as a combined gold standard (Table 1). Reducing the threshold and increasing the inclusion Cq would have led to a sensitivity of close to 100% (gold standard: culture) while the specificity would fall to ~70% (Figure 1), that is unacceptable for both clinical diagnostics and public health investigations.

The defined analysis criteria (Cq ≤ 40; fluorescent threshold: 0.1) were subsequently used in a prospective study to investigate the real-life diagnostic characteristics in a cohort of Lao hospital patients.




FIGURE 1. Sensitivity and specificity characteristics plotted for different threshold values. Real-time quantitative polymerase chain reaction (qPCR) assays were analyzed using three different fluorescent thresholds (0.05, 0.075, 0.1), which were set manually. Receiver operator characteristics (ROC) analysis was used to determine sensitivity and specificity values at different Cq-values. Using (A) culture and (B) both culture and qPCR (blood) as gold standards for the ROC analysis.

TABLE 1
Diagnostic characteristics obtained by ROC analysis for three different fluorescent cutoffs and different gold standard comparators

	Sensitivity (95% CI)	Specificity (95% CI)	Positive/negative*	AUC† (95% CI)
		Threshold: 0.05		
Culture	89.0% (79.5–95.2)	74.6% (62.1–84.7)	73/63	0.94
qPCR	84.2% (60.4–96.6)	25.4% (15.3–37.9)	19/63	0.57
Culture + qPCR	74.7% (64.5–83.3)	74.6% (62.1–84.7)	91/63	0.82
Culture + qPCR + MAT	66.1% (56.4–74.9)	74.6% (62.1–84.7)	109/63	0.76
		Threshold: 0.075		
Culture	87.7% (77.9–94.2)	87.3% (76.5–94.3)	73/63	0.95
qPCR	84.2% (60.4–96.6)	25.4% (15.3–37.9)	19/63	0.57
Culture + qPCR	72.5% (62.2–81.4)	87.3% (76.5–94.3)	91/63	0.85
Culture + qPCR + MAT	64.2% (54.5–73.2)	87.3% (76.5–94.3)	109/63	0.79
		Threshold: 0.1		
Culture	82.2% (71.5–90.2)	95.8% (89.0–99.6)	73/63	0.94
qPCR	89.5% (66.7–98.7)	25.4% (15.3–37.9)	19/63	0.60
Culture + qPCR	65.9% (55.3–75.6)	95.2% (86.7–99.0)	91/63	0.85
Culture + qPCR + MAT	58.7% (48.9–68.1)	95.2% (86.7–99.0)	109/63	0.79

AUC = area under the curve; CI = confidence interval; MAT = microscopical agglutination test; ROC = receiver operator characteristics; qPCR = quantitative real-time polymerase chain reaction.

*Number of positive/negative samples included in the ROC calculation.

†AUC quantifies the overall ability of the test to discriminate between those individuals with the disease and those without the disease (interpretation: 0.9–1 = excellent; 0.8–0.9 = good; 0.7–0.8 = fair; 0.6–0.7 = poor).¹⁹

Prospective evaluation. Over the study period, 363 consecutive patients meeting the inclusion criteria were recruited. The majority was male (217/363; 60%), and the median age was 40 years (range: 1–92) with a median of 6 days (0–90 days) of illness prior to admission. For 352 (97%) patients, BCF and venous blood sample were available and of those 10 (2.8%) patients were positive for *Leptospira* spp. by molecular assays on admission blood samples (buffy coat, serum, or whole blood). When BCF after 24-hour incubation was used as a template for the qPCR, 6/352 (1.7%) patients were positive by qPCR. Out of these six, only three (50%) patients' samples were also positive by qPCR on one or more blood compartments and/or urine (data not shown), whereas seven (70%) patients with positive blood samples (serum, buffy coat, whole blood) were negative by qPCR on BCF. The three patients who were only positive by qPCR on BCF were also positive for other organisms, either by PCR on EDTA blood (*Rickettsia* spp.)²⁰ or blood culture (coagulase-negative *Staphylococcus* and *Burkholderia cepacia*)¹⁶ and their sera were negative by MAT for *Leptospira* spp. infections. The final sensitivity of qPCR on BCF when calculated in comparison to venous blood (buffy coat, serum, whole blood) was 30% with a specificity of 99%. The positive predictive value (PPV) was only 43%, whereas the negative predictive value was very high (98%) (Table 1).

In addition to using the predefined criteria ($C_q \leq 40$, threshold: 0.1), including positives with higher C_q -values was

also explored (Table 2). Although the sensitivity doubled to 50% when including all positives with $C_q \leq 45$,²¹ the specificity dropped to 88%. In this prospective study this would have wrongly identified 41 patients as having leptospirosis, overestimating the leptospira frequency in the hospital population by approximately 4-fold (~3% versus 13%). Only one of those 41 false-positive patients (C_q -cutoff: ≤ 45) had serological evidence for a recent *Leptospira* spp. infection by MAT.

DISCUSSION

The need for improved molecular tools for *Leptospira* spp. diagnosis is reflected in the multitude of new molecular tests that have been recently developed.^{21–24} BCF as a sample for direct detection of leptospirosis has been proposed,¹⁴ but no detailed data on its utility in a clinical laboratory are available. Our study aimed to establish the optimal methodology and subsequently to evaluate the method in febrile inpatients. The initial pilot study showed promising sensitivities of 82%, 90%, 66%, and 59% compared with culture, PCR, culture + PCR, or culture + PCR + MAT, as the respective reference standards. These diagnostic characteristics compare very well with the published sensitivity of the qPCR when used with whole blood (~55%),²⁵ suggesting that BCF might represent a valid and possibly superior blood sample compared with uncultured EDTA blood. However, in the subsequent prospective evaluation in a Lao hospital population,

TABLE 2
Sensitivity and specificity of BCF compared with blood when different inclusion criteria are applied for the BCF qPCR

		Venous blood qPCR		Final diagnostic characteristics for using BCF			
		Positive	Negative	Sensitivity (95% CI)	Specificity (95% CI)	PPV* (%)	NPV* (%)
BCF				BCF: $C_q \leq 40$			
	Positive	3	4	30% (10.3–60.7)	99% (97–100)	43	98
	Negative	7	338				
				BCF: $C_q \leq 42$			
	Positive	3	18	30% (10.3–60.7)	95% (92–97)	14	98
	Negative	7	324				
			BCF: $C_q \leq 45$				
Positive	5	41	50% (23.7–76.3)	88% (84–91)	11	98	
Negative	5	301					

BCF = blood culture fluid; CI = confidence interval; NPV = negative predictive value; PPV = positive predictive value; qPCR = quantitative real-time polymerase chain reaction.

*PPV is the proportion of positive results that are truly positive, and NPV is the proportion of negative results that are truly negative.

the BCF results underestimated leptospiral incidence in comparison to qPCR of uncultured EDTA blood. Seventy percent of *Leptospira* spp. positive patients would have been missed when using only BCF as a sample. Further to highlighting issues with the use of BCF, our results underline the complexity of leptospira diagnosis with none of the qPCR positive patients showing evidence for leptospira infection by the serological gold standard.^{24,25}

The qPCR assay used in this study targets the 16S rRNA gene of *Leptospira* spp., and the ubiquity of this target can result in reduced specificity. A number of studies have described this issue^{9,23,26} and our results confirm those findings as all of the actual false-positive samples had other bacteria present in the blood or blood culture sample, even though two of these are likely to have been contaminants. The specificity and PPV dropped further when the Cq-cutoff was extended to 42 or 45, consistent with the Minimum Information for Publication of Quantitative Real-Time PCR Experiments (MIQE) guidelines that suggest that Cq-values > 40 are suspect and should not be considered positive.¹⁸ This trade-off with the sensitivity has to be accepted, as overestimating disease incidence and initiating the wrong treatment would have adverse consequences for patients.

Using incubated BCF samples seems advantageous for other pathogens (*O. tsutsugamushi*) (S. Dittrich, submitted), but this does not appear to be the case when using a 16S rRNA-based qPCR system for leptospirosis. Blood culture bottles containing glycerol or fatty acids were shown to be conducive to leptospiral growth¹³; hence, future investigations using supplemented blood culture media may be illuminating.

In conclusion, the study shows that, despite the initial promising results, BCF, when combined with a 16S rRNA-targeting assay, does not seem sufficiently sensitive and specific for routine molecular diagnosis of leptospirosis. The results underline the need for increased investment into improved diagnostic tools for this important global pathogen.

Received September 14, 2015. Accepted for publication December 28, 2015.

Published online February 15, 2016.

Acknowledgments: We thank Phouthasen Hyongsithy for technical assistance and we are very grateful to all the doctors and nursing staff of Mahosot Hospital and the staff of the Microbiology Laboratory. We are also very grateful to the Directors of Mahosot Hospital, the Minister of Health, and the Director of the Curative Department, Ministry of Health, for their support in this study.

Financial support: This study was funded by the Wellcome Trust of Great Britain.

Authors' addresses: Sabine Dittrich, Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Molecular Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR, and Center for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom, E-mail: sabine-dittrich@gmx.de. William E. Rudgard, Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Molecular Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR, E-mail: willrudgard@hotmail.co.uk. Kate L. Woods, Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Molecular Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR, and Public Health England, Colindale, London, United Kingdom, E-mail: katewoods79@gmail.com. Joy Silisouk, Weerawat Phuklia, Viengmon Davong, Manivanh Vongsouvath, and Koukeo Phommason, Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Molecular Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR, E-mails: joy@tropmedres.ac, weerawat@tropmedres.ac, viengmon@

tropmedres.ac, manivanh@tropmedres.ac, and koukeo@tropmedres.ac. Sayaphet Rattanavong, Lao-Oxford-Mahosot Hospital-Wellcome Trust Research, Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR, E-mail: sayaphet@tropmedres.ac. Michael Knappik, Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Molecular Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR, and Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine Research Building, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom, E-mail: mikeknappik@hotmail.com. Scott B. Craig, Queensland Health Forensic and Scientific Service, WHO Collaborating Centre for Reference and Research on Leptospirosis, Brisbane, Queensland, Australia, and Faculty of Science Health, Education and Engineering, University of the Sunshine Coast, Sippy Downs, Queensland, Australia, E-mail: scott.craig@health.qld.gov.au. Steven L. Weier, Faculty of Health, Queensland University of Technology, Brisbane, Queensland, Australia, E-mail: s.weier@qut.edu.au. Suhella M. Tulsiani, Department of International Health, Copenhagen Centre for Disaster Research, University of Copenhagen, Copenhagen, Denmark, and Queensland Health Forensic and Scientific Services, WHO Collaborating Centre for Reference and Research on Leptospirosis, Brisbane, Queensland, Australia, E-mail: suhella.tulsiani@gmail.com. David A. B. Dance, Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine Research Building, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom, and Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Molecular Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR, E-mail: david.d@tropmedres.ac. Paul N. Newton, Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine Research Building, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom, and Lao-Oxford-Mahosot Hospital-Wellcome Trust Research, Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao PDR, E-mail: paul@tropmedres.ac.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

REFERENCES

1. Pappas G, Papadimitriou P, Siozopoulou V, Christou L, Akritidis N, 2008. The globalization of leptospirosis: worldwide incidence trends. *Int J Infect Dis* 12: 351–357.
2. Lau CL, Smythe LD, Craig SB, Weinstein P, 2010. Climate change, flooding, urbanisation and leptospirosis: fuelling the fire? *Trans R Soc Trop Med Hyg* 104: 631–638.
3. Forbes AE, Zochowski WJ, Dubrey SW, Sivaprakasam V, 2012. Leptospirosis and Weil's disease in the UK. *QJM* 105: 1151–1162.
4. Taylor AJ, Paris DH, Newton PN, 2015. A systematic review of the mortality from untreated leptospirosis. *PLoS Negl Trop Dis* 9: e0003866.
5. Dittrich S, Rattanavong S, Lee SJ, Panyanivong P, Craig SB, Tulsiani SM, Blacksell SD, Dance DA, Dubot-Peres A, Sengduangphachanh A, Phoumin P, Paris DH, Newton PN, 2015. *Orientia*, *Rickettsia*, and *Leptospira* pathogens as causes of CNS infections in Laos: a prospective study. *Lancet Glob Health* 3: e104–e112.
6. Goris MG, Leeflang MM, Loden M, Wagenaar JF, Klatser PR, Hartskeerl RA, Boer KR, 2013. Prospective evaluation of three rapid diagnostic tests for diagnosis of human leptospirosis. *PLoS Negl Trop Dis* 7: e2290.
7. Levett PN, 2001. Leptospirosis. *Clin Microbiol Rev* 14: 296–326.
8. Picardeau M, Bertherat E, Jancloes M, Skouloudis AN, Durski K, Hartskeerl RA, 2014. Rapid tests for diagnosis of leptospirosis: current tools and emerging technologies. *Diagn Microbiol Infect Dis* 78: 1–8.
9. Thaipadunpanit J, Chierakul W, Wuthiekanun V, Limmathurotsakul D, Amornchai P, Boonslip S, Smythe LD, Limpaboon R, Hoffmaster AR, Day NP, Peacock SJ, 2011. Diagnostic accuracy of real-time PCR assays targeting 16S rRNA and lipL32 genes for human leptospirosis in Thailand: a case-control study. *PLoS One* 6: e16236.
10. Wuthiekanun V, Chierakul W, Limmathurotsakul D, Smythe LD, Symonds ML, Dohnt MF, Slack AT, Limpaboon R,

- Suputtamongkol Y, White NJ, Day NP, Peacock SJ, 2007. Optimization of culture of *Leptospira* from humans with leptospirosis. *J Clin Microbiol* 45: 1363–1365.
11. Agampodi SB, Matthias MA, Moreno AC, Vinetz JM, 2012. Utility of quantitative polymerase chain reaction in leptospirosis diagnosis: association of level of leptospiremia and clinical manifestations in Sri Lanka. *Clin Infect Dis* 54: 1249–1255.
 12. Boonsilp S, Thaipadungpanit J, Amornchai P, Wuthiekanun V, Chierakul W, Limmathurotsakul D, Day NP, Peacock SJ, 2011. Molecular detection and speciation of pathogenic *Leptospira* spp. in blood from patients with culture-negative leptospirosis. *BMC Infect Dis* 11: 338.
 13. Griffith ME, Horvath LL, Mika WV, Hawley JS, Moon JE, Hospenhal DR, Murray CK, 2006. Viability of *Leptospira* in BacT/ALERT MB media. *Diagn Microbiol Infect Dis* 54: 263–266.
 14. Villumsen S, Pedersen R, Krogfelt KA, Jensen JS, 2010. Expanding the diagnostic use of PCR in leptospirosis: improved method for DNA extraction from blood cultures. *PLoS One* 5: e12095.
 15. Mayxay M, Castonguay-Vanier J, Chansamouth V, Dubot-Pérès A, Paris DH, Phetsouvanh R, Tangkhabuanbutra J, Douangdala P, Inthalath S, Souvannasing P, Slesak G, Tongyoo N, Chanthongthip A, Panyanouvong P, Sibounheuang B, Phommason K, Dohnt M, Phonekeo D, Hongvanthong B, Xayadeth S, Ketmayoon P, Blacksell SD, Moore CE, Craig SB, Burns M-A, von Sonnenburg F, Corwin A, de Lamballerie X, González IJ, Christophel EM, Cawthorne A, Bell D, Newton PN, 2013. Causes of non-malarial fever in Laos: a prospective study. *Lancet Glob Health* 1: e46–e54.
 16. Phetsouvanh R, Phongmany S, Soukaloun D, Rasachak B, Soukhaseum V, Soukhaseum S, Frichithavong K, Khounnorath S, Pengdee B, Phiasakha K, Chu V, Luangxay K, Rattanavong S, Sisouk K, Keoulouangkot V, Mayxay M, Ramsay A, Blacksell SD, Campbell J, Martinez-Aussel B, Heuanvongsy M, Bounxouei B, Thamavong C, Syhavong B, Strobel M, Peacock SJ, White NJ, Newton PN, 2006. Causes of community-acquired bacteremia and patterns of antimicrobial resistance in Vientiane, Laos. *Am J Trop Med Hyg* 75: 978–985.
 17. Nic Fhogartaigh C, Dance DA, Davong V, Tann P, Phetsouvanh R, Turner P, Dittrich S, Newton PN, 2015. A novel technique for detecting antibiotic-resistant typhoid from rapid diagnostic tests. *J Clin Microbiol* 53: 1758–1760.
 18. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT, 2009. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 55: 611–622.
 19. Greiner M, Pfeiffer D, Smith RD, 2000. Principles and practical application of the receiver-operating characteristic analysis for diagnostic tests. *Prev Vet Med* 45: 23–41.
 20. Jiang J, Chan TC, Temenak JJ, Dasch GA, Ching WM, Richards AL, 2004. Development of a quantitative real-time polymerase chain reaction assay specific for *Orientia tsutsugamushi*. *Am J Trop Med Hyg* 70: 351–356.
 21. Waggoner JJ, Balassiano I, Abeynayake J, Sahoo MK, Mohamed-Hadley A, Liu Y, Vital-Brazil JM, Pinsky BA, 2014. Sensitive real-time PCR detection of pathogenic *Leptospira* spp. and a comparison of nucleic acid amplification methods for the diagnosis of leptospirosis. *PLoS One* 9: e112356.
 22. Noda AA, Rodriguez I, Rodriguez Y, Govin A, Fernandez C, Obregon AM, 2014. High sensitive PCR method for detection of pathogenic *Leptospira* spp. in paraffin-embedded tissues. *Rev Inst Med Trop Sao Paulo* 56: 411–415.
 23. Villumsen S, Pedersen R, Borre MB, Ahrens P, Jensen JS, Krogfelt KA, 2012. Novel TaqMan(R) PCR for detection of *Leptospira* species in urine and blood: pit-falls of in silico validation. *J Microbiol Methods* 91: 184–190.
 24. Waggoner JJ, Balassiano I, Mohamed-Hadley A, Vital-Brazil JM, Sahoo MK, Pinsky BA, 2015. Reverse-transcriptase PCR detection of leptospira: absence of agreement with single-specimen microscopic agglutination testing. *PLoS One* 10: e0132988.
 25. Limmathurotsakul D, Turner EL, Wuthiekanun V, Thaipadungpanit J, Suputtamongkol Y, Chierakul W, Smythe LD, Day NP, Cooper B, Peacock SJ, 2012. Fool's gold: why imperfect reference tests are undermining the evaluation of novel diagnostics: a reevaluation of 5 diagnostic tests for leptospirosis. *Clin Infect Dis* 55: 322–331.
 26. Ganoza CA, Matthias MA, Saito M, Cespedes M, Gotuzzo E, Vinetz JM, 2010. Asymptomatic renal colonization of humans in the peruvian Amazon by *Leptospira*. *PLoS Negl Trop Dis* 4: e612.