

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Knappik, Michael; Dance, David AB; Rattanaovong, Sayaphet; Pierret, Alain; Ribolzi, Olivier; Davong, Viengmon; Silisouk, Joy; Vongsouvath, Manivanh; Newton, Paul N; Dittrich, Sabine; (2015) Evaluation of Molecular Methods To Improve the Detection of *Burkholderia pseudomallei* in Soil and Water Samples from Laos. *APPLIED AND ENVIRONMENTAL MICROBIOLOGY*, 81 (11). pp. 3722-3727. ISSN 0099-2240 DOI: <https://doi.org/10.1128/AEM.04204-14>

Downloaded from: <http://researchonline.lshtm.ac.uk/4652792/>

DOI: <https://doi.org/10.1128/AEM.04204-14>

Usage Guidelines:

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>

Evaluation of Molecular Methods To Improve the Detection of *Burkholderia pseudomallei* in Soil and Water Samples from Laos

Michael Knappik,^{a,b} David A. B. Dance,^{a,b} Sayaphet Rattanavong,^a Alain Pierret,^c Olivier Ribolzi,^d Viengmon Davong,^a Joy Silisouk,^a Manivanh Vongsouvath,^a Paul N. Newton,^{a,b} Sabine Dittrich^{a,b}

Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit, Microbiology Laboratory, Mahosot Hospital, Vientiane, Lao People's Democratic Republic^a; Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, England, United Kingdom^b; Institute of Ecology and Environmental Science—Paris, Institut de Recherche pour le Développement (IRD), Vientiane, Lao People's Democratic Republic^c; Géosciences Environnement Toulouse (GET), UMR 5563, IRD, Université de Toulouse, UPS (OMP), CNRS, Toulouse, France^d

Burkholderia pseudomallei is the cause of melioidosis, a severe and potentially fatal disease of humans and animals. It is endemic in northern Australia and Southeast Asia and is found in soil and surface water. The environmental distribution of *B. pseudomallei* worldwide and within countries where it is endemic, such as the Lao People's Democratic Republic (Laos), remains unclear. However, this knowledge is important to our understanding of the ecology and epidemiology of *B. pseudomallei* and to facilitate public health interventions. Sensitive and specific methods to detect *B. pseudomallei* in environmental samples are therefore needed. The aim of this study was to compare molecular and culture-based methods for the detection of *B. pseudomallei* in soil and surface water in order to identify the optimal approach for future environmental studies in Laos. Molecular detection by quantitative real-time PCR (qPCR) was attempted after DNA extraction directly from soil or water samples or after an overnight enrichment step. The positivity rates obtained by qPCR were compared to those obtained by different culture techniques. The rate of detection from soil samples by qPCR following culture enrichment was significantly higher (84/100) than that by individual culture methods and all culture methods combined (44/100; $P < 0.001$). Similarly, qPCR following enrichment was the most sensitive method for filtered river water compared with the sensitivity of the individual methods and all individual methods combined. In conclusion, molecular detection following an enrichment step has proven to be a sensitive and reliable approach for *B. pseudomallei* detection in Lao environmental samples and is recommended as the preferred method for future surveys.

The Gram-negative bacterial saprophyte *Burkholderia pseudomallei* is the causative agent of melioidosis and is found in soil and surface water predominantly in regions of Southeast Asia and northern Australia, where the organism is endemic (1–3). It is a common cause of fatal community-acquired bacteremia, pneumonia, and visceral and soft tissue abscesses and poses a significant public health burden (4, 5). Most patients are thought to contract the infection from skin inoculation; other possible routes of transmission are inhalation and ingestion (6). Due to the high mortality from infection with the organism and potential transmission of the organism by aerosols, *B. pseudomallei* is classified as a tier 1 select agent (7). Melioidosis was first reported in the Lao People's Democratic Republic (Laos) in 2001 with the description of two patients with the disease (8). In a randomized soil survey conducted in 2009, the highest isolation frequency was in Saravane Province (southern Laos) (9). Little is known about the true geographical distribution of melioidosis across the country, highlighting the need for a detailed risk map to support empirical patient management. The current “gold standard” for detection of environmental *B. pseudomallei* is culture from soil or water samples (10, 11). Culture methods for soil samples have successfully been employed in many studies, with proposed consensus guidelines being based on a simplified qualitative technique recently published (11). Similar guidelines do not yet exist for the detection of *B. pseudomallei* in water (3, 11, 12). Investigations of *B. pseudomallei* in Laos have highlighted the challenges of detecting the pathogen in different environmental samples (9, 12), hampering the development of a detailed risk map (9). Despite their extensive use, culture methods have their limitations, including the

potential overgrowth of *B. pseudomallei* by other environmental bacterial species, especially when it is present in small numbers. In addition, bacteria in a viable but noncultivable state are not detectable (13), further decreasing the sensitivity of this approach.

To overcome these limitations, molecular methods have successfully been developed and applied to detect *B. pseudomallei* in soil (14, 15). A study in Thailand demonstrated that molecular detection estimated the bacterial load per gram of soil to be 10 times higher than that determined by culture methods (15), and investigations from northern Australia showed that quantitative real-time PCR (qPCR) following an enrichment step yielded a higher positivity rate than culture alone (14). Furthermore, the high specificity of molecular assays allows the specific detection of

Received 24 December 2014 Accepted 16 March 2015

Accepted manuscript posted online 27 March 2015

Citation Knappik M, Dance DAB, Rattanavong S, Pierret A, Ribolzi O, Davong V, Silisouk J, Vongsouvath M, Newton PN, Dittrich S. 2015. Evaluation of molecular methods to improve the detection of *Burkholderia pseudomallei* in soil and water samples from Laos. *Appl Environ Microbiol* 81:3722–3727. doi:10.1128/AEM.04204-14.

Editor: D. W. Schaffner

Address correspondence to Sabine Dittrich, sabine@tropmedres.ac.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AEM.04204-14>.

Copyright © 2015, Knappik et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

doi:10.1128/AEM.04204-14

B. pseudomallei even if closely related bacteria, such as *Burkholderia thailandensis*, *Burkholderia vietnamiensis*, *Burkholderia cepacia*, and other *Burkholderia* and *Ralstonia* species, are present in the same environment (16–18). Although molecular methods have been used for the detection of *B. pseudomallei* from soil, no molecular methods for the detection of the pathogen directly from water samples have been described. As soil and water sampling is crucial for mapping disease risks in the country, updated techniques suitable for large-scale screening are urgently needed.

The aim of this study was to use described and novel molecular methods for the detection of *B. pseudomallei* in soil and water and evaluate those approaches in comparison to different culture techniques.

MATERIALS AND METHODS

Sample collection. (i) Soil. Soil sampling was performed during the dry season (April 2013) in a rice paddy near the village of Ban Nabone, Vientiane Province, Laos (18°22'51.4"N, 102°25'27.8"E; altitude, 195 m). Samples were collected from two depths (30 cm and 60 cm) at 50 random points within a section of the field previously determined to have the highest rates of positivity for *B. pseudomallei* by culture (19), with a minimum distance of 2 m being allowed between sampling sites. Oral informed consent for the removal of soil samples was obtained from the farmers concerned, and written permission was obtained from the relevant authorities. Samples were collected from the field using a hand auger that was disinfected between samplings with 70% alcohol (9). The samples were placed in sterile plastic bags, which were placed in an insulated box that was kept in the shade and maintained at ambient temperature during transport and subsequent manipulation. To ensure that representative subsamples were obtained, the two-dimensional Japanese slab cake method was used, and ~0.5-g (direct extraction from soil), ~10-g (simplified culture method [11]), ~20-g (extraction postenrichment [14]), and ~100-g (conventional culture method [20]) subsamples were obtained (Fig. 1) (21, 22).

Samples were collected within 24 h of each other, and subsampling was performed up to 72 h postsampling. Processing of all samples was started on the same day at 120 h postsampling.

(ii) Water. Water samples (600 ml; $n = 20$) were collected along the course of the Sedone River (Saravane Province, Laos) ($n = 7$) and its main tributaries ($n = 13$) at the onset of the rainy season (June 2013; mean turbidity, 257 nephelometric turbidity units; mean water temperature, 26.3°C; mean pH, 7.1; see Table S2 in the supplemental material). Collection bottles were triple rinsed with river water from the collection site before collecting the sample. The water was mixed thoroughly, and 30-ml subsamples were filtered within 8 h of collection, using 47-mm-diameter membranes. Six subsamples of water were filtered from each site; three were filtered using 0.2- μ m-pore-size filters (30 ml, cellulose acetate membrane filter; Sartorius), and three were filtered using 3.0- μ m-pore-size filters (cellulose nitrate membrane filter; Sartorius) in an attempt to estimate total and attached bacterial loads, respectively (Fig. 1) (23, 24). A manual pump and a 1-liter glass vacuum flask with a stainless steel funnel were used, and membrane supports were sterilized with 70% ethanol between filtrations. In addition, Moore's swab samples ($n = 10$) were collected as described previously (12).

Culture techniques. (i) Soil. Soil was cultured using both plating of soil suspension supernatants on Ashdown's agar as previously described (20) and the simplified broth enrichment culture method recommended in recent guidelines (11). The results of this comparison (unpublished data) will be reported elsewhere.

(ii) Water. Three methods were used concurrently to attempt to culture *B. pseudomallei* from water at each sampling site. First, Moore's swabs ($n = 10$) were used as described previously (12). Second, individual filters from each sampling point ($n = 40$; Fig. 1) were placed in 10 ml modified Ashdown's broth (14) and incubated aerobically without shaking for 7




FIG 1 Schematic representation of the methodologies and environmental samples used in the study. Soil samples were taken from two different depths (30 cm and 60 cm; ~0.5 g for DS DNA extraction, ~10 g for simplified culture, ~20 g for ES DNA extraction, ~100 g for conventional culture) at 50 positions in the rice field, and all samples were processed by direct or enrichment culture, direct DNA extraction, or DNA extraction postenrichment. Water samples were collected at 20 different sites, separate samples were filtered through filters of two different pore sizes (total $n = 40$), and all filters were processed by direct and enrichment culture, while filters were split for molecular methods (direct DNA extraction and DNA extraction postenrichment). In addition, 10 Moore's swabs were collected, and the water from these was cultured directly on solid medium (not shown), as previously described (12). The latitudes/longitudes of the sampling sites were as follows: 15°7'26.22"N/105°48'28.02"E, 15°21'38.81"N/105°49'52.85"E, 15°34'48.01"N/105°48'42.50"E, 15°40'4.62"N/105°54'40.13"E, 15°42'31.58"N/106°4'7.87"E, 15°47'23.56"N/106°17'26.87"E, 15°42'41.46"N/106°25'41.95"E, 15°13'16.88"N/105°44'34.96"E, 15°20'5.86"N/105°58'58.87"E, 15°15'30.33"N/105°55'59.81"E, 15°29'44.72"N/105°45'51.05"E, 15°41'44.84"N/106°16'1.23"E, 15°38'12.39"N/106°22'3.86"E, 15°39'16.84"N/105°50'54.55"E, 15°42'0.52"N/105°58'30.19"E, 15°42'26.27"N/106°8'38.07"E, 15°40'52.12"N/106°25'59.19"E, 15°32'24.51"N/106°15'54.90"E, 15°27'59.66"N/106°10'12.32"E, 15°24'33.56"N/106°5'27.98"E.

days at 40°C. At days 3 and 7, 10- μ l and 100- μ l samples of broth were subcultured onto Ashdown's agar containing gentamicin at 8 mg/liter and incubated at 40°C aerobically for 4 days with daily inspections for colonies resembling *B. pseudomallei* (12). Third, filters from each sampling point ($n = 40$; Fig. 1) were placed directly on Ashdown's agar plates and cultured at 40°C for up to 96 h. *B. pseudomallei* was identified and confirmed as described previously (12). Briefly, all positive cultures were screened by agglutination with a latex agglutination reagent specific for the 200-kDa extracellular polysaccharide of *B. pseudomallei* and tested for susceptibility to amoxicillin-clavulanic acid (co-amoxiclav) and resistance to colistin (8, 14). All suspected isolates were confirmed to be *B. pseudomallei* by qPCR (16), and selected isolates were confirmed to be *B. pseudomallei* by use of an API 20NE system.

Molecular detection. (i) Soil. Direct soil (DS) DNA extraction, in which DNA was extracted from ~0.5 g of soil, was attempted using a kit (PowerSoil DNA isolation kit; Mo Bio) as described previously (14). Enrichment soil (ES) DNA extraction was done as described previously (14) with minor modifications. In brief, soil was homogenized in the modified Ashdown's enrichment broth, shaken for 2 h at 240 rpm, and then incubated at 37°C for 22 h. The liquid phase (~10 ml) was decanted and centrifuged at 700 \times g for 2 min, and the supernatant (~7 ml) was removed and aurintricarboxylic acid (14) was added. After further centrifugation (45 min, 4,000 \times g), DNA was extracted from the soil pellet (~0.5 g) as described previously (14).

(ii) Water. Filters were cut diametrically in half; one half was used to extract DNA directly from the filter (direct filter [DF] extraction), while the other half was subjected to an enrichment step (enrichment filter [EF] extraction) (Fig. 1). For DF extraction, the filter membrane was cut into small pieces using sterile scissors. DNA was extracted from the filter membranes with the PowerSoil DNA isolation kit (Mo Bio), and the detachment of soil particles could be visually observed. After 60 μ l of C1 solution

TABLE 1 Isolates used to test the specificity of the molecular methods

Organism	Source	Location	Strain
International reference strains			
<i>B. cepacia</i>	Environmental	NA ^a	NCTC 10743
<i>B. thailandensis</i>	Environmental	Thailand	E264 (ATCC 700388)
<i>Ralstonia pickettii</i>	Human	Unknown	ATCC 27511
Local isolates			
<i>B. pseudomallei</i> (n = 9)	Human	Laos	Clinical isolate ^b
<i>B. pseudomallei</i> (n = 8)	Environment	Laos	Soil isolate ^b
<i>B. cepacia</i> (n = 9)	Human	Laos	Clinical isolate ^b
<i>B. cepacia</i> (n = 2)	Environmental	Laos	Soil isolate ^b
Local competing soil flora ^c			
<i>Ralstonia</i> spp. (n = 2)	Environmental	Vientiane, Laos	Soil isolate ^{b,d}
<i>Burkholderia</i> spp. (n = 12)	Environmental	Vientiane and Luang Namtha, Laos	Soil isolate ^{b,d}

^a NA, not available.

^b Identities were confirmed by determination of the colony morphology by use of the API 20NE system (bioMérieux, France) and molecular methods (36).

^c Isolates collected during previous environmental surveys.

^d Identities were confirmed by use of the API 20NE system (bioMérieux, France) and 16S rRNA sequencing (GenBank accession numbers KM058066 to KM058079).

was added, DNA extraction was performed according to the manufacturer's instructions (Mo Bio). The second half, used for EF extraction, was incubated in modified Ashdown's broth for 24 h at 37°C (14). Membranes were repeatedly vortexed to release the sediment and bacteria, and after incubation, the filters were removed and free sediment and bacteria were pelleted (45 min, 4,000 × g). The pellet (~0.2 g) was processed for DNA extraction as described previously (14).

qPCR. The PCR targets a 115-bp stretch in *orf2* of the type III secretion system gene cluster (TTS1) of *B. pseudomallei* and was performed, with minor modifications, as described previously (14, 16). In brief, the 25- μ l master mix consisted of 1 U Platinum *Taq* DNA polymerase (Invitrogen), 500 nM each primer, 250 nM probe, 7 mM MgCl₂, 1× PCR buffer, 200 μ M deoxynucleoside triphosphates, and 4 μ l of soil DNA. To reduce the effect of inhibitors, 400 ng/ μ l of bovine serum albumin (BSA; New England Biolabs, USA) was added (15). To control for PCR inhibitors, ~10⁵ copies of inhibitor control plasmid (47 kDa, *Orientia tsutsugamushi* [25, 26]) were amplified alone and in parallel spiked with 4 μ l of sample DNA. Inhibition was monitored as described previously (14). Amplification was performed on a Rotor-Gene 6000 system (Qiagen, Germany) at 94°C for 10 min, and then 45 cycles of 94°C for 15 s and 60°C for 1 min were performed. A standard curve was included in every run, using 1 genome equivalent (GE)/ μ l to 10³ GE/ μ l of *B. pseudomallei* (clinical isolate 1106a from Thailand; assumed genome size, 7.25 Mb). Nontemplate controls were added to each run and were always negative; i.e., no amplification was detected. Extraction controls (n = 10) using molecular-grade water (AccuGENE molecular biology water; Lonza) were used to rule out *B. pseudomallei* contamination of reagents and equipment (27). Samples with threshold cycle (*C_T*) values below 40 were considered positive (28). The limit of detection and the specificity of the assay under the conditions described above were confirmed using *B. pseudomallei* (n = 17), *B. thailandensis* (n = 6), and *B. cepacia* (n = 11) isolates and other competing soil flora (Table 1), artificial soil (29) (n = 4), as well as soil from an area where melioidosis is not endemic (Germany, n = 4; latitude, 48°21'56.29"N; longitude, 10°50'38.32"E; altitude, 514 m).

Data analysis. For the final analysis of the data, the results of all culture methods from soil were combined, and the rate of positivity by any culture method was compared with the rate of positivity by direct DNA extraction or DNA extraction postenrichment. Statistical analysis was performed using Stata/IC (v10) software (StataCorp, College Station, TX, USA). Comparisons were made by the use of McNemar's test (paired samples) or the Mann-Whitney U test, as appropriate. Significance was set at a *P* value of <0.05.

Nucleotide sequence accession numbers. Sequences were analyzed using NCBI-BLAST and subsequently submitted to GenBank and are accessible under accession numbers KM058066 to KM058079 (Table 1).

RESULTS

Comparison of methods for *B. pseudomallei* detection in soil.

The qPCR was highly specific (100%) when tested with a range of reference strains and Lao clinical and environmental isolates (Table 1). The local limit of detection (LOD), which was determined using serial dilutions, was 8 GE/ μ l of soil DNA (equivalent to 32 GE/reaction).

Using the various culture methods, *B. pseudomallei* was detected in 44 out of 100 soil samples (44%; see Table S1 in the supplemental material). *B. pseudomallei* was detected in 6 out of 100 soil samples (6%) using DS DNA extraction, which was a significantly smaller number of soil samples than the number in which *B. pseudomallei* was detected by culture methods (*P* < 0.001). However, when using the ES DNA extraction method followed by qPCR, nearly double the number of samples (84/100, 84%) were identified to be positive compared to the number positive by the culture approach (*P* < 0.001; Fig. 2). Samples positive after direct extraction from soil contained between ~1 and 6 × 10² GE/g soil (range; median, ~1.3 × 10² GE/g soil). Of the 44 culture-positive soil samples, all but 1 were also positive by qPCR following ES DNA extraction (97.7%). No inhibition was observed with any of the extraction methods (change in *C_T* value, \geq 2), suggesting the complete removal of inhibitors using the commercial soil extraction kit. There was no difference in the overall rates of positivity by PCR and/or culture between samples taken at 30 or 60 cm (for samples taken at 60 cm, 42/50 samples [84%] were positive; for samples taken at 30 cm, 43/50 samples [86%] were positive).

Due to the culture step, reliable and accurate quantification of *B. pseudomallei* by qPCR following enrichment was not possible. However, it was observed that culture-positive soil samples had significantly lower *C_T* values after enrichment than culture-negative soil samples (median *C_T* value for culture-positive soil samples = 27.4; interquartile ratio [IQR] of *C_T* values = 23.4 to 30.7;




FIG 2 Comparison of percent positivity by detection type for different environmental sample types. The detection capacities for culture and for enrichment, DNA extraction from soil, and qPCR were significantly different (*, $P < 0.001$). The results obtained by all different culture methods for the different samples were combined.

median C_T value for culture-negative soil samples = 34.3; IQR of C_T values = 32.0 to 45.1; $P = 0.0001$; **Fig. 3**).

Comparison of methods for *B. pseudomallei* detection in river water. Overall, samples taken at 15/20 (75%) sampling points were positive by culture and/or PCR (**Table 2**; see also **Table S2** in the supplemental material). The yield obtained by direct placement of filters on Ashdown's agar (13/20, 65%) was higher than that obtained from broth enrichment cultures (10/20, 50%); the duration of broth enrichment prior to subculture made no difference to the yield. The yield of the 0.2- μm -pore-size filter (12/20, 60%) was higher than that of the 3- μm -pore-size filter (6/20, 30%) when placed on agar; this difference was, however, not significant ($P = 0.070$). No difference in yields from filters of the two pore sizes was observed when broth enrichment was used. Only in one case was culture of the filter directly on Ashdown's agar negative while the enrichment culture was positive with a sample from the same sampling site. In total, 55% (11/20) of sampling points were positive by qPCR when DNA was directly extracted from any filter and 75% (15/20) of sampling points were positive by qPCR when the enrichment approach was used. Samples from nearly all qPCR-positive sites (14/15) were positive with both filter sizes when using the molecular approach with enrichment, with only 1 additional sample being positive with the 3- μm -pore-size filter. No difference in the positivity rates by sampling point was found when the molecular and culture methods were compared (for enrichment and qPCR, 15/20 [75%] samples were positive; for culture, 15/20 [75%] samples were positive). When comparing rates of positivity for individual filters, qPCR postenrichment was more often positive than culture methods (for qPCR postenrichment versus direct plating, $P = 0.003$; for qPCR postenrichment versus broth enrichment culture, $P < 0.001$).

The *B. pseudomallei* bacterial load was estimated to be $\sim 7.5 \times$




FIG 3 C_T values of samples positive or negative by culture methods. Low C_T values indicate a higher bacterial number in the template, while higher C_T values are indicative of lower bacterial numbers. For illustrative purposes, samples negative by qPCR were assigned a C_T value of 50. The median and interquartile ranges are marked. *, statistically significant difference ($P < 0.001$).

10^3 GE/liter river water (median; range, 8×10^2 to 1.3×10^5 GE/liter river water) after direct DNA extraction from filter membranes (pore sizes, 0.2 μm and 3 μm).

DISCUSSION

In order to increase the awareness of health care staff, assist with patient management, and implement public health actions, the availability of a detailed risk map for melioidosis is essential. Our aim was to update and improve detection methods to facilitate further research into the distribution of *B. pseudomallei* in the Lao environment.

Bacterial culture methods have, until now, been the methods of choice to detect *B. pseudomallei* in soil (9, 11, 20, 30). In this investigation, *B. pseudomallei* was detected in 44% of the Lao soil samples tested by culture, while DNA detection following an enrichment step gave a much higher positivity rate for the same sample set (84%). Use of an initial propagation step significantly increased the ability to detect the bacterium in Lao soil, consistent with findings from Australia (14), where DNA detection directly from soil lacked sensitivity. Direct DNA detection from clay-rich soils, like those found in the Lao rice paddy investigated in the present study (~ 30 to 45% clay-size particles; data not shown), can be extremely challenging, as clay particles can reduce the extraction efficiency (14, 31) of commercial kits (32). It is therefore important to include inhibition controls to identify one important cause of false-negative results for samples. Despite the limitations when extracting DNA directly from soil, *B. pseudomallei* DNA has successfully been obtained from sandy loam soils from northeastern Thailand (15) using the SoilMaster DNA extraction kit (Epicentre Biotechnologies, USA). Interestingly, in that investigation the estimated bacterial counts per gram of soil were up to

TABLE 2 Overview of detection methods and positivity frequency for individual water samples and sampling sites^a

Technique	Positivity frequency ^b			
	After filtering through:			
	A 0.2- μ m-pore-size filter	A 3- μ m-pore-size filter	In water	By sample site
Culture techniques				
Moore's swab	NA	NA	2/10 (20)	2/10 (20)
Filtering, direct plating of filter on Ashdown's agar ^c	12/20 (60)	6/20 (30)	NA	13/20 (65)
Filtering, filter placement in Ashdown's broth (3 days) ^{c,d}	6/20 (30)	6/20 (30)	NA	10/20 (50)
Filtering, filter placement in Ashdown's broth (7 days) ^{c,d}	6/20 (30)	6/20 (30)	NA	10/20 (50)
Molecular techniques				
Enrichment, qPCR	14/20 (70)	15/20 (75)	NA	15/20 (75)
Direct qPCR	8/20 (40)	10/20 (50)	NA	11/20 (55)

^a The positivity of individual filters, as well as the overall positivity by sampling site, is listed for the individual techniques.

^b Data represent the number of samples positive/total number of samples tested (percent). NA, not applicable.

^c Combined observations after 48 h, 72 h, and 96 h.

^d Combined results for culture of 10- μ l and 100- μ l supernatants.

140 times higher than those in Lao samples from this study (in Thailand the median is 1.8×10^4 GE/g soil [15]; in Laos the median is 1.3×10^2 GE/g soil). This overall lower *B. pseudomallei* load in Lao soil could also be responsible for the increased difficulties of bacterial DNA detection, and the observed correlation between low C_T values after enrichment and culture positivity might point toward lower *B. pseudomallei* loads in Lao soil.

To reduce the influence of sample variations on the method comparison, we employed a rigorous soil subsampling approach (21, 33, 34). Still, the possibility that variations in results are due to an uneven distribution of the organism within the samples cannot be ruled out (33). Quantitative real-time PCR assays are thought to be highly specific, and the findings of extensive analytical specificity studies performed in this and previous studies (14, 16) support this. In addition, nontemplate and negative controls, used to monitor the potential contamination of reagents and equipment, were consistently negative, supporting the specificity of our results. In future studies, the inclusion of dedicated extraction controls monitoring the DNA extraction efficiency should be considered to further increase the quality of the data.

As with *B. pseudomallei* detection from soil, nucleic acid extraction from filters following an enrichment step proved to be the most sensitive and reliable method to detect the organism in water samples. This report represents the first description of the use of molecular methods to detect *B. pseudomallei* in surface water samples. Water sampling campaigns might represent a promising alternative to large-scale soil sampling campaigns, for example, by using river water as an initial screen to determine whether *B. pseudomallei* is present in the relevant catchment area (12). Despite the overall high positivity rate obtained when using qPCR postenrichment, only the combination of molecular and culture techniques allowed the identification of a *B. thailandensis* isolate that cross-reacted with the *B. pseudomallei* latex reagent (35) due to the presence of a *B. pseudomallei*-like extracellular exopolysaccharide (EPS), emphasizing that one methodology does not fit all research questions.

In conclusion, molecular detection methods using an additional initial enrichment step have proven to be sensitive, specific, and reliable approaches for the detection of *B. pseudomallei* in environmental samples, particularly soil samples. They have the potential for simple scale-up and are less time- and labor-inten-

sive than culture methods. However, the consumables are more expensive (~\$9 per sample) than the consumables used for identification by agglutination assay alone (~\$0.50 per sample), and further follow-up phylogenetic investigations are limited without isolation of the organism.

Additional studies are needed to investigate in particular the water sampling approach and explore the influence of physical and chemical characteristics as well as bacterial loads on the different *B. pseudomallei* detection methods. These data suggest that molecular methods will be an important tool in establishing a melioidosis risk map in Laos and elsewhere.

ACKNOWLEDGMENTS

We are grateful to all staff and students at the Microbiology Laboratory, Mahosot Hospital, Vientiane, Laos, especially P. Panyanouvong, P. Phouminh, M. Seephonelee, D. Sengdetkha, A. Sengduangphachanh, A. Seupsavith, B. Sibounheuang, V. Sihalath, M. Simmalavong, and Ian Ewing, and the farmers who granted permission to dig on their land, as well as Oloth Sengtaheuanghoung (Department of Agriculture Land Management, Lao People's Democratic Republic) for his help in organizing the field mission throughout the Sedone River catchment and liaising with local authorities. We thank Direk Limmathurotsakul from the Mahidol-Oxford-Research Unit for his useful suggestions and fruitful discussions during the analysis and write-up of this study. Further, we are very grateful to the Minister of Health and the director of the Curative Department, Ministry of Health, for their support for this study.

This study was part of the work of the Lao-Oxford-Mahosot Hospital-Wellcome Trust Research Unit funded by the Wellcome Trust of Great Britain. Additional funding was provided by a seed award (Lee Ka Shing Foundation) of the University of Oxford (grant SM40) and the Institut de Recherche pour le Développement (IRD) through the regional pilot program Soils, Waters, Coastal Zones and Societies in Southern and South-east Asia (SELTAR-RPP).

REFERENCES

- Cheng AC, Currie BJ. 2005. Melioidosis: epidemiology, pathophysiology, and management. *Clin Microbiol Rev* 18:383–416. <http://dx.doi.org/10.1128/CMR.18.2.383-416.2005>.
- Wiersinga WJ, Currie BJ, Peacock SJ. 2012. Melioidosis. *N Engl J Med* 367:1035–1044. <http://dx.doi.org/10.1056/NEJMr1204699>.
- Mayo M, Kaesti M, Harrington G, Cheng AC, Ward L, Karp D, Jolly P, Godoy D, Spratt BG, Currie BJ. 2011. *Burkholderia pseudomallei* in unchlorinated domestic bore water, tropical northern Australia. *Emerg Infect Dis* 17:1283–1285. <http://dx.doi.org/10.3201/eid1707.100614>.

4. White NJ. 2003. Melioidosis. *Lancet* 361:1715–1722. [http://dx.doi.org/10.1016/S0140-6736\(03\)13374-0](http://dx.doi.org/10.1016/S0140-6736(03)13374-0).
5. Limmathurotsakul D, Wongratnacheewin S, Teerawattanasook N, Wongsuvan G, Chaisuksant S, Chetchotisakd P, Chaowagul W, Day NP, Peacock SJ. 2010. Increasing incidence of human melioidosis in northeast Thailand. *Am J Trop Med Hyg* 82:1113–1117. <http://dx.doi.org/10.4269/ajtmh.2010.10-0038>.
6. Limmathurotsakul D, Kanoksil M, Wuthiekanun V, Kitphati R, deStavola B, Day NP, Peacock SJ. 2013. Activities of daily living associated with acquisition of melioidosis in northeast Thailand: a matched case-control study. *PLoS Negl Trop Dis* 7:e2072. <http://dx.doi.org/10.1371/journal.pntd.0002072>.
7. National Select Agent Registry. 2014, posting date. National Select Agent Registry: select agents and toxins list. Centers for Disease Control and Prevention, Atlanta, GA. <http://www.selectagents.gov/SelectAgentsandToxinsList.html>.
8. Phetsouvanh R, Phongmany S, Newton P, Mayxay M, Ramsay A, Wuthiekanun V, White NJ. 2001. Melioidosis and Pandora's box in the Lao People's Democratic Republic. *Clin Infect Dis* 32:653–654. <http://dx.doi.org/10.1086/318713>.
9. Rattanavong S, Wuthiekanun V, Langla S, Amornchai P, Sirisouk J, Phetsouvanh R, Moore CE, Peacock SJ, Buisson Y, Newton PN. 2011. Randomized soil survey of the distribution of *Burkholderia pseudomallei* in rice fields in Laos. *Appl Environ Microbiol* 77:532–536. <http://dx.doi.org/10.1128/AEM.01822-10>.
10. Limmathurotsakul D, Wongsuvan G, Aanensen D, Ngamwilai S, Saiprom N, Rongkard P, Thaipadungpanit J, Kanoksil M, Chantratita N, Day NP, Peacock SJ. 2014. Melioidosis caused by *Burkholderia pseudomallei* in drinking water, Thailand, 2012. *Emerg Infect Dis* 20:265–268. <http://dx.doi.org/10.3201/eid2002.121891>.
11. Limmathurotsakul D, Dance DA, Wuthiekanun V, Kaestli M, Mayo M, Warner J, Wagner DM, Tuanyok A, Wertheim H, Yoke Cheng T, Mukhopadhyay C, Puthuchearay S, Day NP, Steinmetz I, Currie BJ, Peacock SJ. 2013. Systematic review and consensus guidelines for environmental sampling of *Burkholderia pseudomallei*. *PLoS Negl Trop Dis* 7:e2105. <http://dx.doi.org/10.1371/journal.pntd.0002105>.
12. Vongphayloth K, Rattanavong S, Moore CE, Phetsouvanh R, Wuthiekanun V, Sengdouangphachanh A, Phouminh P, Newton PN, Buisson Y. 2012. *Burkholderia pseudomallei* detection in surface water in southern Laos using Moore's swabs. *Am J Trop Med Hyg* 86:872–877. <http://dx.doi.org/10.4269/ajtmh.2012.11-0739>.
13. Chantratita N, Wuthiekanun V, Boonbumrung K, TiyaWisutris R, Vesaratchavest M, Limmathurotsakul D, Chierakul W, Wongratnacheewin S, Pukritiyakamee S, White NJ, Day NP, Peacock SJ. 2007. Biological relevance of colony morphology and phenotypic switching by *Burkholderia pseudomallei*. *J Bacteriol* 189:807–817. <http://dx.doi.org/10.1128/JB.01258-06>.
14. Kaestli M, Mayo M, Harrington G, Watt F, Hill J, Gal D, Currie BJ. 2007. Sensitive and specific molecular detection of *Burkholderia pseudomallei*, the causative agent of melioidosis, in the soil of tropical northern Australia. *Appl Environ Microbiol* 73:6891–6897. <http://dx.doi.org/10.1128/AEM.01038-07>.
15. Trung TT, Hetzer A, Gohler A, Topfstedt E, Wuthiekanun V, Limmathurotsakul D, Peacock SJ, Steinmetz I. 2011. Highly sensitive direct detection and quantification of *Burkholderia pseudomallei* bacteria in environmental soil samples by using real-time PCR. *Appl Environ Microbiol* 77:6486–6494. <http://dx.doi.org/10.1128/AEM.00735-11>.
16. Novak RT, Glass MB, Gee JE, Gal D, Mayo MJ, Currie BJ, Wilkins PP. 2006. Development and evaluation of a real-time PCR assay targeting the type III secretion system of *Burkholderia pseudomallei*. *J Clin Microbiol* 44:85–90. <http://dx.doi.org/10.1128/JCM.44.1.85-90.2006>.
17. Rainbow L, Hart CA, Winstanley C. 2002. Distribution of type III secretion gene clusters in *Burkholderia pseudomallei*, *B. thailandensis* and *B. mallei*. *J Med Microbiol* 51:374–384.
18. Winstanley C, Hales BA, Hart CA. 1999. Evidence for the presence in *Burkholderia pseudomallei* of a type III secretion system-associated gene cluster. *J Med Microbiol* 48:649–656. <http://dx.doi.org/10.1099/00222615-48-7-649>.
19. Manivanh L, Pierrat A, Rattanavong S, Buisson Y, Elliott I, Maeght J-L, Xayyathip K, Vongsouvath M, Phetsouvanh R, Newton PN, Dance DAB. 2013. Abstr 7th World Melioidosis Congr 2013, Bangkok, Thailand, poster PPV-03.
20. Wuthiekanun V, Mayxay M, Chierakul W, Phetsouvanh R, Cheng AC, White NJ, Day NP, Peacock SJ. 2005. Detection of *Burkholderia pseudomallei* in soil within the Lao People's Democratic Republic. *J Clin Microbiol* 43:923–924. <http://dx.doi.org/10.1128/JCM.43.2.923-924.2005>.
21. Interstate Technology & Regulatory Council. 2012. Incremental sampling methodology (ISM-1). Interstate Technology & Regulatory Council, Washington, DC.
22. Dittrich S, Knappik MA, Newton PN, Dance DAB. 2013. Abstr 7th World Melioidosis Congr 2013, Bangkok, Thailand, poster PPV-15.
23. Crump BC, Armbrust EV, Baross JA. 1999. Phylogenetic analysis of particle-attached and free-living bacterial communities in the Columbia River, its estuary, and the adjacent coastal ocean. *Appl Environ Microbiol* 65:3192–3204.
24. Suter E, Juhl AR, O'Mullan GD. 2011. Particle association of enterococcus and total bacterial in the lower Hudson River estuary, USA. *J Water Res Prot* 3:715–725. <http://dx.doi.org/10.4236/jwarp.2011.310082>.
25. Jiang J, Chan TC, Temenak JJ, Dasch GA, Ching WM, Richards AL. 2004. Development of a quantitative real-time polymerase chain reaction assay specific for *Orientia tsutsugamushi*. *Am J Trop Med Hyg* 70:351–356.
26. Mayxay M, Castonguay-Vanier J, Chansamouth V, Dubot-Pères A, Paris DH, Phetsouvanh R, Tangkhabuanbutra J, Douangdala P, Inthath S, Souvannasing P, Slesak G, Tongyoo N, Chanthongthip A, Panyanouvong P, Sibounheuang B, Phommason K, Dohnt M, Phonekeo D, Hongvanthong B, Xayadeth S, Ketmayoon P, Blacksell SD, Moore CE, Craig SB, Burns M-A, von Sonnenburg F, Corwin A, de Lamballerie X, González JJ, Christopher EM, Cawthorne A, Bell D, Newton PN. 2013. Causes of non-malarial fever in Laos: a prospective study. *Lancet Global Health* 1:e46–e54. [http://dx.doi.org/10.1016/S2214-109X\(13\)70008-1](http://dx.doi.org/10.1016/S2214-109X(13)70008-1).
27. Peters RP, Mohammadi T, Vandenbroucke-Grauls CM, Danner SA, van Agtmael MA, Savelkoul PH. 2004. Detection of bacterial DNA in blood samples from febrile patients: underestimated infection or emerging contamination? *FEMS Immunol Med Microbiol* 42:249–253. <http://dx.doi.org/10.1016/j.femsim.2004.05.009>.
28. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT. 2009. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 55:611–622. <http://dx.doi.org/10.1373/clinchem.2008.112797>.
29. Ellis RJ. 2004. Artificial soil microcosms: a tool for studying microbial autecology under controlled conditions. *J Microbiol Methods* 56:287–290. <http://dx.doi.org/10.1016/j.mimet.2003.10.005>.
30. Amornchai P, Chierakul W, Wuthiekanun V, Mahakhunkijcharoen Y, Phetsouvanh R, Currie BJ, Newton PN, van Vinh Chau N, Wongratnacheewin S, Day NP, Peacock SJ. 2007. Accuracy of *Burkholderia pseudomallei* identification using the API 20NE system and a latex agglutination test. *J Clin Microbiol* 45:3774–3776. <http://dx.doi.org/10.1128/JCM.00935-07>.
31. Takada-Hoshino Y, Matsumoto N. 2004. An improved DNA extraction method using skim milk from soils that strongly adsorb DNA. *Microbes Environ* 19:13–19. <http://dx.doi.org/10.1264/jmsme.2.19.13>.
32. Young JM, Rawlence NJ, Weyrich LS, Cooper A. 2014. Limitations and recommendations for successful DNA extraction from forensic soil samples: a review. *Sci Justice* 54:238–244. <http://dx.doi.org/10.1016/j.scijus.2014.02.006>.
33. Narizzano R, Riso F, Innocenti R, Mollica V, Tortarolo B. 2008. Soil subsampling in environmental sciences: the role of granulometry. *J Environ Monit* 10:993–997. <http://dx.doi.org/10.1039/b806522p>.
34. Ramsey CA, Suggs J. 2001. Improving laboratory performance through scientific subsampling techniques. *Environ Test Anal* 10:12–16.
35. Sim BM, Chantratita N, Ooi WF, Nandi T, Tewhey R, Wuthiekanun V, Thaipadungpanit J, Tumapa S, Ariyaratne P, Sung WK, Sem XH, Chua HH, Ramnarayanan K, Lin CH, Liu Y, Feil EJ, Glass MB, Tan G, Peacock SJ, Tan P. 2010. Genomic acquisition of a capsular polysaccharide virulence cluster by non-pathogenic *Burkholderia* isolates. *Genome Biol* 11:R89. <http://dx.doi.org/10.1186/gb-2010-11-8-r89>.
36. Ho C-C, Lau CCY, Martelli P, Chan S-Y, Tse CWS, Wu AKL, Yuen K-Y, Lau SKP, Woo PCY. 2011. Novel pan-genomic analysis approach in target selection for multiplex PCR identification and detection of *Burkholderia pseudomallei*, *Burkholderia thailandensis*, and *Burkholderia cepacia* complex species: a proof-of-concept study. *J Clin Microbiol* 49:814–821. <http://dx.doi.org/10.1128/JCM.01702-10>.