



Journal Name

COMMUNICATION

a-2018-02832p

## A Luminescent $[\text{Pd}_4\text{Ru}_8]^{24+}$ Supramolecular Cage

Diego Rota Martir,<sup>a</sup> David B. Cordes,<sup>a</sup> Alexandra M. Z. Slawin,<sup>a</sup> Daniel Escudero,<sup>b</sup> Denis Jacquemin,<sup>b</sup> Stuart L. Warriner,<sup>c</sup> and Eli Zysman-Colman<sup>a</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)

**A phosphorescent cage of the form  $[\text{Pd}_4\text{Ru}_8]^{24+}$  is reported. The cage was formed using the metalloligand  $[\text{Ru}(\text{dtbubpy})_2(\text{qpy})]^{2+}$ , where  $\text{qpy} = 4,4':2'',2''':4'',4''''$ -quaterpyridine and  $\text{dtbubpy} = 4,4'$ -ditert-butyl-2,2'-bipyridine. The cage has been characterised by NMR, ESI-MS, TEM and X-ray diffraction analyses and its emission properties elucidated by steady-state and time-resolved emission spectroscopy.**

Three dimensional metallosupramolecular self-assembly has evolved over the last thirty years as a powerful approach for the construction of structural frameworks, such as cages and capsules, with well-defined shapes and cavities.<sup>1</sup> Due to their ability to encapsulate guest compounds within their cavities, attention has now turned towards the use of these "supramolecules" as functional materials.<sup>2</sup> For instance, cages have been explored in the context of "artificial enzyme" catalysis,<sup>3</sup> for hazardous chemical capture,<sup>4</sup> reactive intermediate stabilisation,<sup>2c</sup> for drug delivery and release,<sup>5</sup> as well as in molecular sensing<sup>6</sup> and in biology.<sup>7</sup> More recently, desirable photophysical properties such as emission tuning,<sup>8</sup> encapsulation-assisted energy and electron transfer,<sup>9</sup> long-lived charge-separated state lifetimes<sup>10</sup> and photoresponsive guest uptake/release<sup>11</sup> have been achieved by integrating emissive moieties such as metalloporphyrins, metal complexes or fluorescent ligands, into the backbone of metallosupramolecular macrocycles and cages.<sup>6a,12</sup>

Ruthenium(II) polypyridyl complexes can generally access multiple oxidation states, and they enjoy a rich history in photocatalysis.<sup>13</sup> However, Ru(II) complexes are general poorly

emissive and their use as luminophores is limited.<sup>14</sup> Many examples have nevertheless been reported where ruthenium(II) complexes are incorporated into polymers,<sup>15</sup> networks,<sup>16</sup> metal-organic frameworks<sup>17</sup> and discrete 2D metallamacrocycles.<sup>18</sup> However, 3D supramolecular cages incorporating ruthenium(II) complexes as structural scaffolds still remain underexplored.<sup>19</sup> Three relevant examples of Ru(II) cages involve a  $\text{Pd}_8\text{Ru}_4$  tetrameric cage<sup>20</sup> formed by the assembly of a Ru(II) metalloligand based on a 2,2',6',2''-terpyridine ligand and  $\text{Pd}^{2+}$  ions, a  $\text{Pd}_6\text{Ru}_8$  cage,<sup>21</sup> in which the eight ruthenium metalloligands and the six  $\text{Pd}^{2+}$  ions are organised into an octahedron, and a  $\text{L}_4\text{Ru}_6$  tetrahedral cage where L is 2,4,6-tri(pyridine-4-yl)1,3,5-triazine (tpt) and Ru is *cis*-bis(2,2'-bipyridine)ruthenium(II).<sup>22</sup> However, these cages as well as most of the supramolecular assemblies incorporating Ru(II) metalloligands are either poorly or non-emissive.

Herein we report, the structural characterisation of a cage of composition  $[(\text{Ruqpy})_8\text{Pd}_4]^{24+}$ , **RuPd**, formed by self-assembly of the ruthenium(II) metalloligand  $[\text{Ru}(\text{dtbubpy})_2(\text{qpy})][\text{PF}_6]_2$ , **Ruqpy**, (where  $\text{dtbubpy}$  is 4,4'-di-*tert*-butyl-2,2'-bipyridine and  $\text{qpy}$  is 4,4':2'',2''':4'',4''''-quaterpyridine) with  $\text{Pd}^{2+}$  ions. The **RuPd** cage has been characterised by NMR spectroscopy, ESI-MS spectrometry, TEM and X-ray crystallography. Notably, single crystal X-ray analysis revealed that **RuPd** has a diagonal distance of 38.4 Å (corresponding to the distance between furthest carbon atoms of opposite *t*-butyl groups), which makes it, to the best of our knowledge, the largest X-ray structure of a Ru(II) cage assembled with  $\text{Pd}^{2+}$  ions reported to date. The emission properties of **RuPd** have been investigated by steady-state and time-resolved photoluminescence spectroscopy and compared with those of the **Ruqpy** metalloligand.

<sup>a</sup> Organic Semiconductor Centre, EaStCHEM School of Chemistry, University of St Andrews, St Andrews, Fife, KY16 9ST, UK, Fax: +44-1334 463808; Tel: +44-1334 463826.


<sup>b</sup> CEISAM UMR CNRS 6230, Université de Nantes, 2 rue de la Houssinière, BP 92208, 44322 Nantes Cedex 3, France.

<sup>c</sup> School of Chemistry, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, UK.

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: characterisation of metalloligand and cage, x-ray structural characterisation (CCDC 1828574 and 1828575), supplementary optoelectronic and computational data. See DOI: 10.1039/x0xx00000x


When two equivalents of **Ruqpy** were reacted with  $[\text{Pd}(\text{NCMe})_4](\text{BF}_4)_2$  in  $\text{DMSO-}d_6$  at  $85^\circ\text{C}$  for 12 h, the proton resonances associated with **Ruqpy** broadened and shifted downfield (Figure 1a). The downfield shift of the signals of **RuPd** compared to **Ruqpy**, is evident for the pyridyl  $\alpha$ -hydrogen atoms ( $\Delta\delta = 0.23$  ppm), and it is characteristic of metal-pyridine complexation. The broadness of the NMR signals of **RuPd** is indicative of the formation of a large structure, the motion of which is very slow on the NMR time scale (Figure S1).  $^1\text{H}$  DOSY NMR spectroscopy clearly documents the formation of a single species in solution, the diffusion coefficient of which, measured in  $\text{DMSO-}d_6$  was found to be  $D = 5.3 \times 10^{-11} \text{ m}^2/\text{s}$ . The magnitude of  $D$  correlates to the presence of a much larger structure than the metalloligand, which has a diffusion coefficient of  $1.3 \times 10^{-10} \text{ m}^2/\text{s}$  in  $\text{DMSO-}d_6$  (Figure 1b). The calculated hydrodynamic radius ( $r_s$ ) of the cage is  $19.7 \text{ \AA}$  (Table S1). The presence of **RuPd** was further corroborated by HR-ESI-MS spectrometry, showing the isotopically resolved peaks for  $[(\text{RuPd})-(\text{BF}_4)_n]^{n+}$  ( $n = 7 - 9$ ) that match with the simulated spectra (Figures S6-S9).




**Figure 1.** a) Self-assembly between **Ruqpy** and  $\text{Pd}^{2+}$  ions yielding the cage **RuPd**, showing the X-ray structure. b)  $^1\text{H}$  DOSY NMR of **Ruqpy** in blue and **RuPd**, in red. The  $^1\text{H}$  DOSY NMR spectra were collected in  $\text{DMSO-}d_6$  at 298 K.

The cubic structure of **RuPd** was unambiguously determined by X-ray crystallography (Figures 1 and 2). Crystals suitable for X-ray analysis were grown by the slow diffusion of a 1:1 mixture of ethyl acetate-hexane into a  $\text{DMSO}$  solution of **RuPd** (25 mM) over 40 days. The topology of **RuPd** resembles that reported by Klein *et al.* wherein the cage structure is constructed such that two ligands doubly bridge adjacent  $\text{Pd}(\text{II})$  centres in a crown-like fashion, disposing the four palladium ions in a square arrangement.<sup>23</sup> This forms a metallomacrocyclic core to the cage. Among the family of  $\text{Pd}_4\text{L}_8$  cages, this specific arrangement is rare with only six examples reported to date.<sup>9c, 23-24</sup> Indeed, this stoichiometry is only possible when the angle between the coordinating 4-pyridyl units is less than  $90^\circ$ ; the X-ray structure of **Ruqpy** revealed that the angle between the coordinating 4-pyridyl units of the qpy ligand

measures  $69^\circ$  (Figure S3). The angle between the coordinating 4-pyridyl units of **Ruqpy** is smaller than that of the previously reported complex  $[\text{Ir}(\text{mesppy})_2(\text{qpy})]\text{PF}_6$ , which showed a coordinating angle of  $78^\circ$ .<sup>25</sup> However, the assembly of both qpy-based complexes with  $\text{Pd}^{2+}$  ions gave rise to cages of the same  $\text{Pd}_4\text{L}_8$  topology.<sup>9c</sup> The **RuPd** cage exhibits a longest diagonal of  $38.4 \text{ \AA}$  (Figures 3, S3), leading to a radius of the cage structure based on this diagonal of  $19.2 \text{ \AA}$ , similar to that obtained by  $^1\text{H}$  DOSY NMR analysis ( $r_s = 19.7 \text{ \AA}$ ). The cage shows a distance between neighboring ruthenium atoms bridging the same  $\text{Pd}\cdots\text{Pd}$  edge of  $15.0 \text{ \AA}$ , an internal diameter of  $27.4 \text{ \AA}$  (corresponding to the longest  $\text{Ru}\cdots\text{Ru}$  distance), and a resulting internal volume of ca.  $4900 \text{ \AA}^3$ . The nanostructures of cage **RuPd** were probed by Transmission Electron Microscopy (TEM) analysis upon deposition of **RuPd** (concentration of  $1 \times 10^{-6} \text{ M}$ ) onto carbon-coated copper grids (Figure 3). The size of the nanostructures observed by TEM are of around  $4.0 - 4.2 \text{ nm}$ , in good agreement with the diagonal distance of  $3.9 \text{ nm}$  observed for the X-ray structure (Figure 3a).



**Figure 2.** Space-filling views of the single crystal X-ray crystal structure of cage **RuPd** viewed down to the crystallographic  $a$ - (left) and  $c$ -axes (right). Hydrogen atoms, solvent molecules and counterions have been omitted for clarity.



**Figure 3.** a) Illustration of the X-ray structure of **RuPd**, showing the diagonal distance across the long axis. b) TEM images of the nanostructures of **RuPd**.


The emission profile of **RuPd** in DCM ( $\lambda_{\text{PL}} = 710 \text{ nm}$ ) is broader and red-shifted compared to that of **Ruqpy** ( $\lambda_{\text{PL}} = 674 \text{ nm}$ ,  $\Phi_{\text{PL}} = 7.3\%$ ), and with a photoluminescence quantum yield,  $\Phi_{\text{PL}}$ , of  $6.9\%$  (Figure 4a). Notably, the  $\Phi_{\text{PL}}$  of **RuPd** is one of the highest reported among ruthenium cages. The red-shifted emission of **RuPd** compared to **Ruqpy** is the result of coordination of the Lewis acidic  $\text{Pd}(\text{II})$  ions to the Ru complex, which essentially stabilizes the  $\pi^*_{\text{qpy}}$  orbital level involved in the emission, and thus lowers the energy of the triplet state.<sup>26</sup> Both **Ruqpy** and **RuPd** exhibit bi-exponential photoluminescence decay kinetics with lifetimes,  $\tau_{\text{PL}}$ , of 324, 1047

ns and 151, 700 ns, respectively (Figure 4b). As a result, similar radiative rate constants,  $k_r$ , of  $6.97 \times 10^4 \text{ s}^{-1}$  and  $9.86 \times 10^4 \text{ s}^{-1}$ , and non-radiative rate constants,  $k_{nr}$ , of  $8.85 \times 10^5 \text{ s}^{-1}$  and  $13.30 \times 10^5 \text{ s}^{-1}$  are obtained for **Ruqpy** and **RuPd**, respectively (Table 1). Thus, the Pd(II) ions do not adversely affect the photophysical properties of this metalloligand. This observation is rather unusual considering that the emissions of the vast majority of metal complexes assembled within cage structures are often partially or fully quenched by the presence of Pd(II) ions,<sup>19b, 20</sup> likely due to the population of low-lying dark states involving the donor and acceptor units. The computational investigation (see details in the ESI) of **RuPd** reveals that the lowest possible transition, from the HOMO to the LUMO (see Figure S12), leads to a Rydberg excited state from d-based Ru orbitals to the Pd units. The presence of these parasitic states explains the slightly lower  $\Phi_{\text{PL}}$  value (along with its increased  $k_{nr}$  value) for **RuPd** as compared to **Ruqpy**. The fact that, in this specific case, the photoluminescence is not completely quenched in the cage indicates that the radiative process is fast enough to compete with internal decay to the lowest-lying dark states.

**Table 1.** Photophysical properties of **Ruqpy** and **RuPd**.<sup>a</sup>

	$\lambda_{\text{PL}} / \text{nm}$	$\Phi_{\text{PL}} / \%$ <sup>b</sup>	$\tau_{\text{PL}} / \text{ns}$ <sup>c</sup>	$k_r \times 10^4 / \text{s}^{-1}$	$k_{nr} \times 10^5 / \text{s}^{-1}$
<b>Ruqpy</b>	674	7.3	324 (0.2) 1047 (0.8)	6.97	8.85
<b>RuPd</b>	710	6.9	151 (0.1) 700 (0.9)	9.86	13.30

<sup>a</sup> Measurements in degassed DCM at 298 K ( $\lambda_{\text{exc}} = 360 \text{ nm}$ ). <sup>b</sup>  $\Phi_{\text{PL}}$  measurements using quinine sulfate as the external reference ( $\Phi_{\text{PL}} = 54.6\%$  in 0.5 M  $\text{H}_2\text{SO}_4$  at 298 K).<sup>27</sup> <sup>c</sup>  $\lambda_{\text{exc}} = 378 \text{ nm}$ ; values in parentheses are pre-exponential weighting factor, in relative % intensity, of the emission decay kinetics.



**Figure 4.** a) Normalised emission spectra of **Ruqpy** (dotted blue line) and **RuPd** (dotted red line) in degassed DCM at 298 K ( $\lambda_{\text{exc}} = 360 \text{ nm}$ ). Photographs of the emissions of **Ruqpy** (left) and **RuPd** (right) are inset to the spectra. b) Emission decays of **Ruqpy** (blue line) and **RuPd** (red line) in degassed DCM at 298 K ( $\lambda_{\text{exc}} = 378 \text{ nm}$ ).

In conclusion a NIR luminescent cage of composition  $[\text{Ru}_8\text{Pd}_4]^{24+}$ , formed through the self-assembly of the ruthenium metalloligand  $[\text{Ru}(\text{dtbubpy})_2(\text{qpy})][\text{PF}_6]_2$  with  $\text{Pd}^{2+}$  ions has been reported. The structure of the ruthenium cage has been elucidated by X-ray diffraction and its emission properties investigated by steady-state and time-resolved emission spectroscopy. Cages and capsules based on ruthenium(II) complexes are very rare and are generally

poorly- or non-emissive. However, **RuPd** exhibits a photoluminescence quantum yield of 6.9%, which is remarkably high, considering the emission maximum of 710 nm. The use of the red-emitting cage **RuPd** as a photosensitiser to induce photochemical transformations of bound guests in water is currently under investigation in our laboratory, the results of which will be reported in due course.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

E.Z.-C. acknowledges the University of St Andrews and EPSRC (EP/M02105X/1) for financial support. We thank the EPSRC UK National Mass Spectrometry Facility at Swansea University for analytical services. We thank Umicore AG for the gift of materials. D.E. thanks funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 700961. D.J. acknowledges the RFI Lumomat for financial support. This research used resources of 1) the GENCI-CINES/IDRIS, 2) Centre de Calcul Intensif des Pays de Loire (CCIPL), and 3) a local Troy cluster.

## Notes and references

- (1). (a) M. M. J. Smulders, I. A. Riddell, C. Browne and J. R. Nitschke, *Chem. Soc. Rev.*, 2013, **42**, 1728-1754; (b) D. Fujita, Y. Ueda, S. Sato, N. Mizuno, T. Kumasaka and M. Fujita, *Nature*, 2016, **540**, 563-566.
- (2). (a) M. Han, D. M. Engelhard and G. H. Clever, *Chem Soc Rev*, 2014, **43**, 1848-1860; (b) W. Wang, Y.-X. Wang and H.-B. Yang, *Chem Soc Rev*, 2016, **45**, 2656-2693; (c) S. Zarra, D. M. Wood, D. A. Roberts and J. R. Nitschke, *Chem Soc Rev*, 2015, **44**, 419-432.
- (3). (a) Y. Nishioka, T. Yamaguchi, M. Yoshizawa and M. Fujita, *J. Am. Chem. Soc.*, 2007, **129**, 7000-7001; (b) Y. Furutani, H. Kandori, M. Kawano, K. Nakabayashi, M. Yoshizawa and M. Fujita, *J. Am. Chem. Soc.*, 2009, **131**, 4764-4768; (c) M. Otte, P. F. Kuijpers, O. Troepfner, I. Ivanovic-Burmazovic, J. N. Reek and B. de Bruin, *Chemistry*, 2014, **20**, 4880-4884.
- (4). (a) I. A. Riddell, M. M. Smulders, J. K. Clegg and J. R. Nitschke, *Chem. Commun.*, 2011, **47**, 457-459; (b) T. K. Ronson, S. Zarra, S. P. Black and J. R. Nitschke, *Chem. Commun.*, 2013, **49**, 2476-2490.
- (5). (a) A. Schmidt, V. Molano, M. Hollering, A. Pothig, A. Casini and F. E. Kuhn, *Chemistry*, 2016, **22**, 2253-2256; (b) A. Ahmedova, R. Mihaylova, D. Momekova, P. Shestakova, S. Stoykova, J. Zaharieva, M. Yamashina, G. Momekov, M. Akita and M. Yoshizawa, *Dalton Trans*, 2016, **45**, 13214-13221; (c) A. Mishra, S. Chang Lee, N. Kaushik, T. R. Cook, E. H. Choi, N. Kumar Kaushik, P. J. Stang and K.-W. Chi, *Chemistry*, 2014, **20**, 14410-14420.
- (6). (a) P. P. Neelakandan, A. Jimenez and J. R. Nitschke, *Chem. Sci.*, 2014, **5**, 908-915; (b) D. P. August, G. S. Nichol and P. J. Lusby, *Angew Chem Int Ed Engl*, 2016, **55**, 15022-15026.
- (7). A. Ahmedova, D. Momekova, M. Yamashina, P. Shestakova, G. Momekov, M. Akita and M. Yoshizawa, *Chem Asian J*, 2016, **11**, 474-477.
- (8). J. B. Pollock, G. L. Schneider, T. R. Cook, A. S. Davies and P. J. Stang, *J Am Chem Soc*, 2013, **135**, 13676-13679.

- (9). (a) J. R. Piper, L. Cletheroe, C. G. P. Taylor, A. J. Metherell, J. A. Weinstein, I. V. Sazanovich and M. D. Ward, *Chem. Commun.*, 2017, **53**, 408-411; (b) S. Durot, J. Taesch and V. Heitz, *Chem Rev*, 2014, **114**, 8542-8578; (c) D. Rota Martir, D. Escudero, D. Jacquemin, D. B. Cordes, A. M. Z. Slawin, H. A. Fruchtl, S. L. Warriner and E. Zysman-Colman, *Chem. Eur. J.*, 2017, **23**, 14358-14366.
- (10). P. D. Frischmann, K. Mahata and F. Wurthner, *Chem Soc Rev*, 2013, **42**, 1847-1870.
- (11). (a) M. Han, R. Michel, B. He, Y.-S. Chen, D. Stalke, M. John and G. H. Clever, *Angew Chem Int Ed Engl*, 2013, **52**, 1319-1323; (b) N. Kishi, M. Akita, M. Kamiya, S. Hayashi, H.-F. Hsu and M. Yoshizawa, *J Am Chem Soc*, 2013, **135**, 12976-12979.
- (12). (a) T. Nakamura, H. Ube and M. Shionoya, *Angew Chem Int Ed Engl*, 2013, **52**, 12096-12100; (b) S. Shanmugaraju, H. Jadhav, Y. P. Patil and P. S. Mukherjee, *Inorg Chem*, 2012, **51**, 13072-13074; (c) J. Wang, C. He, P. Wu, J. Wang and C. Duan, *J Am Chem Soc*, 2011, **133**, 12402-12405.
- (13). C. K. Prier, D. A. Rankic and D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322-5363.
- (14). (a) Q. Sun, S. Mosquera-Vazquez, Y. Suffren, J. Hankache, N. Amstutz, L. M. Lawson Daku, E. Vauthey and A. Hauser, *Coord. Chem. Rev.*, 2015, **282-283**, 87-99; (b) E. Rousset, D. Chartrand, I. Ciofini, V. Marvaud and G. S. Hanan, *Chem. Commun.*, 2015, **51**, 9261-9264; (c) J. D. Slinker, D. A. Bernards, P. L. Houston, H. D. Abruña, S. Bernhard and G. G. Malliaras, *Chem. Commun.*, 2003, **19**, 2392-2399.
- (15). (a) Y. Sun, Z. Chen, E. Puodziukynaite, D. M. Jenkins, J. R. Reynolds and K. S. Schanze, *Macromolecules*, 2012, **45**, 2632-2642; (b) J. E. Beves, E. C. Constable, C. E. Housecroft, C. J. Kepert and D. J. Price, *CrystEngComm*, 2007, **9**, 456-459; (c) J. E. Beves, E. C. Constable, S. Decurtins, E. L. Dunphy, C. E. Housecroft, T. D. Keene, M. Neuburger and S. Schaffner, *CrystEngComm*, 2008, **10**; (d) W. Sun, S. Li, B. Haupler, J. Liu, S. Jin, W. Steffen, U. S. Schubert, H.-J. Butt, X. J. Liang and S. Wu, *Adv Mater*, 2017, **29**; (e) C. Friebe, H. Görls, M. Jäger and U. S. Schubert, *Eur. J. Inorg. Chem.*, 2013, **2013**, 4191-4202.
- (16). (a) C. A. Kent, D. Liu, L. Ma, J. M. Papanikolas, T. J. Meyer and W. Lin, *J Am Chem Soc*, 2011, **133**, 12940-12943; (b) W. A. Maza, R. Padilla and A. J. Morris, *J Am Chem Soc*, 2015, **137**, 8161-8168.
- (17). (a) W. Zhang, B. Li, H. Ma, L. Zhang, Y. Guan, Y. Zhang, X. Zhang, P. Jing and S. Yue, *ACS Appl Mater Interfaces*, 2016, **8**, 21465-21471; (b) R. Chen, J. Zhang, J. Chelora, Y. Xiong, S. V. Kershaw, K. F. Li, P.-K. Lo, K. W. Cheah, A. L. Rogach, J. A. Zapien and C.-S. Lee, *ACS Appl Mater Interfaces*, 2017, **9**, 5699-5708; (c) S. Zhang, L. Li, S. Zhao, Z. Sun and J. Luo, *Inorg Chem*, 2015, **54**, 8375-8379; (d) W. Zhang, M. Kauer, O. Halbherr, K. Epp, P. Guo, M. I. Gonzalez, D. J. Xiao, C. Wiktor, L. I. I. X. FX, C. Woll, Y. Wang, M. Muhler and R. A. Fischer, *Chemistry*, 2016, **22**, 14297-14307.
- (18). (a) G. R. Newkome, T. J. Cho, C. N. Moorefield, P. P. Mohapatra and L. A. Godinez, *Chemistry*, 2004, **10**, 1493-1500; (b) A. Schultz, X. Li, B. Barkakaty, C. N. Moorefield, C. Wesdemiotis and G. R. Newkome, *J Am Chem Soc*, 2012, **134**, 7672-7675; (c) M. Chen, J. Wang, S. Chakraborty, D. Liu, Z. Jiang, Q. Liu, J. Yan, H. Zhong, G. R. Newkome and P. Wang, *Chem. Commun.*, 2017, **53**, 11087-11090; (d) B. Laramée-Milette, F. Nastasi, F. Puntoriero, S. Campagna and G. S. Hanan, *Chem. Eur. J.*, 2017, **23**, 16497-16504.
- (19). (a) A. Valente and M. Garcia, *Inorganics*, 2014, **2**, 96-114; (b) C. Shen, A. D. W. Kennedy, W. A. Donald, A. M. Torres, W. S. Price and J. E. Beves, *Inorg. Chim. Acta*, 2017, **458**, 122-128; (c) J. Y. Ryu, E. H. Wi, M. Pait, S. Lee, P. J. Stang and J. Lee, *Inorg Chem*, 2017, **56**, 5471-5477.
- (20). J. Yang, M. Bhadbhade, W. A. Donald, H. Iranmanesh, E. G. Moore, H. Yan and J. E. Beves, *Chem. Commun.*, 2015, **51**, 4465-4468.
- (21). K. Li, L.-Y. Zhang, C. Yan, S.-C. Wei, M. Pan, L. Zhang and C.-Y. Su, *J Am Chem Soc*, 2014, **136**, 4456-4459.
- (22). C. E. Hauke, A. N. Oldacre, C. R. P. Fulong, A. E. Friedman and T. R. Cook, *Inorg. Chem.*, 2017, **ASAP**, DOI: 10.1021/acs.inorgchem.1027b02657.
- (23). C. Klein, C. Gutz, M. Bogner, F. Topic, K. Rissanen and A. Lutzen, *Angew Chem Int Ed Engl*, 2014, **53**, 3739-3742.
- (24). (a) K. Miki, G. Yoshida, Y. Kai, N. Yasuoka and N. Kasai, *J. Organomet. Chem.*, 1978, **149**, 195-208; (b) K. Suzuki, M. Kawano and M. Fujita, *Angew Chem Int Ed Engl*, 2007, **46**, 2819-2822; (c) D. Kumar Chand, M. Fujita, K. Biradha, S. Sakamoto and K. Yamaguchi, *Dalton Trans.*, 2003, 2750-2756; (d) W. M. Bloch, Y. Abe, J. J. Holstein, C. M. Wandtke, B. Dittrich and G. H. Clever, *J. Am. Chem. Soc.*, 2016, **138**, 13750-13755.
- (25). D. Rota Martir, G. J. Hedley, D. B. Cordes, A. M. Z. Slawin, D. Escudero, D. Jacquemin, T. Kosikova, D. Philp, D. M. Dawson, S. E. Ashbrook, I. D. W. Samuel and E. Zysman-Colman, *Dalton Trans.*, 2016, **45**, 17195-17205.
- (26). D. Rota Martir, M. Averardi, D. Escudero, D. Jacquemin and E. Zysman-Colman, *Dalton Trans.*, 2017, **46**, 2255-2262.
- (27). W. H. Melhuish, *J. Phys. Chem.*, 1961, **65**, 229-235.